

Nanoscale

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Nanoscale

COMMUNICATION

Monocyte Cell Membrane-derived Nanoghosts for Targeted Cancer Therapy †

Received 00th January 20xx,
Accepted 00th January 20xx

S. Krishnamurthy^a, M.K. Gnanasammandhan^b, C. Xie^a, K. Huang^b, M. Y. Cui^a, J. M. Chan^{a*}

DOI: 10.1039/x0xx00000x

www.rsc.org/nanoscale

Core-shell type ‘nanoghosts’ were synthesized with a biodegradable drug-loaded PLGA core and monocyte cell membrane-derived shell. The nanoghosts were monodisperse with an average size <200 nm, and showed good serum stability for 120 h. Drug uptake studies showed greater cellular uptake and cytotoxicity of doxorubicin-loaded nanoghosts compared to non-coated nanoparticle controls in metastatic MCF-7 breast cancer cell lines.

Nanoparticle (NP)-mediated drug delivery systems have contributed significantly to the advancement of clinical therapy.^{1,2} Liposomes and polymeric nanoparticles are the most commonly-used NP drug delivery platforms,³ many of which have entered clinical trials and received US Food and Drug Administration (FDA) approval for a variety of clinical indications.²

In the recent years, however, researchers have explored newer modalities of drug delivery, such as the use of whole biological cells as targeted delivery systems. These whole cell carriers, as they are commonly referred to, have intact cytoplasm and organelles, and are typically chosen for their ability to home towards a site of interest or release a therapeutic payload.⁴ To date, these carriers include red blood cells (RBCs)^{5,6,7} exosomes,^{8,9} mesenchymal stem cells,^{10,11} macrophages,¹² lymphocytes,¹³ and cancer cells^{14,15,16} among others.

Although whole cell carriers have been used successfully in pre-clinical studies, there are a number of limitations to their use. Therapeutic or diagnostic molecules, which are typically loaded by electroporation or osmosis, leak from the cells during and after loading.¹⁷ As these are viable cells, the whole cell carriers have to remain biologically active upon

administration.⁵

To retain the biological targeting properties of the whole cell carriers and concomitantly overcome cell viability and drug loading issues, researchers have designed cell ghosts, which are mammalian cell-surface membranes in the nano- to micro-range devoid of cytoplasm and organelles.^{18,19} This approach of coating naturally-functionalized cell membranes onto preformed nanoparticle surfaces greatly simplifies the current methods of nanoparticle functionalization.²⁰ Moreover, the presence of an additional lipid (cell membrane, in the case of cell ghosts) layer has been shown to provide greater nanoparticle stability and improved drug release profiles,¹ which are advantageous for *in vivo* drug delivery applications. The cell ghosts are produced by cell lysis and serial ultracentrifugation and have a biologically intact bilayer membrane that can be used for both active and passive targeting. Additionally, serial extrusion can give rise to cell ghosts in the 100–200 nm range (‘nanoghosts’)²¹ that can exploit the enhanced permeation and retention (EPR) effect observed in tumors.²² Typically, drugs are conjugated to or encapsulated within the cell ghosts, with glutaraldehyde commonly used as a common chemical cross linker to conjugate drugs such as daunorubicin²³ and doxorubicin (Dox)^{24,25} to RBC cell ghosts.

To increase the drug loading capacity of the nanoghosts and prevent drug leakage, researchers have encapsulated drug-loaded NPs into the nanoghosts instead of free or conjugated drug.^{26,27,28} Zhang *et al.* encapsulated poly(lactic-co-glycolic acid) (PLGA) NPs within RBC cell ghosts by serial extrusion, for use as nanosponges that absorb pore forming toxins²⁹ and for antitoxin vaccination.²⁹ Other NPs that have been encapsulated include gold nanocages for photothermal cancer therapy^{30,31} and gelatin NPs containing vancomycin as a broad-spectrum antibiotic.³² Besides RBCs, researchers have used cancer cells and leukocytes to encapsulate NPs. Zhang *et al.* used B16-F10 mouse melanoma cell coated-polymeric NPs to deliver adjuvants for dendritic cell maturation and elicit a tumor-specific immune response.^{18,33} Leukocyte cell carriers have also been used to evade the immune system and for

^a School of Chemical and Biomedical Engineering and Lee Kong Chian School of Medicine, Nanyang Technological University, Block N1.3, B4-10, 70 Nanyang Drive, Singapore 637457.

^b Department of Biomedical Engineering, National University of Singapore, 9 Engineering Drive 1, Block EA, #03-12, Singapore 117575.

†Electronic Supplementary Information (ESI) available: See DOI: 10.1039/x0xx00000x

payload delivery of nanoporous silicon microparticles³⁴ and Dox-encapsulated silica NPs.³⁵

Previously, we have designed hybrid NP systems comprising of a polymeric core and lipid shell.^{36,37} These hybrid NP systems typically contain a PLGA core surrounded by a lipid monolayer shell consisting of either synthetic lipids or purified lipid populations extracted from soybean. In this study, we design monocyte-derived cell membrane-coated PLGA NPs to actively target cancers that recruit circulating monocytes as a supply of cell-proliferating growth factors.³⁸ Previously, breast cancer cell lines were shown to overexpress cell adhesion molecules such as VCAM-1 that bind to circulating monocytes expressing cell adhesion molecules such as $\alpha 4\beta 1$ integrin.³⁹ PLGA was selected for the NP core due to its excellent biocompatibility and prior FDA approval for clinical use.⁴⁰ Dox was chosen for its anti-proliferative properties and loaded into the PLGA core by nanoprecipitation. Meanwhile, cell membranes from U937 monocytes were purified by hypotonically lysing the cells, homogenization, and subsequent isolation of the membrane fraction by serial ultra-centrifugation (**Figure 1A**). The purified cell membranes were coated onto Dox-loaded PLGA NPs by serial extrusion through polymycarbonate membranes (pore size 400 nm and 200 nm) to form the nanoghosts as shown in **Figure 1B**.

Before cell membrane coating, dynamic light scattering studies showed that the Dox-loaded PLGA NPs were monodisperse with an average hydrodynamic diameter of 157 nm and a zeta potential of -8.3 mV. The size and zeta potential of the pure membrane fraction were 28 nm and -13.6 mV respectively. After coating, the average diameter of the nanoghosts increased by ~20-40 nm and the zeta potential became more negative at -16.5 mV. Transmission electron microscopy (TEM) images showed similar size profiles before and after cell membrane coating (**Figure 2A, B**). The nanoghosts and PLGA NP exhibited good serum stability over a period of 120 h (**Figure 2C**). Dox-loading of PLGA NPs was high at 21% (drug to polymer weight) and remained at similar levels after extrusion to form the nanoghosts. The nanoghosts released Dox in a sustained fashion over 72 h, similar to PLGA NP controls, but drug release was slower initially (< 30 h) and plateaued at 60 h as shown in **Figure 2D**. The slower release from nanoghosts could be due to an additional cell membrane bilayer that poses a diffusional barrier.

To confirm the post-purification retention and integrity of CD49d, which is a part of the heterodimeric integrin $\alpha 4\beta 1$ that binds to VCAM-1 on target cells, the nanoghost membrane protein profile was compared to the U937 cell membrane extract by SDS-PAGE and western blotting. As shown in **Figure 3A**, protein expression patterns were similar between U937 cell membranes and nanoghosts, including integrin $\alpha 4\beta 1$ and control cell membrane marker proteins sodium-potassium ATPase and GAPDH, suggesting that the U937 cell membrane coated onto the PLGA NPs is biologically relevant.

The intracellular uptake of nanoghosts into metastatic breast cancer cells was tested using MCF-7 cell lines. MCF-7 cells were treated with FITC-loaded nanoghosts or FITC-loaded PLGA NP controls for 2, 4 and 6 h and NP uptake was quantified by flow cytometry. At each time point, the percentage of FITC-positive cells in the parent population was analyzed. The number of FITC-positive cells increased over time in both samples, and the percentage of FITC-positive cells was significantly higher at all timepoints in the nanoghost samples, in comparison to PLGA NP control samples (**Figure 3B**). Similarly, the uptake of these nanoparticles in primary mouse skeletal myocytes that have negligible expression of cell adhesion molecules like VCAM-1, ICAM-1 and ELAM-1 was carried out as a control (**Figure 3C**). Nanoparticle uptake was similar for both the nanoghosts and PLGA NP samples at all time points and reached a maximum of ~10%. We hypothesize that the higher cellular uptake observed with the nanoghosts in cancer cells is due to the monocyte cell membrane coating that provides enhanced cancer cell targeting, cell receptor-specific binding and intracellular uptake.

Finally, the cytotoxicity of Dox-loaded nanoghosts was evaluated using MCF-7 cells. The cytotoxicity of blank nanoghosts and PLGA NPs was assessed to rule out non-specific toxicity. Both formulations showed no observable toxicity towards MCF-7 cells up to the tested concentrations of 2 mg/mL and for a treatment period of 72 h (**Figure 4A**). Next, MCF-7 cells were treated with free Dox, Dox-loaded nanoghosts or Dox-loaded PLGA NPs for 6 h, followed by 66 hours of incubation. Using a standard MTS cytotoxicity assay, free Dox was shown to have a half maximal inhibitory concentration (IC50) of 0.5 μ M, while nanoghosts and PLGA NP had respective IC50 of 4 μ M and 12 μ M. (**Figure 4B**).

The difference in cytotoxicity between free drug and NP samples could be because of the fundamentally different routes of uptake. Comparing the two formulations, it can be seen that the nanoghosts had a significantly lesser IC50 value compared to PLGA NPs. Therefore, these results demonstrate the significance of using biological cell membrane coatings for efficient nanoparticle-mediated drug delivery applications.

In summary, the nanoghosts combine both synthetic and biological features for enhanced target specificity and efficacy. Further studies planned include optimizing the ratio of PLGA NPs and cell membranes to minimize the presence of free cell membranes that might cause physical aggregation. *In vivo* experiments aimed at investigating the pharmacokinetic and pharmacodynamic profiles, tumor-specific accumulation and tumor reduction of the nanoghosts will be performed in MCF-7 xenograft mice models. We believe that cell membrane-coated NP systems such as our monocyte cell membrane-derived nanoghosts hold great promise as biomimetic applications for targeted cancer therapy, and that further research will only help advance its clinical relevance.

Acknowledgements

JMC acknowledges start-up grant funding from Nanyang Technological University (NTU) and the Lee Kong Chian School of Medicine (LKCmedicine). We thank Ronnie Fang and Liangfang Zhang for experimental advice.

References

- S. Krishnamurthy, R. Vaiyapuri, L. Zhang and J. M. Chan, *Biomater. Sci.* 2015.
- L. Zhang, F. Gu, J. Chan, A. Wang, R. Langer and O. Farokhzad, *Clin. Pharmacol. Ther.* 2007, **83**, 761-769.
- S. Krishnamurthy, X. Ke and Y. Y. Yang, *Nanomedicine*, 2015, **10**, 143-160.
- C. Latorre-Romero, M. R. Marin-Yaseli, C. Belmar-Lopez, R. del Moral, P. C. Marijuan, M. Quintanilla and P. Martin-Duque, *Clin. Transl. Oncol.* 2011, **13**, 10-17.
- M. Hamidi, A. Zarrin, M. Foroozesh and S. Mohammadi-Samani, *J. Control. Release*, 2007, **118**, 145-160.
- A. Fraternali, A. Casabianca, C. Orlandi, A. Cerasi, L. Chiarantini, G. Brandi and M. Magnani, *Antiviral Res.* 2002, **56**, 263-272.
- R. Luo, S. Mutukumaraswamy, S. S. Venkatraman and B. Neu, *J. Mater. Sci. Mater. Med.* 2012, **23**, 63-71.
- C. Théry, M. Ostrowski and E. Segura, *Nat. Rev. Immunol.* 2009, **9**, 581-593.
- A. J. O'Loughlin, C. A. Woffindale and M. J. A. Wood, *Curr. Gene Ther.* 2012, **12**, 262-274.
- C. D. Porada and G. Almeida-Porada, *Adv. Drug Deliv. Rev.* 2010, **62**, 1156-1166.
- N. E. Toledano Furman, Y. Lupu-Haber, T. Bronshtein, L. Kaneti, N. Letko, E. Weinstein, L. Baruch and M. Machluf, *Nano Lett.* 2013, **13**, 3248-3255.
- N. Doshi, A. J. Swiston, J. B. Gilbert, M. L. Alcaraz, R. E. Cohen, M. F. Rubner and S. Mitragotri, *Adv. Mater.* 2011, **23**, H105-H109.
- C. H. June, B. R. Blazar and J. L. Riley, *Nat. Rev. Immunol.* 2009, **9**, 704-716.
- K. Iguchi, F. Sakurai, K. Tomita, K. Katayama, T. Yamaguchi, K. Kawabata, M. Tagawa, M. Kawabata, T. Shirakawa and H. Mizuguchi, *Cancer Gene Ther.* 2012, **19**, 118-125.
- T. Zhang, K. Hamada, M. Hyodo, H. Itoh, K. Tani, H. Goda, K. I. Nakashiro and H. Hamakawa, *Oncol. Rep.* 2011, **25**, 795-802.
- A. T. Power, J. H. Wang, T. J. Falls, J. M. Paterson, K. A. Parato, B. D. Lichty, D. F. Stojdl, P. A. J. Forsyth, H. Atkins and J. C. Bell, *Mol. Ther.* 2007, **15**, 123-130.
- S. Ravilla, B. R. Chandu, S. Nama, B. Nagaveni, *Journal of Applied Pharmaceutical Science* 2012, **2**, 166-176.
- R. H. Fang, C. M. J. Hu, B. T. Luk, W. W. Gao, J. A. Copp, Y. Y. Tai, D. E. O'Connor and L. F. Zhang, *Nano Lett.* 2014, **14**, 2181-2188.
- C.-M. J. Hu, L. Zhang, S. Aryal, C. Cheung, R. H. Fang and L. Zhang, *Proc. Natl. Acad. Sci. U.S.A.* 2011, **108**, 10980-10985.
- S. A. McCarthy, G.-L. Davies, Y. K. Gun'ko, *Nature protocols* 2012, **7**, 1677-1693
- C. M. J. Hu, R. H. Fang and L. F. Zhang, *Adv. Healthc. Mater.* 2012, **1**, 537-547.
- H. Maeda, J. Wu, T. Sawa, Y. Matsumura and K. Hori, *J. Control. Release* 2000, **65**, 271-284.
- A. Lejeune, M. Moorjani, C. Gicquaud, J. Lacroix, P. Poyet and R. C. Gaudreault, *Anticancer Res.* 1994, **14**, 915-919.
- A. Alachi and M. Boroujerdi, *Drug. Dev. Ind. Pharm.* 1990, **16**, 2199-2219.
- A. Alachi and M. Boroujerdi, *Drug. Dev. Ind. Pharm.* 1990, **16**, 1325-1338.
- A. C. Anselmo, V. Gupta, B. J. Zern, D. Pan, M. Zakrewsky, V. Muzykantov and S. Mitragotri, *ACS Nano* 2013, **7**, 11129-11137.
- E. Chambers and S. Mitragotri, *J. Control. Release* 2004, **100**, 111-119.
- E. Chambers and S. Mitragotri, *Exp. Biol. Med.* 2007, **232**, 958-966.
- C. M. J. Hu, R. H. Fang, B. T. Luk and L. F. Zhang, *Nat. Nanotechnol.* 2013, **8**, 933-938.
- W. Gao, C. M. J. Hu, R. H. Fang, B. T. Luk, J. Su and L. Zhang, *Adv. Mater.* 2013, **25**, 3549-3553.
- J.-G. Piao, L. Wang, F. Gao, Y.-Z. You, Y. Xiong and L. Yang, *ACS Nano*, 2014, **8**, 10414-10425.
- L. L. Li, J. H. Xu, G. B. Qi, X. Z. Zhao, F. Q. Yu and H. Wang, *ACS Nano*, 2014, **8**, 4975-4983.
- D. J. Schwartzenruber, D. H. Lawson, J. M. Richards, R. M. Conry, D. M. Miller, J. Treisman, F. Gailani, L. Riley, K. Conlon, B. Pockaj, K. L. Kendra, R. L. White, R. Gonzalez, T. M. Kuzel, B. Curti, P. D. Leming, E. D. Whitman, J. Balkissoon, D. S. Reintgen, H. Kaufman, F. M. Marincola, M. J. Merino, S. A. Rosenberg, P. Choyke, D. Vena and P. Hwu, *N. Engl. J. Med.* 2011, **364**, 2119-2127.
- A. Parodi, N. Quattrocchi, A. L. van de Ven, C. Chiappini, M. Evangelopoulos, J. O. Martinez, B. S. Brown, S. Z. Khaled, I. K. Yazdi and M. V. Enzo, *Nat. Nanotechnol.* 2013, **8**, 61-68.
- M. Xuan, J. Shao, L. Dai, Q. He and J. Li, *Adv. Healthc. Mater.* 2015.
- L. Zhang, J. M. Chan, F. X. Gu, J.-W. Rhee, A. Z. Wang, A. F. Radovic-Moreno, F. Alexis, R. Langer and O. C. Farokhzad, *ACS Nano*, 2008, **2**, 1696-1702.
- J. M. Chan, L. Zhang, R. Tong, D. Ghosh, W. Gao, G. Liao, K. P. Yuet, D. Gray, J.-W. Rhee and J. Cheng, *Proc. Natl. Acad. Sci. U.S.A.* 2010, **107**, 2213-2218.
- D. Laoui, E. Van Overmeire, P. De Baetselier, J. A. Van Ginderachter and G. Raes, *Front. Immunol.* 2014, **5**.
- Q. Chen, X. H.-F. Zhang and J. Massagué, *Cancer cell*, 2011, **20**, 538-549.
- R. A. Jain, *Biomaterials*, 2000, **21**, 2475-2490.

FIGURE LEGENDS

Figure 1. (A) Illustration of cell membrane purification and (B) nanoghost preparation. (A) U937 monocyte cell membranes were extracted by hypotonic lysis, homogenization and serial ultracentrifugation. (B) Subsequently, nanoghosts were prepared by serial extrusion with pre-formed PLGA NP and U937 cell membrane fractions.

Figure 2. (A) TEM characterization of PLGA NPs and (B) nanoghosts. Scale bar: 200 nm. (C) Dynamic light scattering studies on nanoghosts and PLGA NPs. (D) Cumulative drug release from nanoghosts and PLGA NPs in PBS at 37 °C.

Figure 3. (A) Western blot studies on U937 cell membrane extract and nanoghosts. Percentage of FITC-positive MCF-7 (B) and mouse skeletal myocyte (C) cells in cell populations treated with either FITC-loaded nanoghosts or PLGA NPs for 2, 4 or 6 h. Statistical analysis was performed using the student's t-test. Differences are considered statistically significant when probability $P < 0.05$. *, $P < 0.05$ vs. PLGA.

Figure 4. (A) Cytotoxicity of 'blank' nanoghosts and PLGA NPs against MCF-7 cells. (B) Cytotoxicity of Dox-loaded nanoghosts and PLGA NPs against MCF-7 cells. Statistical analysis was performed using the student's t-test. Differences are considered statistically significant when probability $P < 0.05$. *, $P < 0.05$ vs. PLGA at all Dox concentrations from 2 μM .




Figure 1. (A) Illustration of cell membrane purification and (B) nanoghost preparation. (A) U937 monocyte cell membranes were extracted by hypotonic lysis, homogenization and serial ultracentrifugation. (B) Subsequently, nanoghosts were prepared by serial extrusion with pre-formed PLGA NP and U937 cell membrane fractions.
88x52mm (220 x 220 DPI)




Figure 2. (A) TEM characterization of PLGA NPs and (B) nanoghosts. Scale bar: 200 nm. (C) Dynamic light scattering studies on nanoghosts and PLGA NPs. (D) Cumulative drug release from nanoghosts and PLGA NPs in PBS at 37 °C.
80x153mm (220 x 220 DPI)




Figure 3. (A) Western blot studies on U937 cell membrane extract and nanoghosts. Percentage of FITC-positive MCF-7 (B) and mouse skeletal myocyte (C) cells in cell populations treated with either FITC-loaded nanoghosts or PLGA NPs for 2, 4 or 6 h. Statistical analysis was performed using the student's t-test. Differences are considered statistically significant when probability $P < 0.05$. *, $P < 0.05$ vs. PLGA. 159x380mm (300 x 300 DPI)




Figure 4. (A) Cytotoxicity of 'blank' nanoghosts and PLGA NPs against MCF-7 cells. (B) Cytotoxicity of Dox-loaded nanoghosts and PLGA NPs against MCF-7 cells. Statistical analysis was performed using the student's t-test. Differences are considered statistically significant when probability $P < 0.05$. *, $P < 0.05$ vs. PLGA at all Dox concentrations from 2 μM .

99x163mm (300 x 300 DPI)