

# Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

**Synthesis of 6,8-9 Poly-substituted Purine Analogue Libraries as Pro-Apoptotic Inducers of Human Leukemic Lymphocytes and DAPK-1 Inhibitors.**

AUTHORS: Maria J. Pineda de las Infantas<sup>a\*</sup>, Sara Torres-Rusillo<sup>b</sup>, Juan Diego Unciti-Broceta<sup>c</sup>, Pablo Fernandez-Rubio<sup>b</sup>, Maria Angelica Luque-Gonzalez<sup>a,c</sup>, Miguel A. Gallo<sup>a</sup>, Asier Unciti-Broceta<sup>c</sup>, Ignacio J. Molina<sup>b,e</sup>, and Juan J. Díaz-Mochón<sup>a,c,e</sup>

AFFILIATIONS: <sup>a</sup>*Departamento de Química Farmacéutica y Orgánica. Facultad de Farmacia, Universidad de Granada, Campus de Cartuja s/n, 18011 Granada, Spain.*

<sup>b</sup>*Instituto de Biopatología y Medicina Regenerativa, Centro de Investigación Biomédica, Universidad de Granada, Parque Tecnológico de Ciencias de la Salud (PTS), Avda. del Conocimiento s/n, 18016 Armilla, Granada, Spain.*

<sup>c</sup>*Centro Pfizer - Universidad de Granada - Junta de Andalucía de Genómica e Investigación Oncológica (GENYO), Parque Tecnológico de Ciencias de la Salud (PTS), Avenida de la Ilustración 114, 18016 Granada, Spain.*

<sup>d</sup>*Edinburgh Cancer Research UK Centre, MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Crewe Road South, Edinburgh EH4 2XR, UK;*


<sup>e</sup>*IJM and JJD-M share credit for senior authorship.*

KEYWORDS: Purine synthesis, DAPK1, kinase inhibitors, target-blinded phenotypic screening, apoptosis, human leukemic lymphocytes.

\*To whom the correspondence should be addressed. Email: [juanjose.diaz@genyo.es](mailto:juanjose.diaz@genyo.es); [imolina@ugr.es](mailto:imolina@ugr.es); [mjpineda@ugr.es](mailto:mjpineda@ugr.es)

## Table of contents entry

6,8,9-substitued purines are presented as interesting candidates to study the role of DAPK1 kinase in triggering apoptosis in cancer cells.



## ABSTRACT

A 18-member library of 6,8,9-poly-substituted purines was prepared from pyrimidines, primary alcohols and N,N-dimethylamides under basic conditions via a novel one-pot synthetic pathways controlled by amide sizes and the novel analogues were tested against two leukemia cell lines: Jurkat (acute T cell leukemia) and K562 (chronic erythroleukemia) cells. Compounds having a benzyloxy group at C6 position of the aromatic ring exhibited antiproliferative activity in Jurkat cells whereas all compounds induced a lower effect on K562 cells. Analysis of cell cycle, Annexin-V staining and cleavage of initiator caspases assays showed that the active purine analogues induce cell death by apoptosis. Based on these results, a new purine derivative was synthesized, 6-benzyloxy-9-tert-butyl-8-phenyl-9H-purine (**6d**), which displayed the highest activity of the series against Jurkat cell lines. Finally,  $^{33}\text{P}$ -radiolabeled kinase assays using 96 recombinant human kinases known to be involved in apoptotic events were performed. Just one of the kinases tested, DAPK-1, was inhibited 50% or more by the phenotypic hits at 10  $\mu\text{M}$ , suggesting that the inhibition of this target could be responsible for the

induction of cell death by apoptosis. In agreement with the phenotypic results, the most active antiproliferative agent, **6d**, displayed also the lowest IC<sub>50</sub> value against recombinant DAPK1 (2.5  $\mu$ M), further supporting the potential role of this protein on the observed functional response. DAPK-1 inhibition led by **6d** together with its pro-apoptotic properties against Jurkat line makes it an interesting candidate for further investigate the role of DAPK1 kinase in triggering apoptosis in cancer cells, a role which is attracting recent interest.

## 1. Introduction

The new paradigm of anticancer therapy is the development of new drugs that enable personalized medicine. Targeted therapy, which encompasses therapies with antibodies and with small-molecule drugs, has that idyllic aim. In contrast to classic cytotoxic chemotherapies, which killed rapidly dividing cells, targeted therapies are focused to tackle essential biochemical pathways, acting against the progression and proliferation of tumour cells.<sup>8</sup> Biomolecules, which participate in these essential pathways, are also present in normal tissues, but they are either mutated or overexpressed in tumour tissues if compared with normal ones. This makes them excellent molecular candidates to be targeted. In this particular area, protein kinases can be considered cornerstones of many of these cancer molecular mechanisms.<sup>26</sup> Protein kinase networks control cellular signalling pathways and are key elements in tumour development and progression which use ATP, a purine derivative, to transfer phosphate groups.<sup>14</sup> Purine derivatives are synthetic compounds that mimic their endogenous counterparts acting as antimetabolites. These kinds of antimetabolites interrupt cellular processes by interfering with essential enzymes involved in the intracellular signalling, the tyrosine kinases<sup>10</sup>.

Tackling kinases by small-molecules is therefore a very attractive way to both develop new antitumor drugs and to elucidate new molecular pathways involved in cancer.<sup>25</sup> In fact, at least fourteen small-molecules which act as kinase inhibitors have been approved over the last few years by the FDA for its use as antitumor drugs in humans.<sup>19</sup>

A total of 518 proteins have been characterized up to date as kinases, forming the so-called human kinome.<sup>15</sup> This means that there is still room for finding both drug targets and biological pathways related to cancer which remain unknown. To confront this challenge, new small-molecule libraries need to be prepared and used to interrogate kinase activity in cellular and biochemical systems. As most small-molecule kinase inhibitors target the conserved adenosine triphosphate (ATP)-binding site, purine libraries decorated with different substituents are expected to have a high probability of yielding bioactive compounds and kinase inhibitors<sup>9</sup>, and consequently, purine-derived compounds have been subject to vast exploration in both heterocyclic and medicinal chemistry.


The aim of our research was to develop a drug discovery workflow process divided step-wise into: i) synthesis of poly-substituted purine libraries via novel synthetic pathways developed by our group; ii) in vitro cell-based assays using human erythroleukemic cells and leukemic T lymphocytes to select hits based on their antiproliferative activity and to determine the mechanism(s) by which cell proliferation is affected; and iii) target deconvolution by searching potential molecular targets known to be involved in pathways leading to the observed antiproliferative effects.

## 2. Results and discussion

### 2.1 Chemistry


Our group has developed new one-pot synthetic ways to prepare 6,8,9-substituted purines from 6-chloro-4,5-diamino-pyrimidines using alcohols and N,N-dimethylamides under basic and metal-free conditions (see ref. 2 and Pineda de las Infantas et al, submitted). Depending on the size of the R1 group of the amides employed to carry out the chemical reactions (Scheme 1), two different synthetic pathways can take place, resulting in purine analogues whose substituent at C8 comes either from the amide (Route A) or the alcohol's carbon chains (Route B). We have just reported these findings elsewhere to explain these competing reactions (Pineda de las Infantas et al., submitted).

Briefly, Route A proceeds via in-situ generation of alkoxyiminium species, which are created by reaction between amides and alkoxides, leading to purine derivatives with either H or methyl in C8 when using N,N-dimethylformamide or N,N-dimethylacetamide, respectively. When steric hindrance of amides increase with the use of N,N-dimethylbenzamide or N,N-dimethylpropionamide, Route A is then impeded and a metal-free tandem alcohol oxidation/annulation reaction occurs giving rise to Route B products. In this case, the purine's C8 substituent originates from the primary alcohols used in the reactions. Therefore, competition between alkoxyiminium formation and metal-free oxidative coupling of primary alkoxides and diaminopyrimidines with Schiff base formation and subsequent annulation can be controlled. In both routes, SNAr of the chloro atom at C6 by alkoxides take place and, along with the R<sup>3</sup> group of the starting pyrimidine, implement structural diversity in a one-pot reaction that overcome the low yields found in some of the reactions. Figure 1 shows the set of 18 purines presented here which were prepared by either Route A or B. Therefore, this synthetic platform allows creating a diversity of purine analogues in a



Scheme 1. Synthetic routes to generate purine types "A" or "B". Size of R1 controls which synthetic mechanism is followed under the same reaction conditions.  
83x88mm (300 x 300 DPI)


## Route A Purines



5 a-n

- 5a:** R<sup>3</sup>= *i*Pr, R<sup>2</sup>= Et, R<sup>1</sup>= H  
**5b:** R<sup>3</sup>= *i*Pr, R<sup>2</sup>= Et, R<sup>1</sup>= Et  
**5c:** R<sup>3</sup>= *i*Pr, R<sup>2</sup>= Et, R<sup>1</sup>= Me  
**5d:** R<sup>3</sup>= *i*Pr, R<sup>2</sup>= Bn, R<sup>1</sup>= H  
**5e:** R<sup>3</sup>= *i*Pr, R<sup>2</sup>= Bn, R<sup>1</sup>= Me  
**5f:** R<sup>3</sup>= Bn, R<sup>2</sup>= Et, R<sup>1</sup>= H  
**5g:** R<sup>3</sup>= Bn, R<sup>2</sup>= Et, R<sup>1</sup>= Me  
**5h:** R<sup>3</sup>= Bn, R<sup>2</sup>= Et, R<sup>1</sup>= Et  
**5i:** R<sup>3</sup>= Bn, R<sup>2</sup>= Bn, R<sup>1</sup>= H  
**5j:** R<sup>3</sup>= *i*Pr, R<sup>2</sup>= *i*Pr, R<sup>1</sup>= H  
**5k:** R<sup>3</sup>= *i*Pr, R<sup>2</sup>= *i*Pr, R<sup>1</sup>= Me  
**5l:** R<sup>3</sup>= *i*Pr, R<sup>2</sup>= *t*Bu, R<sup>1</sup>= H  
**5m:** R<sup>3</sup>= *i*Pr, R<sup>2</sup>= Me, R<sup>1</sup>= H  
**5n:** R<sup>3</sup>= *t*Bu, R<sup>2</sup>= Et, R<sup>1</sup>= H

## Route B Purines



6 a-d

- 6a:** R<sup>3</sup>= *i*Pr; R<sup>2</sup>=Ph (with propionamide)  
**6b:** R<sup>3</sup>= *i*Pr; R<sup>2</sup>=Et (with benzamide)  
**6c:** R<sup>3</sup>= *i*Pr; R<sup>2</sup>=*i*Pr (with benzamide)  
**6d:** R<sup>3</sup>= *t*Bu; R<sup>2</sup>=Ph (with propionamide)

Fig. 1. Set of 18 purines prepared by either Route A or B (Scheme 1).  
85x96mm (300 x 300 DPI)






parallel and straightforward fashion using a variety of amides and alcohols with different pyrimidines under the same reaction conditions.

## 2.2 Biological Activity

*2.2.1. Antiproliferative activity of compounds 5a-n and 6a-c.* The panel of small molecules was incubated with two haematopoietic tumour cell lines: Jurkat (acute T cell leukaemia) and K562 (chronic erythroleukaemia). Cells were cultured for 48 h in the presence of increasing concentrations of each compound: 50  $\mu$ M and 100  $\mu$ M (SI Figure 1) and 25  $\mu$ M and 200  $\mu$ M (not shown) and cell viability assessed by the MTT colorimetric screening assay. Results shown in SI Figure 1 are expressed as percentage of inhibition of cell proliferation calculated over the values obtained with untreated cells. Proliferation of Jurkat T cells was inhibited at levels close or superior to 50% by a number of the analogues (SI Figure 1), whereas the K562 erythroleukaemic cells were significantly more resistant to the action of the compounds (not shown). From this initial set of data, a sub-pool of compounds which inhibited cell proliferation by 50% or more were chosen for further phenotypic characterization (**5e**, **5i**, and **6a**; Figure 2). Interestingly, the three of them presented a benzoxy group at purine's C6 position.

Dose-response curves in Jurkat and K562 cell lines were determined for the chosen derivatives and their IC<sub>50</sub> values calculated (Table 1), highlighting compound **6a** as the strongest antiproliferative hit. These IC<sub>50</sub> demonstrate that K562 cells are more resistant than Jurkat cells to the antiproliferative activity of the assayed compounds, thus confirming the general tendency of this library.

**5e****6a****5i**


79x43mm (300 x 300 DPI)

**Table 1.** IC<sub>50</sub> values of indicated compounds assayed by inhibition of cell proliferation

Compound	Jurkat ( $\mu\text{M}$ )	K562 ( $\mu\text{M}$ )
<b>5e</b>	205	>500
<b>5i</b>	152	>500
<b>6a</b>	63	154

IC<sub>50</sub> values were calculated over a lineal regression line of specific inhibition of proliferation values in a MTT assay of cells treated for 48 h with increasing amounts of compounds.

*2.2.2 Study of the mechanism responsible for inhibiting cell proliferation.* Cell cycle analyses were performed to determine whether the antiproliferative effect was a consequence of cell cycle arrest, induction of apoptosis or death by necrosis. Jurkat and K562 cells were treated for 48 h with each compound (**5e**, **5i** and **6a**) at different concentrations and the cell cycle profile was determined by flow cytometry after staining of nuclear DNA with propidium iodide. Treatment of Jurkat cells with the chosen sub-library of compounds resulted in accumulation of cells within the sub-G1 region<sup>18</sup> due to DNA fragmentation and, consequently, loss of nuclear DNA content, which is consistent with apoptotic populations. The three compounds induced cell death in Jurkat cells in a dose-dependent manner (Figure 3). Interestingly, the G0/G1 peak was greatly diminished after treatment of cells with our compounds, thus suggesting that no cell cycle arrest was involved.



Effect of compounds 5e, 5i and 6a on cell cycle. A representative experiment showing Jurkat cells incubated for 48 h in the presence of indicated compounds and doses. Cell cycle subpopulations were analyzed by flow cytometry after permeabilisation and staining of cells with Propidium iodide. Additional controls containing equivalent amounts of DMSO used as drug vehicle were included, showing no toxicity in any case (not shown). Percentage of cells within the sub-G1 region are indicated in the plots.  
254x190mm (96 x 96 DPI)

Flow cytometry profiles of K562 cells (SI Figure 2) showed a lower cell population accumulating within the sub-G1 region when comparing the effect induced by each compound to those observed on Jurkat cells, suggesting that the former are more resistant to death induced by compounds **5e**, **5i** and **6a**.

To further analyze the cell death mechanism(s) we used Annexin V, a calcium-dependent phospholipid-binding protein with a high affinity for phosphatidylserine (PS), which is a distinct feature of early stages of apoptosis<sup>12</sup>. Figure 4 shows a representative flow cytometry dot plot of double stained cells with Propidium iodide and Annexin-V. As expected, untreated Jurkat cells (upper left-hand panel) presented a negligible apoptotic cell population (right-hand gates).

Etoposide was used as a positive inductor of apoptosis (lower left-hand panel), and two populations of cells arise after exposure to this agent: early apoptotic cells (Annexin-V<sup>+</sup>/PI<sup>-</sup>) and late apoptotic cells (Annexin-V<sup>+</sup>/PI<sup>+</sup>). In Jurkat cells treated with **5e**, **5i** and **6a** at the indicated doses, transition behaviour from early to late apoptotic stages was observed, confirming that these compounds induce death cell by apoptosis in a dose-dependent manner (Figure 4).

To address whether or not our compounds had a preferential activity on tumour cells, we directly compared their action on two cell lines belonging to the same lineage: Jurkat and N1, both CD4<sup>+</sup> T cells. The latter cells are a long-term allospecific primary cell line established in our laboratory by weekly allostimulation of peripheral blood lymphocytes with the Class II<sup>+</sup> Raji B cell line in the presence of rIL-2.<sup>24</sup> We treated both cells with our compounds and assessed their effect on the cell cycle. Although these compounds, particularly **6a**, have a discreet cytotoxic effect on primary T lymphocytes at high doses, the overall activity of our compounds is significantly lower on primary lymphocytes




Figure 4. Membrane phosphatidylserine expression after treatment of Jurkat cells with compounds 5e, 5i and 6a. Jurkat cells were treated for 48 h with the indicated compounds and doses and the percentages of apoptotic cells was measured by flow cytometry after double staining with propidium iodide and Annexin-V-Fluos. Dot plots represent the results of a typical experiment. Untreated cells and cells exposed to Etoposide were included as negative and positive controls.

73x54mm (300 x 300 DPI)

than that exerted over tumour cells (Figure 5). Since primary T cells remain viable at low and intermediate doses, this could be of importance in the event of a future therapeutic development of this family of compounds.

To gather further direct evidence that treatment of Jurkat cells with the selected compounds resulted in programmed cell death by apoptosis, caspase activation events were analysed by western blot.

Caspases are a series of cysteine-proteases that cleave certain proteins after specific aspartic acid residues.<sup>23</sup> Caspases are expressed as inactive proenzymes that become activated after being cleaved by their preceding caspase in a cascade-like mechanism.<sup>16</sup> Initiator caspases (8 and 9) trigger effector caspases (3, 6 and 7) and, therefore, cleavage of initiator caspases is an early event of the biochemical events leading to apoptosis. The appearance of cleavage products (Figure 6) of caspases 8 and 9 indicates that apoptosis was triggered in cells treated with compound **6a**. Similar effects were observed with compounds **5e** and **5i** (not shown).

*2.2.3. Synthesis and biological evaluation of compound 6d.* The pro-apoptotic activity of compound **6a** prompted us to design and synthesize a new derivative, compound **6d** (Figure 7), by replacing the isopropyl group found at N9 position of the aromatic ring for a tert-butyl group. These groups are known to bind lipophilic pockets of enzymes, improving generally inhibiting activities of small-molecules. With this structure in mind, we decided to apply the novel one-pot synthetic route developed (Scheme 1, Route B) using benzyl alcohol, *N,N*-dimehtylpropanamide and 5-amino-4-tert-butylamino-6-chloropyrimidine in the presence of NaH. Following synthesis, purification and characterization, Jurkat and K562 cells were incubated for 48h with compound **6d** and cell viability assessed by MTT. Evident dose-dependent curves of




Figure 5. Effect of indicated compounds on primary lymphocytes and Jurkat cells. Primary cell line N1 and Jurkat cells were treated for 48h at the indicated doses of compounds 5e, 5i and 6a and cell cycles were analysed by flow cytometry after staining with Propidium Iodide. Graphs represent the mean $\pm$ S.D. of cells gated within the sub-G1 region obtained from three independent experiments. UT: Untreated.  
254x108mm (96 x 96 DPI)






Figure 6. Study of early signaling events of apoptosis induced by compound 6a in Jurkat cells. Western blot analysis of total lysates of Jurkat cells after treatment with either compound 6a (200  $\mu$ M, left-hand lanes) or equivalent amount of vehicle (DMSO, right-hand lanes) for 5, 6 or 7 h. Membranes were probed with antibodies against initiator caspases 8 or 9, using actin as loading control. Cells treated with Etoposide were included as positive control. Arrows indicate cleavage products (43/41 kD bands for caspase 8 and 37/< kD bands for caspase 9), which are detected after caspase activation whereas upper bands correspond to native proteins. Similar results were obtained after treating cells with compounds 5e and 5i (not shown). NT: Not

treated  
56x35mm (300 x 300 DPI)




Figure 7. Structure of compound 6-benzyloxy-9-tert-butyl-8-phenyl-9H-purine 6d.  
32x36mm (300 x 300 DPI)

inhibition of cell proliferation were detected after exposure of cells to purine **6d**, allowing calculation of its IC<sub>50</sub> values against Jurkat (Figure 8, solid points) and K562 cells on a 11-point 1:2 serial dilution, yielding values of 29±2.4 μM for Jurkat and 120±10 μM for K562.

To determine the effects of the new compound, cell cycle analyses of Jurkat (Figure 9, panel A) and K562 cells (SI Figure 3) were carried out after exposure of cells to purine **6d**. Treatment of Jurkat cells for 24 h resulted in cell death induction which was determined by the percentage of cells accumulated in the sub-G1 fraction (Figure 9, panel A). Consistent with results obtained with compounds **5e**, **6a** and **5i**, K562 cells showed increased resistance to induction of cell death after treatment with **6d** (SI Figure 3), which also agrees with the lower antiproliferative effect observed for this cell line.

As described above for other compounds, the mechanism by which **6d** induces death in Jurkat cells was further studied. Again, Annexin-V flow cytometric analysis of cells treated with compound **6d** (Figure 9, panel B) showed a stronger apoptotic-induced effect than those triggered by compounds **5e**, **5i** and **6a** (Figure 4).

The capacity of compound **6d** to induce activation of caspases was then assessed by Western Blotting (Figure 9, panel C). Treatment of Jurkat cells with **6d** for 2, 4 and 6 h yielded similar results to those shown in Figure 5, clearly showing the expected cleavage products of both caspases 8 and 9 as early as 2 h of treatment.

To further confirm that the cytotoxic effect exerted by compound **6d** was strictly dependent upon caspase activation, we pre-treated Jurkat cells with the pan-caspase inhibitor Q-VD-OPh, which is known to be a potent and irreversible blocker of cell death mediated by apoptosis with reduced toxicity.<sup>3</sup> Figure 10 shows that pre-treatment of Jurkat cells with the pan-caspase inhibitor inhibits almost completely apoptosis of




Figure 8.- IC<sub>50</sub> value of compound 6d in Jurkat cells as determined by inhibition of cell proliferation. A 11-point 1:2 serial dilution of compound 6d, starting at 800 μM, was used to determine its IC<sub>50</sub> value by the MTT assay after treatment of Jurkat cells for 48 h (solid circles). Equivalent amounts of DMSO used as vehicle in each dilution were set as negative controls (open circles). Results indicate percentages of cell proliferation over the results obtained with untreated cells. Data are expressed as mean ± SEM.  
273x178mm (300 x 300 DPI)




Figure 9. A) Effect of compound 6d on the cell cycle of Jurkat cells. Jurkat cells were treated for 48 h with 6d at indicated concentrations or their equivalent amounts of DMSO used as vehicle. Cells gated on the sub-G1 region after staining with propidium iodide are indicated. B) Membrane phosphatidylserine expression after treatment of Jurkat cells with compound 6d. Jurkat cells were treated for 24 h with compound 6d and the indicated doses and the percentages of apoptotic cells was measured by flow cytometry after double staining with propidium iodide and Annexin-V-Fluos. Dot plots represent the results of a typical experiment. Untreated cells and cells exposed to Etoposide were included as negative and positive controls. C) Percentages of apoptotic cells after treatment of Jurkat cells with compound 6d. Graph summarizes values of percentages of apoptotic cells and SEM obtained from three independent experiments. UT: Untreated. D) Activation of initiator caspases 8 and 9 after cell treatment with compound 6d. Western blot analysis of total lysates from Jurkat cells after treatment with either compound 6d (200 μM, right-hand lanes) or vehicle (DMSO, left-hand lanes) for 2, 4 and 6 h. Membranes were probed with antibodies against initiator caspases 8 or 9, using actin as loading control. Arrows indicate cleavage products (43/41 kD bands for caspase 8 products and 37/< kD bands for caspase 9 ones) after caspase activation whereas upper bands correspond to native proteins (caspase 8 correspond to 55/57 kD band and caspase 9 to 45 kD band)

97x67mm (300 x 300 DPI)




Figure 10.- Pre-treatment of Jurkat cells with the pan-caspase inhibitor Q-VD-OPh abrogates cell death mediated by compound 6d. A) Jurkat cells were pre-treated for 1 h with 50  $\mu$ M of the Q-VD-OPh pan-caspase inhibitor and cultured thereafter for 24 h in the presence of 200  $\mu$ M of compound 6d (bottom left panel). Controls include untreated cells (upper left panel); cells treated with Q-VD-OPh alone (upper right panel) or only with compound 6d (bottom left panel). Cells were stained with propidium iodide and cell cycles analysed by flow cytometry. Percentages of cells within the sub-G1 region are indicated in the plots. B) Western Blot analysis of Caspase-8 activation. The same cultures of cells examined in Panel A were collected 8 h after treatment with compound 6d and cytosolic lysates were resolved by SDS-PAGE and membranes probed with the anti-caspase 8 mAb. Actin was used as loading control. UT: Untreated.

194x190mm (96 x 96 DPI)

Jurkat cells induced by compound 6d, as analysed by cell cycle staining with propidium iodide (Figure 10 A, bottom panels). Furthermore, this effect is clearly specific as Western Blot analysis shows that the Caspase-8 cleavage products strongly observed after treatment of Jurkat cells with compound 6d are absent in those cells pre-treated with the pan-caspase inhibitor (Figure 10, panel B). As expected, cells treated with the Q-VD-OPh alone showed no toxicity (Figure 10 A, upper right panel)

These results are clear evidence that activation of caspases is an absolute requirement for the cytotoxic effect mediated by compound 6d, thereby confirming that cell death occurs by apoptosis.

*2.2.4. Kinase screening assay.* Since both cell cycle studies and caspase 8/9 activation demonstrated that the chosen sub-family of compounds triggered apoptosis, a panel of kinases from the human kinome were selected. The selection criteria was based on their known involvement in apoptotic processes. 5e and 6a purines were chosen as representative compounds to perform the assay in order to see if this family of purines, presenting a benzoxy group at C6 position, were capable of inhibiting kinases. Hence, 96 recombinant kinases were used to run a <sup>33</sup>P-radiolabeled kinase assay profiling using a single-dose duplicate mode at a 10 μM purine concentration with 10 μM ATP. Surprisingly, just one kinase out of the 96 tested showed a significant reduction in kinase activity by both compounds. This kinase was Death Associated Protein Kinase-1 (DAPK1) which presented a percentage of activity of 50.9% and 44.3% when incubated with compounds 5e and 6a, respectively. Compounds 5e, 5i, 6a and 6d purines were then tested against DAPK1 via a <sup>33</sup>P-radiolabeled kinase assay using 10-doses to determine their IC<sub>50</sub> (DAPK1) values (Table 2).

---

**Table 2.** IC<sub>50</sub> values of indicated compounds calculated as inhibition of DAPK1 enzymatic activity.

---

Compound	DAPK1 (μM)
<b>5e</b>	23
<b>5i</b>	5.9
<b>6a</b>	ND*
<b>6d</b>	2.5

---

\*The IC<sub>50</sub> curve for **6a** against DAPK1 presented a Gaussian profile, which suggested inconsistent solubility of this compound and therefore impeded us a precise calculation of the IC<sub>50</sub> dose in this system (See SI Figure 4-7). Whereas both **6a** and **6d** present very similar clogP values, the added steric behavior of *tert*-butyl groups was expected to also influence its solubility properties. Furthermore, the sigmoid curve obtained on the MTT assay (Figure 6) is a strong indication of the improved solubility of compound **6d**.

---



This assay revealed that our compounds inhibited DAPK1 at the low micromolar level. This result is remarkable since an elegant analysis where 178 small-molecules kinase inhibitors were assayed against 300 recombinant protein kinases<sup>1</sup> showed that DAPK1 has 0.03 as Selective Score 50, one of the lowest for any of the kinases included in that study. This meant that just 5 out of the 178 compounds tested at 10  $\mu$ M by this group against DAPK1 inhibited this kinase more than 50%, evidence of how unusual is to hit this target. Moreover, these 5 compounds were very promiscuous, targeting many other kinases. While more studies are needed in order to link their pro-apoptotic features with their DAPK1 inhibition, our results indicate that these derivatives do not inhibit other kinases known to be involved in apoptotic events. Furthermore, our data supports recent findings by Schneider-Stockemil<sup>4</sup> that for the first time reported that selective inhibition of DAPK1, an enzyme that plays a critical role in interferon- $\gamma$ -mediated apoptosis<sup>5-7,27</sup>, might impinge upon the biology of tumour cells via STAT3 activation. The relevant role of DAPK1 in pro-apoptotic pathways is underlined by the fact that many chronic lymphocytic leukaemia cells downregulate the expression of the protein to favour tumour progression.<sup>11,20</sup> Our findings, as well as those of Schneider-Stockemil<sup>4</sup>, may thus be surprising if we only view DAPK1 as a pro-apoptotic kinase. The pro-apoptotic activity of the enzyme requires the integrity of the Death Domain<sup>6</sup>, but growing evidence indicates that post-transcriptional modifications may confer opposing functions to the protein. For instance, autophosphorylation restrains the apoptotic activity of members of the DAPK family<sup>22</sup> and disruption of the cathepsin B-DAPK1 interaction breaks a multiprotein complex whose perturbation sensitizes cells to apoptosis.<sup>13</sup> Since our compounds have a preferential activity on Jurkat cells when compared to K562, a cell line with low expression of DAPK-1, it can be speculated that the mechanism of action for the observed pro-apoptotic effects after inhibiting DAPK1

activity may be related to posttranscriptional mechanisms resulting in sensitizing treated cells to death by apoptosis. The precise mechanisms involved, however, need further investigations which will answer the question of whether targeting DAPK1 is a sound strategy for the treatment of certain types of tumours.

### 3. Conclusions

To overcome the disadvantages of target-centric drug discovery strategies and promote a more empirical, evidence-led, approach to anticancer drug discovery, novel derivatives were directly tested in disease relevant leukemia cell lines. In order to obtain purine analogues, novel chemistry pathways developed by our group and based on one-pot synthesis were carried out. The novel chemistry allows to obtain 6,8-9 poly-substituted purines starting from diamino-pyrimidines and a range of different primary alcohols and amides. Following this approach, a series of 18 purine analogues exhibiting antiproliferative activity and pro-apoptotic properties on the acute T cell leukemia Jurkat cell line were identified, having all of them a benzyloxy group at C6 position and being 9-*tert*-butyl-6-(benzyloxy)-8-phenyl-9*H*-purine (**6d**) the most active one displaying a  $IC_{50}$  value of 29  $\mu$ M on Jurkat cells. Cell cycle analyses, Annexin-V staining, cleavage of initiator caspases and blocking of the cytotoxic effects after pre-incubation of cells with a pan-caspase inhibitor are cumulative evidence clearly showing that cell death was induced by apoptosis. Finally, to assess if their pro-apoptotic activities were due to inhibition of multiple kinases, 96 kinases known to be involved in apoptosis processes were assayed against these purines. Surprisingly, just one kinase, DAPK1, was inhibited at significant levels, with **6d** purine compound being the one showing the lowest  $IC_{50\_DAPK1}$  (2.5  $\mu$ M). Compound **6d** is therefore the first selective inhibitor of DAPK1. Since all other available inhibitors target multiple protein kinases, this makes it difficult to dissect precisely the role of DAPK1 on certain

biological processes. Among these, growing evidence of the role of DAPK1<sup>28</sup> on cancer processes is appearing and our results tend to support this. Therefore, the use of a selective inhibitor could shed light on the exact participation of DAPK1 in tumor development. Furthermore, the fact that primary T lymphocytes are relatively refractory to cytotoxicity induced by our compounds make it worth to pursue further studies aiming to explore DAPK1 as a potential therapeutic target in certain tumors.

#### 4. Experimental procedures

##### 4.1. Synthesis and Characterization of Compounds

###### 4.1.1. General

Reaction courses and products mixtures were routinely monitored by TLC on silica gel Merck 60-200 mesh silica gel. Melting points were determined on a Stuart Scientific SMP3 apparatus and are uncorrected. <sup>1</sup>H-NMR spectra were obtained in CDCl<sub>3</sub>, CD<sub>3</sub>OD solutions on a Varian Inova Unity (300 MHz) and Varian Direct Drive (400 MHz and 500 MHz). Chemical shifts (δ) are given in ppm upfield from tetramethylsilane. <sup>13</sup>C-NMR spectra were obtained in CDCl<sub>3</sub>, CD<sub>3</sub>OD solutions on a Varian Direct Drive (125 MHz). All products reported showed <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectra in agreement with the assigned structures. Mass spectra were obtained by electrospray (ESY) with a LCT Premier XE Micromass Instrument (High resolution mass spectrometry).

###### 4.1.2. General procedure for the preparation of compounds 4 (a-c):

To commercially available 4,6-dichloro-5-nitropyrimidine 1 (10 mmol, 1 equiv.), in anhyd. THF (15 ml) was added NaHCO<sub>3</sub> (11 mmol, 1.1 equiv.). The mixture was then heated to 55°C and at this temperature the appropriate amine (11 mmol, 1.1 equiv.)

dissolved in THF was added dropwise. After 1 h, the mixture was cooled at room temperature, filtered and evaporated to give a crude product, which was directly used for the next step without purification.

This crude solution (5 mmol, 1 equiv.) in EtOH (10 ml) containing SnCl<sub>2</sub>·2H<sub>2</sub>O (25 mmol, 5 equiv.), was refluxed for 1 h, monitoring the reaction by TLC. The mixture was then cooled at room temperature and NaHCO<sub>3</sub> was added until pH 8 was reached. After two extractions with EtOAc (10 ml for each one), the organic phase was washed with an aq. saturated solution of NaCl (2x25 mL) and dried on Na<sub>2</sub>SO<sub>4</sub>. Evaporation under vacuum gave a solid that was purified by column chromatography eluting with ethyl acetate/petroleum ether solutions.

*4.1.2.1. 5-Amino-6-chloro-4-isopropylaminopyrimidine (4a).* White solid, yield 73%; mp 185-186°C.  $\delta_{\text{H}}$ (400.58 MHz, CDCl<sub>3</sub>): 8.08 (1H, s, NCHN), 4.68 (1H, bs, NH), 4.29 (1H, m, CH(CH<sub>3</sub>)<sub>2</sub>), 3.33 (2H, bs, NH<sub>2</sub>), 1.27 (6H, d, J=8, CH(CH<sub>3</sub>)<sub>2</sub>).

*4.1.2.2. 5-Amino-4-tert-butylamino-6-chloropyrimidine (4b).* White solid; yield 75%; mp: 135°C.  $\delta_{\text{H}}$ (499.79 MHz, CDCl<sub>3</sub>): 8.03 (1H, s, NCHN), 4.87 (1H, bs, NH), 3.45 (2H, bs, NH<sub>2</sub>), 1.47 (9H, s, C(CH<sub>3</sub>)<sub>3</sub>). ES+HRMS: Calculated M+H = 201.0907. C<sub>8</sub>H<sub>14</sub>N<sub>4</sub>Cl. Obtained: 201.0909.

*4.1.2.3. 5-Amino-4-benzilamino-6-chloropyrimidine (4c).* White solid; yield 65%; mp 205-206°C.  $\delta_{\text{H}}$ (499.79 MHz, CD<sub>3</sub>OD): 7.78 (1H, s, NCHN), 7.35-7.22 (5H, m, Ph), 4.69 (2H, s, CH<sub>2</sub>Ph).  $\delta_{\text{C}}$ (125.68 MHz, CDCl<sub>3</sub>): 150.13, 142.76, 137.79, 132.02, 131.18, 130.72, 127.49, 48.51. ES+HRMS: Calculated M+H = 235.0750. C<sub>11</sub>H<sub>12</sub>N<sub>4</sub>Cl. Obtained: 235.0753.

#### 4.1.3. General procedure for the preparation of compounds 5 (a-n). Synthesis A:

A suspension of NaH (50%) in mineral oil (3.7 mmol, 10 equiv.) was dissolved in a mixture cooled at 0° constituted both from the appropriate alcohol (3.7 mmol, 10 equiv.) and amide (18.5 mmol, 50 equiv.). The solution was stirred at room temperature for 30 min and then at 90°C for the same time. At 90°C compounds 4 (a-c) (0.37 mmol, 1 equiv.), dissolved in the appropriate amide (18.5 mmol, 50 equiv.) were added dropwise and the mixture heated for 24 h. The reaction mixture quenched to pH 7 with an aq. saturated NH<sub>4</sub>Cl solution, was then extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x15 mL), the combined organic extracts dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent evaporated. The residue was chromatographed on silica gel using ethyl acetate/petroleum ether solutions.

*4.1.3.1.1 6-Ethoxy-9-isopropyl-9H-purine (5a).* Light yellow solid; yield 44%.  $\delta_{\text{H}}$ (499.79 MHz, CDCl<sub>3</sub>): 8.49 (1H, s, NCHN), 7.96 (1H, s, NHCHN), 4.88 (1H, m, CH(CH<sub>3</sub>)<sub>2</sub>), 4.64 (2H, q, J=5 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.60 (6H, d, J=5 Hz, NCH(CH<sub>3</sub>)<sub>2</sub>), 1.49 (3H, t, J=5 Hz, OCH<sub>2</sub>CH<sub>3</sub>).  $\delta_{\text{C}}$ (125.68 MHz, CDCl<sub>3</sub>): 160.81, 151.70, 139.53, 121.83, 62.98, 47.33, 22.60, 14.51. ES+HRMS: Calculated M+H = 207.1246. C<sub>10</sub>H<sub>15</sub>N<sub>4</sub>O. Obtained: 207.1245.

*4.1.3.1.2 6-Ethoxy-8-ethyl-9-isopropyl-9H-purine (5b).* Yellow viscous oil; yield 10%.  $\delta_{\text{H}}$ (300.20 MHz, CDCl<sub>3</sub>): 8.47 (1H, s, NCHN), 4.72 (1H, m, CH(CH<sub>3</sub>)<sub>2</sub>), 4.67 (2H, q, J=6 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 2.96 (2H, q, J=6 Hz, CH<sub>2</sub>CH<sub>3</sub>), 1.74 (6H, d, J=6 Hz, NCH(CH<sub>3</sub>)<sub>2</sub>), 1.53 (3H, t, J=6 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.47 (3H, t, J=6 Hz, -CH<sub>2</sub>CH<sub>3</sub>). ES+HRMS: Calculated M+H = 235.1559. C<sub>12</sub>H<sub>19</sub>N<sub>4</sub>O. Obtained: 235.1557.

*4.1.3.2 6-Ethoxy-9-isopropyl-8-methyl-9H-purine (5c).* Yellow viscous oil; yield 10%.  $\delta_{\text{H}}$ (499.79 MHz, CDCl<sub>3</sub>): 8.44 (1H, s, NCHN), 4.72 (1H, m, CH(CH<sub>3</sub>)<sub>2</sub>), 4.62 (2H, q, J=10 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 2.64 (3H, s, CH<sub>3</sub>), 1.68 (6H, d, J=5 Hz, NCH(CH<sub>3</sub>)<sub>2</sub>), 1.50 (3H,

t, J=10 Hz, OCH<sub>2</sub>CH<sub>3</sub>).  $\delta_{\text{C}}$ (125.68 MHz, CDCl<sub>3</sub>): 159.74, 153.15, 151.75, 150.53, 120.71, 62.67, 48.31, 21.24, 15.22, 14.58. ES+HRMS: Calculated M+H = 221.1402. C<sub>11</sub>H<sub>17</sub>N<sub>4</sub>O. Obtained: 221.1396.

4.1.3.1.3. 6-(Benzyloxy)-9-isopropyl-9H-purine (**5d**). White solid; yield 38%; mp 167-169°C.  $\delta_{\text{H}}$ (499.79 MHz, CDCl<sub>3</sub>): 8.56 (1H, s, NCHN), 8.00 (1H, s, NHCHN), 7.56-7.30 (5H, m, Ph), 5.69 (2H, s, OCH<sub>2</sub>Ph), 4.91 (1H, m, CH(CH<sub>3</sub>)<sub>2</sub>), 1.64 (6H, d, J=10 Hz, NCH(CH<sub>3</sub>)<sub>2</sub>).  $\delta_{\text{C}}$ (125.68 MHz, CDCl<sub>3</sub>): 160.53, 151.63, 139.77, 136.27, 130.03, 128.41, 128.29, 128.05, 68.29, 47.43, 22.63. ES+HRMS: Calculated M+H = 269.1402. C<sub>15</sub>H<sub>17</sub>N<sub>4</sub>O. Obtained: 269.1402.

4.1.3.1.4. 6-(Benzyloxy)-9-isopropyl-8-methyl-9H-purine (**5e**). White solid; yield 16%.  $\delta_{\text{H}}$ (499.79 MHz, CDCl<sub>3</sub>): 8.47 (1H, s, NCHN), 7.54-7.30 (5H, m, Ph), 5.65 (2H, s, OCH<sub>2</sub>Ph), 4.75 (1H, m, CH(CH<sub>3</sub>)<sub>2</sub>), 2.65 (3H, s, CH<sub>3</sub>), 1.69 (6H, d, J=10 Hz, NCH(CH<sub>3</sub>)<sub>2</sub>).  $\delta_{\text{C}}$ (125.68 MHz, CDCl<sub>3</sub>): 159.41, 153.57, 150.69, 150.41, 136.42, 128.43, 128.35, 127.98, 120.73, 68.06, 48.35, 21.23, 15.22. ES+HRMS: Calculated M+H = 283.1559. C<sub>16</sub>H<sub>19</sub>N<sub>4</sub>O. Obtained: 283.1558.

4.1.3.1.5. 9-Benzyl-6-ethoxy-9H-purine (**5f**). White solid; yield 62%; mp 133-135°C.  $\delta_{\text{H}}$ (499.79 MHz, CDCl<sub>3</sub>): 8.55 (1H, s, NCHN), 7.88 (1H, s, NHCHN), 7.34-7.28 (5H, m, Ph), 5.41 (2H, s, CH<sub>2</sub>Ph), 4.66 (2H, q, J=10 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.51 (3H, t, J=10 Hz, OCH<sub>2</sub>CH<sub>3</sub>).  $\delta_{\text{C}}$ (125.68 MHz, CDCl<sub>3</sub>): 160.93, 152.33, 141.81, 135.34, 129.08, 128.27, 127.75, 121.39, 63.16, 47.42, 14.53. ES+HRMS: Calculated M+H = 255.1246. C<sub>14</sub>H<sub>15</sub>N<sub>4</sub>O. Obtained: 255.1252.

4.1.3.1.6. 9-Benzyl-6-ethoxy-8-methyl-9H-purine (**5g**). White solid; yield 41%; mp 56-58°C.  $\delta_{\text{H}}$ (499.79 MHz, CDCl<sub>3</sub>): 8.50 (1H, s, NCHN), 7.32-7.13 (5H, m, Ph), 5.40 (2H, s, CH<sub>2</sub>Ph), 4.65 (2H, q, J=10 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 2.51 (3H, s, CH<sub>3</sub>), 1.51 (3H, t, J=10 Hz,

OCH<sub>2</sub>CH<sub>3</sub>). $\delta_C$ (125.68 MHz, CDCl<sub>3</sub>): 159.83, 153.55, 151.51, 151.42, 135.49, 128.98, 128.10, 126.92, 120.33, 62.95, 46.08, 14.58, 14.39. ES+HRMS: Calculated M+H = 269.1402 C<sub>15</sub>H<sub>17</sub>N<sub>4</sub>O. Obtained: 269.1396.

4.1.3.1.7. *9-Benzyl-6-ethoxy-8-ethyl-9H-purine (5h)*. White viscous oil; yield 2%.  $\delta_H$ (499.79 MHz, CDCl<sub>3</sub>): 8.50 (1H, s, NCHN), 7.31-7.12 (5H, m, Ph), 5.43 (2H, s, CH<sub>2</sub>Ph), 4.68 (2H, q, J=10 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 2.80 (2H, q, J=10 Hz, CH<sub>2</sub>CH<sub>3</sub>), 1.53 (3H, t, J=10 Hz, OCH<sub>2</sub>CH<sub>3</sub>), 1.27 (3H, t, J=10 Hz, CH<sub>2</sub>CH<sub>3</sub>). ES+HRMS: Calculated M+H = 283.1559 C<sub>16</sub>H<sub>19</sub>N<sub>4</sub>O. Obtained: 283.1560.

4.1.3.1.8. *9-Benzyl-6-(benzyloxy)-9H-purine (5i)*. White solid; yield 30%.  $\delta_H$ (499.79 MHz, CDCl<sub>3</sub>): 8.59 (1H, s, NCHN), 7.90 (1H, s, NHCHN), 7.57-7.28 (10H, m, Ph), 5.70 (2H, s, OCH<sub>2</sub>Ph), 5.43 (2H, s, CH<sub>2</sub>Ph).  $\delta_C$ (125.68 MHz, CDCl<sub>3</sub>): 161.51, 153.09, 142.91, 137.06, 136.17, 129.98, 129.72, 129.40, 129.31, 129.23, 128.99, 128.65, 127.84, 122.32, 69.31, 48.35. ES+HRMS: Calculated M+H = 317.1402. C<sub>19</sub>H<sub>17</sub>N<sub>4</sub>O. Obtained: 317.1407.

4.1.3.1.9. *6-Isopropoxy-9-isopropyl-9H-purine (5j)*. Yellow viscous oil; yield 10%.  $\delta_H$ (300.20 MHz, CDCl<sub>3</sub>): 8.54 (1H, s, NCHN), 7.99 (1H, s, NHCHN), 5.70 (1H, m, OCH(CH<sub>3</sub>)<sub>2</sub>), 4.91 (1H, m, NCH(CH<sub>3</sub>)<sub>2</sub>), 1.66 (6H, d, J=6 Hz, OCH(CH<sub>3</sub>)<sub>2</sub>), 1.51 (6H, d, J=6 Hz, NCH(CH<sub>3</sub>)<sub>2</sub>).  $\delta_C$ (125.68 MHz, CDCl<sub>3</sub>): 160.84, 151.87, 150.40, 139.60, 121.83, 70.68, 47.55, 22.89, 22.05. ES+HRMS: Calculated M+H = 221.1402. C<sub>11</sub>H<sub>17</sub>N<sub>4</sub>O. Obtained: 221.1402.

4.1.3.1.11. *6-Isopropoxy-9-isopropyl-8-methyl-9H-purine (5k)*. Yellow viscous oil; yield 12%.  $\delta_H$ (300.20 MHz, CDCl<sub>3</sub>): 7.99 (1H, s, NCHN), 5.70 (1H, m, OCH(CH<sub>3</sub>)<sub>2</sub>), 4.91 (1H, m, NCH(CH<sub>3</sub>)<sub>2</sub>), 2.07 (3H, s, CH<sub>3</sub>), 1.71 (6H, d, J=6 Hz, OCH(CH<sub>3</sub>)<sub>2</sub>), 1.49

(6H, d, J=6 Hz, NCH(CH<sub>3</sub>)<sub>2</sub>). ES+HRMS: Calculated M+H = 235.1559.C<sub>12</sub>H<sub>19</sub>N<sub>4</sub>O.Obtained: 235.1553.

4.1.3.1.12. *6-tert-butoxy-9-isopropyl-9H-purine (5l)*. Yellow viscous oil; yield 15%.  $\delta_{\text{H}}$ (300.20 MHz, CDCl<sub>3</sub>): 8.51 (1H, s, NCHN), 7.95 (1H, s, NHCHN), 4.90 (1H, m, NCH(CH<sub>3</sub>)<sub>2</sub>), 1.78 (9H, s, C(CH<sub>3</sub>)<sub>3</sub>), 1.65 (6H, d, J=6 Hz, OCH(CH<sub>3</sub>)<sub>2</sub>).ES+HRMS: Calculated M+H = 235.1559.C<sub>12</sub>H<sub>19</sub>N<sub>4</sub>O.Obtained: 235.1563.

4.1.3.1.13. *9-Isopropyl-6-methoxy-9H-purine (5m)*. White viscous oil; yield 10%.  $\delta_{\text{H}}$ (400.57 MHz, CDCl<sub>3</sub>): 8.53 (1H, s, NCHN), 7.98 (1H, s, NHCHN), 4.89 (1H, m, CH(CH<sub>3</sub>)<sub>2</sub>), 4.18 (3H, s, OCH<sub>3</sub>), 1.63 (6H, d, J=4 Hz, NCH(CH<sub>3</sub>)<sub>2</sub>). $\delta_{\text{C}}$ (125.68 MHz, CDCl<sub>3</sub>): 161.19, 152.25, 151.86, 139.89, 122.07, 54.26, 47.63, 22.78. ES+HRMS: Calculated M+H :193.1089. C<sub>9</sub>H<sub>13</sub>N<sub>4</sub>O.Obtained: 193.1087.

4.1.3.1.14. *9-tert-butyl-6-ethoxy-9H-purine (5n)*. White solid; yield 57%.  $\delta_{\text{H}}$ (300.20 MHz, CDCl<sub>3</sub>): 8.49 (1H, s, NCHN), 7.97 (1H, s, NHCHN), 4.64 (2H, q, OCH<sub>2</sub>CH<sub>3</sub>), 1.80 (9H, s, NC(CH<sub>3</sub>)<sub>3</sub>), 1.50 (3H, t, OCH<sub>2</sub>CH<sub>3</sub>). $\delta_{\text{C}}$ (75.49 MHz, CDCl<sub>3</sub>): 161.22, 151.20, 139.87, 123.16, 63.10, 29.30, 14.80. ES+HRMS: Calculated M+H = 221.1402. C<sub>11</sub>H<sub>17</sub>N<sub>4</sub>O. Obtained: 221.1400.

#### 4.1.3.2. General procedure for the preparation of compounds 6 (a-d). Synthesis B

A suspension of NaH (50%) in mineral oil (3.7 mmol, 10 equiv. ) was dissolved in a mixture cooled at 0° constituted both for appropriate alcohol (3.7 mmol, 10 equiv.) and appropriate N,N-dimethylamide corresponding (18.5 mmol, 50 equiv.), when the amide is liquid, or the mixture is dissolved in dioxane, when the amide is solid. The solution was stirred at room temperature for 30 min and then at 90°C for the same time. At 90°C compound 4 (a-b) (0.37 mmol, 1 equiv.), dissolved in the amide (18.5 mmol, 50 equiv.)



or in dioxane were added dropwise and the mixture heated for 24 h. The reaction mixture quenched to pH 7 with an aq. saturated  $\text{NH}_4\text{Cl}$  solution, was then extracted with  $\text{CH}_2\text{Cl}_2$  (3x15 mL), the combined organic extracts dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent evaporated. The residue was chromatographed on silica gel using ethyl acetate/petroleum ether solutions.

*4.1.3.2.1. 6-Benzyloxy-9-isopropyl-8-phenyl-9H-purine (6a)*: White solid; yield 24%.  $\delta_{\text{H}}$ (300.20 MHz,  $\text{CDCl}_3$ ): 8.57 (1H, s, NCHN), 7.70- 7.38 (10H, m, Ph,  $\text{CH}_2\text{Ph}$ ), 5.72 (2H, s,  $\text{OCH}_2\text{Ph}$ ), 4.80 (1H, m,  $\text{CH}(\text{CH}_3)_2$ ), 1.76 (6H, d,  $J=6$  Hz,  $\text{NCH}(\text{CH}_3)_2$ ).  $\delta_{\text{C}}$ (125.68 MHz,  $\text{CDCl}_3$ ): 160.54, 153.14, 151.01, 136.66, 130.40, 129.80, 128.95, 128.76, 128.61, 128.27, 128.05, 122.14, 68.42, 50.04, 21.51. ES+HRMS: Calculated  $\text{M}+\text{H}=345.1715$ .  $\text{C}_{21}\text{H}_{21}\text{N}_4\text{O}$ . Obtained: 345.1716.

*4.1.3.2.2. 8-Ethyl-9-isopropyl-6-propoxy-9H-purine (6b)*: Yellow viscous oil; yield 20%.  $\delta_{\text{H}}$ (300.20 MHz,  $\text{CDCl}_3$ ): 8.43 (1H, s, NCHN), 4.69 (1H, m,  $\text{CH}(\text{CH}_3)_2$ ), 4.52 (2H, t,  $J=9$  Hz,  $\text{OCH}_2\text{CH}_2\text{CH}_3$ ), 2.94 (2H, q,  $J=6$  Hz,  $\text{CH}_2\text{CH}_3$ ), 1.92 (2H, m,  $\text{OCH}_2\text{CH}_2\text{CH}_3$ ), 1.70 (6H, d,  $J=10$  Hz,  $\text{NCH}(\text{CH}_3)_2$ ), 1.43 (3H, t,  $J=6$  Hz,  $\text{CH}_2\text{CH}_3$ ), 1.05 (3H, t,  $J=9$ ,  $\text{OCH}_2\text{CH}_2\text{CH}_3$ ).  $\delta_{\text{C}}$ (125.68 MHz,  $\text{CDCl}_3$ ): 160.28, 155.28, 150.65, 150.31, 124.41, 68.65, 48.57, 21.24, 23.13, 22.49, 21.51, 12.49, 10.66. ES+HRMS: Calculated  $\text{M}+\text{H}=249.1715$ .  $\text{C}_{13}\text{H}_{21}\text{N}_4\text{O}$ . Obtained: 249.1721.

*4.1.3.2.4. 6-Isobutoxy-8,9-diisopropyl-9H-purine (6c)*: White solid; yield 22%.  $\delta_{\text{H}}$ (300.20 MHz,  $\text{CDCl}_3$ ): 8.44 (1H, s, NCHN), 4.74 (1H, m,  $\text{NCH}(\text{CH}_3)_2$ ), 4.36 (2H, d,  $\text{OCH}_2\text{Pr}^i$ ,  $J=9$  Hz), 3.28 (1H, m,  $\text{CCH}(\text{CH}_3)_2$ ), 2.29 (1H, m,  $\text{CH}_2\text{CH}(\text{CH}_3)_2$ ), 1.72 (6H, d,  $J=9$  Hz,  $\text{NCH}(\text{CH}_3)_2$ ), 1.47 (6H, d,  $J=9$  Hz,  $\text{CCH}(\text{CH}_3)_2$ ), 1.06 (6H, d,  $J=9$  Hz,  $\text{CH}_2\text{CH}(\text{CH}_3)_2$ ). ES+HRMS: Calculated  $\text{M}+\text{H}=277.2028$ .  $\text{C}_{15}\text{H}_{25}\text{N}_4\text{O}$ . Obtained: 277.2036.

4.1.3.2.3. *9-tert-butyl-6-(benzyloxy)-8-phenyl-9H-purine (6d)*: White solid; yield 45%.  $\delta_{\text{H}}$ (300.20 MHz,  $\text{CDCl}_3$ ): 8.59 (1H, s, *NCHN*), 7.58- 7.34 (10H, m, *Ph*,  $\text{CH}_2\text{Ph}$ ), 5.69 (2H, s,  $\text{OCH}_2\text{Ph}$ ), 1.70 (9H, s,  $\text{NC}(\text{CH}_3)_3$ ).  $\delta_{\text{C}}$ (125.68 MHz,  $\text{CDCl}_3$ ): 160.62, 154.52, 153.37, 150.46, 136.58, 134.97, 130.11, 129.72, 128.85, 128.59, 128.27, 128.09, 121.76, 68.43, 61.06, 31.15. ES+HRMS: Calculated  $\text{M}+\text{H} = 359.1872$ .  $\text{C}_{22}\text{H}_{23}\text{N}_4\text{O}$ . Obtained: 359.1880.

## 4.2 Cells and culture

Jurkat (acute T cell leukemia) and K562 (chronic erythroleukaemia) cell lines were cultured at 37° C in a humidified atmosphere of 5%  $\text{CO}_2$  in RPMI 1640 medium (BioWhittaker, Verviers, Belgium) supplemented with 10% Fetal Bovine Serum (Gibco, Auckland, New Zealand) and 1% Glutamax (Gibco) and Penicillin-Streptomycin (BioWhittaker). The N1 cell line are long-term primary  $\text{CD4}^+$  T lymphocytes derived from a normal individual<sup>24</sup> and expanded by weekly stimulation of Mytomicin-C treated Class II<sup>+</sup> Raji B lymphoma cells in the presence of 50 U.I./ml of rIL-2 (kindly provided by the National Institutes of Health's AIDS Reagent program, Bethesda, MD) as described in detail elsewhere<sup>17</sup>.

### 4.2.1 Cell proliferation assay

To analyze **5e**, **5i** and **6a** activity, cells were cultured in 96-well microtiter plates (50,000 cells/well) and treated for 48 h with the library of compounds at increasing concentrations (25, 50, 100 and 200  $\mu\text{M}$ ). Cell proliferation was measured with the MTT (3-[4-5 dimethylthiazol-2-7P]-2,5-diphenyl tetrazolium bromide:thiazolyl blue)

(Sigma) colorimetric assay as described in detail elsewhere<sup>21</sup>, and results are expressed as percentage of inhibition calculated over the values obtained with untreated cells and corrected for the non-specific effect of the amounts of DMSO contained in each sample of the compounds. Data are represented as mean  $\pm$  standard deviation of the different experiments.

To analyze **6d** activity, each well of a 96-well tissue culture plate containing 50  $\mu$ L from serial doubling dilutions of **6d** compound was inoculated with 50  $\mu$ L of previously prepared cell culture at 20,000 cells/mL, with the exception of two rows, which received medium only. Eleven different final concentrations of each compound, ranging from 0.8–800  $\mu$ M, were tested in the assay. Each concentration point was assayed in three replicates, and experiments were repeated at least three times. The same volume of DMSO, **6d** solvent, used in each concentration point was assayed in parallel as control. The IC<sub>50</sub> value was calculated using GraphPad Prism5 software.

#### **4.3. Analysis of cell cycle and Western Blotting**

For the analysis of cell cycle, 250,000 cells/well were cultured in 24-well plates and incubated with the chosen compounds at the indicated doses for 48 h, as described elsewhere<sup>21</sup>. After treatment with the compounds, cells were collected, fixed with 70% cold ethanol for 5 min at 4° C and then incubated with DNA extraction buffer (0.2 M Na<sub>2</sub>HPO<sub>4</sub> and 0.004 M citric acid) at 37° C for 10 min. Subsequently, cells were resuspended in PBS containing 100  $\mu$ g/ml RNase and 40  $\mu$ g/ml propidium iodide (PI) and incubated at 37° C for 30 min in the dark. Quantitative analysis of cells with sub-G1 DNA content was carried out in a FACScan flow cytometer using the Cell Quest software (BD Biosciences). Data are expressed as mean  $\pm$  Standard Error Mean (SEM) of three different experiments. For the analysis of caspase activation, cells were treated

with compounds at 200  $\mu\text{M}$  for the indicated times and lysed in 150 mM NaCl, 50 mM Tris-Cl pH 8.0, 1% NP-40 and protease inhibitors for 30 min. Lysates were resolved by SDS-PAGE as described, blotted onto PVDF membranes, blocked for 1 h and hybridized overnight with pre-determined optimal concentrations of monoclonal antibodies against human caspase-9 (R&D Systems; Minneapolis, MN) and caspase-8 (Alexis Biochemicals; San Diego, CA). Membranes were washed, incubated for 1 h with an HRP-labelled goat anti-mouse antibody and revealed by chemiluminescence (ECL Advance Western Blotting Detection Kit, Amersham, GE Healthcare UK). Light emission was detected with a digital imaging system (Fujifilm Image Analyzer LAS-4000, Tokyo, Japan) and analyzed with the Multi Gauge software. Where indicated, cells were pre-treated with the Pan Caspase OPH Inhibitor Q-VD (Q-Val-Asp(non-methylated)-OPh) (R&D systems, Minneapolis, MN) for 1 h prior adding compound 6d and maintained throughout the culture.

#### ***4.4. Annexin-V staining.***

For the analysis of phosphatidylserine exposure, 250,000 cells/well were cultured in 24-well plates and incubated with the chosen compounds at the indicated doses for 48 h, as described above. Etoposide was used as positive control and untreated cells were used as control of live cells. After treatment, cells were collected and washed with annexin V binding buffer (10 mM HEPES pH 7.4, 140 mM NaCl, 2.5 mM  $\text{CaCl}_2$ ) and resuspended in 100  $\mu\text{l}$  of this buffer. Then, 2.5  $\mu\text{l}$  of Annexin V-FLUOS solution (Roche, Mannheim, Germany) were added to each sample, followed by incubation for 20 min in the dark at room temperature. After a final wash, the pellet was resuspended in annexin V binding buffer containing 10  $\mu\text{l}$  of PI at 10  $\mu\text{g}/\text{ml}$ . Stained samples were immediately analyzed by flow cytometry using a FACSCalibur flow cytometer. Data were analyzed with the FlowJo software.

#### 4.5 *In vitro* kinase screening assay.

A panel of 96 recombinant kinases (see Supporting Information) was used to run a  $^{33}\text{P}$ -radiolabeled kinase assay profiling against compounds **5e**, **6a** and **5i** by Reaction Biology Corp. (Malvern, PA) according to their established procedures. Compounds were tested in a single-dose duplicate mode at a concentration of 10  $\mu\text{M}$ . Reactions were carried out at 10  $\mu\text{M}$  ATP. Compounds were tested in a 10-dose  $\text{IC}_{50}$  mode with 2-fold serial dilution starting at 200  $\mu\text{M}$  using the same  $\text{P}^{33}$ -radiolabeled kinase assay as before to determine their  $\text{IC}_{50}$  values against DAPK1 and DAPK2.

#### Supporting Information

Experimental synthetic protocols and characterization data, including NMR data ( $^1\text{H}$  and  $^{13}\text{C}$ ) and mass spectrometry data (HRMS) of compounds are presented in SI file. Raw data of the  $^{33}\text{P}$  radiolabeled kinase assay containing the list of the 96 kinases used.  $\text{IC}_{50\_DAPK1}$  graphs.

#### Acknowledgements

We thank undergraduate students Alvaro Lorente Macias and Pilar Aguilar Martinez for their help with basic laboratory techniques. JJDM thanks the Spanish Ministerio de Economia y Competitividad for funding a Ramon y Cajal Fellowship. AUB thanks Medical Research Council UK - Institute of Genetics and Molecular Medicine for funding an Academic Fellowship. We acknowledge the generous supply of rIL-2 provided by the National Institute's of Health AIDS Reagent Program. IJM is grateful for the support provided to his laboratory by grant 12UDG01-ATF from Sparks, The Children's Medical Charity. London, UK.

#### Author Contributions

MJPI, AUB and JJDM designed the compounds and their syntheses. MJPI synthesized and characterized the compounds. STR, PFD and JDUB carried out or contributed to the biological experiments. MAG contributed with reagents. MALG, MPI, AUB and JJDM analyzed the chemical data. AUB, JJD and IJM interpreted the biological data and wrote the paper. IJM and JJDM directed the work.

**FIGURE LEGENDS**

**Scheme 1.** Synthetic routes to generate purine types “A” or “B”. Size of R1 controls which synthetic mechanism is followed under the same reaction conditions.

**Figure 1.** Set of 18 purines prepared through either Route A or B (see Scheme 1).

**Figure 2.** Structure of purine analogues **5e**, **6a** and **5i**, which were chosen for detailed functional screening.

**Figure 3.** Effect of compounds **5e**, **5i** and **6a** on cell cycle. A representative experiment showing Jurkat cells incubated for 48 h in the presence of indicated compounds and doses. Cell cycle subpopulations were analyzed by flow cytometry after permeabilisation and staining of cells with Propidium iodide. Additional controls containing equivalent amounts of DMSO used as drug vehicle were included, showing no toxicity in any case (not shown). Percentage of cells within the sub-G1 region are indicated in the plots.

**Figure 4.** Membrane phosphatidylserine expression after treatment of Jurkat cells with compounds **5e**, **5i** and **6a**. Jurkat cells were treated for 48 h with the indicated compounds and doses and the percentages of apoptotic cells was measured by flow cytometry after double staining with propidium iodide and Annexin-V-Fluos. Dot plots represent the results of a typical experiment. Untreated cells and cells exposed to Etoposide were included as negative and positive controls.

**Figure 5.** Effect of indicated compounds on primary lymphocytes and Jurkat cells. Primary cell line N1 and Jurkat cells were treated for 48h at the indicated doses of compounds **5e**, **5i** and **6a** and cell cycles were analysed by flow cytometry after staining

with Propidium Iodide. Graphs represent the mean $\pm$ S.D. of cells gated within the sub-G1 region obtained from three independent experiments. UT: Untreated.

**Figure 6.** Study of early signaling events of apoptosis induced by compound **6a** in Jurkat cells. Western blot analysis of total lysates of Jurkat cells after treatment with either compound **6a** (200  $\mu$ M, left-hand lanes) or equivalent amount of vehicle (DMSO, right-hand lanes) for 5, 6 or 7 h. Membranes were probed with antibodies against initiator caspases 8 or 9, using actin as loading control. Cells treated with Etoposide were included as positive control. Arrows indicate cleavage products (43/41 kD bands for caspase 8 and 37/< kD bands for caspase 9), which are detected after caspase activation whereas upper bands correspond to native proteins. Similar results were obtained after treating cells with compounds **5e** and **5i** (not shown). NT: Not treated

**Figure 7.** Structure of compound 6-benzyloxy-9-tert-butyl-8-phenyl-9H-purine **6d**.

**Figure 8.-** IC<sub>50</sub> value of compound **6d** in Jurkat cells as determined by inhibition of cell proliferation. A 11-point 1:2 serial dilution of compound **6d**, starting at 800  $\mu$ M, was used to determine its IC<sub>50</sub> value by the MTT assay after treatment of Jurkat cells for 48 h (solid circles). Equivalent amounts of DMSO used as vehicle in each dilution were set as negative controls (open circles). Results indicate percentages of cell proliferation over the results obtained with untreated cells. Data are expressed as mean $\pm$ SEM.

**Figure 9.** A) Effect of compound **6d** on the cell cycle of Jurkat cells. Jurkat cells were treated for 48 h with **6d** at indicated concentrations or their equivalent amounts of DMSO used as vehicle. Cells gated on the sub-G1 region after staining with propidium iodide are indicated. B) Membrane phosphatidylserine expression after treatment of Jurkat cells with compound **6d**. Jurkat cells were treated for 24 h with compound **6d** and the indicated doses and the percentages of apoptotic cells was measured by flow



cytometry after double staining with propidium iodide and Annexin-V-Fluos. Dot plots represent the results of a typical experiment. Untreated cells and cells exposed to Etoposide were included as negative and positive controls. C) Percentages of apoptotic cells after treatment of Jurkat cells with compound **6d**. Graph summarizes values of percentages of apoptotic cells and SEM obtained from three independent experiments. UT: Untreated. D) Activation of initiator caspases 8 and 9 after cell treatment with compound **6d**. Western blot analysis of total lysates from Jurkat cells after treatment with either compound **6d** (200  $\mu$ M, right-hand lanes) or vehicle (DMSO, left-hand lines) for 2, 4 and 6 h. Membranes were probed with antibodies against initiator caspases 8 or 9, using actin as loading control. Arrows indicate cleavage products (43/41 kD bands for caspase 8 products and 37/35 kD bands for caspase 9 ones) after caspase activation whereas upper bands correspond to native proteins (caspase 8 correspond to 55/57 kD band and caspase 9 to 45 kD band)


**Figure 10.-** Pre-treatment of Jurkat cells with the pan-caspase inhibitor Q-VD-Oph abrogates cell death mediated by compound **6d**. A) Jurkat cells were pre-treated for 1 h with 50  $\mu$ M of the Q-VD-Oph pan-caspase inhibitor and cultured thereafter for 24 h in the presence of 200  $\mu$ M of compound **6d** (bottom left panel). Controls include untreated cells (upper left panel); cells treated with Q-VD-Oph alone (upper right panel) or only with compound **6d** (bottom right panel). Cells were stained with propidium iodide and cell cycles analysed by flow cytometry. Percentages of cells within the sub-G1 region are indicated in the plots. B) Western Blot analysis of Caspase-8 activation. The same cultures of cells examined in Panel A were collected 8 h after treatment with compound **6d** and cytosolic lysates were resolved by SDS-PAGE and membranes probed with the anti-caspase 8 mAb. Actin was used as loading control. UT: Untreated.

## References

- 1 T. Anastassiadis, S. W. Deacon, K. Devarajan, H. Ma, and J. R. Peterson, *Nat. Biotechnol.*, 2011, **29**, 1039-1045.
- 2 P. G. Baraldi, A. Unciti-Broceta, M. J. Pineda de las Infantas, J. J. Díaz-Mochón, A. Espinosa, and R. Romagnoli, *Tetrahedron*, 2002, **58**, 7607-7611.
- 3 T. M. Caserta, A. N. Smith, A. D. Gultice, M. A. Reedy, and T. L. Brown, *Apoptosis.*, 2003, **8**, 345-352.
- 4 S. Chakilam, M. Gandesiri, T. T. Rau, A. Agaimy, M. Vijayalakshmi, J. Ivanovska, R. M. Wirtz, J. Schulze-Luehrmann, N. Benderska, N. Wittkopf, A. Chellappan, P. Ruemmele, M. Vieth, M. Rave-Frank, H. Christiansen, A. Hartmann, C. Neufert, R. Atreya, C. Becker, P. Steinberg, and R. Schneider-Stock, *Am. J. Pathol.*, 2013, **182**, 1005-1020.
- 5 O. Cohen, E. Feinstein, and A. Kimchi, *EMBO J.*, 1997, **16**, 998-1008.
- 6 O. Cohen, B. Inbal, J. L. Kissil, T. Raveh, H. Berissi, T. Spivak-Kroizaman, E. Feinstein, and A. Kimchi, *J. Cell Biol.*, 1999, **146**, 141-148.
- 7 L. P. Deiss, E. Feinstein, H. Berissi, O. Cohen, and A. Kimchi, *Genes Dev.*, 1995, **9**, 15-30.
- 8 M. Hojjat-Farsangi, *Int. J. Mol. Sci.*, 2014, **15**, 13768-13801.
- 9 H. Huang, J. Ma, J. Shi, L. Meng, H. Jiang, J. Ding, and H. Liu, *Bioorg. Med. Chem.*, 2010, **18**, 4615-4624.
- 10 L. P. Jordheim, D. Durantel, F. Zoulim, and C. Dumontet, *Nat. Rev. Drug Discov.*, 2013, **12**, 447-464.

- 11 J. L. Kissil, E. Feinstein, O. Cohen, P. A. Jones, Y. C. Tsai, M. A. Knowles, M. E. Eydmann, and A. Kimchi, *Oncogene*, 1997, **15**, 403-407.
- 12 G. Koopman, C. P. Reutelingsperger, G. A. Kuijten, R. M. Keehnen, S. T. Pals, and M. H. van Oers, *Blood*, 1994, **84**, 1415-1420.
- 13 Y. Lin, C. Stevens, and T. Hupp, *J. Biol. Chem.*, 2007, **282**, 16792-16802.
- 14 Y. Liu and N. S. Gray, *Nat. Chem. Biol.*, 2006, **2**, 358-364.
- 15 G. Manning, D. B. Whyte, R. Martinez, T. Hunter, and S. Sudarsanam, *Science*, 2002, **298**, 1912-1934.
- 16 D. R. McIlwain, T. Berger, and T. W. Mak, *Cold Spring Harb. Perspect. Biol.*, 2013, **5**, a008656.
- 17 I. J. Molina, D. M. Kenney, F. S. Rosen, and E. Remold-O'Donnell, *J. Exp. Med.*, 1992, **176**, 867-874.
- 18 I. Nicoletti, G. Migliorati, M. C. Pagliacci, F. Grignani, and C. Riccardi, *J. Immunol. Methods*, 1991, **139**, 271-279.
- 19 Z. O'Brien and M. Moghaddam, *Exper. Opin. Drug Metab. Toxicol.*, 2013, **9**, 1597-1612.
- 20 A. Raval, S. M. Tanner, J. C. Byrd, E. B. Angerman, J. D. Perko, S. S. Chen, B. Hackanson, M. R. Grever, D. M. Lucas, J. J. Matkovic, T. S. Lin, T. J. Kipps, F. Murray, D. Weisenburger, W. Sanger, J. Lynch, P. Watson, M. Jansen, Y. Yoshinaga, R. Rosenquist, P. J. de Jong, P. Coggill, S. Beck, H. Lynch, C. A. de la, and C. Plass, *Cell*, 2007, **129**, 879-890.


- 21 C. Ruiz-Ruiz, G. K. Srivastava, D. Carranza, J. A. Mata, I. Llamas, M. Santamaria, E. Quesada, and I. J. Molina, *Appl. Microbiol. Biotechnol.*, 2011, **89**, 345-355.
- 22 G. Shani, S. Henis-Korenblit, G. Jona, O. Gileadi, M. Eisenstein, T. Ziv, A. Admon, and A. Kimchi, *EMBO J.*, 2001, **20**, 1099-1113.
- 23 N. A. Thornberry and Y. Lazebnik, *Science*, 1998, **281**, 1312-1316.
- 24 M. G. Toscano, C. Frecha, K. Benabdellah, M. Cobo, M. Blundell, A. J. Thrasher, E. Garcia-Olivares, I. J. Molina, and F. Martin, *Hum. Gene Ther.*, 2008, **19**, 179-197.
- 25 L. Xi, J. Q. Zhang, Z. C. Liu, J. H. Zhang, J. F. Yan, Y. Jin, and J. Lin, *Org. Biomol. Chem.*, 2013, **11**, 4367-4378.
- 26 A. Zamecnikova, *Expert. Opin. Drug Discov.*, 2014, **9**, 77-92.
- 27 J. Zhang, M. M. Hu, H. B. Shu, and S. Li, *Cell Mol. Immunol.*, 2014, **11**, 245-252.
- 28 Huang, L. Chen, L. Guo, T.R. Hupp and Y. Lin, *Apoptosis*, 2014, **19**, 371-386.




1 Antiproliferative Screening

2 Kinase Inhibition Screening

3 Structure Refining



### Inductor of Apoptosis & DAPK1 Inhibitor



6,8,9-substitued purines are presented as interesting candidates to study the role of DAPK1 kinase in triggering apoptosis in cancer cells.  
84x45mm (300 x 300 DPI)