

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

Importance of topology for glycoclusters binding to *Pseudomonas aeruginosa* and *Burkholderia ambifaria* bacterial lectins

Caroline Ligeour,^a Lucie Dupin,^b Anthony Angeli,^a Gérard Vergoten,^d Sébastien Vidal,^c Albert Meyer,^a Eliane Souteyrand,^b Jean-Jacques Vasseur,^a Yann Chevolut^{b*} and François Morvan^{a*}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Pseudomonas aeruginosa (PA) and *Burkholderia ambifaria* (BA) are two opportunistic gram negative bacteria and major infectious agents involved in lung infection of cystic fibrosis patients. Both bacteria can develop resistance to conventional antibiotherapies. An alternative strategy consists in targeting virulence factors and in particular lectins with high affinity ligands such as multivalent glycoclusters. LecA (PA-IL) and LecB (PA-III) are two tetravalent lectins from PA that recognise galactose and fucose respectively. BamBL lectin from BA is trimeric with 2 binding sites per monomer and is also specific for fucose. These three lectins are potential therapeutic targets in an anti-adhesive anti-bacterial approach. Herein, we report the synthesis of 18 oligonucleotide pentofuranose-centered or mannitol-centered glycoclusters leading to tri-, penta- or decaivalent clusters with different topologies. The linker arm length between the core and the carbohydrate epitope was also varied leading to 9 galactocusters targeting LecA and to 9 fucocusters targeting both LecB and BamBL. Their dissociation constants (K_d) were determined using a DNA-based carbohydrate microarray technology. Trivalent xylo-centered galactocuster and ribo-centered fucocuster exhibited the best affinity for LecA and LecB respectively while the mannitol-centered deca-fucocuster displayed the best affinity to BamBL. This data demonstrated that topology and nature of linkers were the predominant factors for achieving high affinity rather than valency.

INTRODUCTION

Pseudomonas aeruginosa (PA) and *Burkholderia ambifaria* (BA) are opportunistic pathogens involved in nosocomial diseases and especially affect cystic fibrosis patients. PA has two soluble lectins LecA and LecB involved in its virulence.¹ Both lectins are homotetrameric with one binding site per monomer and one or two calcium ions involved in the binding site respectively. LecA is galactose specific with association constant (K_a) of $3.4 \times 10^4 \text{ M}^{-1}$ while LecB is fucose specific with K_a of $1.6 \times 10^6 \text{ M}^{-1}$. LecA has a cobblestone shape where the binding sites are located in the corner of the rectangle (32 and 70 Å away on the width and length, respectively).^{2, 3} Due to this configuration, several studies suggest that galactocusters can reach simultaneously two binding sites along the small side of

one tetramer leading to a glycoside cluster effect.^{3, 4} LecB has more or less a tetrahedral spatial distribution of the binding sites that are more difficult to reach simultaneously and very few glycoside cluster effects have been reported so far with LecB.^{5, 6} Furthermore, its affinity for monovalent fucose is already high with a K_d in the micromolar range.^{1, 2} BA expresses a fucose-binding lectin called BamBL. Similarly to LecB, it has a very high affinity for fucose (K_d of $0.96 \mu\text{M}$).⁷ It is crown shape trimeric lectin with two binding sites per monomer. The six binding sites are on the same side of the protein with a inter binding site distance ranking between 18 and 36 Å. So far very little fucocusters targeting BamBL have been synthesized and no glycoside cluster effect has been reported.^{8, 9} Nevertheless, its topology is favorable to the glycoside cluster effect with binding sites that can be reached by multivalent ligands with a topology in close relation with the AB₅ shiga toxin complexes.

The use of glycoclusters exhibiting high affinity for the targeted lectins has emerged as a promising strategy to fight against such pathogens.^{5, 10} Since there is no pressure on the bacteria this strategy should not induce a resistance phenomenon. However, the design of good candidates remains empirical since the precise mode of recognition of multivalent glycoclusters by their bacterial lectin partners is still not fully understood.

^aInstitut des Biomolécules Max Mousseron (IBMM), UMR 5247 CNRS, Université de Montpellier, ENSCM, place Eugène Bataillon, CC1704, 34095 Montpellier Cedex 5, France. E-mail: morvan@univ-montp2.fr

^bUniversité de Lyon, Institut des Nanotechnologies de Lyon UMR CNRS 5270, Ecole Centrale de Lyon, 36 avenue Guy de Collongue, 69134 Ecully, France E-Mail: yann.chevolut@ec-lyon.fr

^cInstitut de Chimie et Biochimie Moléculaires et Supramoléculaires (ICBMS), Laboratoire de Chimie Organique 2 – Glycochimie, UMR 5246, CNRS, Université Claude Bernard Lyon 1, 43 Boulevard du 11 Novembre 1918, 69622 Villeurbanne, France.

^dUnité de Glycobiologie Structurale et Fonctionnelle (UGSF), UMR 8576 Université de Lille 1, Cité Scientifique Avenue Mendeleiev, Bat C9, 59655 Villeneuve d'Ascq cedex, France.

Electronic Supplementary Information (ESI) available: NMR data and HPLC of glycoclusters. See DOI: 10.1039/x0xx00000x

RESULTS AND DISCUSSION

We have previously reported the syntheses of glycoclusters with a hexose-platform exhibiting either galactose¹¹⁻¹³ or fucose^{12, 14} epitopes for interactions with LecA and LecB respectively. To gain more insight on the effect of topology and valency on the affinity of the corresponding glycoclusters toward LecA and LecB, we have designed two new families of glycoclusters. The first one used furanose-based platforms such as arabinose, ribose and xylose (**3a-c**, Fig. 1) and the second one used an acyclic azido-mannitol (**10**, Fig. 1). The furanose platforms exhibit three hydroxyl functions which are differently orientated in the space according to the nature of the furanose to study the effect of the orientation of the 2'- and 3'-hydroxyl groups of the furanose moiety. The mannitol core **10** exhibits five hydroxyl functions on a flexible linear scaffold. The impact of rigidity and multivalence on the affinity for LecA and LecB can therefore be evaluated using these different scaffolds. The affinity of the fucocusters toward BambL was also evaluated since this lectin from *Bulkhoderia ambifaria* recognizes fucose moieties.

According to our strategy, the polyhydroxylated platforms **3a-c** and **10** were immobilized on solid support and each hydroxyl was phosphitylated using phosphoramidite chemistry with an alkyne linker leading to scaffolds bearing three or five alkynes. In addition, two alkyne functions were introduced on each hydroxyl of the reduced mannosyl platform leading to a decaalkyne scaffold. Finally, the conjugation between the polyalkyne scaffolds and the carbohydrate azides (galactoside **16** or fucoside **17**) occurred through a copper-catalyzed azide-alkyne cycloaddition (CuAAC) affording the expected glycoclusters exhibiting from three to ten galactose or fucose residues. Finally, a DNA sequence was incorporated for an immobilization on a DNA microarray of the final glycocluster leading to a glycoarray. K_d values of the glycoclusters toward LecA, LecB or BambL were determined using this glycoarray.





Figure 1: Structure of building blocks used for the synthesis of the glycoclusters.


Building blocks synthesis

The different building blocks required for their further assembly to gain the desired glycoclusters were synthesized: the three ribo, arabino and xylo-furanoside propargyl platforms **3a-c**, the 6-azido-6-deoxymannitol **10**, azide **11**¹⁵ and alkyne **12**¹⁶ solid supports, three different alkyne phosphoramidites exhibiting one **13**¹⁵ and **14**¹² or two **15**¹⁷ alkynes and β -D-galactoside **16**¹⁸ or α -L-fucoside **17**¹⁹ azide derivatives (Figure 1).

For the synthesis of the three different furanose scaffolds, we used ribo-, arabino- and xylo-furanosyl uracil since it is easy to introduce an alkyne function on the N^3 of uracil by a selective alkylation.²⁰ Commercially available ribo- **1** and arabino- **2** uracils were treated with propargyl bromide and potassium carbonate in DMF at 60 °C overnight to give compounds **3a** and **3b** respectively exhibiting N^3 -propargyl for subsequent immobilization on the azide solid support **11** (Scheme 1). Since xylo-furanosyl uracil is not commercial, N^3 -propargyl xylo-furanosyl uracil **3c** was synthesized in two steps by glycosylation of the 1,2,3,5-tetra-*O*-acetyl-D-xylofuranose with uracil, using bis(trimethylsilyl)acetamide and trimethylsilyl trifluoromethanesulfonate in acetonitrile. In accord with Baker's rule,²¹ the direct condensation of the suitably protected 2'-*O*-acetyl-D-xylofuranose **4** and uracil led to the formation of the β (trans-1',2') nucleoside **5**. The resulting peracetyl xylofuranosyl uracil **5** was alkylated as described above. The deprotection of the acetyl groups on the xylofuranoside was achieved by addition of methanol in the basic medium to give **3c**.



Scheme 1: Synthesis of ribo-, arabino- and xylo-furanosyl uracils **3a-c**. a) $\text{BrCH}_2\text{CCH}_3$, K_2CO_3 , DMF, 60 °C, overnight; b) *N,N*-bis(trimethylsilyl)acetamide, TMSOTf, CH_3CN , reflux, 4h30; c) MeOH, rt.




Scheme 2: Synthesis of 6-azido-6-deoxymannitol **10**. a) TsCl, pyridine, 22h, 0°C, 65%; b) NaN₃, DMF, 2h, 80°C, 84%; c) HCl, H₂O, dioxane, 2h, reflux, 46%; d) NaBH₄, EtOH, 1h, rt, 82%.

The linear platform **10** was obtained from methyl α -D-mannopyranoside **6** (Scheme 2). The primary 6-hydroxyl group was selectively tosylated in pyridine. Then, the tosyl group was replaced by an azide function by treatment with NaN₃ to afford **8**.²² The acetal **8** was converted to the corresponding hemiacetal **9** in acid conditions and reduction using sodium borohydride²³ afforded the 6-azido-6-deoxy-D-mannitol **10**.

Synthesis of glycocluster oligonucleotide conjugates.

Ribo-, arabino-, xylo-furanosyl uracil-centered glycol-oligonucleotides were synthesized using the building blocks reported in Figure 1. The first step was the immobilization of scaffolds **3a-c** on azide solid support **11** by CuAAC under microwaves assistance (Scheme 3). The resulting tri-hydroxyl scaffolds **18a-c** were then phosphorylated with alkyne phosphoramidites **13** or **14** using a DNA synthesizer with a triple coupling step. Then, the oligonucleotide was elongated

and labelled with a fluorescent dye (Cy3) using phosphoramidite chemistry. The resulting trialkyl furanoside oligonucleotide conjugates **19a-c** and **20a-c** were deprotected and released from the solid support by ammonia treatment leading to the trialkyl furanoside oligonucleotide conjugates **21a-c** and **22a-c** exhibiting different furanoside scaffolds (i.e. ribo-, arabino- or xylo) with alkyne linkers of different lengths (pent-4-ynyl phosphodiester **21** or propargyl diethyleneglycol phosphodiester **22**) in solution. Crude materials were analyzed and characterized by HPLC and MALDI-TOF spectrometry. Around 100 nmol of crude material was conjugated with either galactoside **16** or fucoside **17** azide by CuAAC using copper nanopowder²⁴ under microwaves assistance. Finally, a short ammonia treatment was applied to remove the acetyl groups. The twelve resulting glycoclusters exhibiting three galactosides (**G1a-c** and **G2a-c**) or three fucosides (**F1a-c** and **F2a-c**) with different scaffolds (ribo-, arabino- and xylo-furanose) and with short (pro, **G1**) or long (EG₂M, **G2**) linkers were purified by C₁₈ HPLC.




Scheme 3. Synthesis of the triglycocluster oligonucleotides **G1a-c**, **G2a-c**, **F1a-c** and **F2a-c**. a) **3a-c**, CuSO₄, Na Ascorbate, MeOH/H₂O (1:1), MW, 60°C, 1h; b) Solid Phase Oligonucleotide Synthesis (SPOS) with **13** or **14**, then SPOS with A, T, C, G, Cy3; c) NH₄OH; d) Cu⁰, **16** or **17**.

For the mannitol-centered oligonucleotides, the synthetic strategy was changed since difficulties in characterizing the final glycocluster oligonucleotide conjugates were encountered due to their high molecular weights. A short DNA sequence was synthesized on the alkyne solid support **12** and then the mannitol azide **10** was immobilized by CuAAC with

copper sulfate and sodium ascorbate under microwaves assistance (Scheme 4). The resulting pentahydroxyl compound **28** was phosphorylated with the three different phosphoramidites **11-13** with a triple coupling step to get respectively **29a-c**. Both galactoside and fucoside azide derivatives were conjugated on solid support by CuAAC using

CuSO₄ and sodium ascorbate to afford **30a-c** and **31a-c** respectively. A few beads were withdrawn and treated with ammonia for deprotection and release. After evaporation, the sample was analyzed by HPLC and MALDI-TOF showing the efficient introduction of carbohydrate moieties on the

polyalkyne scaffolds. Then, the rest of the oligonucleotide was elongated and labeled with Cy3. A final treatment with ammonia led to the four pentaglycoclusters **G3a-b**, **F3a-b** and the two decaglycoclusters **G3c**, **F3c**.



Scheme 5. Synthesis of the penta- and deca-glycocluster oligonucleotides **G3a-c** and **F3a-c**. a) SPOS with A, T, C; b) **10**, CuSO₄, Na Ascorbate, MeOH/H₂O (1:1), MW, 65°C, 1h; c) SPOS with 3; d) **16** or **17**, CuSO₄, Na Ascorbate, dioxane/H₂O (1:1), MW, 70°C, 1h; e) SPOS with A, T, C, G, Cy3, ii) NH₄OH.

Table 1: K_d values of the DNA-glycoclusters towards LecA, LecB or BamB1 determined by DNA Directed Immobilization glycoarray.²⁵

Glyco- Clusters	K_d (nM)			
	LecA	LecB	BamB1	
G1a	riboPro	66	/	/
G2a	riboEG ₂ M	54	/	/
G1b	araPro	64	/	/
G2b	araEG ₂ M	62	/	/
G1c	xyloPro	54	/	/
G2c	xyloEG ₂ M	49	/	/
G3a	mannipro	66	/	/
G3b	manniEG ₂ M	78	/	/
G3c	manni2	50	/	/
F1a	riboPro	/	56	20.1
F2a	riboEG ₂ M	/	73	21.2
F1b	araPro	/	81	25.1
F2b	araEG ₂ M	/	76	29.6
F1c	xyloPro	/	62	20.5
F2c	xyloEG ₂ M	/	93	26.5
F3a	manniPro	/	91	22.8

F3b	manniEG ₂ M	/	95	18.2
F3c	manni2	/	84	13.8

^aName of the core (ribose, arabinose, xylose or mannitol) with the nature of linker L (Pro or EG₂M) and manni2 for mannitol with two glycosides per hydroxyl.

The affinity (K_d) of the eighteen new glycoclusters displaying 3, 5 or 10 galactose or fucose moieties with different topologies and conjugated to different DNA sequences (Table S1) were determined by microarray (Table 1). The 9 galactocusters can only target LecA while the 9 fucocusters can target both LecB and BamB1. The microarray was elaborated by DNA Directed Immobilization (DDI).^{26, 27} Each glycocluster labelled with Cy3 and bearing a specific DNA sequence was hybridized specifically with an immobilized complementary DNA sequence on the array.^{28, 29} Fluorescent scanning of the Cy3 signal allows verifying glycocluster immobilization with a deviation lower than 14% illustrating similar surface densities for all immobilized glycoclusters. Microstructured glass slides, bearing 40 microwells, were used as an array of 64 spots/microwell allowing 40 independent experiments on one

single slide. In each microwell, two glycoclusters to be tested were immobilized and incubated with increasing concentrations (one concentration per well) of Alexa-647-labelled lectin (LecA, LecB or BambL). Scanning at 635 nm provided the binding isotherms for each glycocluster/lectin binding event. K_d values were obtained using the linear regression described in the experimental section.³⁰ The background noise of the Alexa-647 fluorescence signal was very low (below 20 a.u.) illustrating that Alexa-647-labelled lectin was not adsorbed nonspecifically on the surface.

The 9 galactoclusters tested exhibited the same order of magnitude of binding strength with K_d values between 49 and 78 nM for LecA. For each trimeric cluster, the hydrophilic and flexible EG₂M linker (**G2a-c**) seems slightly preferred to the short hydrophobic Pro linker (**G1a-c**) especially for the ribose core. The opposite was observed for mannitol core (**G3a** vs **G3b**). It was previously observed that the length of a linker could have a strong effect on the binding. Thus dimeric galactoclusters with a linker differing from only four methylenes exhibit a four-fold difference of binding.³¹

Regardless of the linker or the valence (3 or 5), the xylose core seems preferred. Then, with a slightly lower affinity, arabinose and ribose scaffolds showed comparable binding properties. The decavalent glycocluster **G3c** displayed similar K_d value (50 nM) than the best trimeric glycocluster **G2c** (49 nM). This data could be explained by some steric hindrances leading to a lower accessibility of the ten galactoside motifs. Furthermore, we have previously observed that mannose-centered galactoclusters exhibiting four phenyl galactoside motifs with Pro and EG₂M linkers have a K_d value of 57 and 39 nM³² respectively, showing also the benefit of a longer linker. Thus, the affinity of the xylose centered trigalactocluster **G2c** with the EG₂M linker exhibiting a K_d value of 49 nM is not so different, taking into account that only three motifs are involved in the construct.

This data showed that the "aromatic effect" has a strong consequence on the high binding of glycoclusters, as already observed by others and we.^{13, 32-38} Nevertheless, the topology brings an additional benefit allowing a high binding with only trivalent clusters.

In the case of LecB, glycoclusters with ribo and xylo cores exhibited the best affinities (56 and 62 nM). A stronger affinity was found with the shortest Pro linker (**F1a** and **F1c**) than with the EG₂M linker (**F2a** and **F2c**). Surprisingly, for the glycoclusters with an arabinose (**F1b-F2b**) or with the mannitol core (**F3a-F3b**), the effect of the linker on the binding was not significant. Interestingly, the glycoclusters with an arabinose were less prone to bind LecB. Finally the decavalent cluster (**F3c**) bound LecB better than the pentavalent ones but lower than the trivalent clusters. One can imagine that the rigidity of the furanose cores has an entropic benefit on the interaction of the corresponding clusters leading to a better binding.

For BambL, all fucoclusters displayed a strong binding with only few differences between them (K_d from 13.8 to 29.6 nM). The increase of the length of the linker from Pro to EG₂M was slightly detrimental for the furanose cores in contrast to the

mannitol one. Ribo and xylo-centered clusters were found better ligands to BambL than arabino-centered ones. Interestingly, the decafucocluster (**F3c**) displayed the strongest affinity ($K_d = 13.8$ nM) this effect can be explained by the fact that the CRD of BambL are localized on the same side leading to an easy and simultaneous (multivalent) access of several fucoses of the cluster in a "glycoside cluster effect" fashion.

In order to understand why glycoclusters with an arabinofuranose core have a higher dissociation constant (K_d) against LecA and LecB compared to ribofuranose and xylofuranose, preliminary molecular modeling studies have been performed. For each lectin, the glycoclusters exhibiting the best affinity have been chosen: **G2a**, **G2b**, **G2c** and **F1a**, **F1b**, **F1c**. Quantum semi empirical calculations using the PM6 Hamiltonian³⁹ and the solvation model SM5.4/A⁴⁰ showed that β -D-ribofuranose, β -D-arabinofuranose and β -D-xylofuranose exhibit a very similar heat of formation both in vacuum (ΔH_f) and in aqueous solution (ΔH_{aq}) (Table 2). From a structural point of view, the main differences among the three sugars stand in the chiralities of carbon 2 and carbon 3 of the furanose ring (Table 2) while that of carbons 1 and 4 is the same (R).

Table 2: ΔH_f and ΔH_{aq} and absolute configuration of carbon 2 and 3 obtained by quantum semi empirical calculations using PM6 Hamiltonian and SM5.4/A solvation model

furanose	ΔH_f kcal/mol	ΔH_{aq} kcal/mol	Absolute configuration	
			C2	C3
β -D-ribo	-231.8	-252.0	R	S
β -D-arabino	-230.9	-251.2	S	S
β -D-xylo	-229.7	-251.9	R	R

According to these findings, the differences in glycocluster interactions with LecA and LecB are supposed to be related to branches originating from C2 and C3. Accordingly, *in silico* molecular modelling was performed and the following empirical potential energies of interaction were obtained (Table 3). They clearly show that arabinofuranose derivatives (**G2b** and **F1b**) are the weakest ligands as found from their K_d values. When docking was applied with the epitopes on carbons 4 and 2 involved in the CRD, the empirical potential energies of interaction (ΔE) were not in agreement with the K_d values. This data strongly suggest that interaction with lectin occurred with epitopes on carbons C2 and C3.

Table 3: Empirical potential energy of interaction (ΔE) and experimental dissociation constants (K_d)

Complex	ΔE (kcal/mol)	K_d (nM)
LecA- G2a	-255.8	54
LecA- G2b	-198.9	62
LecA- G2c	-244.4	49
LecB- F1a	-232.7	56

LecB-F1b	-192.0	81
LecB-F1c	-222.0	62

In the case of LecA interaction, the difference of the energies of interaction could be tentatively explain by the position of the galactose with the linker on carbon 4 that brought an additional stabilization by interaction with the lectin. For LecA-G2a, the free ending galactose interacts with the R48 residue of LecA. In the case of LecA-G2c (figure 2a), the interaction takes place with P38 and T39 (figure 2b). Finally for LecA-G2b, no interaction occurs (figure 2c):




Figure 2. LecA in complex with a) G2a, b) G2c and c) G2b

The situation of LecB is slightly different. In the case of F1a and F1c, the free fucose, on the C4 branch, is located at equal distance from the two binding sites. F1a (figure 3a) exhibits strong intramolecular hydrogen bonds, a pseudo stacking interaction with the uracil ring and some interactions with residues A1, Q3 and L76. The free fucose in F1c (figure 3b) also takes a central position and hence displays strong interactions with N34, K62 and Q64. In the case of F1b, the free fucose is oriented towards the triazole moiety of branch on C2 leading to some interactions with P73 and S74 (figure 3c).




Figure 3. LecB in complex with a) F1a, b) F1c and c) F1b

CONCLUSION

In the present report, 18 glycoclusters were synthesized in order to target two lectins of PA (LecA and LecB) and one from BA (BambL). These three lectins are virulence factors involved in the adhesion and/or in biofilm formation. Their binding to the 18 glycoclusters was evaluated using DDI-based glycocluster array allowing for the determination of their K_d values which were below 100 nM. In silico molecular modeling study suggested that, for furanose-centered glycoclusters, epitopes born on carbons 2 and 3 are involved in the CRD of LecA or LecB while the free epitope on carbon 4 for ribose and xylose interacts with the lectin leading to an additional stabilization. Topology and nature of linkers were the predominant factors for achieving high affinity rather than valency. LecB and BambL showed different bonding profiles toward the set of 9 fucosylated clusters suggesting that the topology of the binding sites leads an additional (additional with respect to the saccharide residues) selectivity toward the clusters structure (topology and linker). These selectivity profiles can be viewed as a fingerprint of the lectin. These

results demonstrated that high avidity of a glycocluster to a lectin is a delicate fine-tuning of its topology.

Experimental

General methods. All reagents for synthesis were commercial and used without further purification. Solvents were commercial and used without further purification. All moisture sensitive reactions were performed under an argon atmosphere. NMR solvents were purchased from Euriso-Top. NMR spectra were recorded at 293 K using a 200, 300, 400 or a 600 MHz spectrometer (Bruker). Shifts (δ) are referenced relative to deuterated solvent residual peaks and expressed in parts per million (ppm). Coupling constants are expressed in Hertz (Hz). The following abbreviations are used to explain the observed multiplicities: s (singlet), d (doublet), t (triplet), q (quartet), dd (doublet of doublets), m (multiplet), sbr (broad singlet). High-resolution (HR-ESI-QToF) mass spectra were recorded using a Q-ToF Micromass spectrometer. MALDI-TOF mass spectra were recorded on a Voyager mass spectrometer (Perspective Biosystems, Framingham, MA) equipped with a nitrogen laser. Thin-layer chromatography (TLC) was carried out on aluminum sheets coated with silica gel 60 F₂₅₄ (Merck). TLC plates were inspected by UV light ($\lambda=254$ nm) and developed by treatment with a mixture of 10% H₂SO₄ in EtOH/H₂O (1:1 v/v), vanillin, KMnO₄ or phosphomolybdic acid followed by heating. Silica gel column chromatography was performed with silica gel Si 60 (40-63 mm). Reverse phase chromatography was performed with C₁₈ flash column.

Glycooligonucleotides synthesis. The oligonucleotides synthesis on solid support was performed on a DNA synthesizer Applied Biosystems (381A or 394A DNA synthesizer). Reactions under microwaves activation were performed on a Biotage Initiator system or Anton Paar Monowave 300 system. Solutions of Cap A, Cap B and iodide were purchased from link technologies as well as the commercial solid-supports. Solutions of TCA and CH₃CN for DNA synthesis were purchased from Biosolve. Cy₃-amidite was purchased from GE Healthcare. All glycooligonucleotides were purified and analysed by C₁₈ reversed-phase HPLC (Macherey-Nagel, Nucleodur 4.6×75 mm, 3 mM) on a Dionex Ultimate 3000 system with a Reodyn injector and a detector UV DAD 3000. Oligonucleotides were dosed by UV-Vis spectrophotometry at 260 nm on a VARIAN Cary 300 Bio UV-Vis spectrophotometer.

General procedure for N³-alkylation of uracyl 1 and 2. A solution of propargyl bromide (1.18 eq.) was added to a solution of furanosyl uracyl 1 or 2 (1.0 eq.) and potassium carbonate (1.08 eq.) in DMF, and the system was stirred at 60 °C overnight. The reaction mixture was filtered and evaporated to dryness. The crude product was filtered on silica gel column chromatography (5% MeOH in AcOEt) to afford the desired compound 3a-b.

1-(β -D-Ribofuranosyl)-3-(prop-2-yn-1-yl)uracil 3a. Obtained as a pale yellow solid (702 mg, 83%) following the general

procedure for alkylation: Uridine 1 (730 mg, 3.0 mmol), propargyl bromide (420 mg, 3.5 mmol), potassium carbonate (450 mg, 3.2 mmol), DMF (30 mL). ¹H NMR (300 MHz, CD₃OD) δ 8.08 (d, $J=8.1$ Hz, 1H, H₆), 5.95 (d, $J=4.2$ Hz, 1H, H₁), 5.82 (d, $J=8.1$ Hz, 1H, H₅), 4.68 (dd, $J=2.4, 1.6$ Hz, 2H, CH₂C \equiv CH), 4.22-4.16 (m, 2H, H₂, H₃), 4.06-4.03 (m, 1H, H₄), 3.76-3.92 (dd AB system, $J=9.0, 2.1$ Hz, 2H, H₅), 2.56 (t, $J=2.4$ Hz, 1H, C \equiv CH). ¹³C NMR (75 MHz, CD₃OD) δ 163.9, 151.9, 141.3, 102.0, 91.7, 86.4, 79.1, 75.9, 71.9, 71.1, 62.2, 31.1. HR-ESI-QToF MS (positive mode): m/z calcd. for C₁₂H₁₅N₂O₆ [M+H]⁺ 283.0930 found 283.0926.

1-(β -D-Arabinofuranosyl)-3-(prop-2-yn-1-yl)uracil 3b. Obtained as a pale yellow solid (761 mg, 90%) following the general procedure for alkylation: arauridine 2 (730 mg, 3.0 mmol), propargyl bromide (420 mg, 3.5 mmol), potassium carbonate (450 mg, 3.2 mmol), DMF (30 mL). ¹H NMR (300 MHz, CD₃OD) δ 7.88 (d, $J=8.1$ Hz, 1H, H₆), 6.16 (d, $J=4.2$ Hz, 1H, H₁), 5.76 (d, $J=8.1$ Hz, 1H, H₅), 4.66 (dd, $J=2.4, 1.2$ Hz, 2H, CH₂C \equiv CH), 4.19 (dd, $J=4.2, 2.8$ Hz, 1H, H₂), 4.08 (t, $J=3.3$ Hz, 1H, H₃), 3.97-3.93 (m, 1H, H₄), 3.86-3.76 (dd J_{AB}, $J=11.7, 3.0$ Hz, 2H, H₅), 2.56 (t, $J=2.4$ Hz, 1H, C \equiv CH). ¹³C NMR (75 MHz, CD₃OD) δ 162.9, 152.3, 141.5, 95.1, 87.3, 85.4, 77.9, 76.4, 75.8, 70.7, 61.3, 29.8. HR-ESI-QToF MS (positive mode): m/z calcd. for C₁₂H₁₅N₂O₆ [M+H]⁺ 283.0930 found 283.0930.

1-(2',3',5'-Tri-O-acetyl- β -D-xylofuranosyl)uracil 5. Uracil (262 mg, 2.3 mmol, 1.2 eq.) and *N,O*-bis(trimethylsilyl)acetamide (1.1 mL, 4.5 mmol, 2.2 eq.) were added to a solution of 1,2,3,5-tetra-*O*-acetyl- β -D-xylofuranose 4 (637 mg, 2.0 mmol, 1.0 eq.) in CH₃CN (25 mL) and refluxed until the suspension became a clear solution. Then, reaction mixture was cooled to 0 °C and TMSOTf (511 mg, 2.3 mmol, 1.2 eq.) was added dropwise, followed by reflux during 4.5 h. CH₃CN was evaporated under reduced pressure. To the residue was added saturated NH₄Cl solution and the mixture was extracted with CH₂Cl₂. The organic layer was dried with Na₂SO₄, evaporated and nucleoside 5 was purified by silica gel column chromatography (AcOEt:cyclohexane 8:2) to afford the title compound 5 (488 mg, 65%). ¹H NMR (300 MHz, CDCl₃) δ 9.82 (s, 1H, NH), 7.46 (d, $J=8.1$ Hz, 1H, H₆), 5.96 (d, $J=2.4$ Hz, 1H, H₁), 5.74 (d, $J=8.1$ Hz, 1H, H₅), 5.30 (dd, $J=3.9, 1.5$ Hz, 1H, H₃), 5.13 (t, $J=2.1$ Hz, 1H, H₂), 4.46 (q, $J=9.6, 5.4$ Hz, 1H, H₄), 4.27 (d, $J=5.7$ Hz, 2H, H₅). ¹³C NMR (100 MHz, CDCl₃) δ 169.4, 168.1, 168.0, 162.3, 149.3, 138.1, 101.8, 87.4, 78.6, 77.7, 73.3, 59.9, 20.0, 19.7, 19.6. HR-ESI-QToF MS (positive mode): m/z calcd. for C₁₅H₁₉N₂O₉ [M+H]⁺ 371.1091 found 371.1092.

1-(β -D-Xylofuranosyl)-3-(prop-2-yn-1-yl)uracil 3c. A solution of propargyl bromide (76 mg, 0.64 mmol, 1.18 eq.) was added to a solution of 2',3',5'-tri-*O*-acetyl xylofuranosyluridine 5 (200 mg, 0.54 mmol, 1.00 eq.) and potassium carbonate (373 mg, 2.7 mmol, 5.00 eq.) in DMF (5 mL). The system was stirred at 60 °C overnight and then methanol was added (300 mL). The reaction mixture was filtered and evaporated to dryness. The crude product was purified on silica gel column chromatography (AcOEt:cyclohexane 8:2) to afford the title

compound **3c** (98 mg, 65%). ^1H NMR (300 MHz, CD_3OD) δ 7.86 (d, $J=8.4$ Hz, 1H, H_6), 5.69 (d, $J=3.3$ Hz, 1H, H_1), 5.67 (d, $J=8.1$ Hz, 1H, H_5), 4.55 (d, $J=2.4$ Hz, 2H, $\text{CH}_2\text{C}\equiv\text{CH}$), 4.22 (m, 1H, H_4), 4.04-3.99 (m, 2H, H_2 , H_3), 3.84 (d, $J=5.4$ Hz, 2H, H_5), 2.45 (t, $J=2.4$ Hz, 1H, $\text{C}\equiv\text{CH}$). ^{13}C NMR (75 MHz, CD_3OD): δ 30.87, 61.13, 71.87, 76.42, 79.08, 82.46, 85.49, 94.16, 100.71, 141.72, 151.73, 164.03. HR-ESI-QToF MS (positive mode): m/z calcd. for $\text{C}_{12}\text{H}_{15}\text{N}_2\text{O}_6$ $[\text{M}+\text{H}]^+$ 283.0930 found 283.0930.

6-Azido-6-deoxy-D-mannose 9. To a solution of methyl 6-azido-6-deoxy- α -D-mannopyranoside **8**²² (379 mg, 1.7 mmol, 1 eq.) in H_2O /dioxane (15 mL, 1:2) was added a solution of HCl 12N (4 mL). The mixture was stirred under reflux for 4 h and then a saturated solution of sodium carbonate was added to increase the pH to 7. The crude product was concentrated *in vacuo*, diluted with MeOH and filtered. The filtrate was concentrated and purified by silica gel column chromatography (EtOAc) to afford the desired compound **9** as a a:b mixture (252 mg, 72%) as a pale yellow oil. Analyses were in agreement with the literature.⁴¹

6-Azido-6-deoxy-D-mannitol 10. To a solution of 6-azido-6-deoxy-D-mannose **9** (60 mg, 0.29 mmol, 1 eq.) in EtOH (8 mL) was added NaBH_4 (66 mg, 1.75 mmol, 6 eq.). The mixture was stirred for 2 h at room temperature and ion exchange resin (DOWEX-50W X8, H^+ form) was then added to decrease the pH to 3. The resin was then removed by filtration. The filtrate was concentrated and co-evaporated with MeOH to give the desired compound **10** after silica gel chromatography (49 mg, 88%). ^1H NMR (300 MHz, CD_3OD) δ 3.83-3.61 (m, 5H), 3.55 (dd, $J=12.6$, 2.7 Hz, 1H), 3.39 (dd, $J=6.6$, 12.6 Hz, 1H). ^{13}C NMR (100 MHz, D_2O) δ 73.0, 71.8, 71.5, 71.0 ($4\times\text{CHOH}$), 65.1 (CH_2OH), 55.8 (CH_2N_3). HR-ESI-QToF MS (negative mode): m/z calcd for $\text{C}_6\text{H}_{12}\text{N}_3\text{O}_5$ $[\text{M}-\text{H}]^-$: 206.0777 found 206.0777.

General procedure for elongation of DNA sequences and labelling with Cy3. The DNA sequences were synthesized on the solid-supported scaffolds (LCAA-CPG, 500Å) at the 1 mmol scale on a DNA synthesizer (ABI 394) by standard phosphoramidite chemistry. For the coupling step, benzylmercaptotetrazole was used as activator (0.3 M in anhydrous CH_3CN), and commercially available nucleosides phosphoramidites (0.1 M in anhydrous CH_3CN) were introduced with a 20 s coupling time and Cy3 amidite (0.06 M in anhydrous CH_3CN) with a 180 s coupling time. The capping step was performed with acetic anhydride using commercial solution (Cap A, Ac₂O/pyridine/THF, 10:10:80; Cap B, 10 % *N*-methylimidazole in THF) for 15 s. Each oxidation was performed with commercial solution of iodide (0.1 M in THF/pyridine/ H_2O , 78:20:2) for 15 s. Detritylation was performed with 3% TCA in CH_2Cl_2 for 35 s.

Immobilization on solid supports 11 or 12 of compounds 3a-c or 10 by CuAAC. An aqueous solution of modified carbohydrates **3a-c** or **10** (0.1 M, 50 mL, 5 eq.), freshly prepared aqueous solutions of CuSO_4 (0.1 M, 4 mL, 0.4 eq.) and sodium ascorbate (0.5 M, 4 mL, 2 eq.), TEAAC buffer (50

mL), water (17 mL) and MeOH (125 mL) were added to 1 mmol of solid support **11** or **12**. The resulting mixture was heated at 65 °C for 1 h. The solution was removed, and CPG beads were washed with H_2O (3 \times 2 mL), MeOH (3 \times 2 mL) and CH_3CN (3 \times 2 mL), and dried.

General procedure for introduction of phosphoramidites 13-15 on furanoside and reduced mannoside scaffolds. Solid-supported furanoside **18a-c** and mannitol **28** scaffolds (1 mmol) were treated by phosphoramidite chemistry, on a DNA synthesizer, with phosphoramidites **13**, **14** or **15**. Only coupling and oxidation steps were performed. For the coupling step, benzylmercaptotetrazole was used as activator (0.3 M in anhydrous CH_3CN) and phosphoramidites **13-15** (0.3 M with platform **18a-c** and 0.5 M with platform **28** in anhydrous CH_3CN) were introduced three times with a 180 s coupling time. Oxidation was performed with 0.1 M commercial solution of iodide for 15 s.

General procedure for deprotection of solid-supported alkyne oligonucleotides. The CPG beads bearing modified oligonucleotides were transferred to a 4 mL screw top vial and treated with 2 mL of concentrated aqueous ammonia for 15 h at room temperature and warmed to 55 °C for 2 h. The supernatants were withdrawn and evaporated.

General procedure for introduction of galactoside 16 or fucoside 17 derivatives by CuAAC reaction.

In solution. To a solution of 5'-fluorescent-3'-alkyne oligonucleotide **21a-c** and **22a-c** (100 nmol) were added galactoside **16** or fucoside **17** derivatives (3 eq per alkyne function, 0.1 M in MeOH or dioxane for **16**), 1 mg of $\text{Cu}^{(0)}$ nanopowder, triethylammonium acetate buffer 0.1 M, pH 7.7 (25 mL), water and MeOH or dioxane to obtain a final volume of 250 mL (water/MeOH, 1:1 v/v). The resulting mixture was heated at 65 °C for 1 h. EDTA (0.1 M, 100 mL) was added to the mixtures, and after centrifugation, the supernatants were withdrawn to eliminate $\text{Cu}(0)$ and desalted by size exclusion chromatography on NAP10. After evaporation, the 5'-fluorescent 3'-acetyl-glycomimetic oligonucleotides were treated with concentrated aqueous ammonia for 2 h at room temperature to remove acetyl groups and evaporated. Final compounds were purified by C_{18} reverse phase HPLC.

On solid support. A solution of galactoside **16** or fucoside **17** derivatives (5 eq per alkyne function, 0.1 M in MeOH or dioxane for **16**), freshly prepared aqueous solutions of CuSO_4 (0.1 M, 0.4 eq) and sodium ascorbate (0.5 M, 2 eq), TEAAC buffer (50 mL) were added to the solid supported oligonucleotide **29a-c** (1 mmol). The resulting mixture was heated at 70 °C for 1 h. The solution was removed, and CPG beads were washed with H_2O (3 \times 2 mL), MeOH (3 \times 2 mL) and CH_3CN (3 \times 2 mL), and dried. Finally, the DNA sequence was elongated and labelled with Cy3.

Microarray production and K_d microarray assays:

The procedure has been described in Goudot *et al.*²⁵: 3'-Amino-modified oligonucleotides were purchased from

Eurogentec and chemicals from Sigma-aldrich. Alexa Fluor® 647 Microscale Protein Labelling Kit was purchased from Invitrogen. Fluorescent scanning was performed using the Axon microarray scanner, GenePix 4100A software package.

Microarray production

Nexterion Glass D slides from Schott AG were used. The slides were structured by adapting the protocol of Mazurczyk et al.⁴² leading to slides bearing 40 microwells (square 3×3 mm, depth: 102 ± 1 μm, with 4.5 mm spacing between each wells).

The resulting glass slides were silanized according to the gas phase protocol⁴³ leading to tert-butyl ester functionalized surface. Deprotection of the acid function was accomplished in glacial formic acid for 7 h. The slides were washed in dichloromethane and water (10 min, under ultrasound). Activation of the carboxylic function was performed using diisopropylcarbodiimide and *N*-hydroxysuccinimide solution at 0.1 M in tetrahydrofuran, overnight at room temperature. The slides were washed in THF followed by dichloromethane (10 min, ultrasounds).

Next, four amino-modified oligonucleotides (25 μM, PBS 10x pH 8.5) were covalently immobilized in an array fashion using a Scienion sciFLEXARRAYER s3 system piezoelectric. Amino-modified oligonucleotides were spotted at 25 μM for LecB and at 0.5 μM for LecA and BambL in PBS 10X (pH 8.5). The immobilized sequences were 5'-GTG AGC CCA GAG GCA GGG-(CH₂)₇-NH₂, 5'-GTG GAG GCA CCA AGC TTT-(CH₂)₇-NH₂, 5'-CCA AGC GAG GTG GCA TTT-(CH₂)₇-NH₂ and 5'-CCA GCA GTG GAG AGC TTT-(CH₂)₇-NH₂. In each microwell, 64 spots of two sequences (2×32 spots) were spotted in a water saturated atmosphere. Water was allowed to evaporate gently overnight. Then, slides were washed for 30 min at 70 °C in SDS 0.1% and rinsed with water.

Blocking non-specific adsorption was performed using Bovine Serum Albumin (BSA). This step was performed before DDI of the glycoclusters and repeated before incubation with the lectins. The slides were incubated with a BSA solution 4%; in PBS 1X, pH 7.4 for 2 h. The slides were washed in 0.05% PBS1X-Tween20 (3×3 min), in PBS 1X (3×3 min) and rinsed with water.

Immobilization of the glycoclusters was accomplished using DNA directed Immobilization (DDI).²⁶ Two μL of a mixture containing two different glycoclusters (1 μM per glycocluster in PBS 1X pH 7.4) was poured at the bottom of each microwell and incubated (3 h, 37 °C, in a water saturated chamber). The slides were washed in SSC 2X, SDS 0.1% at 51 °C for 1 min, SSC 2X for 5 min and rinsed with water. Correct immobilization of the Cy3 labeled glycoclusters was insured by scanning the slides at 532 nm (excitation of Cy3). The fluorescent signal of each glycocluster was determined as the average of the mean fluorescence signal of 32 spots. The surface densities of the glycoclusters deviated by less than 14%.

Lectins LecA, LecB and BambL (a gift from Dr. Anne Imberty, CERMAV) were labeled with Alexa 647 according to the supplier protocol. The degrees of labeling per lectin (multimeric form) were 0.30, 0.34 and 0.24 moles of Alexa-647 per mole of lectin respectively.

K_d determination

Solutions of BSA_{2%}, 7.5 μM CaCl₂ and labeled lectin ranging from 0.01 nM to 2.0 μM for LecA and BambL (or 1 nM to 2.4 μM for LecB) were prepared in PBS 1X (pH 7.4) in order to draw isotherm curves of 20 discrete points. In each microwell, 2 μL of solution was added and incubated for 3 h at 37 °C in a water saturated chamber. Then, slides were washed for 5 min with 0.02% PBS1X-Tween20 and rinsed with water. An average of 32 spots per glycocluster of the mean fluorescence signal at 635 nm (FI) was calculated. For each glycocluster, *K_d* values were determined by the intercept with the y-axis using the linear regression: [LecA]/FI = 1/FI_{max} × [LecA] + *K_d*/FI_{max}.

In silico molecular docking

The three dimensional structures of LecA and LecB were retrieved from the Protein Data Bank (www.rcsb.org) under the PDB codes 2VXJ and 1UVZ respectively. As an example, the building procedure of the lectin (LecA)-ligand (**G2b**) complex is depicted as follows: the galactose endings of the two glycocluster branches linked on C2 and C3 of the arabinofuranose ring are brought closed to the crystallographic galactose ligands (figure S1). The terminal sugars of the two glycocluster branches are removed and the chemical bond with galactose moieties of LecA are built (figure S2). LecA and its two galactose groups are considered as aggregates and the complex is optimized (figure S2). A classical Monte Carlo conformational searching procedure is then performed as described in the BOSS software.⁴⁴ The constraints have been removed and the whole structure relaxed. In the minimization procedures, the spectroscopic empirical energy function SPASIBA and the corresponding parameters are used.⁴⁵⁻⁴⁷ A typical lowest energy structure for the complex LecA-**G2b** is shown in figure S3.

In the same way, an empirical potential energy of interaction Δ*E* for the lectin-ligand complexes using the simple expression:

$$\Delta E_{\text{interaction}} = E_{\text{complex}} - E_{\text{protein}} - E_{\text{ligand}}$$

was evaluated using the same force field.

Acknowledgements

This work was financially supported by ANR-12-BSV5-0020 "GLYCOMIME". Plateforme NanoLyon is acknowledged for its technical support. C. L and A. A. thank University of Montpellier for a research student ship. F.M. is member of Inserm.

References

1. N. Gilboa-Garber, *Methods Enzymol.*, 1982, **83**, 378-385.
2. A. Imberty, M. Wimmerova, E. P. Mitchell and N. Gilboa-Garber, *Microbes and Infection*, 2004, **6**, 221-228.
3. D. Sicard, S. Cecioni, M. Iazykov, Y. Chevolut, S. E. Matthews, J. P. Praly, E. Souteyrand, A. Imberty, S. Vidal and M. Phaner-Goutorbe,

- Chemical Communications*, 2011, **47**, 9483-9485.
4. S. Cecioni, R. Lalor, B. Blanchard, J. P. Praly, A. Imberty, S. E. Matthews and S. Vidal, *Chemistry-a European Journal*, 2009, **15**, 13232-13240.
 5. S. Cecioni, A. Imberty and S. Vidal, *Chemical Reviews*, 2015, **115**, 525-561.
 6. K. Buffet, E. Gillon, M. Holler, J. F. Nierengarten, A. Imberty and S. P. Vincent, *Organic & Biomolecular Chemistry*, 2015, **13**, 6482-6492.
 7. A. Audfray, J. Claudinon, S. Abounit, N. Ruvoen-Clouet, G. Larson, D. F. Smith, M. Wimmerova, J. Le Pendu, W. Romer, A. Varrot and A. Imberty, *J. Biol. Chem.*, 2012, **287**, 4335-4347.
 8. C. Ligeour, A. Audfray, E. Gillon, A. Meyer, N. Galanos, S. Vidal, J. J. Vasseur, A. Imberty and F. Morvan, *Rsc Adv*, 2013, **3**, 19515-19524.
 9. B. Richichi, A. Imberty, E. Gillon, R. Bosco, I. Sutkeviciute, F. Fieschi and C. Nativi, *Organic & Biomolecular Chemistry*, 2013, **11**, 4086-4094.
 10. R. J. Pieters, *Organic & Biomolecular Chemistry*, 2009, **7**, 2013-2025.
 11. G. Pourceau, A. Meyer, Y. Chevolut, E. Souteyrand, J. J. Vasseur and F. Morvan, *Bioconjugate Chem.*, 2010, **21**, 1520-1529.
 12. B. Gerland, A. Goudot, G. Pourceau, A. Meyer, V. Dugas, S. Cecioni, S. Vidal, E. Souteyrand, J. J. Vasseur, Y. Chevolut and F. Morvan, *Bioconjugate Chem.*, 2012, **23**, 1534-1547.
 13. B. Gerland, A. Goudot, C. Ligeour, G. Pourceau, A. Meyer, S. Vidal, T. Gehin, O. Vidal, E. Souteyrand, J. J. Vasseur, Y. Chevolut and F. Morvan, *Bioconjugate Chem.*, 2014, **25**, 379-392.
 14. B. Gerland, A. Goudot, G. Pourceau, A. Meyer, S. Vidal, E. Souteyrand, J.-J. Vasseur, Y. Chevolut and F. Morvan, *J. Org. Chem.*, 2012, **77**, 7620-7626.
 15. G. Pourceau, A. Meyer, J. J. Vasseur and F. Morvan, *J. Org. Chem.*, 2009, **74**, 6837-6842.
 16. J. Lietard, A. Meyer, J. J. Vasseur and F. Morvan, *J. Org. Chem.*, 2008, **73**, 191-200.
 17. C. Ligeour, A. Meyer, J. J. Vasseur and F. Morvan, *European Journal of Organic Chemistry*, 2012, 1851-1856.
 18. S. Cecioni, J. P. Praly, S. E. Matthews, M. Wimmerova, A. Imberty and S. Vidal, *Chemistry-a European Journal*, 2012, **18**, 6250-6263.
 19. F. Morvan, A. Meyer, A. Jochum, C. Sabin, Y. Chevolut, A. Imberty, J. P. Praly, J. J. Vasseur, E. Souteyrand and S. Vidal, *Bioconjugate Chem.*, 2007, **18**, 1637-1643.
 20. M. Nakane, S. Ichikawa and A. Matsuda, *J. Org. Chem.*, 2008, **73**, 1842-1851.
 21. B. R. Baker, in *The Ciba Foundation Symposium on the Chemistry and Biology of the Purine*, eds. G. E. W. Wolstenholme and C. M. O'Connor, John Wiley & Sons, Chichester, UK, 1957, pp. 120-133.
 22. J. J. Reina, O. S. Maldonado, G. Tabarani, F. Fieschi and J. Rojo, *Bioconjugate Chem.*, 2007, **18**, 963-969.
 23. L. Chaveriat, I. Stasik, G. Demailly and D. Beaupere, *Tetrahedron-Asymmetr.*, 2006, **17**, 1349-1354.
 24. note, *Since the catalytic form of copper is the Cu(I), the Cu(0) was used as a nanopowder to ensure a large specific surface area allowing its oxidation by oxygen contained in solvents.*
 25. A. Goudot, G. Pourceau, A. Meyer, T. Gehin, S. Vidal, J. J. Vasseur, F. Morvan, E. Souteyrand and Y. Chevolut, *Biosens. Bioelectron.*, 2013, **40**, 153-160.
 26. Y. Chevolut, C. Bouillon, S. Vidal, F. Morvan, A. Meyer, J. P. Cloarec, A. Jochum, J. P. Praly, J. J. Vasseur and E. Souteyrand, *Angew. Chem., Int. Ed.*, 2007, **46**, 2398-2402.
 27. R. Wacker and C. M. Niemeyer, *ChemBiochem*, 2004, **5**, 453-459.
 28. J. Zhang, G. Pourceau, A. Meyer, S. Vidal, J.-P. Praly, E. Souteyrand, J.-J. Vasseur, F. Morvan and Y. Chevolut, *Biosens. Bioelectron.*, 2009, **24**, 2515-2521.
 29. J. Zhang, G. Pourceau, A. Meyer, S. Vidal, J. P. Praly, E. Souteyrand, J. J. Vasseur, F. Morvan and Y. Chevolut, *Chemical Communications*, 2009, 6795-6797.
 30. S. Park and I. Shin, *Org. Lett.*, 2007, **9**, 1675-1678.
 31. F. Pertici, N. J. de Mol, J. Kemmink and R. J. Pieters, *Chemistry-a European Journal*, 2013, **19**, 16923-16927.
 32. F. Casoni, L. Dupin, G. Vergoten, A. Meyer, C. Ligeour, T. Gehin, O. Vidal, E. Souteyrand, J. J. Vasseur, Y. Chevolut and F. Morvan, *Organic & Biomolecular Chemistry*, 2014, **12**, 9166-9179.
 33. N. Garber, U. Guempel, A. Belz, N. Gilboa-Garber and R. J. Doyle, *Biochim. Biophys. Acta*, 1992, **1116**, 331-333.

34. R. U. Kadam, M. Bergmann, M. Hurley, D. Garg, M. Cacciarini, M. A. Swiderska, C. Nativi, M. Sattler, A. R. Smyth, P. Williams, M. Camara, A. Stocker, T. Darbre and J. L. Reymond, *Angew. Chem., Int. Ed.*, 2011, **50**, 10631-10635.
35. J. Rodrigue, G. Ganne, B. Blanchard, C. Saucier, D. Giguere, T. C. Shiao, A. Varrot, A. Imberty and R. Roy, *Organic & Biomolecular Chemistry*, 2013, **11**, 6906-6918.
36. R. U. Kadam, D. Garg, J. Schwartz, R. Visini, M. Sattler, A. Stocker, T. Darbre and J. L. Reymond, *ACS Chem. Biol*, 2013, **8**, 1925-1930.
37. M. L. Gening, D. V. Titov, S. Cecioni, A. Audfray, A. G. Gerbst, Y. E. Tsvetkov, V. B. Krylov, A. Imberty, N. E. Nifantiev and S. Vidal, *Chemistry-a European Journal*, 2013, **19**, 9272-9285.
38. C. Ligeour, L. Dupin, A. Marra, G. Vergoten, A. Meyer, A. Dondoni, E. Souteyrand, J. J. Vasseur, Y. Chevolot and F. Morvan, *European Journal of Organic Chemistry*, 2014, 7621-7630.
39. J. J. P. Stewart, *J Mol Model*, 2007, **13**, 1173-1213.
40. G. D. Hawkins, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem.*, 1996, **100**, 19824-19839.
41. D. C. M. Kong and M. von Itzstein, *Carbohydr. Res.*, 1997, **305**, 323-329.
42. R. Mazurczyk, G. E. Khoury, V. Dugas, B. Hannes, E. Laurenceau, M. Cabrera, S. Krawczyk, E. Souteyrand, J. P. Cloarec and Y. Chevolot, *Sens. Actuators, B*, 2008, **128**, 552-559.
43. M. Phaner-Goutorbe, V. Dugas, Y. Chevolot and E. Souteyrand, *Mater. Sci. Eng., C*, 2011, **31**, 384-390.
44. W. L. Jorgensen and J. Tirado-Rives, *J. Comput. Chem.*, 2005, **26**, 1689-1700.
45. P. Derreumaux and G. Vergoten, *J. Chem. Phys.*, 1995, **102**, 8586-8605.
46. G. Vergoten, I. Mazur, P. Lagant, J. C. Michalski and J. P. Zanetta, *Biochimie*, 2003, **85**, 65-73.
47. P. Lagant, D. Nolde, R. Stote, G. Vergoten and M. Karplus, *J Phys Chem A*, 2004, **108**, 4019-4029.