

Organic & Biomolecular Chemistry

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.


Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Graphical Abstract

The reactivity of a selection of azulenium salts, derived from photochromic dihydroazulenes (DHAs), towards a nucleophile is reported, and one regioisomer product is tautomerised into a new photochromic DHA that undergoes a reversible ring-opening reaction to a vinylheptafulvene (VHF).



ARTICLE

Azulenium chemistry – towards new derivatives of photochromic dihydroazulenes

Cite this: DOI: 10.1039/x0xx00000x

Anne Ugleholdt Petersen,^a Martyn Jevric,^a Jonas Elm,^a Stine T. Olsen,^a Christian G. Tortzen,^a Anders Kadziola,^a Kurt V. Mikkelsen,^a and Mogens Brøndsted Nielsen^{a,*}

Received 00th January 2012,
Accepted 00th January 2012


DOI: 10.1039/x0xx00000x

www.rsc.org/

Here we present the preparation of a selection of azulenium ions by hydride abstraction from photochromic 1,8a-dihydroazulenes (1,8a-DHAs) incorporating two cyano groups at C-1. The reactivity of the electrophilic tropylium ring of these molecules towards lithium triisopropylsilylacetylide was investigated. The position of attack by the nucleophile depended on the substitution pattern of the azulenium cation but was in general found to occur preferentially at positions C-4, C-5, and C-6, and to a minor extent at positions C-7 and C-8. The outcome was a mixture of non-photochromic, regioisomeric DHAs. One of these compounds containing the ethynyl substituent at position C-4 was partly tautomerized to the photochromic 1,8a-dihydroazulene, which was isolated and its switching properties were investigated by UV-Vis absorption spectroscopy. Upon irradiation, it undergoes a ring-opening reaction to form a vinylheptafulvene (VHF), which in time returns to the original DHA. The half-life of this reaction was significantly smaller than for a derivative with the alkynyl substituent placed at C-7. This fast switching behavior was according to calculations explained by an enhancement in the stability of the reactive *s-cis* conformer of the VHF relative to the, still more stable, *s-trans* conformer, and by a smaller activation energy for this *s-cis* conformer to undergo ring-closure.

Introduction

Dihydroazulene **1** (DHA) is a photoswitch that upon exposure to light undergoes a ring-opening to vinylheptafulvalene **2** (VHF).¹ VHF returns in time to DHA (Scheme 1) with a rate of ring-closure that is solvent dependent and is believed to proceed via a zwitterionic transition state. We have recently focused on exploiting these reversible switching properties in molecular electronics devices² and further efforts are currently underway to employ the DHA/VHF for development of solar thermal energy storage systems and other advanced materials such as photoresponsive liquid crystals.³



Scheme 1. DHA-VHF photo/thermoswitch and numbering of the DHA core.

In order to tune the DHA-VHF properties, in particular in regard to the forward and backward switching events, the DHA and VHF absorption maxima, the energy storage capacity of the metastable VHF, and the possibility for liquid crystallinity, it is desirable to achieve reliable synthetic protocols for regioselectively placing substituents in all positions of the DHA core. So far, it has been possible to selectively brominate the 3- and the 7-positions,⁴ the latter giving an appropriate functional handle for further transformations using palladium cross-coupling methodologies.⁵ In fact, it has been found that the electronic character of substituent groups at position 7 strongly influences the rate of the thermal VHF to DHA back-reaction.^{5a,b,6}

Rather than subjecting DHA **1** to further functionalization, an alternate approach employed phenyltropylium as an

electrophilic starting material for constructing a substituted DHA core.⁷ This method gave, however, a mixture of regioisomers with the phenyl substituent at different positions in the seven-membered ring. From this reaction, small amounts of the 5-Ph substituted isomer could be isolated pure, but the lack of selectivity made this strategy rather inconvenient. Recently, it has been shown that DHA **1** could be oxidized by means of hydride abstraction using tritylium cation to affect the formation of the azulonium salt **3** in good yield (Figure 1).⁸ The same method has been used to form azulonium cations from other dihydroazulenes.⁹ We became interested to prepare a large selection of such azulonium cations with aryl substituents at various positions (**4-7**) from DHAs **8-11** and to investigate the scope of nucleophilic additions to such species. This approach was stimulated by the fact that some regioselectivity was previously observed for reaction of aryltropylium species with nucleophiles,¹⁰ depending on the bulkiness of the nucleophile and the electronic character of the aryl group. The parent 2*H*-azulonium cation has previously been shown to have a clear vinyltropylium ion character and with the charge predominantly localized on C-3a, C-5, C-7, C-8a in the seven-membered ring and on C2 and C3 in the five-membered ring.¹¹ Synthetic protocols for donor- (OMe) and acceptor- (CN) substituted DHAs **8** and **9** have previously been reported^{5d} as has protocols for the 2,3-diphenyl DHA **10**.¹² Synthesis of the new 2,3,7-triphenyl substituted DHA **11** will be presented below.





Figure 1. Retrosynthetic route to azulonium salts **3-7** by hydride abstraction from DHA precursors **1, 8-11**.

Results and Discussion

Synthesis

Synthesis of DHA **11** was performed according to Scheme 2. First, DHA **10** was subjected to the bromination-elimination protocol providing the intermediate **12**, which was then subjected to a Suzuki reaction with phenylboronic acid yielding DHA **11** in reasonable yield over three steps. In addition to **11**, azulenes **13** and **14** were isolated from the reaction mixture, a result of loss of hydrogen cyanide. Single crystals of both **11** and **14** were subjected to X-ray crystallographic analysis, which confirmed the identities of both structures (Figure 2).



Scheme 2. Synthesis of 2,3,7-triphenyl-substituted DHA **11**. RuPhos = 2-dicyclohexylphosphino-2',6'-diisopropoxybiphenyl.




Figure 2. Top: Molecular structure of **11** (CCDC 1441529). The crystals were grown from dichloromethane/ethanol. Bottom: Molecular structure of **14** (CCDC 1441530). The crystals were grown from THF/heptane. The blue spheres represent nitrogen atoms.

Subjecting DHA **11** to light produced the corresponding VHF as evidenced by a redshift in the absorption maximum in the UV-Vis absorption spectrum (Figure 3). The VHF underwent a very fast back-reaction with a half-life of ca. 11 s at 25 °C in acetonitrile. The half-life is thus halved relative to that of the VHF corresponding to DHA **10**, for which a half-life of 20 s at 25 °C has been reported.¹²





Figure 3. UV-Vis absorption spectra of pure DHA **11** and the corresponding VHF in MeCN at 25 °C. On account of the fast VHF to DHA conversion, the spectrum of VHF contains overlapping DHA absorptions.

Our next objective was to study the possibility of converting DHAs **1** and **8-11** to the corresponding azulonium ions. Hydride abstraction of these DHAs was accomplished by treatment with $[\text{Ph}_3\text{C}]\text{BF}_4$ in refluxing 1,2-dichloroethane (DCE) with reaction times varying from one night to two days depending upon the substitution pattern of the DHA (Scheme 3, Table 1). Noteworthy, generation of the azulonium ion required a long reaction time in the presence of the electron-withdrawing cyano substituent (R^1). The salts **3-7** were conveniently precipitated by the addition of diethyl ether to the cooled reaction mixtures. All these salts were generally poorly soluble in most organic solvents although it was possible to obtain full NMR spectroscopic analyses using CD_3CN as solvent, despite a limited stability in this solvent. The yield of the salt **3** was slightly improved in comparison to our previous report⁸ (from 86 to 93%) by performing the reaction under inert atmosphere.



Scheme 3. Synthesis of azulonium tetrafluoroborate salts **3-7**. DCE = 1,2-dichloroethane.

Table 1. Synthesis of azulonium salts – reaction times and yields.


DHA – Starting material	R^1	R^2	R^3	Reaction time (h)	Azulenium salt – Product	Yield (%)
1	H	H	H	17	3	93
8	OMe	H	H	15	4	85
9	CN	H	H	38	5	71
10	H	Ph	H	24	6	69
11	H	Ph	Ph	39	7	78

The absorption profiles for the pure azulonium tetrafluoroborates **3-7** were measured in both CH_3CN and CH_2Cl_2 (see ESI). Table 2 lists the longest-wavelength absorption maxima. When going from CH_3CN to CH_2Cl_2 the salts experience a redshift in the absorption maximum of ca. 35 nm.

Table 2. Longest-wavelength absorption maxima of azulonium salts.

Azulenium salt	$\lambda_{\text{max}}(\text{CH}_3\text{CN})$ (nm)	$\lambda_{\text{max}}(\text{CH}_2\text{Cl}_2)$ (nm)
3	435	468
4	509	545
5	438	475
6	438	473
7	463	501

Azulenium salts **3-7** were then subjected to nucleophilic attack with lithium triisopropylsilylacetylide. This nucleophile was chosen as we found that other nucleophiles such as alcohols tended to give labile products. In addition, molecules with potential for subsequent acetylenic coupling reactions (after desilylation) are desirable. The product outcomes are summarized in Scheme 4 and Table 3. The general reaction conditions involved the addition of a THF solution of the nucleophile at -78 °C to a suspension of the azulonium salt also held at -78 °C in THF. In these cases, low temperature was used to avoid the formation of azulene products through an elimination of hydrogen cyanide, a side reaction which typically can occur under basic conditions.^{4a} The addition product ratios were determined after the reaction mixture was passed through a silica gel column to remove any unwanted impurities and all these mixtures were determined to be analytically pure. In all cases, some of the regioisomers could be separated from their respective mixtures using subsequent flash column chromatography, which allowed for the assignment of some of the products.

Scheme 4. Addition of triisopropylsilylacetylide anion to azulonium ions **3-7**.Table 3. Product distributions from nucleophilic attack of lithium triisopropylsilylacetylide on cations **3-7**.


Entry	Starting material	Product	Overall yield (%)	% of isomers ^{a)}				
				a	b	c	d	e
1	3	15	73	29	47	9	10	5
2	4	16	57	28	28	27	12	5
3	5	17	40	19	30	39	9	3
4	6	18	44	6	51	20	16	7
5	7	19	25	25	21 ^{b)}	54	-	-

^{a)} Determined by ¹H NMR spectroscopic analysis of the pure regioisomeric mixture (obtained after chromatographic purification). ^{b)} Compound **20**.


It was found that the reaction of the acetylide with **3** (Entry 1) gave a mixture of isomers, with the addition pattern predominating in the 5- (**15b**) and 4-positions (**15a**). Using flash column chromatography, it was possible to isolate and characterize these two main isomers. All addition products have proton connectivity spanning the 7-membered ring from position 4 to 7, where the point of attack is sp³ hybridized. Coupling constants could be used to distinguish the nature of

the protons upon the adjacent carbons, as *cis*-alkene couplings (ca. 10 Hz) were distinguishable from couplings between protons upon adjacent “alkene units” (ca. 7 Hz), which assisted in elucidating the structural framework. The orientation relative to the 5-membered ring could be verified by a NOESY correlation from the proton at the 3-position. The reactions with the azulonium **4** bearing an electron-donating OMe group and the azulonium **5** bearing an electron-withdrawing CN group showed minor changes to the addition pattern (Entries 2 and 3), but unfortunately these reactions appeared to be less regioselective. It was envisaged that the phenyl substituent in position 3 of azulonium **6** could potentially block attack at position 4 (**18a**). Indeed, it was found that attack on position 4 did not occur in the treatment of **6** with the acetylide anion (Entry 4), and the substitution at the 5-position (**18b**) was the predominant product. Nevertheless, the yield for this reaction was significantly lower than for the others. The addition of triisopropylsilylacetylide to the highly substituted system gave **19a** and **19c** in addition to an unexpected product **20**. In this case addition to the 5-position had occurred but with formation of a norcaradiene product. This type of product¹³ has previously been observed in the reaction of 1,3-diphenyltropylium with nucleophiles.¹⁴

None of the products formed are photoactive in regard to a ring-opening reaction. They all require a double bond migration to move a hydrogen to the 8a position in order to generate the photoactive DHA. A 1,5-sigmatropic shift has been reported to occur in alkynyl substituted cycloheptatriene in warm DMF.¹⁵ As the photoactive DHA has the longest conjugation pathway, one could imagine sigmatropic shifts to terminate at the presumably thermodynamically most stable, photoactive DHA. Due to the tedious purification procedure in separating the isomers from the addition reaction, the double bond isomerization reaction was examined on only two derivatives, **15a** and **15b** featured in Scheme 5. It was found that heating the 4-isomer **15a** was very destructive but resulted in both formation of the photoactive compound **21** (in 11%) and the corresponding azulene **22** (in 16%), arising from loss of hydrogen cyanide. Purification of the DHA was made possible by preparative TLC, where the plate was continuously irradiated using several TLC lamps (355 nm), as the VHF form of **21**, formed under the illumination, is more polar allowing for an easier purification. On the other hand, thermolysis of the 5-substituted isomer **15b** in DMF did not yield any photoswitching products, instead significant conversion to the azulene **23** was obtained, the structure of which was confirmed by X-ray crystallography (Figure 4). Heating of both **15a** and **15b** in other solvents was investigated, but they did not facilitate the formation of neither photoactive species nor azulene by-products and due to such low yields of both the addition products and the isomerization, this was not further pursued.






Scheme 5. Tautomerization of dihydroazulene products.

Figure 4. Top: Molecular structure of **23** (CCDC 1441531). The crystals were grown from evaporating ethyl acetate. The blue sphere represents nitrogen and the yellow sphere silicon.

Switching studies of 4-alkynyl substituted DHA **21**

With the first, pure 4-substituted 1,8a-DHA **21** in hand, its photoswitching properties were studied using UV-vis absorption spectroscopy. Upon irradiation of **21** with light (ca. 375 nm), the corresponding VHF **24** was formed with a characteristic absorption maximum at 473 nm (Figure 5); the VHF returned in the dark to **21** with an identical absorption spectrum as that measured before photolysis and with spectral evolution with clean isobestic points, suggesting no conversion to the congested 8a-isomer **25** (with a Z-VHF as open form, Scheme 6). In contrast, it was previously found that a ring-opening/closing cycle of a 7-substituted DHA results in the formation of a mixture of 6- and 7-substituted DHAs accompanied by a red-shift in the characteristic DHA absorption, arising from the contributions from the 6-isomer.^{4a} This isomerism has also been observed in experiments upon 5-phenyl substituted DHA (5-/8-substituted isomers), and this study was complimented by NMR spectroscopy. The lack of isomerization of the 4-substituted isomer is in agreement with the actual purification procedure that was performed where the compound was constantly irradiated. DFT calculations also show that DHA **25** is significantly higher in Gibbs free energy than the isomer **21**, by 30.6-38.8 kJ/mol, depending on the DFT

functional utilized (see ESI). This clearly indicates that there should be no conversion to **25** in an opening-closure cycle since it is thermochemically disfavored.

Scheme 6. Ring-opening/closure of DHA **21**.Figure 5. Top: Absorption spectra of DHA **21** and its corresponding VHF **24** in MeCN at 25 °C. Bottom: Arrhenius plot for the thermal VHF to DHA ring-closure. $E_a = 95.8 \pm 1.9 \text{ kJ/mol}$ and $A = 7.10 \times 10^{12} \text{ s}^{-1}$.Figure 6. Conformations of VHF **E-24** and **Z-26**.

The VHF **24** to DHA **21** conversion is readily followed by the first-order decay of the VHF absorption at 473 nm. The back-reaction was followed at five different temperatures in order to make an Arrhenius plot for the kinetics of the ring-closure. This plot provided an activation energy for the back-reaction of $E_a = 95.8 \pm 1.9$ kJ/mol and a pre-exponential factor of $A = 7.10 \times 10^{12}$ s⁻¹. The rate constant at 25 °C was determined to 1.165×10^{-4} s⁻¹, corresponding to a half-life of 99.2 min. In comparison, the VHF **26** (Figure 6) of the isomeric 7-triisopropylsilylethynyl substituted DHA was previously found to undergo ring-closure with a half-life of 670 min at 25 °C.^{4a} Thus, VHF **24** undergoes a significantly faster ring-closure. It is known from previous studies of the ring-closure of substituted phenyl VHF that electron-withdrawing or donating groups can affect the rate of the thermal back reaction (Hammett correlations),⁶ yet in the present case the position of the substituent on the heptafulvene core seems of importance. The VHF has two conformers, *s-cis* and *s-trans*, which exist in equilibrium through a single bond rotation (Figure 6). It is only *s-cis* that can close back to the DHA, yet the *s-trans* is usually more stable. If the energy difference between the two conformers is lower, then the VHF equilibrium lies closer to the reactive *s-cis* form, and ring-closure should then occur faster. To shed further light on this, we subjected the VHF isomers to DFT calculations (at different levels, see ESI). In all cases the *s-trans* conformation is found more stable than the corresponding *s-cis* conformation. The *s-trans* conformation of **Z-26** is found to be 10-11 kJ/mol lower in Gibbs free energy than the corresponding *s-cis* conformation. For **E-24**, the Gibbs free energy difference is found somewhat lower, with the *s-trans* conformation being now only 2.9 to 4.1 kJ/mol lower in Gibbs free energy than the *s-cis* conformation. This supports that VHF **E-24** closes faster due to a lower energy difference between the *s-cis* and *s-trans* conformations, shifting the thermal equilibrium more towards the reactive *s-cis* conformation. The reactivity of the *s-cis* conformers of **24** and **26** may also depend on the position of the alkyne unit. For example, conjugation effects could play a role. Therefore, we calculated the transition state (TS) energies (in MeCN) of the related *s-cis* VHF where the *i*-Pr groups were substituted with Me groups (for calculational reasons). The corresponding TS structures are shown in ESI, and we obtain differences in Gibbs free energies of activation at 25 °C for “**24**-SiMe₃” and “**26**-SiMe₃”, which vary between 1.0 and 3.4 kJ mol⁻¹ depending on the functional used. At the CAM-B3LYP/6-311+G(d) level, free energies of activation of 91.5 (“**24**-SiMe₃”) and 94.9 kJ mol⁻¹ (“**26**-SiMe₃”) were obtained, which correspond to a factor of 4 in rate constants (using the Eyring equation), while that observed experimentally is ca. 6.8. The influence was, however, smaller at the M06-2X level (difference in TS energies of +1.0 kJ mol⁻¹). In all, it seems that not only does the larger content of *s-cis* conformer of **24** relative to that of **26** play a role for its faster back-reaction, but the *s-cis* conformer of **24** seems also to be more reactive.

Conclusions

It has been possible to synthesize a host of azulenium salts from their respective DHAs. Addition of lithium triisopropylsilylacetylide to these salts has shown that the positions 4, 5, and 6 are the most predominant points of attack in an otherwise reaction of low selectivity. The electronic influence of either donating or withdrawing groups upon the *para*-position of a phenyl ring at C-2 did little to enhance the selectivity of this addition, yet the placement of a phenyl blocking group in the 3-position of the azulenum acted as to prevent attack in the 4-position. The 4-substituted derivative was found to undergo a 1,5-sigmatropic shift to form a photoactive DHA (with a hydrogen on position 8a), which allowed switching studies of this 4-substituted DHA/VHF system. An opening-closure cycle gave none of the isomeric 8a-substituted isomer, which according to DFT calculations is of significantly higher energy. Interestingly, the VHF to DHA ring-closure rate was significantly enhanced compared to the previously studied 7-substituted analogue. For the ring-opened VHF a lower energy difference between *s-cis* and *s-trans* conformations was calculated (but still in favor of *s-trans*) than for the VHF obtained from the 7-substituted DHA. More molecules are thus in the reactive *s-cis* conformation on account of the sterical constraints imposed by the bulky triisopropylsilyl group. The calculations also indicate that the *s-cis* conformer is more reactive by itself when the alkyne substituent is next to the exocyclic double bond.

While previous experiments have shown that fast VHF to DHA ring-closure can be promoted by *i*) suitable donor-acceptor substitution,⁶ *ii*) by having a phenyl substituent at position 3,^{1b, 12} or *iii*) by locking the VHF in its *s-cis* conformation via a covalent bridge (fused ring system),¹⁶ the current study has shown that a bulky substituent at position 4 also provides a means of achieving fast switching.

Experimental

General Methods. NMR spectra were acquired with a Bruker Avance UltraShield 500 Plus instrument equipped with a cryoprobe or a Bruker 500MHz Avance III system with a broadbandprobe probe (frequencies of 500 and 125 MHz for ¹H and ¹³C NMR, respectively). All chemical-shift values in ¹H and ¹³C NMR spectra were referenced to the residual solvent peak (CDCl₃: δ_H = 7.26 ppm, δ_C = 77.16 ppm; CD₃CN: δ_H = 1.96 ppm, δ_C = 1.94 ppm; C₆D₆: δ_H = 7.16 ppm, δ_C = 128.06 ppm). For compounds **19a**, **19c**, and **20**, two ¹³C signals were observed for the TIPS CH₃ carbon nuclei at 300 K in CDCl₃, which we ascribe to slow conversion between rotamers. Thus, recording a sample of **19a** at 323 K caused coalescence of these two signals into one signal. All starting materials were either purchased from Sigma-Aldrich, Combiblocks or synthesized; compounds **1**,¹⁷ **8**,^{5d} **9**,^{5d} and **10**¹² were prepared according to literature procedures. Palladium catalysed cross couplings and all reactions involving the formation and use of azulenum salts were performed under an argon atmosphere. HPLC grade solvents were used as received with the exception of anhydrous THF and diethyl ether which were freshly distilled from a

sodium/benzophenone couple. Flash column chromatography was performed on 40–63 μm SiO_2 , and TLC was performed on commercially available precoated silica plates and examined under UV light. All manipulations of DHA in solution were done so in the dark and the vessels were covered with aluminium foil. Melting points were measured with a Büchi apparatus and are uncorrected. Microanalyses were performed either at the University of Copenhagen, Department of Chemistry, or at London Metropolitan University. Electrospray ionization (ESI) mass spectra were acquired with a Bruker MicroTOF-QII-system, whereas matrix-assisted laser desorption ionization (MALDI) mass spectra were acquired with a Bruker FT-ICR instrument equipped with a 7 T magnet. Prior to the experiments, the instrument was calibrated by using NaTFA cluster ions. UV-Vis absorption spectroscopy measurements were performed using a 1 cm-path-length cuvette. UV-Vis absorption spectra were obtained by scanning the wavelength from 800 to 200 nm in a 1 cm-path-length cuvette using a Cary 50 Bio UV-visible spectrophotometer. Photoswitching experiments were performed using a 150-W Xenon arc lamp equipped with a monochromator. The thermal back-reaction was performed by heating the sample (cuvette) by a Peltier unit in the UV-Vis spectrophotometer.

2,3-Diphenylazulene-1,1(8aH)-dicarbonitrile (10): Compound **10** was made according to the literature procedure.¹² ^1H NMR (500 MHz, CDCl_3) δ = 7.39–7.31 (m, 5H), 7.30–7.26 (m, 3H), 7.20–7.12 (m, 2H), 6.56 (dd, J = 11.1, 6.0 Hz, 1H), 6.51 (dd, J = 11.1, 5.7 Hz, 1H), 6.34 (ddd, J = 10.0, 5.7, 2.0 Hz, 1H), 6.07 (br d, J = 6.0 Hz, 1H), 5.82 (dd, J = 10.0, 4.0 Hz, 1H), 3.73 (ddd, J = 4.0, 2.0, 2.0 Hz, 1H) ppm. ^{13}C NMR (125 MHz, CDCl_3) δ = 145.7, 140.1, 136.2, 132.6, 131.7, 131.2, 130.9, 129.4, 129.4, 129.0, 129.0, 129.0, 128.8, 127.7, 121.1, 119.7, 115.8, 113.1, 50.2, 47.4 ppm.

2,3,7-Triphenylazulene-1,1(8aH)-dicarbonitrile (11): To a stirring solution of **10** (1.07 g, 3.22 mmol) in CH_2Cl_2 (30 mL) at -78°C was added dropwise a solution of Br_2 (4.15 mL, 0.78M in CH_2Cl_2 , 3.24 mmol) and stirring was continued for 1 h. The solvent was removed under reduced pressure to give the dibromide as an off-orange solid. The dibromide was dissolved in dry THF (50 mL), under an argon atmosphere and cooled to 0°C , after which time a solution of LiHMDS (3.5 mL, 1.0M in toluene, 3.5 mmol) was slowly added and the contents stirred for 2 h, where the temperature was allowed to slowly reach rt. The reaction was quenched with saturated aqueous NH_4Cl (20 mL) and the contents diluted with water (80 mL). The mixture was extracted with a THF/ Et_2O mixture (1:9; 2 x 100 mL). The combined organics were dried over MgSO_4 , filtered and the solvent was removed in vacuo. The residue was taken up in toluene (100 mL) and water (20 mL). To this degassed biphasic mixture containing **12** were sequentially added potassium phosphate (1.99 g, 9.38 mmol), phenylboronic acid (516 mg, 4.23 mmol), RuPhos (156 mg, 0.334 mmol) and $\text{Pd}(\text{OAc})_2$ (39 mg, 0.174 mmol), and the resulting mixture was heated to 55°C for 16 h. Since the reaction was not done, the reaction mixture was degassed again and additional potassium phosphate (1.07 g, 5.04 mmol), phenylboronic acid (264 mg, 2.17 mmol), RuPhos (88 mg, 0.189 mmol) and $\text{Pd}(\text{OAc})_2$ (25 mg, 0.111 mmol) were added, and the reaction mixture was heated to 65°C for 8 h. The contents of the vessel were diluted with water (100 mL) and the phases separated. The aqueous phase was extracted with CH_2Cl_2 (2 x 100 mL) and the

combined organics dried over MgSO_4 . Filtration and removal of the solvent in vacuo gave a crude residue which was subjected to flash column chromatography (CH_2Cl_2 /heptane 1:1) to afford pure **11** (579 mg, 44%) as a yellow solid. The reaction also yielded **13** (78 mg, 8%) as a purple solid and **14** (102 mg, 8%) as a green solid.

(11): M.p. = $172\text{--}175^\circ\text{C}$. R_f = 0.38 (1:1 CH_2Cl_2 /heptane). ^1H NMR (500 MHz, CDCl_3) δ = 7.47–7.27 (m, 13H), 7.22–7.16 (m, 2H), 6.84–6.79 (m, 2H), 6.12 (ddd, J = 4.9, 1.5, 1.5 Hz, 1H), 6.03 (br d, J = 4.9 Hz, 1H), 3.83 (dd, J = 4.9, 1.5 Hz, 1H) ppm. ^{13}C NMR (125 MHz, CDCl_3) δ = 145.1, 142.0, 139.9, 139.8, 136.8, 132.7, 132.4, 131.9, 131.7, 129.5, 129.4, 129.1, 129.0, 129.0, 128.9, 128.7, 128.2, 127.9, 120.4, 116.3, 115.9, 113.2, 50.2, 47.3 ppm. HRMS (MALDI +ve) calcd for $\text{C}_{30}\text{H}_{20}\text{N}_2$ [M^+]: m/z = 408.1616; found m/z = 408.1630. Anal. calcd (%) for $\text{C}_{30}\text{H}_{20}\text{N}_2$ (408.50): C 88.21, H 4.94, N 6.86; found C 88.23, H 4.85, N 6.90.

2,3-Diphenylazulene-1-carbonitrile (13): M.p. = $201\text{--}216^\circ\text{C}$ (decomposes). R_f = 0.26 (CH_2Cl_2 /heptane 1:1). ^1H NMR (500 MHz, CDCl_3) δ = 8.47 (d, J = 9.7 Hz, 1H), 8.04 (d, J = 9.8 Hz, 1H), 7.60 (d, J = 7.7 Hz, 2H), 7.14–7.04 (m, 8H), 7.03–6.87 (m, 1H), 6.71 (t, J = 9.7 Hz, 1H), 6.65 (t, J = 9.8 Hz, 1H) ppm. ^{13}C NMR (125 MHz, CDCl_3) δ = 151.3, 144.3, 140.1, 139.1, 137.2, 136.4, 135.5, 134.6, 131.6, 130.9, 129.9, 128.8, 128.8, 128.4, 127.5, 127.4, 127.3, 117.4, 97.8 ppm. HRMS (MALDI +ve) calcd for $\text{C}_{23}\text{H}_{15}\text{N}$ [M^+]: m/z = 305.1199; found m/z = 305.1203.

2,3,7-Triphenylazulene-1-carbonitrile (14): M.p. > 230°C . R_f = 0.31 (CH_2Cl_2 /heptane 1:1). ^1H NMR (500 MHz, C_6D_6) δ = 9.02 (d, J = 1.8 Hz, 1H), 8.04 (d, J = 9.8 Hz, 1H), 7.71–7.56 (m, 2H), 7.44 (dt, J = 10.4, 0.9 Hz, 1H), 7.39–7.29 (m, 2H), 7.29–7.07 (m, 10H), 7.06–6.95 (m, 1H), 6.74 (t, J = 10.1 Hz, 1H) ppm. ^{13}C NMR (125 MHz, C_6D_6) δ = 151.8, 143.7, 143.6, 141.1, 140.0, 139.5, 136.9, 136.1, 135.5, 134.6, 131.6, 130.9, 129.6, 129.3, 128.9, 128.8, 128.6, 128.4, 127.4, 127.1, 117.5, 98.6 ppm, 1C masked. HRMS (MALDI +ve) calcd for $\text{C}_{29}\text{H}_{19}\text{N}$ [M^+]: m/z = 381.1512; found m/z = 381.1522. Anal. calcd (%) for $\text{C}_{29}\text{H}_{19}\text{N}$ (381.48): C 91.31, H 5.02, N 3.67; found C 91.22, H 5.14, N 3.59.

1,1-Dicyano-2-phenyl-1H-azulenium tetrafluoroborate (3): Prepared from a modified literature procedure.⁸ A solution of **1** (1.56 g, 6.08 mmol) and $[\text{Ph}_3\text{C}]\text{BF}_4$ (3.01 g, 9.13 mmol) in $\text{CH}_2\text{ClCH}_2\text{Cl}$ (60 mL) was heated to reflux point for 17 h under argon atmosphere, after which time, the reaction mixture was allowed to cool to rt and precipitation was induced by the addition of dry Et_2O (40 mL). The precipitate was isolated by filtration and washed with dry Et_2O (3 x 20 mL) yielding **3** (1.94 g, 93%) as a yellow solid. M.p. = 178°C (decomposes). ^1H NMR (300 MHz, CD_3CN) δ = 9.44 (dd, J = 8.9, 1.5 Hz, 1H), 9.18–9.09 (m, 2H), 9.08–8.97 (m, 2H), 8.29 (s, 1H), 8.23–8.16 (m, 2H), 7.86–7.73 (m, 3H) ppm. ^{13}C NMR (75 MHz, CD_3CN) δ = 170.9, 157.6, 157.5, 155.5, 153.6, 153.1, 148.0, 147.7, 136.8, 135.7, 131.4, 129.8, 128.6, 110.4, 46.2 ppm. MS (FAB +ve); m/z = 255 [M^+]. Anal. calcd (%) for $\text{C}_{18}\text{H}_{11}\text{N}_2\text{BF}_4$ (342.10): C 63.04, H 3.00, N 8.14; found C 63.18, H 3.24, N 8.19.

1,1-Dicyano-2-(4-methoxyphenyl)-1H-azulenium tetrafluoroborate (4): To a solution of **8** (1.05 g, 3.67 mmol) in $\text{CH}_2\text{ClCH}_2\text{Cl}$ (60 mL) was added $[\text{Ph}_3\text{C}]\text{BF}_4$ (1.45 g, 4.40

mmol), and the resulting solution was heated to reflux point for 15 h under argon atmosphere. To the cooled reaction mixture was added dry Et₂O (40 mL) and the resulting precipitate was collected by suction filtration and washed with dry Et₂O (3 x 80 mL) giving **4** (1.16 g, 85%) as golden brown microcrystals. M.p. = 209–220 °C (decomposes). ¹H NMR (500 MHz, CD₃CN) δ = 9.20 (dd, *J* = 8.7, 1.4 Hz, 1H), 8.94–8.83 (m, 2H), 8.82–8.68 (m, 2H), 8.17–8.11 (m, 2H), 8.04 (s, 1H), 7.31–7.23 (m, 2H), 3.99 (s, 3H) ppm. ¹³C NMR (125 MHz, CD₃CN) δ = 171.8, 166.8, 156.8, 156.7, 156.3, 151.9, 151.4, 146.7, 146.1, 133.9, 132.8, 121.3, 117.3, 110.6, 57.1, 45.6 ppm. MS (ESP+): *m/z* = 258 [M⁺]. Anal. calcd (%) for C₁₉H₁₃N₂OBF₄ (372.13): C 61.32, H 3.52, N 7.53. found C 61.65, H 3.07, N 7.42.

1,1-Dicyano-2-(4-cyanophenyl)-1H-azulenium tetrafluoroborate (5): A solution of **9** (1.00 g, 3.58 mmol) and [Ph₃C]BF₄ (1.52 g, 4.60 mmol) in CH₂ClCH₂Cl (60 mL) was heated to reflux point for 14 h argon atmosphere. [Ph₃C]BF₄ (792 mg, 2.40 mmol) was added and the mixture was refluxed for a further 8 h, and extra [Ph₃C]BF₄ (892 mg, 2.70 mmol) was added, and the mixture was refluxed for a further 16 h. The contents of the reaction vessel were cooled to rt and the product was precipitated by addition of dry Et₂O (40 mL). The precipitate was collected by filtration and washed with dry Et₂O (3 x 20 mL) and a dry Et₂O / CH₂Cl₂ mixture (1:1, 3 x 15 mL) giving **5** (931 mg, 71%) as a yellow powder. M.p. = 208–217 °C (decomposes). ¹H NMR (500 MHz, CD₃CN) δ = 9.53 (d, *J* = 9.2 Hz, 1H), 9.37–8.79 (m, 4H), 8.39 (s, 1H), 8.26 (d, *J* = 8.5 Hz, 2H), 8.07 (d, *J* = 8.5 Hz, 2H) ppm. ¹³C NMR (125 MHz, CD₃CN) δ = 169.7, 158.1, 158.1, 154.7, 154.3, 152.3, 148.8, 148.8, 139.7, 134.9, 132.3, 130.0, 118.6, 117.6, 109.9, 46.4 ppm. HRMS (ESP +ve) calcd for C₁₉H₁₀N₃: *m/z* = 280.0867; found *m/z* = 280.0869. Anal. calcd (%) for C₁₉H₁₀N₃BF₄ (367.11): C 62.16, H 2.75, N 11.45; found C 62.42, H 2.51, N 10.92.

1,1-Dicyano-2,3-diphenyl-1H-azulenium tetrafluoroborate (6): A solution of **10** (1.56 g, 4.70 mmol) and [Ph₃C]BF₄ (2.33 g, 7.05 mmol) in CH₂ClCH₂Cl (60 mL) was heated to reflux point for 15 h argon atmosphere, after which extra [Ph₃C]BF₄ (1.23 g, 3.73 mmol) was added, and the mixture was refluxed for a further 9h. Then the reaction mixture was allowed to cool to rt and precipitation was induced by the addition of dry Et₂O (40 mL). The precipitate was isolated by filtration and washed with dry Et₂O (3 x 50 mL) affording **6** (1.36 g, 69%) as a yellow powder. M.p. = 208–217 °C (decomposes). ¹H NMR (500 MHz, CD₃CN) δ = 9.55 (dd, *J* = 9.3, 1.0 Hz, 1H), 9.21–8.98 (m, 3H), 8.79–8.65 (m, 1H), 7.70–7.63 (m, 5H), 7.62–7.58 (m, 1H), 7.53–7.46 (m, 4H) ppm. ¹³C NMR (125 MHz, CD₃CN) δ = 171.2, 157.7, 157.3, 154.2, 154.0, 150.2, 149.9, 148.1, 147.0, 133.9, 131.8, 131.2, 130.7, 130.7, 130.6, 130.0, 129.9, 110.4, 48.1 ppm. HRMS (MALDI +ve) calcd for C₂₄H₁₅N₂: *m/z* = 331.1230; exp = 331.1228. Anal. calcd (%) for C₂₄H₁₅N₂BF₄ (418.20): C 68.93, H 3.62, N 6.70; found C 69.02, H 3.59, N 6.79.

1,1-Dicyano-2,3,7-triphenyl-1H-azulenium tetrafluoroborate (7): A solution of **11** (1.06 g, 2.59 mmol) and [Ph₃C]BF₄ (1.28 mg, 3.89 mmol) in CH₂ClCH₂Cl (60 mL) was heated to reflux point for 16 h argon atmosphere, after which extra [Ph₃C]BF₄ (669 mg, 2.03 mmol) was added, and the reaction mixture was refluxed for a further 23 h. Then the reaction mixture was allowed to cool to rt and precipitation was induced by the addition of dry Et₂O (50 mL). The precipitate was isolated by

filtration and washed with a dry Et₂O / CH₂Cl₂ mixture (1:1; 2 x 20 mL). The title compound **7** was isolated as an orange powder (1.00 g, 78%). M.p. >230 °C. ¹H NMR (500 MHz, CD₃CN) δ = 9.68 (d, *J* = 2.1 Hz, 1H), 9.19 (ddd, *J* = 10.2, 2.1, 0.8 Hz, 1H), 8.97 (t, *J* = 10.2 Hz, 1H), 8.58 (dd, *J* = 10.2, 0.8 Hz, 1H), 8.07–8.05 (m, 2H), 7.95–7.74 (m, 3H), 7.72–7.56 (m, 6H), 7.56–7.45 (m, 4H) ppm. ¹³C NMR (125 MHz, CD₃CN) δ = 168.0, 167.1, 156.1, 155.4, 152.2, 149.7, 149.1, 146.5, 144.9, 139.4, 134.5, 133.6, 131.8, 131.7, 131.2, 131.0, 130.7, 130.7, 130.6, 130.1, 110.7, 48.6 ppm, 1C masked. HRMS (MALDI +ve) calcd for C₃₀H₁₉N₂ [M⁺]: *m/z* = 407.1543; found *m/z* = 407.1543. Anal. calcd (%) for C₃₀H₁₉N₂BF₄ (494.30): C 72.90, H 3.87, N 5.76; found C 72.24, H 3.65, N 5.49

Reaction of 3 with lithium triisopropylsilylacetylide: To a degassed solution of triisopropylsilylacetylene (1.2 mL, 5.35 mmol) at -40 °C in Et₂O (25 mL) was added LiHMDS (3.2 mL, 1.0 M in toluene, 3.2 mmol) and the vessel allowed to warm to -20 °C. The reaction contents were cooled to -78 °C and added via cannula to a vessel containing a degassed suspension of **3** (978 mg, 2.86 mmol) in THF (50 mL) at -78 °C. The reaction mixture was stirred for 1 h at -78 °C, quenched with saturated aqueous NH₄Cl (50 mL) and allowed to warm to ambient temperature. The reaction mixture was diluted with Et₂O (100 mL) and water (50 mL) and the phases separated. The aqueous component was extracted with Et₂O (50 mL) and the combined organic extracts were dried over MgSO₄ and filtered. The solvent was removed under reduced pressure and the crude residue was purified by flash column chromatography (toluene/heptane 2:3) to give a regioisomeric mixture of **15** (914 mg, 73%) as a yellow oil. Anal. calcd (%) for C₂₉H₃₂N₂Si (436.67): C 79.77, H 7.39, N 6.42; found C 79.73, H 7.47, N 6.49. Mixture **15** could be subject to further flash column chromatography (2:23 EtOAc/heptane) to give pure samples of the following products.

2-Phenyl-4-[(triisopropylsilyl)ethynyl]-1,4-dihydroazulene-1,1-dicarbonitrile (15a): A yellow solid. M.p. = 98–100 °C. *R_f* = 0.49 (2:23 EtOAc/heptane). ¹H NMR (500 MHz, CDCl₃) δ = 7.86–7.65 (m, 2H), 7.53–7.45 (m, 2H), 7.43–7.37 (m, 1H), 7.30 (s, 1H), 6.95 (d, *J* = 11.2 Hz, 1H), 6.84 (dd, *J* = 11.2, 6.1 Hz, 1H), 6.29 (ddd, *J* = 9.7, 6.1, 1.2 Hz, 1H), 5.46 (dd, *J* = 9.7, 5.0 Hz, 1H), 3.46 (dd, *J* = 5.0, 1.2 Hz, 1H), 1.16–1.14 (m, 21H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ = 142.3, 139.3, 133.5, 132.7, 132.6, 130.2, 129.8, 129.6, 127.2, 125.5, 125.0, 122.4, 112.1, 112.0, 105.2, 85.2, 43.4, 32.2, 18.8, 11.4 ppm. HRMS (MALDI +ve) calcd for C₂₉H₃₁N₂Si [(M-H)⁺]: *m/z* = 435.2251; found *m/z* = 435.2258. Anal. calcd (%) for C₂₉H₃₂N₂Si (436.67): C 79.77, H 7.39, N 6.42; found C 79.35, H 7.40, N 6.34.

2-Phenyl-5-[(triisopropylsilyl)ethynyl]-1,5-dihydroazulene-1,1-dicarbonitrile (15b): A yellow solid. M.p. = 110–112 °C. *R_f* = 0.45 (2:23 EtOAc/heptane). ¹H NMR (500 MHz, CDCl₃) δ = 7.73–7.68 (m, 2H), 7.50–7.46 (m, 2H), 7.45–7.40 (m, 1H), 7.18 (d, *J* = 6.5 Hz, 1H), 7.01 (s, 1H), 6.35 (ddd, *J* = 9.4, 6.5, 1.4 Hz, 1H), 5.56–5.53 (m, 2H), 3.22 (ddd, *J* = 5.4, 5.4, 1.4 Hz, 1H), 1.11–1.08 (m, 21H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ = 139.3, 138.4, 136.7, 132.0, 130.3, 130.0, 129.5, 126.1, 125.8, 123.6, 117.9, 114.1, 113.4, 107.7, 82.2, 43.9, 32.4, 18.8, 11.3 ppm, 1C masked. HRMS (MALDI +ve) calcd for C₂₉H₃₁N₂Si [(M-H)⁺]: *m/z* = 435.2251; found *m/z* = 435.2262. Anal. calcd

(%) for C₂₉H₃₂N₂Si (436.67): C 79.77, H 7.39, N 6.42; found C 79.54, H 7.41, N 6.34.

2-Phenyl-6-[(triisopropylsilyl)ethynyl]-1,5-dihydroazulene-1,1-dicarbonitrile (15c): A yellowish oil. *R_f* = 0.40 (2:23 EtOAc/heptane). ¹H NMR (500 MHz, CDCl₃) δ = 7.75-7.72 (m, 2H), 7.50-7.47 (m, 2H), 7.44-7.41 (m, 1H), 7.14 (s, 1H), 6.75 (dd, *J* = 9.3, 1.4 Hz, 1H), 6.51 (dd, *J* = 9.3, 1.4 Hz, 1H), 5.60-5.54 (m, 2H), 2.92 (dddd, *J* = 5.6, 5.6, 1.4, 1.4 Hz, 1H), 1.10-1.09 (m, 21H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ = 146.2, 139.6, 137.3, 133.5, 129.9, 129.9, 129.6, 126.9, 125.9, 125.0, 121.6, 119.7, 112.4, 112.3, 107.8, 82.2, 44.3, 33.3, 18.8, 11.3 ppm. HRMS (MALDI +ve) calcd for C₂₉H₃₁N₂Si [(M-H)⁺]: *m/z* = 435.2251; found *m/z* = 435.2249.

2-Phenyl-7-[(triisopropylsilyl)ethynyl]-1,5-dihydroazulene-1,1-dicarbonitrile (15d): A yellowish oil. *R_f* = 0.31 (2:23 EtOAc/heptane). ¹H NMR (500 MHz, CDCl₃) δ = 7.71-7.69 (m, 2H), 7.50-7.47 (m, 2H), 7.45-7.41 (m, 1H), 7.14 (s, 1H), 6.70 (d, *J* = 6.4 Hz, 1H), 6.34 (ddd, *J* = 9.9, 6.4, 1.2 Hz, 1H), 6.20 (d, *J* = 5.4 Hz, 1H), 5.44 (dd, *J* = 9.9, 5.4 Hz, 1H), 3.40 (ddd, *J* = 5.4, 5.4, 1.2 Hz, 1H), 1.09-1.08 (m, 21H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ = 141.1, 137.2, 136.0, 134.9, 130.3, 130.0, 129.5, 126.7, 126.4, 126.3, 126.2, 124.1, 113.9, 113.4, 106.9, 82.6, 42.4, 32.2, 18.8, 11.3 ppm. HRMS (MALDI +ve) calcd for C₂₉H₃₁N₂Si [(M-H)⁺]: *m/z* = 435.2251; found *m/z* = 435.2256.

2-Phenyl-8-[(triisopropylsilyl)ethynyl]-1,5-dihydroazulene-1,1-dicarbonitrile (15e): A yellowish oil. *R_f* = 0.26 (2:23 EtOAc/heptane). ¹H NMR (500 MHz, CDCl₃) δ = 7.70-7.68 (m, 2H), 7.47-7.44 (m, 2H), 7.40-7.37 (m, 1H), 7.03 (s, 1H), 6.78 (dd, *J* = 10.6, 6.2 Hz, 1H), 6.71 (d, *J* = 10.6 Hz, 1H), 6.22 (br dd, *J* = 9.2, 6.2 Hz, 1H), 5.38 (dd, *J* = 9.2, 6.2 Hz, 1H), 3.73 (dd, *J* = 6.2, 1.0 Hz, 1H), 1.13 (br s, 21H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ = 141.1, 138.8, 133.6, 132.6, 130.1, 129.5, 129.5, 129.4, 125.9, 124.1, 112.1, 112.0, 87.0, 43.7, 29.0, 18.8, 11.4 ppm, 3C masked. HRMS (MALDI +ve) calcd for C₂₉H₃₁N₂Si [(M-H)⁺]: *m/z* = 435.2251; found *m/z* = 435.2257.

Reactions of 4-7 with lithium triisopropylsilylacetylide were performed in an analogous manner; for details, see ESI.

Tautomerisation of 15a: A solution of **15a** (106 mg, 0.243 mmol) in freshly distilled degassed DMF (7 mL) was heated to 85 °C for 3.5 h, after which time the volatiles were removed under reduced pressure. The crude material was subjected to flash column chromatography (toluene/heptane 1:1), which gave **22** (16.3 mg, 16%), as a dark green glassy solid, in addition to an impure component, which contained **21**. This solution was concentrated and irradiated at 353 nm for 10 h. The then formed VHF was purified by preparative TLC (toluene/heptane 3:1) whilst the plate was subjected to continuous irradiation at 353 nm. To remove the last small impurities the residue was subjected to flash column chromatography (toluene/heptane 1:1), which furnished **21** (12 mg, 11%) as a yellow solid.

1,1-Dicyano-2-phenyl-4-triisopropylsilylacetyl-1,8a-dihydroazulene (21): M.p. = 115-118 °C. *R_f* = 0.41 (toluene/heptane 1:1). ¹H NMR (500 MHz, CDCl₃) δ = 7.81-7.67 (m, 2H), 7.61-7.40 (m, 3H), 7.24 (s, 1H), 6.64 (d, *J* = 11.4 Hz, 1H), 6.47 (dd, *J* = 11.4, 5.9 Hz, 1H), 6.28 (ddd, *J* = 10.1, 5.9, 2.1 Hz, 1H), 5.89 (dd, *J* = 10.1, 4.0 Hz, 1H), 3.82 (dd, *J* =

4.0, 2.1 Hz, 1H), 1.13-1.15 (m, 21H) ppm. ¹³C NMR (125 MHz, CDCl₃) δ = 143.7, 141.6, 133.5, 131.7, 130.7, 130.6, 130.5, 129.5, 127.6, 126.5, 121.1, 115.4, 115.0, 112.7, 104.0, 96.6, 51.0, 45.3, 18.9, 11.4 ppm. HRMS (ESP +ve) calcd for C₂₉H₃₃N₂Si: *m/z* = 437.2424; found *m/z* = 437.2407.

2-Phenyl-4-[(triisopropylsilyl)ethynyl]azulene-1-carbonitrile (22): *R_f* = 0.16 (toluene/heptane 1:1). ¹H NMR (500 MHz, C₆D₆) δ = 8.33 (d, *J* = 9.5 Hz, 1H), 8.24 (dd, *J* = 8.3, 1.0 Hz, 2H), 8.08 (s, 1H), 7.27 (d, *J* = 10.2 Hz, 1H), 7.24-7.21 (m, 2H), 7.11-7.07 (m, 1H), 6.82 (dd, *J* = 10.2, 10.0 Hz, 1H), 6.63 (ddd, *J* = 10.0, 9.5, 1.5 Hz, 1H), 1.22-1.19 (m, 21H) ppm. ¹³C NMR (125 MHz, C₆D₆) δ = 152.5, 145.1, 142.1, 136.5, 135.8, 134.9, 131.7, 131.3, 129.8, 129.5, 129.0, 127.6, 117.6, 117.5, 108.0, 101.5, 96.5, 18.9, 11.7 ppm. HRMS (ESP +ve) calcd for C₂₈H₃₂NSi [(M+H)⁺]: *m/z* = 410.2300; found *m/z* = 410.2299.

Tautomerisation of 15b: A solution of **15b** (102 mg, 0.234 mmol) in distilled degassed DMF (10 mL) was heated to 90 °C for 5.5 h. The solution was allowed to cool to rt, and the DMF was removed under reduced pressure. The crude mixture was purified by flash column chromatography (toluene/heptane 1:1) to give **23** (57 mg, 60%) as a blue solid.

2-Phenyl-5-[(triisopropylsilyl)ethynyl]azulene-1-carbonitrile (23): M.p. = 125-128 °C. *R_f* = 0.21 (toluene/heptane 1:1). ¹H NMR (500 MHz, C₆D₆) δ = 8.26 (ddd, *J* = 1.6 Hz, 1H), 8.18 (ddd, *J* = 9.5, 0.9, 0.9 Hz, 1H), 8.05-8.03 (m, 2H), 7.48 (ddd, *J* = 10.5, 1.6, 0.9 Hz, 1H), 7.23-7.19 (m, 2H), 7.13-7.10 (m, 1H), 6.91 (s, 1H), 6.50 (dd, *J* = 10.5, 9.6 Hz, 1H), 1.25-1.24 (m, 21H) ppm. ¹³C NMR (125 MHz, C₆D₆) δ = 152.8, 145.2, 141.3, 140.8, 140.8, 135.6, 134.4, 129.8, 129.4, 129.0, 126.4, 122.7, 118.3, 117.4, 110.5, 96.9, 92.1, 18.9, 11.7 ppm. HRMS (ESP +ve) calcd for C₂₈H₃₁NNa [(M+Na)⁺]: *m/z* = 432.2118, found *m/z* = 432.2118. Anal. calcd (%) for C₂₈H₃₁NSi (409.65): C 82.10, H 7.63, N 3.42; found C 81.58, H 7.34, N 3.74.

Acknowledgements

University of Copenhagen and the Danish Council for Independent Research | Technology and Production Sciences (#12-126668) are acknowledged for financial support. We thank Dr. Søren L. Broman for generous donation of starting materials.

Notes and references

^a Department of Chemistry, University of Copenhagen, Universitetsparken 5, DK-2100 Copenhagen Ø, Denmark. E-mail: mbn@chem.ku.dk

[†] Electronic Supplementary Information (ESI) available: Protocols for reactions of **4-7** with triisopropylsilyl acetylide, NMR and UV-Vis spectra and calculational data. See DOI: 10.1039/b000000x/

- (a) J. Daub, T. Knöchel and A. Mannschreck, *Angew. Chem. Int. Ed. Engl.* 1984, **23**, 960; (b) T. Mrozek, A. Ajayaghosh and J. Daub, in *Molecular Switches* (Ed. B. L. Feringa), Wiley-VCH, Weinheim, Germany, 2001, pp. 63–106. (c) S. L. Broman and M. B. Nielsen, *Phys. Chem. Chem. Phys.*, 2014, **16**, 21172.
- (a) S. Lara-Avila, A. V. Danilov, S. E. Kubatkin, S. L. Broman, C. R. Parker and M. B. Nielsen, *J. Phys. Chem. C*, 2011, **115**, 18372; (b) S.

- L. Broman, S. Lara-Avila, C. L. Thisted, A. D. Bond, S. Kubatkin, A. Danilov and M. B. Nielsen, *Adv. Funct. Mater.*, 2012, **22**, 4249; (c) T. Li, M. Jevric, J. R. Hauptmann, R. Hviid, Z. Wei, R. Wang, N. E. A. Reeler, E. Thyraug, S. Petersen, J. A. S. Meyer, N. Bovet, T. Vosch, J. Nygård, X. Qui, W. Hu, Y. Liu, G. C. Solomon, H. G. Kjaergaard, T. Bjørnholm, M. B. Nielsen, B. W. Laursen and K. Nørgaard, *Adv. Mater.*, 2013, **25**, 4164.
- 3 (a) S. T. Olsen, J. Elm, F. E. Storm, A. N. Gejl, A. S. Hansen, M. H. Hansen, J. R. Nikolajsen, M. B. Nielsen, H. G. Kjaergaard and K. V. Mikkelsen, *J. Phys. Chem. A*, 2015, **119**, 896; (b) M. Cacciarini, A. B. Skov, M. Jevric, A. S. Hansen, J. Elm, H. G. Kjaergaard, K. V. Mikkelsen and M. B. Nielsen, *Chem. Eur. J.*, 2015, **21**, 7454; (c) A. U. Petersen, M. Jevric, R. J. Mandle, E. J. Davis, S. J. Cowling, J. W. Goodby and M. B. Nielsen, *RSC Adv.*, 2015, **5**, 89731.
- 4 (a) M. Å. Petersen, S. L. Broman, A. Kadziola, K. Kilså and M. B. Nielsen, *Eur. J. Org. Chem.*, 2009, 2733; (b) V. Mazzanti, M. Cacciarini, S. L. Broman, C. R. Parker, M. Schau-Magnussen, A. D. Bond and M. B. Nielsen, *Beilstein J. Org. Chem.*, 2012, **8**, 958; (c) M. Cacciarini, S. L. Broman and M. B. Nielsen, *Arkivoc*, 2014, **i**, 249.
- 5 (a) S. L. Broman, M. Å. Petersen, C. G. Tortzen, A. Kadziola, K. Kilså and M. B. Nielsen, *J. Am. Chem. Soc.*, 2010, **132**, 9165; (b) M. Å. Petersen, S. L. Broman, K. Kilså, A. Kadziola and M. B. Nielsen, *Eur. J. Org. Chem.*, 2011, 1033; (c) M. Jevric, S. L. Broman and M. B. Nielsen, *J. Org. Chem.*, 2013, **78**, 4348; (d) S. L. Broman, M. Jevric, A. D. Bond and M. B. Nielsen, *J. Org. Chem.*, 2014, **79**, 41.
- 6 S. L. Broman, M. Jevric and M. B. Nielsen, *Chem. Eur. J.*, 2013, **19**, 9542.
- 7 L. Skov, M. Å. Petersen, S. L. Broman, A. D. Bond and M. B. Nielsen, *Org. Biomol. Chem.*, 2011, **9**, 6498.
- 8 S. L. Broman, A. U. Petersen, C. G. Tortzen, J. Vibenholt, A. D. Bond and M. B. Nielsen, *Org. Lett.*, 2012, **14**, 318.
- 9 (a) S. Itoh, N. Morita, R. Miyatake and S. Kuroda, *Chem. Lett.*, 1997, 1011; (b) M. Oda, A. Sakamoto, T. Uchiyama, T. Kajioka, R. Miyatake and S. Kuroda, *Tetrahedron Lett.*, 1999, **40**, 3595; (c) M. Oda, H. Kainuma, T. Uchiyama, R. Miyatake and S. Kuroda, *Tetrahedron*, 2003, **59**, 2831; (d) M. Oda, N. Nakajima, N. C. Thanh, T. Kajioka and S. Kuroda, *Tetrahedron*, 2006, **62**, 8177; (e) M. Oda, N. Nakajima, N. C. Thanh, K. Kitahara, R. Miyatake and S. Kuroda, *Eur. J. Org. Chem.*, 2008, 5301.
- 10 (a) W. Abraham, K. Buck, S. Neubert and D. Kreysig, *Z. Chem.*, 1979, **19**, 97; (b) M. Balci, *Chim. Acta Turica*, 1983, **11**, 31.
- 11 T. Okazaki and K. K. Laali, *Org. Biomol. Chem.*, 2003, **1**, 3078.
- 12 T. Mrozek, H. Görner and J. Daub, *Chem. Eur. J.*, 2001, **7**, 1028.
- 13 For a review on the cycloheptatriene-norcaradiene conversion, see: O. C. McNamara and A. R. Maguire, *Tetrahedron*, 2011, **67**, 9.
- 14 (a) T. Mukai, H. Kubota and T. Toda, *Tetrahedron Lett.*, 1967, **37**, 3581; (b) S. W. Staley, M. A. Fox and A. Cairncross, *J. Am. Chem. Soc.*, 1977, **99**, 4524.
- 15 M. Tamm, T. Bannenber, K. Baum, R. Frolich, T. Steiner, T. Meyer-Friederichsen and J. Heck, *Eur. J. Inorg. Chem.*, 2000, **6**, 1161.
- 16 S. L. Broman, O. Kushnir, M. Rosenberg, A. Kadziola, J. Daub and M. B. Nielsen, *Eur. J. Org. Chem.*, 2015, 4119.
- 17 S. L. Broman, S. L. Brand, C. R. Parker, M. Å. Petersen, C. G. Tortzen, A. Kadziola, K. Kilså and M. B. Nielsen, *Arkivoc*, 2011, **ix**, 51.