

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.


Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

RSC Author Templates - ChemDraw (CHM) - Graphical Abstracts

All text and images must be placed within the frame.



71x62mm (300 x 300 DPI)

COMMUNICATION

DDQ-mediated synthesis of functionalized unsymmetrical disulfanes

Cite this: DOI: 10.1039/x0xx00000x

Mateusz Musiejuk, Tomasz Klucznik, Janusz Rachon and Dariusz Witt*

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

We developed a simple and efficient method for the synthesis of functionalized unsymmetrical disulfanes under mild conditions in good yields. The designed method is based on the reaction of bis(5,5-dimethyl-2-thioxo-1,3,2-dioxaphosphorinan-2-yl)disulfane with thiols in the presence of 2,3-dichloro-5,6-dicyanobenzoquinone (DDQ). The developed method allows the preparation of unsymmetrical disulfanes bearing additional hydroxy, carboxy, or amino functionalities.

Compounds with the structure R-S-S-R, where the R group can be alkyl, vinyl or aryl are termed symmetrical disulfides when the R groups are the same. The wide range of unsymmetrical disulfides in which the R groups are different is also well known. These compounds are often termed organic disulfides in the literature; however, the IUPAC recommended nomenclature is disulfanes.¹ The name disulfide should only be applied to ionic compounds, such as sodium disulfide (Na₂S₂). Moreover, the term disulfane is more widely applicable than disulfide because it facilitates naming, even when the R groups are acyl and/or phosphoryl.

The synthesis of unsymmetrical disulfanes is an important transformation in organic synthesis and medicinal chemistry.²⁻⁵ Recent developments in disulfide bond formation have been reviewed.⁶⁻⁹ Although many different methods exist for the preparation of unsymmetrical disulfanes, the most prevalent approach involves substitution of a sulfenyl derivative with a thiol or its derivative. To date, the most commonly utilized sulfenyl derivatives are the following: sulfenyl chlorides,¹⁰⁻¹² S-alkyl thiosulfates and S-aryl thiosulfates (Bunte salts),¹³⁻¹⁴ S-alkylsulfanyl-isothioureas,¹⁵ benzothiazol-2-yl disulfanes,¹⁶⁻¹⁷ benzotriazolyl-sulfanes,¹⁸ dithioperoxyesters,¹⁹ (alkylsulfanyl)dialkylsulfonium salts,²⁰ 2-pyridyl disulfanes and derivatives,²¹⁻²² N-alkyltetrazolyl disulfanes,²³ sulfenamides,²⁴ sulfenyldimesylamines,²⁵ sulfenylthiocyanates,²⁶ 4-nitroarenesulfenylidides,²⁷ thiosulfonates and thiosulfonates,²⁸⁻³¹ sulfanyl-sulfonamides,³² thionitrites,³³ sulfenylthiocarbonates,³⁴

thioimides,³⁵⁻³⁷ and thiophosphonium salts.³⁸ Other practical procedures involve the reaction of a thiol with a sulfinylbenzimidazole,³⁹ the rhodium-catalyzed disulfide exchange,⁴⁰⁻⁴¹ an electrochemical method,⁴² ring opening of aziridines using tetrathiomolybdate in the presence of symmetrical disulfanes,⁴³⁻⁴⁴ and the use of diethyl azodicarboxylate (DEAD)⁴⁵ or a solid support⁴⁶ in a sequential coupling of two different thiol groups. Recently, the oxidation of a mixture of two different thiols by 2,3-dichloro-5,6-dicyanobenzoquinone (DDQ) to produce an unsymmetrical disulfane has also been reported.⁴⁷⁻⁴⁸

Disulfanes have been used for the preparation of self-assembled monolayers (SAMs)⁴⁹⁻⁵⁰ and monolayer-protected clusters (MPCs) with a number of versatile properties.⁵¹⁻⁵² Compounds containing the disulfide linkage have also been used for the preparation of dynamic combinatorial libraries,⁵³ catenanes,⁵⁴⁻⁵⁵ macrocycles,^{56,5} carceplexes,⁵⁷ dendrimers,⁵⁸ rotaxanes, micelles,⁵⁹⁻⁶⁰ and a wide range of chemosensors and pro-drugs.⁶¹ These species illustrate the wide applications of disulfanes and show that the synthesis of the disulfide bond is a critical transformation in organic chemistry.⁶⁻⁹

We have previously demonstrated the preparation of functionalized unsymmetrical molecules, such as dialkyl disulfanes,⁶² alkyl-aryl disulfanes,⁶³ 'bioresistant' disulfanes,⁶⁴ the unsymmetrical disulfanes of L-cysteine and L-cystine,⁶⁵ and diaryl disulfanes,⁶⁶ based on the readily available 5,5-dimethyl-2-thioxo-1,3,2-dioxaphosphorinane-2-disulfanyl derivatives **1**. These disulfanyl derivatives **1** of phosphorodithioic acid were also convenient for the preparation of α -sulfenylated carbonyl compounds,⁶⁷ functionalized phosphorothioates,⁶⁸ as well as symmetrical⁶⁹⁻⁷⁰ and unsymmetrical⁷¹⁻⁷² trisulfanes (Figure 1).

As part of our continued interest in the preparation of functionalized unsymmetrical disulfanes, in this study, we report an efficient and convenient synthesis of unsymmetrical disulfanes **1** directly from phosphorodithioic acid disulfane **2a** and functionalized thiols **3**.

Wang and co-workers have developed a new method for the synthesis of unsymmetrical disulfanes from simple aliphatic and aromatic thiols





Figure 1 Synthetic applications of unsymmetrical disulfanes 1.

using DDQ as the oxidant.⁴⁷ This method is particularly interesting due to its apparent selectivity for the exclusive formation of unsymmetrical disulfanes, despite the presence of two different thiols in the reaction mixture in a 1:1 ratio before the addition of DDQ. We expected that this method might also be applicable to the synthesis of unsymmetrical disulfanes **1**. As a test reaction, 5,5-dimethyl-2-sulfanyl-2-thioxo-1,3,2-dioxaphosphorinane and dodecane-1-thiol **3a** (1:1 ratio) were mixed and then treated with DDQ (0.5 equivalent) following Wang's procedure.⁴⁷ Unfortunately, the yield of unsymmetrical disulfane **1a** after separation was moderate, and the observed ratio of products (**1a**:**2a**:**4**) was typical for the oxidation of most thiol mixtures⁸ (Figure 2).

Figure 2 Oxidation of phosphorodithioic acid and thiol **3a** mixture by DDQ.

Although Wang and co-workers did not discuss the mechanism of the formation of the unsymmetrical disulfanes, it can be speculated that DDQ converts the more easily oxidized thiol to the symmetrical disulfane and the second thiol to the alkylthio radical (thiyl radical RS^{\cdot}) which then reacts with symmetrical disulfane to produce the unsymmetrical product. To verify this hypothesis, we performed a reaction of bis-(5,5-dimethyl-2-thioxo-1,3,2-dioxaphosphorinane-2-yl) disulfane **2a** (variable amount) and dodecane-1-thiol **3a** (1 equivalent) in the presence of DDQ (0.5 equivalent) (Table 1).

Table 1 Reaction of disulfane **2a** with thiol **3a** and DDQ^a

Entry	2a (equiv.)	Yield ^b (%)	
		CH ₂ Cl ₂	CH ₃ CN
1	6 ^c	100	100
2	2	100	99
3	1.5	100	98
4	1	99	96
5	0.6	81	78
6	0.5	72	69

^a Conditions: disulfane **2a** (0.5-6.0 mmol), thiol **3a** (1 mmol), DDQ (0.5 mmol), 4.0 mL of solvent. ^b Isolated yields. ^c 10 mL of solvent was used.

As the data in Table 1 demonstrate, the excess symmetrical disulfane **2a** improved the yield of unsymmetrical product **1a**. The reactions using 1 equivalent of disulfane **2a** in CH₂Cl₂ and CH₃CN produced unsymmetrical product **1a** in a 99% and 96% yield, respectively (entry 4). Further studies were performed with 1 equivalent of disulfane **2a** because the yield of product **1a** was high when the excess amount of **2a** was acceptable.

Under the optimized conditions, the scope and limitations of this new unsymmetrical disulfane formation reaction were investigated, and the results are summarized in Table 2.

Table 2 Synthesis of unsymmetrical disulfanes **1** from functionalized thiols **3**^a.

Entry	RSH	Product	Yield ^b (%)	
			CH ₂ Cl ₂	CH ₃ CN
1	HS-(CH ₂) ₁₁ CH ₃ 3a	1a	99	96
2	HS-(CH ₂) ₁₁ OH 3b	1b	88	89
3	HS-(CH ₂) ₁₀ CO ₂ H 3c	1c	77	82
4	HS-(CH ₂) ₁₁ N ₃ 3d	1d	93	96
5	HS-(CH ₂) ₁₀ CONHS 3e	1e	88	89
6	HS-(CH ₂) ₁₀ CO-Ferrocene 3f	1f	82	85
7	HS-Ph 3g	1g	99	98
8	HS-C ₆ H ₄ -4-CH ₃ 3h	1h	98	97
9	HS-C ₆ H ₄ -2-CO ₂ H 3i	1i	96	95
10	HS-(CH ₂) ₉ -CH=CH ₂ 3j	1j	87	92
11	Ac-Cys-OH 3k	1k	-	89
12	HS-(CH ₂) ₁₁ NH ₂ ·HCl 3l	1l	0	0


^a Conditions: disulfane **2a** (1 mmol), thiol **3** (1 mmol), DDQ (0.5 mmol), 4.0 mL of solvent. ^b Isolated yields

A broad range of thiols reacted smoothly under the optimized reaction conditions. The reaction was tolerant of various functional groups, including the hydroxy, carboxy, azide, ferrocene, active ester

and carbon-carbon double bond groups. Aromatic thiols (**3g-i**) underwent disulfide bond formation to furnish the desired products (**1g-i**) with excellent yields (95-99%, entries 7-9, Table 2). Steric hindrance does not appear to affect the progress of the reaction (entry 9, Table 2). The aliphatic thiols also provided unsymmetrical disulfanes **1** in high yields (77-99%, entries 1-6, Table 2). Note that the aliphatic thiol **3j** underwent the reaction in the presence of a carbon-carbon double bond (entry 10, Table 2), and the hydroxy and carboxy groups did not require protection (entries 2-3, Table 2). In the case of *N*-acetyl-L-cysteine **3k**, the reaction did not occur in CH₂Cl₂ due to the low solubility of the starting material in the solvent. However, using acetonitrile as the solvent allowed the preparation of the corresponding unsymmetrical disulfane **1k** in high yield (89%, entry 11, Table 2).

Unfortunately, an unprotected amino group was not tolerated under the developed conditions. Although thiol **3l** was consumed during the reaction, the unsymmetrical disulfane was not produced (entry 12, Table 2). DDQ most likely reacted with an amino group to produce a complex reaction mixture without the formation of unsymmetrical disulfane **1**. Similar unsuccessful results have also been observed by Wang and co-workers in the case of amine-substituted thiols.⁴⁷ To overcome that limitation, we decided to use protected amino-thiols under the developed conditions (Table 3).

Table 3. Reaction of disulfane **2a** with amino-thiol and DDQ^a



Entry	RSH	Product	Yield ^b (%)	
			CH ₂ Cl ₂	CH ₃ CN
1	HS-(CH ₂) ₁₁ NH ₂ ·HCl 3l	1l	0	0
2	HS-(CH ₂) ₁₁ NHBoc 3m	1m	53	56
3	HS-(CH ₂) ₁₁ NBoc ₂ 3n	1n	94	96


^a Conditions: disulfane **2a** (1 mmol), thiol **3** (1 mmol), DDQ (0.5 mmol), 4.0 mL of solvent. ^b Isolated yields.

Our data show that the protection of the amino group plays a vital role in the yield of the unsymmetrical disulfanes **1m** and **1n**. When the amino group is protected by a single Boc group, the corresponding unsymmetrical disulfane **1m** is produced in moderate yield (53-56%, entry 2, Table 3). However, using thiol **3n** with a double-protected amino group provided unsymmetrical product **1n** in high yields (94-96%, entry 3, Table 2). The appropriate protection of the amino group appears to facilitate the preparation of unsymmetrical disulfanes **1m** and **1n** under the developed reaction conditions.

We were curious to continue our investigation with other symmetrical disulfanes, such as 2,2'-dipyridyl disulfane **2b** and diphenyldisulfane **2c** (Table 4).

As demonstrated, the formation of unsymmetrical disulfanes **1** strongly depends on the structure of the starting material **2**. In the case of disulfane **2a** and **2b**, the unsymmetrical product **1** was obtained in 99% and 23% yield, respectively. However, the reaction of diphenyldisulfane **2c** with dodecane-1-thiol **3a** in the presence of DDQ did not produce an unsymmetrical product, and exclusive formation of

Table 4 Reaction of symmetrical disulfane **2** with thiol **3a** and DDQ^a



Entry	R	Yield ^b (%)	
		1	4
1		99	0
2		23	70
3	Ph- 2c	0	99

^a Conditions: disulfane **2** (1 mmol), thiol **3a** (1 mmol), DDQ (0.5 mmol), 4.0 mL of CH₂Cl₂. ^b Isolated yields.

didodecyl disulfane **4** was observed. The determination of the scope of disulfanes **2** capable of producing unsymmetrical disulfanes under the developed conditions is under investigation.

The suggested mechanism to explain this transformation involves the initial formation of the thiyl radical (RS[•]) through the oxidation of thiol RSH **3** by DDQ. The resulting thiyl radical reacts with the symmetrical disulfane **2a** to yield the unsymmetrical product **1** and the phosphorodithioic acid radical, which undergoes recombination to produce symmetrical disulfane **2a** (Figure 3).




Figure 3 Plausible reaction mechanism The phosphorodithioic acid radical cannot extract hydrogen from the starting thiol because in this case, the catalytic amount of DDQ would be sufficient to produce unsymmetrical disulfane **1** in high yield. When a smaller amount of DDQ was used, a lower yield of product was obtained. In addition, we did not observe the formation of phosphorothioic acid in the reaction mixture. Moreover, the suggested mechanism explains why the excess of disulfane **2** improved the yield of the reaction. The initially formed thiyl radical (RS[•]) can undergo either recombination or

reaction with product **1** to produce symmetrical disulfane RSSR. These side reactions can be avoided when an excess of disulfane **2** is present in the reaction mixture. The reactivity of symmetrical disulfane **2** with the thiyl radical appears to play a vital role in the formation of unsymmetrical disulfane **1**. Most likely, 2,2'-dipyridyl disulfane **2b** and diphenyldisulfane **2c** are less reactive toward thiyl radicals than the produced unsymmetrical disulfanes **1**, which resulted in the formation of symmetrical disulfane **4** (entries 2-3, Table 4).

Conclusions

In summary, we developed an efficient and convenient method for the preparation of unsymmetrical disulfanes **1** directly from the disulfane of phosphorodithioic acid **2a** and functionalized thiols **3** in the presence of DDQ. A wide range of functional groups is tolerated, including the hydroxy, carboxy, azido, ferrocene, protected amino, and carbon-carbon double bond groups. Reactions of **2a** with a variety of functionalized thiols **3** in the presence of DDQ in CH₂Cl₂ or CH₃CN at 0°C were generally complete within 5 minutes and gave unsymmetrical disulfanes **1** exclusively in good or very good yield after isolation. The simplicity and good yields render this method one of the most attractive approaches to the preparation of functionalized unsymmetrical disulfanes, especially derivatives of phosphorodithioic acid with versatile synthetic applications.

General procedure for the preparation of disulfanyl derivatives **1** and representative analytical data

A thiol **3** (1.0 mmol) and bis-(5,5-dimethyl-2-thioxo-1,3,2-dioxaphosphorinan-2-yl) disulfane **2a** 394 mg (1.0 mmol) were dissolved in solvent (2.0 mL, dichloromethane or acetonitrile) and cooled to 0 °C in the ice bath. Then a solution of DDQ 114 mg (0.5 mmol) in solvent (2.0 mL, dichloromethane or acetonitrile) was added slowly to the reaction mixture and stirred for 5 min at 0 °C. The reaction was monitored by TLC analysis. Solvent was removed under reduced pressure and the residue was directly purified by column chromatography (SiO₂) to afford the desired products (**1a–1n**).

Acknowledgments

We gratefully acknowledge the National Science Centre (NCN) for financial support (grant no. 2013/09/B/ST5/01261)

Notes and references

^aDepartment of Organic Chemistry, Chemical Faculty, Gdansk University of Technology, Narutowicza 11/12, 80-233 Gdansk, Poland

[†]Electronic Supplementary Information (ESI) available: [experimental details and spectroscopic data for all new compounds **1**]. See DOI: 10.1039/c000000x/

1. R. Steudel, *Chem. Rev.*, 2002, **102**, 3905.
2. R. Cremlyn and J. An, *Introduction to Organosulfur Chemistry*; Wiley: New York, 1996.
3. S. Oae, *Organic Sulfur Chemistry: Structure and Mechanism*; CRC Press: Boca Raton FL, 1991.
4. V. M. Vrudhula, J. F. MacMaster, L. Zhengong, D. E. Kerr and P. D. Senter, *Bioorg. Med. Chem. Lett.*, 2002, **12**, 359.

5. Y. Mu, M. Nodwell, J. L. Pace, J. P. Shaw and J. K. Judice, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 735.
6. I. Shcherbakova and A. F. Pozharskii, In *Comprehensive Organic Functional Group Transformations II*; A. R. Katritzky, R. Taylor and Ch. Ramsden, Eds.; Pergamon: Oxford, 2004; vol. 2, pp. 177–187.
7. R. Sato and T. Kimura, In *Science of Synthesis*; N. Kambe, J. Drabowicz and G. A. Molander, Eds.; Thieme: Stuttgart-New York, 2007; vol. 39, pp. 573–588.
8. D. Witt, *Synthesis*, **2008**, 2491.
9. B. Mandal and B. Basu, *RSC Adv.*, 2014, **4**, 13854.
10. T. Endo, H. Tasai and T. Ishigami, *Chem. Lett.* **1975**, 813.
11. D. N. Harpp, B. T. Friedlander, C. Larsen, K. Steliou and A. Stockton, *J. Org. Chem.* 1978, **43**, 3481.
12. C. Brown and G. R. Evans, *Tetrahedron Lett.*, 1996, **37**, 9101.
13. J. M. Swan, *Nature* 1957, **180**, 643.
14. P. Hiver, A. Dicko and D. Paquer, *Tetrahedron Lett.*, 1994, **35**, 9569.
15. K. Sirakawa, O. Aki, T. Tsujikawa and T. Tsuda, *Chem. Pharm. Bull.*, 1970, **18**, 235.
16. A. L. Ternay, C. Cook and E. Brzezinska, *Phosphorus, Sulfur Silicon Relat. Elem.*, 1994, **95**, 351.
17. A. L. Ternay and E. Brzezinska, *J. Org. Chem.*, 1994, **59**, 8239.
18. R. Hunter, M. Cairra and N. Stellenboom, *J. Org. Chem.*, 2006, **71**, 8268.
19. C. Lriverend and P. Metzner, *Synthesis* **1994**, 761.
20. P. Dubs and R. Stuessi, *Helv. Chim. Acta* 1976, **59**, 1307.
21. D. H. R. Barton, C. Chen and M. G. Wall, *Tetrahedron* 1991, **47**, 6127.
22. D. H. R. Barton, A. C. O'Sullivan and M. M. Pechet, *J. Org. Chem.*, 1991, **56**, 6697.
23. M. Ohtani and N. Narisada, *J. Org. Chem.*, 1991, **56**, 5475.
24. M. Bao and M. Shimizu, *Tetrahedron* 2003, **59**, 9655.
25. A. Blaschette and M. Naveke, *Chem.-Zig.* 1991, **115**, 61.
26. R. G. Hiskey and B. F. Jr. Ward, *J. Org. Chem.*, 1970, **35**, 1118.
27. L. Benati, P. C. Montecocchi and P. Spagnolo, *Tetrahedron Lett.*, 1986, **27**, 1739.
28. R. Cragg, J. P. N. Husband and A. F. Weston, *J. Chem. Soc., Chem. Commun.*, **1970**, 1701.
29. D. A. Armitage, M. J. Clark and C. C. Tsao, *J. Chem. Soc., Perkin Trans. 1* **1972**, 680.
30. G. Capozzi, A. Capperucci, A. Degl'Innocenti, R. DelDuce and S. Menichetti, *Tetrahedron Lett.*, 1989, **30**, 2995.
31. A. Rajca and M. Wiessler, *Tetrahedron Lett.*, 1990, **31**, 6075.
32. I. V. Koval, *Russ. J. Org. Chem. (Engl. Transl.)* 2002, **38**, 232.
33. S. Oae, Y. H. Kim, D. Fukushima and K. Shinham, *J. Chem. Soc., Perkin Trans. 1* **1978**, 913.
34. S. J. Brois, J. F. Pilot and H. W. Barnum, *J. Am. Chem. Soc.*, 1970, **92**, 7629.
35. K. S. Boustang and A. B. Sullivan, *Tetrahedron Lett.*, 1970, **11**, 3547.
36. D. H. Harpp, D. K. Ash, T. G. Beck, J. G. Gleason, B. A. Orwig, W. F. VanHorn and J. P. Snyder, *Tetrahedron Lett.*, 1970, **11**, 3551.
37. J. Klose, C. B. Reese and Q. Song, *Tetrahedron* 1997, **53**, 14411.
38. M. Masui, Y. Mizuki, K. Sakai, C. Ueda and H. Ohmori, *J. Chem. Soc., Chem. Commun.*, **1984**, 843.
39. D. R. Graber, R. A. Morge and J. C. Sih, *J. Org. Chem.*, 1987, **52**, 4620.
40. M. Arisawa and M. Yamaguchi, *J. Am. Chem. Soc.*, 2003, **125**, 6624.
41. K. Tanaka and K. Ajiki, *Tetrahedron Lett.*, 2004, **45**, 5677.

42. Q. T. Do, D. Elothmani, G. Le Guillanton and J. Simonet, *Tetrahedron Lett.*, 1997, **38**, 3383.
43. D. Sureshkumar, V. Ganesh, R. S. Vidyarini and S. Chandrasekaran, *J. Org. Chem.*, 2009, **74**, 7958.
44. D. Sureshkumar, S. M. Koutha and S. Chandrasekaran, *J. Am. Chem. Soc.*, 2005, **127**, 12760.
45. T. Mukaiyama and K. Takahashi, *Tetrahedron Lett.*, 1968, **9**, 5907.
46. A. K. Galawde and A. F. Spatola, *Org. Lett.*, 2003, **5**, 3431.
47. J. K. Vandavasi, W.-P. Hu, Ch.-Y. Chen and J.-J. Wang, *Tetrahedron* 2011, **67**, 8895.
48. R. Smith, X. Zeng, H. Müller-Bunz and X. Zhu, *Tetrahedron Lett.*, 2013, **54**, 5348.
49. A. Ulman, *Chem. Rev.*, 1996, **96**, 1533.
50. D. Witt, R. Klajn, P. Barski and B. A. Grzybowski, *Curr. Org. Chem.*, 2004, **8**, 1763.
51. L. A. Jr. Porter, D. Ji, S. L. Westcott, M. Graupe, R. S. Czernuszewicz, N. J. Halas and T. R. Lee, *Langmuir*, 1998, **14**, 7378.
52. Y. S. Shon, C. Mazzitelli and R. W. Murray, *Langmuir* 2001, **17**, 7735.
53. K. R. West, K. D. Bake and S. Otto, *Org. Lett.*, 2005, **7**, 2615.
54. W. Wang, L. Q. Wang, B. J. Palmer, G. J. Exarhos and A. D. Q. Li, *J. Am. Chem. Soc.*, 2006, **128**, 11150.
55. L. Raehm, Ch. Hamann, J. M. Kern and J. P. Sauvage, *Org. Lett.*, 2000, **2**, 1991.
56. S. W. Tam-Chang, J. S. Stehouwer and J. Hao, *J. Org. Chem.*, 1999, **64**, 334.
57. Ch. Naumann, S. Place and J. C. Sherman, *J. Am. Chem. Soc.*, 2002, **124**, 16.
58. A. P. Umali and E. E. Simanek, *Org. Lett.*, 2003, **5**, 1245.
59. Y. Furusho, T. Oku, T. Hasegawa, A. Tsuboi, N. Kihara and T. Takata, *Chem. Eur. J.*, 2003, **9**, 2895.
60. S. Ghosh, K. Irvin and S. Thayumanavan, *Langmuir* 2007, **23**, 7916.
60. M. H. Lee, Z. Yang, Ch. W. Lim, Y. H. Lee, S. Dongbang, Ch. Kang and J. S. Kim, *Chem. Rev.*, 2013, **113**, 5071.
62. S. Lach, S. Demkowicz and D. Witt, *Tetrahedron Lett.*, 2013, **54**, 7021.
63. S. Antoniow and D. Witt, *Synthesis* **2007**, 363.
64. J. Kowalczyk, P. Barski, D. Witt and B. A. Grzybowski, *Langmuir* 2007, **23**, 2318.
65. M. Szymelfejnik, S. Demkowicz, J. Rachon and D. Witt, *Synthesis* **2007**, 3528.
66. S. Demkowicz, J. Rachon and D. Witt, *Synthesis* **2008**, 2033.
67. E. Okragla, S. Demkowicz, J. Rachon and D. Witt, *Synthesis* **2009**, 1720.
68. S. Lach and D. Witt, *Synthesis* **2011**, 3975.
69. A. Kertmen, S. Lach, J. Rachon and D. Witt, *Synthesis* **2009**, 1459.
70. S. Lach and D. Witt, *Heteroatom Chem.*, 2014, **25**, 10.
71. S. Lach, M. Sliwka-Kaszynska and D. Witt, *Synlett* **2010**, 2857.
72. S. Lach and D. Witt, *Synlett* 2013, **24**, 1927.