

RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Bis(imidazolium) salts derived from amino acids as receptors and transport agents for chloride anion

Cite this: DOI: 10.1039/x0xx00000x

Laura González-Mendoza,^a Belén Altava,^{*a} M. Isabel Burguete,^a Jorge Escorihuela,^b Elsa Hernando,^c Santiago V. Luis,^{*a} Roberto Quesada^c and Cristian Vicent^d

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/


The binding properties of bis(imidazolium) hosts **1a-c** derived from amino acids towards different anions have been studied by ¹H NMR titration experiments in 95:05 CD₃CN:H₂O at 303 K, ESI-MS and theoretical calculations. Among this family, the salt **1c** showed a strong and high selectivity for chloride anion. Transmembrane chloride transport activity has also been studied for the three bis(imidazolium) based transporters in POPC liposome models, and compound **1a** was identified as an active chloride/nitrate exchanger.

Introduction

Considerable attention continues to be shown in the design and synthesis of anion receptors possessing high affinity and selectivity.¹ In this regard, anionic species are of biological and environmental importance,² chloride being the most abundant anion in living organisms.³ The proper design, synthesis and study of abiotic hosts allows a minimalistic approach to the binding phenomena, largely simplifying the deep understanding of the corresponding processes.⁴ Recently, numerous efforts have been devoted to the study of imidazolium receptors, including benzene-based tripodal systems, cyclophane and caliximidazolium, fluorescent imidazolium, ferrocenyl imidazolium, cavitand and calixarene, and polymeric imidazolium systems.^{5,6} In contrast to the well-known N–H–anion or N⁺–H–anion hydrogen bonding, which is often found in amide, pyrrole, urea, ammonium or guanidinium groups, the imidazolium group can interact with anions through the involvement of (C–H)⁺–anion hydrogen bonds and, usually, the recognition is dominated by charge–charge electrostatic interactions.^{7,8}

Furthermore, the transport of chloride across cell membranes is important to numerous biological processes, and is often mediated by proteins that act as anion pores, anion exchangers or cation-dependent co-transporters.⁹ Malfunction of these proteins, through genetic mutation or otherwise, can lead to diseases such as cystic fibrosis.¹⁰ With this in mind, much focus is being placed on the development of small synthetic molecules that can mimic the role of such proteins.¹¹ Optimizing their potential to bind or encapsulate ions, such synthetic “carriers” could be developed for therapeutic applications.¹² In this context, the biocompatibility of the imidazole ring makes it suitable for biomimetic applications,¹³ and it has previously been incorporated in the structure of transmembrane carriers by Gale *et al.*¹⁴ and more recently by Schmitzer *et al.*,¹⁵ representing a new class of potential anion transporters.

In this regard, our research group has been involved in the preparation of imidazolium based receptors derived from enantiomerically pure materials obtained from the chiral pool.¹⁶ We have recently reported the synthesis of new ditopic dicationic receptors derived from amino acids with the general structure shown in Fig. 1, as receptors for the recognition of the chiral anions of dicarboxylic acids.¹⁷ As an extension of this work, here we report their use of as receptors for inorganic anions such as the spheric chloride, bromide and iodide anions and the trigonal NO₃[−] anion. The corresponding anion complexation properties have been investigated by ¹H NMR spectroscopy, ESI-MS and computational studies. Besides, these receptors have been studied as transmembrane transporters for chloride anion.



- 1a:** R = Bn; R¹ = Bn
1b: R = CH(CH₃)₂; R¹ = Bn
1c: R = CH(CH₃)₂; R¹ = (CH₂)₁₁CH₃

Fig. 1 General structure of receptors **1a-c**, prepared as their bis(trifluoromethylsulfonil)imide salts, X[−] = NTF₂[−].

Results and discussion

Chiral bis(imidazolium) compounds (**1a-c**) derived from different amino acids with variable N-amide substitution were prepared following the procedure described previously by our group.¹⁷

In preliminary recognition studies the interaction of the dicationic receptors **1a-c** with chloride was studied by ¹H NMR titrations with the corresponding tetrabutylammonium (TBA)

salt using CDCl_3 (5 % DMSO) as the solvent at 303 K. The corresponding binding constants could not be calculated due to the strong association with chloride in this medium. Thus, we studied the interaction between receptors **1a-c** and chloride using a more competitive solvent like aqueous CD_3CN . Fig. 2 shows the chemical shift variation of the different protons of receptor **1c** during titration experiments with TBACl in CD_3CN (5 % D_2O). Most of the signals of the receptor were affected by the presence of the chloride anion, although the overlapping of some of them precluded their accurate quantitative analysis. The signals more affected by the interaction were those corresponding to Ha, NH, Hb, He and Hd (see Fig. 1 for the identification of the corresponding hydrogen atoms). These variations suggested a strong electrostatic and H-bonding interaction between **1c** and the anion. The same trends in the chemical shifts variations were obtained with receptors **1a** and **1b** (see ESI†). It is interesting to note that the signal corresponding to the proton He of the central aromatic spacer is also significantly shifted to lower fields in the presence of Cl^- along with the signals corresponding to the hydrogen atoms of the imidazolium subunit (including Hb and Hc, although at a lower extent). Interestingly, the diastereotopic methylene Hf,f' hydrogen atoms become less equivalent as chloride is added, which is more evident when the less polar solvent (CDCl_3) is used (see ESI†).¹⁷




Fig. 2 Partial ^1H NMR spectra (500 MHz, 5 % aqueous CD_3CN at 303 K) of **1c** (5mM) in the presence of different amounts of TBACl.

The stoichiometry of the resulting host-guest complex was then investigated by ^1H NMR using the Job plot method.¹⁸ The Job-plot for the complexation of **1c** with the chloride anion in CD_3CN (5 % D_2O) exhibit a maximum at $X_L = 0.5$ for the different proton signals of compound **1c** analysed, which indicates that the 1:1 complex is predominant in solution (see ESI†).

The data obtained from the titration experiments using compound **1c** are shown in Fig. 3, displaying the corresponding isotherms for the binding of chloride anion. As can be observed, the most affected proton signal is the NH for which $\Delta\delta_{\text{max}} > 1$ ppm.

The simultaneous fitting of the variation of the chemical shifts of the five ^1H NMR proton signals of the different receptors rendered the corresponding binding constants which were determined fitting the data to a 1:1 and 1:2 receptor:anion binding isotherm using the HypNMR program (Table 1).¹⁹ The receptor **1c** showed a strong interaction with chloride even in

this competitive medium, and formed a stable 1:1 complex ($K_a 10^{3.2}$) with an additional weaker interaction with a second chloride anion ($K_a 10^{1.4}$). Interestingly, the binding constant for **1c** is eight times higher than for **1b** for the 1:1 complex (Table 1). We also determined the BC_{50}^0 parameter for a suitable comparison between ligands.²⁰ This parameter is defined here as the concentration of anion needed to bind 50 % of the dication and, thus, the lower the BC_{50}^0 value the stronger the supramolecular host-guest interaction. As expected, this parameter also shows with clarity the same trends detected in the values of the association constants, with the BC_{50}^0 value for chloride being significantly lower for **1c** (0.6 mM), the valine derivative containing additional long aliphatic tails. When comparing the role of the amino acid residue (**1a** against **1b**), the results suggest that this structural parameter does not have a strong influence on the anion recognition (Table 1, entries 1, 2), while the nature of R^1 seems to have a bigger effect (Table 1, entries 2, 3). Most likely, according to modeling studies on the receptors and complexes, R^1 substituents play an important role in defining conformational preferences, in particular the nature of folding, which could explain this trend.




Fig. 3 ^1H NMR (500 MHz, 5 % aqueous CD_3CN at 303 K) titration plot of **1c** (5mM) versus the equiv. of TBACl added. Symbols represent the experimental chemical shifts; lines correspond to the calculated ones for the best fitting.

Table 1. Binding constants for the interaction of **1a-c** with chloride, as its TBA salt, determined by ^1H NMR in 5 % D_2O in CD_3CN at 500 MHz and 303 K.

Entry	Receptor ^[a]	$\log \beta^{[b]}$ (receptor/anion)	K_a	BC_{50}^0 [mM]
1	1a	2.16 ± 0.02 (1:1) ^[c]	145 (1:1)	(6.7 ± 0.3)
		2.61 ± 0.09 (1:2) ^[d]	3 (1:2)	
2	1b	2.28 ± 0.02 (1:1) ^[c]	191 (1:1)	(5.0 ± 0.2)
		3.0 ± 0.1 (1:2) ^[d]	5 (1:2)	
3	1c	3.20 ± 0.08 (1:1) ^[c]	1585 (1:1)	(0.6 ± 0.1)
		4.6 ± 0.1 (1:2) ^[d]	25 (1:2)	

[a] As the bis(trifluoromethylsulfonyl)imide salt. [b] Obtained by the simultaneous non-linear regression fitting of all the available ^1H NMR signals using the HypNMR program. [c] β (1:1) is defined as the formation constant for the 1:1 receptor/substrate complex in M^{-1} . [d] In this case β (1:2) is the cumulative formation constant for the corresponding 1:2 receptor/substrate complex in M^{-2} .

As a function of these initial results, the interaction of the receptor **1c** with other spherical anions, such as bromide and iodide, and the trigonal (NO_3^-), was also studied. The results obtained showed that the corresponding binding constants were more than one order of magnitude smaller than that for chloride

(Table 2), indicating the weaker interaction of these anions with the bis(imidazolium) receptor. According to these results, the interaction for the different anions is $\text{Cl}^- \gg \text{Br}^- > \text{I}^- \sim \text{NO}_3^-$. As observed, in the case of the halogen anions, the interaction becomes smaller as the anion radius increases.

Table 2. Binding constants for the interaction of **1c** with different anions, as their TBA salts, determined by ^1H NMR in 5% D_2O in CD_3CN at 500 MHz and 303 K.

Entry	Anion	$\log \beta^{[a]}$ (receptor/anion)	K_a	BC_{50}^0 [mM]
1	Cl^-	3.20 ± 0.08 (1:1) ^[b]	1585 (1:1)	(0.6 ± 0.1)
		4.6 ± 0.1 (1:2) ^[c]	25 (1:2)	
2	Br^-	2.20 ± 0.02 (1:1) ^[b]	159 (1:1)	(5.9 ± 0.3)
		2.96 ± 0.08 (1:2) ^[c]	6 (1:2)	
3	I^-	1.92 ± 0.03 (1:1) ^[b]	83 (1:1)	(11.2 ± 0.2)
		2.5 ± 0.1 (1:2) ^[c]	4 (1:2)	
4	NO_3^-	1.95 ± 0.04 (1:1) ^[b]	89 (1:1)	(10.3 ± 0.8)
		2.6 ± 0.1 (1:2) ^[c]	5 (1:2)	

[a] Obtained by the simultaneous non-linear regression fitting of all the available ^1H NMR signals using the HypNMR program. [b] β (1:1) is defined as the formation constant for the 1:1 receptor/substrate complex in M^{-1} . [c] In this case β (1:2) is the cumulative formation constant for the corresponding 1:2 receptor/substrate complex in M^{-2} .

ESI-MS represents a versatile tool to evaluate both qualitatively and quantitatively binding properties in molecular recognition processes.²¹ In this work we carried out an ESI-MS study aimed to estimate the distinctive binding affinities of compound **1c** towards different anions. A qualitative agreement between NMR titration experiments and ESI-MS data were found only when MeCN was used as the mobile phase using equimolar solutions (1×10^{-5} M) of $[\mathbf{1c}](\text{NTf}_2)_2$ and $(\text{TEA})\text{Cl}$. We are aware that comparison of adducts with Cl^- and NTf_2^- certainly represents two extreme situations in terms of ion size and hydrophilic character; however, we envisage that for ion adducts displaying similar size, shape and properties, such as those comprising the $[\mathbf{1c} \text{X}]^+$ adducts where X stands for Cl^- , Br^- , I^- and NO_3^- anions, competitive ESI experiments might provide a good match between ion abundances in the ESI mass spectrum and solution concentration. To validate this hypothesis, we performed a competitive ESI mass analysis on an equimolar solution of $[\mathbf{1c}](\text{NTf}_2)_2$ and TEAX ($\text{X} = \text{Cl}^-$, Br^- , I^- and NO_3^-) in MeCN (see Fig. 4).




Fig. 4 Competitive ESI mass spectrum of an equimolar mixture of $[\mathbf{1c}](\text{NTf}_2)_2$ and TEAX ($\text{X} = \text{Cl}^-$, Br^- , I^- and NO_3^-) in MeCN. The inset shows the expanded region in the m/z 750–1150 range illustrating the ion abundances of the species $[\mathbf{1c} \text{X}]^+$.

On the basis of the relative intensities from the ESI mass spectrum, and normalizing the ion abundances of $[\mathbf{1c} \text{Cl}]^+$ to 100%, the corresponding adducts follow the trend $[\mathbf{1c} \text{Cl}]^+(100) > [\mathbf{1c} \text{Br}]^+(37) > [\mathbf{1c} \text{NO}_3]^+(20) > [\mathbf{1c} \text{I}]^+(16)$ in qualitative agreement with the trend observed by NMR titration methods. We also detected a minor peak at m/z 974 that formally correspond to the

$[\mathbf{1cCl}_2+\text{TEA}]^+$ cation (marked in blue in Fig. 4). After diluting $\times 10$ and $\times 100$ the sample solutions, this species was still present, thus suggesting that adducts $[\mathbf{1c}:\text{Cl}_2]$ with a 1:2 stoichiometry are present in the solution rather than being formed due to non-specific interactions during the ESI process.²² It must be noted that the use of ESI-MS to estimate IL:anion interaction strengths based on i) the propensity to aggregate into $[\text{IL}_x\text{Anion}_{(x-1)}]^+$ in single stage ESI mass spectra or ii) CID of mixed $[\text{IL}_1\text{IL}_2\text{A}]^+$ adducts (the same applies for anions), is well-documented.^{23, 24} However, in our particular case, single stage ESI mass spectra of **1c** displayed poor aggregation. For example, for the ESI-MS of **1c** in MeCN, the $[\mathbf{1c}:\text{anion}_3]^+$ or the $[\mathbf{1c}\text{anion}_3]^+$ ions were barely detected, and accordingly the methodology based on i) and ii) criterions could not be satisfactorily applied.

Theoretical studies

According with the predominant 1:1 host–guest complex observed in solution by ^1H NMR and ESI-MS, theoretical calculations were carried out at the B3LYP/6-31G(d,p) level to predict the structure of the positively charged receptor–anion complexes when considering the interaction with only one chloride anion. Since the supramolecular complexes can display certain flexibility in solution, we subjected the corresponding host–guest structures to Monte Carlo conformational searches without restrictions followed by MMFF minimizations, in order to fully map all the possible conformations and anion–cation relative dispositions. To this aim, 10000 structures were stochastically generated and minimized with the MMFF molecular force field using the Monte Carlo conformational search available in Spartan^{14, 25}. Then, the lowest 100 minima were analysed and ordered attending to their relative energies. The resulting minimum energy conformer was optimized at the B3LYP/6-31+G*level of theory using the Gaussian 09 software,²⁶ being the corresponding global minima shown in Fig. 5 for a 1:1 model complex and in Fig. 6 for the corresponding dichloride salts of **1a-c**. For 1:1 complexes, the dicationic bis(imidazolium) salt with methyl substituents ($\text{R} = \text{R}' = \text{CH}_3$) was used as a model compound to simplify the high level calculations. The global minima optimized structure is illustrated in Fig. 5 and shows the ability of this receptor to form a very stable complex displaying multiple H-bonding interactions. In this 1:1 structure the receptor can efficiently fold over the chloride anion in such a way that the anion is able to form strong hydrogen bonds with the acidic hydrogen atoms of the receptor and be shielded from the influence of the solvent. In this case, the chloride anion is hydrogen bonded, simultaneously, to the hydrogen atoms of both amide NH groups and the two Ha atoms of the imidazolium rings. The anion is also at short distances from both Hd atoms and from He. The calculated $\text{Cl}\cdots\text{H}$ distances are in very good agreement with the trends observed for $\Delta\delta$ values in ^1H NMR upon addition of TBACl. For Hd the observed shift in the corresponding signals is similar to that observed for Ha in NMR experiments (ca. 0.6 ppm) although the calculated distance to the anion is larger, but in this case it must be taken into consideration that the effect of the interaction of chloride with Ha also affects to the shifts of the other two imidazolium hydrogen atoms Hb and Hd ($\Delta\delta$ values ≈ 0.17 ppm, similar to those for He).

We carried out also simple molecular modelling calculations for the 1:2 complexes and in this case we considered the full set of structural parameters to analyse their effects and to rationalize the experimental data for chloride binding.




Fig. 5 Global minima optimized structure for the supramolecular 1:1 complex formed between dicationic bis(imidazolium) salt with methyl substituents ($R = R' = \text{CH}_3$) and Cl^- .




Fig. 6 Global minima optimized structures for the supramolecular complexes formed between bis(imidazolium) cations **1a-c** and two Cl^- anions. For clarity, we only show the polar hydrogen atoms implicated in H-bonding interactions (shown as red dashed lines).

The minima display different conformations with a different connectivity of H-bonds implicating the chloride anions. For dication **1a**, the receptor adopts a cleft conformation in which the N-benzyl groups are located closing the gate of the cleft. This geometry is stabilized by a number of electrostatic H-bonds between the chloride anions and the acidic hydrogens of the receptor. In this regard, each chloride anion is H-bonded to the hydrogen of one amide NH group and to one imidazolium Ha, with two eight membered chelated rings, being formed at each pseudopeptidic chain. However, for **1b** and **1c** the receptor adopts a cleft conformation in which the N-substituents do not close the cleft gate but are folded backwards. In this case, the intramolecular H-bonds involve each chloride anion with the amide hydrogen of one of the chains and the imidazolium Ha of the other chain. The observed conformations correlate well with the ^1H NMR shifts observed during the titration experiments. The calculated H-bond distances are shown in Table 3. Again, although the NH—anion distances are the shortest ones, very short distances are observed from the chloride to other hydrogen atoms, besides the imidazolium Ha, like Hd, He and Hf.

Table 3. Cl-H distances (Å) for the optimized structures^[a]

Entry	Receptor	NH-Cl	Ha-Cl	Hd-Cl	He-Cl	Hf-Cl
1	1a	2.207	2.962	2.554	3.291	2.683
2	1b	2.151	2.950	2.595	2.624	2.695
3	1c	2.206	2.930	2.569	2.831	2.718

[a] Calculated for the bis(imidazolium) cations **1a-c** and two Cl^- anions.

Chloride transport studies

The complexation properties of these receptors prompted us to study them as transport agents across lipid bilayer membranes, in particular for a biologically very relevant species like chloride. Since transmembrane transport experiments were carried out in the presence of nitrate, ^1H NMR anion-binding studies were also conducted for receptors **1b-c** with nitrate anions (Table 4) in order to provide the appropriate basic data for chloride transport studies. The corresponding stability constants, K_a , were determined again by following the shift of the ^1H NMR signals of five protons from the receptor and fitting the data to 1:1 and 1:2 receptor:anion binding isotherms using HypNMR. As is shown by the data gathered in Table 4, the receptors interact weakly with nitrate anions, displaying stability constants lower than 113 M^{-1} . For receptors **1b** and **1c** a good selectivity for chloride is observed as reflected in the BC_{50}^0 values (5.0 vs. 16.5 for **1b** and 0.6 vs. 10.3 for **1c**), while the selectivity is smaller for the receptor **1a** (6.7 vs. 8.9 for BC_{50}^0 values).

Table 4. Binding constants for the interaction with NO_3^- as its TBA salt determined by ^1H NMR in 5% D_2O in CD_3CN at 500 MHz and 303 K.

Entry	Receptor	$\log \beta^{[a]}$ (receptor/anion)	K_a	BC_{50}^0 [mM]
1	1a	2.05 ± 0.02 (1:1) ^[b]	113 (1:1)	(8.9 ± 0.4)
		2.65 ± 0.06 (1:2) ^[c]	4 (1:2)	
2	1b	1.78 ± 0.03 (1:1) ^[b]	61 (1:1)	(16.5 ± 0.9)
		2.2 ± 0.1 (1:2) ^[c]	2 (1:2)	
3	1c	1.95 ± 0.04 (1:1) ^[b]	89 (1:1)	(10.3 ± 0.8)
		2.6 ± 0.1 (1:2) ^[c]	5 (1:2)	

[a] Obtained by simultaneous non-linear regression fitting of all the available ^1H NMR signals. [b] β (1:1) is defined as the formation constant for the 1:1 receptor/substrate complex in M^{-1} . [c] In this case β (1:2) is the cumulative formation constant for the corresponding 1:2 receptor/substrate complex in M^{-2} .

The transmembrane anion transport properties of receptors **1a**, **1b** and **1c** were explored using vesicle based methods.²⁷ Unilamellar POPC vesicles encapsulating 488 mM NaCl buffered to pH 7.2 with 5 mM sodium phosphate salts were suspended in an isotonic NaNO_3 solution. The studied compounds were added as DMSO solutions using a minimal amount of solvent to avoid disturbance of the liposomes. The chloride efflux from the interior of the vesicles was monitored using a chloride ion selective electrode (ISE). After a fixed time (300 s) detergent was added to lyse the vesicles and induce the efflux of all encapsulated chloride. The final electrode reading was used to calibrate the 100 % chloride release. The chloride efflux induced by compounds **1a-c** (15 % molar carrier to phospholipid loading) is shown in Fig. 7. Under these conditions compound **1a** showed a good activity whereas **1b** and **1c** induced a more modest chloride efflux. By carrying out

a Hill analysis of the chloride efflux induced by **1a** at different concentrations, an EC_{50} value of 10.5 % molar **1a**:POPC was calculated (see Figs. S35-S36).²⁸ EC_{50} is defined as the concentration needed for releasing 50 % of encapsulated chloride under the conditions described above for this assay. The Hill parameter was found at 3.8 suggesting cooperativity in the transport activity exerted by **1a**. It can be seen that the receptor with a lower chloride affinity presents the higher transport activity. This is not surprising as transmembrane ion transport is an extremely complicated process involving many equilibria and several steps and thus, the affinity for the guest is not the single parameter determining the transport efficiency.²⁹ In this regard, for instance, differences in lipophilicity (**1c** presents high lipophilicity) can be as important as those in binding constants. The transport activity is defined by a delicate balance between different parameters for which binding constants in different solvent media and lipophilicity are just two of the involved parameters.




Fig. 7 Chloride efflux promoted by compounds **1a-c** (75 μ M, 15 % mol carrier to lipid concentration) in unilamellar POPC vesicles. Vesicles loaded with 489 mM NaCl buffered at pH 7.2 with 5 mM phosphate dispersed in 489 mM $NaNO_3$ buffered at pH 7.2. Each trace represents the average of at least different three trials.

In order to investigate the impact of the external anion in the chloride release mediated by the receptors, another set of experiments using POPC vesicles containing 451 mM NaCl buffered to pH 7.2 with 20 mM sodium phosphate salts suspended in 150 mM Na_2SO_4 buffered to pH 7.2 was performed. The results are shown in Fig. 8. Under these conditions chloride efflux promoted by **1a** was greatly reduced whereas the transmembrane transport activity observed for **1b** and **1c** was essentially maintained. These results could indicate that anion exchange mechanism is dominant in the case of receptor **1a**. Thus chloride efflux is more efficient in the presence of nitrate, a relatively lipophilic anion, whereas the high hydrophilicity of the SO_4^{2-} anion makes its transport much more difficult reducing the overall transport activity.

Further experiments to investigate the transmembrane transport activity of these receptors were performed using fluorescence based assays. We decided to use the 5(6)-carboxyfluorescein leakage test.³⁰ Carboxyfluorescein (CF) is self-quenching at high concentration. Thus, if CF is encapsulated at high concentrations in the interior of the liposomes, little fluorescence is detected. On the other hand,

leakage of entrapped CF into the much larger volume of the cuvette is readily signalled by the characteristic highly fluorescence of the diluted CF solutions. CF is a membrane impermeable dye and thus CF release from the interior of the vesicle is indicative of disruption of the membrane integrity or formation of membrane pores large enough to allow the leaking of this anion. The CF leakage test was performed on vesicles containing CF and NaCl, dispersed in a Na_2SO_4 solution (Fig. 9). Illustrative results are shown in Figs. S40-S46. No indications of leakage were observed. After addition of compounds **1a-c** very little fluorescence was detected prior to the addition of detergent to the vesicle suspension. These results ruled out the possibility of an unspecific detergent action exerted by these derivatives.




Fig. 8 Chloride efflux promoted by compounds **1a-c** (75 μ M, 15 % mol carrier to lipid concentration) in unilamellar POPC vesicles. Vesicles loaded with 450 mM NaCl buffered at pH 7.2 with 20 mM phosphate dispersed in 150 mM Na_2SO_4 buffered at pH 7.2. Each trace represents the average of at least different three trials.




Fig. 9 CF leakage induced by **1a-c** (15 % mol carrier to lipid) from unilamellar POPC vesicles (0.05 mM) loaded with 50 mM CF and 451 mM NaCl, buffered to pH 7.2 with 20 mM sodium phosphate salts. The vesicles were suspended in 150 mM Na_2SO_4 buffered to pH 7.2 with 20 mM sodium phosphate salts. At $t = 60$ s, a DMSO solution of the transporter was added to start the experiment. At the end of the experiment ($t = 360$ s), detergent was added to lyse the vesicles and induce the release of all encapsulated CF.

Conclusions

The results here presented demonstrate that bis(imidazolium) salts can be selective, when properly designed, for the binding of chloride anion in solution as analysed by NMR spectroscopy and ESI-MS. ^1H NMR titration experiments reveal that compound **1c** shows binding constants more than two orders of magnitude larger for chloride than for any other halide or anion studied. The binding affinities of **1c** with different anions were also investigated by competitive ESI-MS experiments. Qualitative agreement between binding affinities estimated by ESI-MS and NMR data was found by considering the relative abundances of the 1:1 adducts $[\mathbf{1c X}]^+$ ($X = \text{Cl}^-, \text{Br}^-, \text{I}^-$ and NO_3^-). In order to establish a proper correlation between ESI-MS and NMR experiments, several experimental factors need to be considered: i) the selection of acetonitrile as the solvent is important to minimize the distinctive effect of the surface activity of the anions, following recent observations from McInnes's group on closely related IL's,^{22,23} ii) because of the large size of **1c**, the dominant effect manifested in the individual ion abundances for the observed $[\mathbf{1c X}]^+$ ($X = \text{Cl}^-, \text{Br}^-, \text{I}^-$ and NO_3^-) cations must be the binding energy rather than surface activity differences. Regarding the transmembrane chloride transport studies, compound **1a** posses a reasonable activity, favouring the transport via an exchange mechanism, while compounds **1b-c** display a low level of chloride transport activity. Importantly, these derivatives do not exert an unspecific detergent action in the lipid bilayer. Although the level of activity found for these compounds is yet far from optimal, the tunability of the design holds promise for developing improved transmembrane transporters based on ionic liquids derived from aminoacids.

Experimental

General: Reagents and solvents were purchased from commercial suppliers (Aldrich, Fluka or Merck) and were used without further purification.

NMR spectroscopy: The NMR experiments were carried out on a Varian INOVA 500 spectrometer (500MHz for ^1H and 125 MHz for ^{13}C). Chemical shifts are reported in ppm by using TMS as a reference. Titration experiments followed by ^1H NMR spectroscopy were performed in $\text{CD}_3\text{CN}/\text{D}_2\text{O}$ 95:5 at 303 K and data fitting was carried out with the HypNMR 2008 version 4.0.71 software.

Molecular modelling.

The supramolecular complexes between **1a-c** and the guests were generated manually and submitted to conformational analysis by Monte Carlo conformational searches as implemented in Spartan'14 software. To this aim, 10000 structures were stochastically generated and minimized with the MMFF molecular force field. Then, the lowest 100 minima were analysed and ordered attending to their relative energies. Different starting geometries and anion-cation dispositions were used in order to verify that the systems converged to the same final minima. The obtained minima were subsequently fully minimized at the B3LYP/6-31G(d,p) level of theory. The stationary points have been characterized as true minima by the calculation of the normal vibration modes, being all the values positive

Membrane transport assays.

Preparation of Phospholipid Vesicles: A chloroform solution (20 mg/mL) of 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC) (Sigma-Aldrich) was evaporated *in vacuo* using a rotary evaporator. The lipid film thus obtained was dried under high vacuum for at least 4 hours and then it was rehydrated by addition of a sodium chloride solution (489 mM NaCl and 5 mM phosphate buffer, pH 7.2) followed by vortexing. The lipid suspension was then subjected to nine freeze-thaw cycles and twenty-nine extrusions through a 200 nm polycarbonate Nucleopore membrane using a LiposoFast Basic extruder (Avestin, Inc.). The resulting unilamellar vesicles were dialyzed against a NaNO_3 solution (489 mM NaNO_3 and 5 mM phosphate buffer, pH 7.2) or Na_2SO_4 solution (150 mM Na_2SO_4 and 20 mM phosphate buffer, pH 7.2) to remove unencapsulated chloride.

Ion selective electrode (ISE) transport assays.

Chloride/Nitrate Transport Assays: Unilamellar POPC vesicles containing NaCl (prepared as described above) were suspended in a 489 mM NaNO_3 solution buffered to pH 7.2 with sodium phosphate salts (5 mM buffer). The final lipid concentration per sample was 0.5 mM and the total volume of 5 mL. A DMSO solution of the carrier molecule, typically 10-20 μL to avoid the influence of the solvent molecule in the assay, was added and the chloride release out of vesicles was monitored using a chloride selective electrode for 5 minutes. Then, the vesicles were lysed by adding 40 μL of Triton-X (10 % dispersion in water). The initial value was set at 0 % chloride efflux and the final one was set as 100 % chloride efflux. All other data were calibrated to these points. All traces represent an average of at least three independent experiments.

Ion selective electrode (ISE) transport assays.

Chloride/Sulfate Transport Assays: The experiments were conducted as described above using unilamellar POPC vesicles (100 nm mean diameter) containing an encapsulated solution of 451 mM NaCl and 20 mM phosphate buffer, pH 7.2, were suspended in a solution 150 mM Na_2SO_4 and 20 mM phosphate buffer, pH 7.2, for a final lipid concentration of 0.5 mM and a total volume of 5 mL. All traces represent an average of at least three independent experiments.

Fluorescence assays: 200 nm unilamellar vesicles encapsulating 5(6)-carboxyfluorescein (CF) dissolved in a sodium chloride solution (451 mM NaCl, 50 mM 5(6)-carboxyfluorescein (CF), 20 mM phosphate buffer at pH 7.2) were prepared as described above. Unencapsulated 5(6)-carboxyfluorescein was removed by size exclusion chromatography on a Sephadex G-50 column using a sodium sulfate solution as eluent (150 mM Na_2SO_4 , 20 mM phosphate buffer at pH 7.2). The vesicles were suspended in a solution of 150 mM Na_2SO_4 and 20 mM phosphate buffer, pH 7.2 for a final lipid concentration of 0.025 mM or 0.05 mM.

The fluorescence emission of the vesicle suspension (2.5 mL samples) at 520 nm after excitation at 490 nm was recorded for 8 minutes (a measure every 10 seconds under continuous stirring) using a HITACHI F-7000 Fluorescence spectrophotometer. At $t = 60$ s, a solution of the carrier molecule in DMSO (15 % molar carrier to lipid) was added. At the end of the experiment ($t = 360$ s) the vesicles were lysed by addition of detergent (20 μL of Triton-X (10 % dispersion in water)). The extent of the CF leakage from the interior of the

vesicles can be estimated from the variation of the fluorescence intensity.

$$\text{CF leakage} = \left[\frac{I_t - I_0}{I_{\text{max}} - I_0} \right]_{\lambda=520\text{nm}}$$

I_0 = fluorescence intensity at time 0

I_t = fluorescence intensity at time t

I_{max} = maximum fluorescence intensity observed after addition of detergent.

Acknowledgements

Financial support from Ministerio de Economía y Competitividad (CTQ2011-28903-C02-01), Generalitat Valenciana (PROMETEO 2012/020) and Universitat Jaume I (P11B2013-38), Consejería de Educación de la Junta de Castilla y León (Project BU340U13) and Fundació la Marató de TV3 is acknowledged.

Notes and references

^a Departamento de Química Inorgánica y Orgánica, Universitat Jaume I, Av. Sos Baynat, s/n, E-12071 Castellón (Spain).

Fax: (+34) 964728214, E-mail: luiss@uji.es, altava@uji.es

^b Wageningen UR, Laboratory of Organic Chemistry, Dreijenplein 8, 6703 HB Wageningen, The Netherlands.

^c Departamento de Química, Facultad de Ciencias, Universidad de Burgos, 09001 Burgos (Spain).

^d Serveis Centrals d'Instrumentació Científica, Universitat Jaume I, Avda. Sos Baynat, s/n, 12071 Castellón (Spain).

† Electronic Supplementary Information (ESI) available: ¹H NMR titration experiments and HypNMR fitting data, ESI-MS experiments and chloride transport studies and theoretical data. See DOI: 10.1039/b000000x/.

- Anion receptors: (a) J. L. Sessler, P. A. Gale, W.-S. Cho, Anion Receptor Chemistry; RSC: Cambridge, 2006; (b) F. P. Schmidtchen, M. Berger, *Chem. Rev.* 1997, **97**, 1609; (c) P. D. Beer, P. A. Gale, *Angew. Chem., Int. Ed.* 2001, **40**, 486; (d) R. Martínez-Manez, F. Sancenon, *Chem. Rev.* 2003, **103**, 4419; (e) P. A. Gale, *Coord. Chem. Rev.* 2003, **240**, 1; (f) P. A. Gale, R. Quesada, *Coord. Chem. Rev.* 2006, **250**, 3219; (g) P. A. Gale, S. E. García-Garrido, J. Garric, *Chem. Soc. Rev.* 2008, **37**, 151; (h) C. Caltagirone, P. A. Gale, *Chem. Soc. Rev.* 2009, **38**, 520; (i) M. Cametti, K. Rissanen, *Chem. Commun.* 2009, 2809. (j) N. H. Evans, P. D. Beer, *Angew. Chem. Int. Ed.* 2014, **53**, 11716.
- (a) J. L. Sessler, P. A. Gale, W. S. Cho, Anion Receptor Chemistry, RSC Publishing, Cambridge, 2006; (b) Supramolecular Chemistry of Anions (Eds.: A. Bianchi, K. Bowman-James, E. García-España), Wiley-VCH, Weinheim, 1997; (c) Special Issue: Anion Coordination Chemistry II (Eds.: A. B. P. Lever, P. A. Gale), *Coord. Chem. Rev.* 2006, **250**, 2917.

- (a) M. Wenzel, J. R. Hiscock, P. A. Gale, *Chem. Soc. Rev.* 2012, **41**, 480; (b) A. P. Davis, D. N. Sheppard, B. D. Smith, *Chem. Soc. Rev.* 2007, **36**, 348.
- (a) H. T. Chifotides, I. D. Giles, K. R. Dunbar, *J. Am. Chem. Soc.* 2013, **135**, 3039; (b) L. Trembleau, T. A. D. Smith, M. H. Abdelrahman, *Chem. Commun.* 2013, **49**, 5850; (c) A. Basu, G. Das, *Chem. Commun.* 2013, **49**, 3997; (d) E. García-España, P. Díaz, J. M. Llinares, A. Bianchi, *Coord. Chem. Rev.* 2006, **250**, 2952.
- (a) J. Yoon, S. K. Kim, N. J. Singh, K. S. Kim, *Chem. Soc. Rev.* 2006, **35**, 355. (b) Z. Xu, S. K. Kim, J. Yoon, *Chem. Soc. Rev.* 2010, **39**, 1457. (c) Q.-S. Lu, L. Dong, J. Zhang, J. Li, L. Jiang, Y. Huang, S. Qin, C.-W. Hu, X.-Q. Yu, *Org. Lett.* 2009, **11**, 669.
- (a) A. H. Flood, Y. Liu, J. F. Stoddart, In Modern Cyclophane Chemistry; Gleiter, R., Hopf, H., Eds.; Wiley-VCH: Weinheim, 2004; (b) Y. Yuan, G. Gao, Z.-L. Jiang, J.-S. You, Z.-Y. Zhou, D.-Q. Yuan, R.-G. Xie, *Tetrahedron* 2002, **58**, 8993; (c) J. Yoon, S. K. Kim, N. J. Singh, J. W. Lee, Y. J. Yang, K. Chellappan, K. S. Kim, *J. Org. Chem.* 2004, **69**, 581. (d) P. P. Neelakandan, M. Hariharan, D. Ramaiah, *Org. Lett.* 2005, **7**, 5765; (e) P. P. Neelakandan, M. Hariharan, D. Ramaiah, *J. Am. Chem. Soc.* 2006, **128**, 11334; (f) P. P. Neelakandan, D. Ramaiah, *Angew. Chem., Int. Ed.* 2008, **47**, 8407. (g) V. K. Khatri, S. Upreti, P. S. Pandey, *Org. Lett.* 2006, **8**, 1755; (h) V. K. Khatri, M. Chahar, K. Pavani, P. S. Pandey, *J. Org. Chem.* 2007, **72**, 10224.
- (a) P. A. Gale, P. Anzenbacher, J. L. Sessler, *Coord. Chem. Rev.* 2002, **222**, 57; (b) M. D. Best, S. L. Tobey, E. V. Anslyn, *Coord. Chem. Rev.* 2003, **240**, 3; (c) J. M. Llinares, D. Powell, K. Bowman-James, *Coord. Chem. Rev.* 2003, **240**, 57; (d) C. R. Bondy, S. J. Loeb, *Coord. Chem. Rev.* 2003, **240**, 77; (e) P. Anzenbacher, R. Nishiyabu, M. A. Palacios, *Coord. Chem. Rev.* 2006, **250**, 2929; (f) S. O. Kang, M. A. Hossain, K. Bowman-James, *Coord. Chem. Rev.* 2006, **250**, 3038; (g) C. Schmuck, *Coord. Chem. Rev.* 2006, **250**, 3053.
- H. Ihm, S. Yun, H. G. Kim, J. K. Kim, K. S. Kim, *Org. Lett.* 2002, **4**, 2897.
- (a) B. Hille, Ionic Channels of Excitable Membranes, Sinauer Associates, Sunderland, 3rd edn, 2001; (b) J. T. Davis, O. Okunola and R. Quesada, *Chem. Soc. Rev.*, 2010, **39**, 3843.
- (a) F. M. Ashcroft, *Nature*, 2006, **440**, 440. F. M. Ashcroft; (b) Ion Channels and Disease; Academic Press: San Diego, 2000.
- (a) S. Hussain, P. R. Brotherhood, L. W. Judd and A. P. Davis, *J. Am. Chem. Soc.*, 2011, **133**, 1614; (b) S. J. Moore, M. Wenzel, M. E. Light, R. Morley, S. J. Bradberry, P. Gomez-Iglesias, V. Soto-Cerrato, R. Perez-Tomas and P. A. Gale, *Chem. Sci.*, 2012, **3**, 2501; (c) S. Rastogi, E. Marchal, I. Uddin, B. Groves, J. Colpitts, S.A. McFarland, J. T. Davis and A. Thompson, *Org. Biomol. Chem.*, 2013, **11**, 3834; (d) Louise E. Karagiannidis, Cally J. E. Haynes, Katie J. Holder, Isabelle L. Kirby, Stephen J. Moore, Neil J. Wells, Philip A. Gale, *Chem. Commun.*, 2014, **50**, 12050.
- (a) P. A. Gale, R. Pérez-Tomás, R. Quesada, *Acc. Chem. Res.*, 2013, **56**, 2801; (b) I. Alfonso and R. Quesada, *Chem. Sci.*, 2013, **4**, 3009; (c) N. Busschaert and P. A. Gale, *Angew. Chem., Int. Ed.*, 2013, **52**, 1374. (d) S.-K. Ko, S. K. Kim, A. Share, V. M. Lynch, J. Park, W. Namkung, W. Van Rossom, N. Busschaert, P. A. Gale, J. L. Sessler and I. Shin, *Nat. Chem.*, 2014, **6**, 885.
- E. B. Anderson, T. E. Long, *Polymer*, 2010, **51**, 2447.

- 14 P. A. Gale, J. Garric, M. E. Light, B. A. McNally, B. D. Smith, *Chem. Commun.*, 2007, 1736.
- 15 (a) C.-R. Elie, N. Noujeim, C. Pardin, A. R. Schmitzer, *Chem. Commun.* 2011, **47**, 1788; (b) C.-R. Elie, M. Charbonneau and A. R. Schmitzer, *MedChemComm*, 2012, **3**, 1231; (c) M. Vidal, A. Schmitzer, *Chem. Eur. J.*, 2014, **20**, 9998.
- 16 (a) L. González, J. Escorihuela, B. Altava, M. I. Burguete, S. V. Luis, *Eur. J. Org. Chem.* **2014**, 5356; (b) L. González, B. Altava, M. Bolte, M. I. Burguete, E. García-Verdugo, S. V. Luis, *Eur. J. Org. Chem.* **2012**, 4996; (c) B. Altava, D. S. Barbosa, M. I. Isabel Burguete, J. Escorihuela, S. V. Luis, *Tetrahedron: Asymmetry*, 2009, **2**, 999.
- 17 L. González-Mendoza, J. Escorihuela, B. Altava, M. I. Burguete, S. V. Luis, *Org. Biomol. Chem.*, **2015**, DOI: 10.1039/C5OB00348B.
- 18 (a) V. M. S. Gil, N. C. Oliveira, *J. Chem. Ed.*, 1990, **67**, 473; (b) P. Job, *Ann. Chim. Fr.*, 1928, **9**, 113.
- 19 (a) C. Frassinetti, L. Alderighi, P. Gans, A. Sabatini, A. Vacca, S. Ghelli, *Anal. Bioanal. Chem.* 2003, **376**, 1041; (b) C. Frassinetti, S. Ghelli, P. Gans, A. Sabatini, M. S. Moruzzi, A. Vacca, *Anal. Biochem.* 1995, **231**, 374.
- 20 (a) A. Vacca, O. Francesconi, S. Roelens, *Chem. Rec.* 2012, **12**, 544; (b) C. Nativi, O. Francesconi, G. Gabrielli, A. Vacca, S. Roelens, *Chem. Eur. J.* 2011, **17**, 4814; (c) S. Roelens, A. Vacca, C. Venturi, *Chem. Eur. J.* 2009, **15**, 2635; (d) S. Roelens, A. Vacca, O. Francesconi, C. Venturi, *Chem. Eur. J.* 2009, **15**, 8296.
- 21 R. Bini, O. Bortolini, C. Chiappe, D. Pieraccini, C. Siciliano, *J. Phys. B.* 2007, **111**, 598.
- 22 Z. Fei, D.-R. Zhu, Yan, N., R. Scopelliti, S. A. Katsuba, G. Laurencyzy, D. M. Chisholm, J. S. McIndoe, K. R. Seddon, P. J. Dyson, *Chem. Eur. J.* 2014, **20**, 4273.
- 23 J. Pape, K. L. Vikse, E. Janusson, N. Taylor, J. S. McIndoe, *Int. J. Mass Spectrom.* 2014, **373**, 66.
- 24 V. Gabelica, N. Galic, E. De Pauw, *J. Am. Soc. Mass Spectrom.* 2002, **13**, 946.
- 25 Spartan '14, version 1.1.0 (Sept 27 2013), Wavefunction Inc.: Irvine CA, 2014.
- 26 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, GAUSSIAN 09 (Revision D.1), Gaussian, Inc., Wallingford CT, 2009.
- 27 (a) B. D. Smith and T. N. Lambert, *Chem. Commun.*, 2003, 2261; (b) J. T. Davis, P. A. Gale, O. A. Okunola, P. Prados, J. C. Iglesias-Sanchez, T. Torroba and R. Quesada, *Nat. Chem.*, 2009, **1**, 138.
- 28 (a) A. V. Hill, *Biochemistry*, 1913, **7**, 471; (b) S. Bhosale and S. Matile, *Chirality*, 2006, **18**, 849.
- 29 E. Hernando, V. Soto-Cerrato, S. Cortés-Arroyo, R. Pérez-Tomás and R. Quesada, *Org. Biomol. Chem.*, 2014, **12**, 1771.
- 30 E. E. Ambroggio, F. Separovic, J. H. Bowie, G. D. Fidelio and L. A. Bagatolli, *Biophys. J.*, 2005, **89**, 1874.