

RSC Advances




This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.





Journal Name

COMMUNICATION

Modulation of the stability of *i*-motif structures using an acyclic threoninol cytidine derivative†

Sonia Pérez-Rentero,^a Raimundo Gargallo,^b Carlos González^c and Ramon Eritja^{*a}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

The effect of α TNA (acyclic threoninol nucleic acids) units on the stability of intramolecular *i*-motifs was investigated by spectroscopic techniques. The replacement of selected positions of the C-core can modulate the stability at different pH ranges.

The *i*-motif is a secondary structure found in C-rich DNA sequences, which is stabilised at acidic. The structure comprises two strands arranged in a parallel-type duplex with hemiprotonated CCH⁺ base pairs, and two of these duplexes are held together in an antiparallel fashion by intercalation of each CCH⁺ base pairs (Fig. 1a).¹

From the biological point of view, the presence of *i*-motif-forming sequences in telomeric and promoter regions supports the hypothesis that they may be functionally relevant.² Recently, the role of these structures as modulators of gene expression has been assessed.³ Additionally, these structures are interesting in the field of nanotechnology, since for instance they have been used to control the assembly of supramolecular structures⁴ or in the design of molecular motors⁵ and sensing systems such as nanoswitches driven by pH changes to monitor biological processes.⁶ Desirable properties regarding this last application points towards the necessity of nuclease resistant *i*-motifs and the availability of different pH-sensors suited to different pH regimes.^{6a} The stability of *i*-motif structures is highly dependent of the base sequence.⁷ It has been reported that simple manipulations of the C-core⁸ or the loops⁹ can modulate the pH sensitivity and also the introduction of chemical modifications.² In particular, sugar modifications of the cytidine core⁴ have pointed the relevance of the sugar-sugar contacts in the stability of *i*-motifs.¹⁰

Threoninol is an acyclic scaffold that has been used to design new artificial nucleic acids (α TNA), capable of forming exceptionally stable homo-duplexes.¹¹




Fig. 1 a) Schematic representation of a potential folding topology of the intermolecular *i*-motif studied in this work. b) Chemical structure of C_{LT}.

Moreover, α TNA's can form stable hetero-duplexes with DNA and RNA in a parallel manner,¹² and also they are compatible with parallel G-quadruplex structures.¹³ Recently, this acyclic scaffold has been used to design functional RNA/ α TNA chimeras with improved nuclease resistance.¹⁴ To date the effect of this modification in the formation of *i*-motifs has not been tested. In sight of these results, we are interested in study the effect of this sugar replacement on the thermal and pH stability.

The present work is based on a well described C-rich fragment of the vertebrate telomere (C₃TA₂)₃C₃T, which folds intramolecularly, where four cytidine stretches connected by three TAA loops form six CCH⁺ base pairs.¹⁵ In this work the loop bases were substituted by thymines and then, different positions of the C-core stretches of the 24 nt fragment (C₃T₃)₄ (Lt₀) were systematically replaced by C_{LT} units (Fig 1b), (Table 1). For detailed information see ESI[†] (sections 6-8). Lt₀ was modified by a single C_{LT} substitution at various positions of the 5'-terminal stretch (Lt₁, Lt₂ and Lt₃) and more internal positions (Lt₁₃, Lt₁₄ and Lt₁₅). Additionally, oligonucleotides containing two C_{LT} modifications involved in the same CCH⁺ base pair (Lt₁/13, Lt₂/14 and Lt₃/15) were studied. Finally, Lt₁/2/3 with three consecutive C modifications in the 5'-terminal stretch was included in the analysis. We assessed the thermodynamic stability of the modified 24 nt fragments by monitoring the absorbance at 295 nm as a function of temperature (Fig. S1, ESI[†]) and compared with the non-modified version (Table 1). For additional information see ESI[†] (section 2).

^a Institute for Advanced Chemistry of Catalonia (IQAC-CSIC), CIBER-BBN, Jordi Girona 18-26, E-08034 Barcelona, Spain. E-mail: recqma@cid.csic.es

^b Department of Analytical Chemistry, University of Barcelona, Martí i Franqués 1-11, E-08028 Barcelona, Spain

^c Institute of Physical Chemistry "Rocasolano", CSIC, Serrano 119, E-28006 Madrid, Spain.

† Electronic Supplementary Information (ESI) available: Experimental methods, synthesis and characterizations of C_{LT} monomers and oligonucleotides. See DOI: 10.1039/c000000x

Table 1. Thermal stabilities and thermodynamic parameters of Lt_0 and modified oligonucleotides at pH 5.3.^a

Name ^b	T_m	ΔT_m	ΔG_{298}^0	$\Delta\Delta G_{298}^0$	ΔH^0	$T\Delta S_{298}^0$
Lt_0	47.8	-----	-5.7	-----	-79.3	-73.7
Lt_1	39.5	-8.3	-2.9	2.8	-61.7	-58.8
Lt_2	29.1	-18.7	-0.6	5.1	-45.6	-45.0
Lt_3	38.4	-9.4	-2.6	3.1	-60.9	-58.4
Lt_13	37.4	-10.4	-2.7	3.0	-67.4	-64.8
Lt_14	29.3	-18.5	-0.8	4.9	-56.3	-55.5
Lt_15	37.9	-9.9	-2.7	3.0	-65.1	-62.4
Lt_1/13	35.4	-12.4	-2.3	3.4	-67.5	-65.2
Lt_2/14	32.4	-15.4	-1.6	4.1	-66.4	-64.8
Lt_3/15	34.7	-13.1	-2.1	3.6	-65.8	-63.7
Lt_1/2/3	35.6	-12.2	-2.4	3.3	-69.3	-67.0

^a T_m is expressed in °C and ΔG_{298}^0 , ΔH^0 and $T\Delta S_{298}^0$ in kcal/mol. Experimental errors are shown in Table S1 (see ESI[†]). ^bThe numbers indicate the positions substituted. $C_{\text{oligonucleotide}} = 2.7 \mu\text{M}$.

At pH 5.3 the UV melting analyses reveal that all the C_{LT} substitutions reduce significantly the T_m , and the effect is highly dependent of the position. Single C_{LT} substitutions near the loops are less destabilizing than those in the middle of C-tracts. Similar effect has been observed with other acyclic backbone modifications.^{10d} The effect of C_{LT} substitutions is not additive. When the two cytidines involved in a $C\text{CH}^+$ base pair were replaced, the T_m is slightly lower than in single substitutions. The effect is opposite when the $C\text{CH}^+$ is located in the center of the C-tracts (Lt_2/14). Finally, Lt_1/2/3 shows a higher thermal stability than some of the oligonucleotides containing one or two modifications. In all cases the values of Gibbs's free energies follow the same pattern observed with the T_m values as expected because depend on the equilibrium constant. Analysing the enthalpic term for oligonucleotides containing one or two C_{LT} substitutions near the loops, it can be observed that the ΔH^0 exhibit a similar decrease indicating that the destabilization is mostly of enthalpic origin. The influence of the enthalpic change is compensated by opposite change in the entropy term leaving the influence on Gibbs's energy rather modest. A single replacement in central positions of a cytidine core causes a higher drop of ΔH^0 , especially when is implicated the 5'-terminal stretch, suggesting that the $C\text{CH}^+$ base pair is affected and additionally the distortion may be affecting the neighbouring base pairing. Surprisingly, when both modifications occupy the central part of the core, it leads to a more stable structure than the ones obtained with Lt_2 and Lt_14. Taking into account that in this case the enthalpic term is more favourable, it can be reasoned that the pairing dC- C_{LT} is less tolerated than the $C_{\text{LT}}-C_{\text{LT}}$ pair. Finally, Lt_1/2/3 has the highest enthalpic term, but at the same time the most unfavourable entropic term.

Also we investigated the *i*-motif formation by CD spectroscopy. In Fig. 2 there are depicted a selection of some CD spectra. For detailed information see ESI[†] (Fig. S2 and Table S2). Lt_0 has the characteristic *i*-motif signature,¹⁶ with positive and negative peaks around 287 and 262 nm respectively and as well all modified sequences show this specific profile. Comparing with Lt_0, in

general the modifications produce blue shifts of the positive and/or negative bands and a decrease in the molar ellipticity to a greater or lesser extent. The most affected is the version containing three C_{LT} residues, followed by the oligonucleotides containing a substitution in an internal position of the 5'-terminal stretch. As CD spectroscopy is sensitive to structural changes,¹⁸ it can be reasoned that the deviations in the CD spectra from the natural sequence would be more pronounced as higher is the degree of distortion introduced in the *i*-motif structure.

Since structural changes may affect the pH stability, we decided to compare the pH dependence of two modified oligonucleotides with the unmodified version. We selected the one showing the most altered CD spectrum (Lt_1/2/3) and one showing moderate changes (Lt_3). The unfolding of the *i*-motif with pH at 25 °C was monitored at 295 nm. By plotting the unfolded fraction against pH and adjusting to a sigmoidal (see section 4 ESI[†]) we obtained the titration curves showed in Fig. 3. The pH transition midpoint ($\text{pH}_{1/2}$) and the transitional range (R_T , range between 10 and 90% unfolded structure) were calculated from the adjusted curves (Table 2). Comparing Lt_3 with Lt_0, the R_T has been slightly widened and also shifted toward acidic pH values, showing a $\text{pH}_{1/2}$ around pH 6.00. Lt_1/2/3 as well shows a decrease in the $\text{pH}_{1/2}$, but the R_T is almost 3-fold wider. The presence of three consecutive C_{LT} in the 5'-external stretch introduces higher flexibility.




Fig. 2 CD spectra of Lt_0 and modified oligonucleotides at pH 5.3.




Fig. 3 pH titration of Lt_0, Lt_3 and Lt_1/2/3.


Table 2. pH transition midpoint and transitional range for Lt_0, Lt_3 and Lt_1/2/3.

Name	pH _T	pH ₁₀	pH ₉₀	R _T
Lt_0	6.54±0.02	6.05±0.02	6.71±0.02	0.66±0.04
Lt_3	6.00±0.08	5.42±0.02	6.20±0.07	0.78±0.09
Lt_1/2/3	6.16±0.02	4.80±0.02	6.70±0.02	1.90±0.04

These effects may lead to the formation of several misfolded conformations provoking a decreased cooperativity.

Next, we performed 1D ¹H NMR experiments with Lt_0, Lt_3 and Lt_1/2/3 at different temperatures to obtain more detailed structural information (see Fig. S4, ESI[†]). Fig. 4 shows for each oligonucleotide the experiments recorded at 5 °C. In the three cases, the NMR spectra are consistent with *i*-motif formation. The characteristic imino signals of protonated cytosines are clearly observed at 15.0-16.0 ppm,^{7,15} as well as the thymine imino protons in the loop regions (around 11.0 ppm), in agreement with previous studies reported in the literature.¹⁵ The differences in the exchangeable protons spectra between the unmodified and Lt_3 and Lt_1/2/3 are significant, indicating the effect of the acyclic threoninol backbone propagates up to the base pairs, affecting the core of the *i*-motif structure. Such effect is particularly pronounced in the case of Lt_1/2/3. The unfolding of the structure with increasing temperature can be followed in Fig S4 in ESI[†]. In the unmodified oligonucleotide, the imino protons are observed up to 45 °C. In agreement with UV melting experiments, the exchangeable protons signals of Lt_3 and Lt_1/2/3 disappear at lower temperatures than the unmodified oligonucleotide.

In summary, we evaluated for the first time the effect of deoxy-sugar replacements by C_{LT} substitutions in the ability to form intramolecular *i*-motifs. Different positions of the C-core were replaced and the thermal and thermodynamic stability have been studied. In general the C_{LT} modifications reduced significantly the stability of the resulting structures and the effect is dependent of the position and the type of pair implicated, being the pairing dC-C_{LT} less favourable than the C_{LT}-C_{LT} pair. When a single substitution was introduced, the positions with a lesser decrease in stability are the positions near the loops.

**Fig. 4** NMR spectra at 5 °C and pH = 4.2 of a) Lt_0, b) Lt_3 and c) Lt_1/2/3.

In addition the pH stability of Lt_3 and Lt_1/2/3 was studied. In comparison with Lt_0, a single C_{LT} substitution produced a decrease of pH_T of about 0.54 units and maintained a comparable range of response. The substitution of three consecutive cytidines led to a less cooperative transition. Recently it has been described that the L-threoninol nucleic acids are more stable to nuclease degradation than unmodified oligonucleotides [14]. For this reason the possibility of modulating the pH stability of *i*-motifs by selected replacements of cytidines by C_{LT} modifications may allow the design of nuclease resistant *i*-motif structures suited for different pH ranges.

Acknowledgements

This work was supported by Spanish Ministry of Economy and Competitiveness (grants CTQ2014-52588-R and CTQ2012-38616-C02-02), the Generalitat de Catalunya (grant 2014 SGR 1106; 2014 SGR 187) and the Communities MULTIFUN (contract NMP4-LA-2011-262943). CIBER-BBN is an initiative funded by the VI National R&D&I Plan 2008–2011, Iniciativa Ingenio 2010, Consolider Program, CIBER Actions and financed by the Instituto de Salud Carlos III with assistance from the European Regional Development Fund.

Notes and references

- 1 K. Gehring, J. L. Leroy and M. Guéron, *Nature*, 1993, **363**, 561-565.
- 2 (a) S. Benabou, A. Aviñó, R. Eritja, C. González; and R. Gargallo, *RSC Adv.*, 2014, **4**, 26956-26980; (b) H. A. Day, P. Pavlou and Z. A. E. Waller, *Bioorg. Med. Chem.*, 2014, **22**, 4407-4418.
- 3 H.-J. Kang, S. Kendrick, S. M. Hecht and L. H. Hurley, *J. Am. Chem. Soc.*, 2014, **136**, 4172-4185.
- 4 (a) H. B. Goedke, R. Krishnan, K. Vignesh, G. Kumar, C. Narayana and Y. Krishnan, *Angew. Chem. Int. Ed.*, 2007, **11**, 2700-2703; (b) E. Guittet, D. Renciuik and J.-L. Leroy, *Nucleic Acids Res.*, 2012, **40**, 11, 5162-5170.
- 5 D. Liu and S. Balasubramanian, *Angew. Chem. Int. Ed.*, 2003, **42**, 5734-36.
- 6 (a) S. Modi, C. Nizak, S. Surana, S. Halder and Y. Krishnan, *Nat. Nanotechnol.*, 2013, **8**, 459-467; (b) S. Surana, J. M. Bhat, S. P. Koushika and Y. Krishnan, *Nat. Commun.*, 2011, **2**, 339-345.
- 7 J.-L. Mergny, L. Lacroix, X. Han, J.-L. Leroy and C. Hélène, *J. Am. Chem. Soc.*, 1995, **6**, 8887-8898.
- 8 I. V. Nesterova and E. E. Nesterov, *J. Am. Chem. Soc.*, 2014, **136**, 8843-8846.
- 9 S. P. Gurung, C. Schwarz, J. P. Hall, C. J. Cardin and J. A. Brazier, *Chem. Commun.*, 2015, **51**, 5630-5632.
- 10 (a) C. P. Fenna, V. J. Wilkinson, J. R. P. Arnold and R. Cosstick, *Chem. Commun.*, 2008, 3567-3569. (b) N. Kumar, I. Petersen and S. Maiti, *Chem. Commun.*, 2009, 1532-1534; (c) S. Modi, A. H. Wani, and Y. Krishnan, *Nucleic Acids Res.*, 2006, **34**, 16, 4354-4363; (d) A. Pasternak and J. Weng, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 752-755.
- 11 H. Asanuma, T. Toda, K. Murayama, X. Liang and H. Kashida, *J. Am. Chem. Soc.*, 2010, **132**, 14702-14703.
- 12 K. Murayama, H. Kashida and H. Asanuma, *Chem. Commun.*, 2015, **51**, 6500-6503.
- 13 J. Zhou, K. Murayama, S. Amrane, F. Rosu, H. Kashida, J. Bourdoncle, H. Asanuma, and J.-L. Mergny, *Chem. Sci.*, 2013, **4**, 3693-3698.

COMMUNICATION

Journal Name

- 14 (a) A. Alagia, M. Terrazas and R. Eritja, *Molecules*, 2014, **19**, 17872-17896. (b) A. Alagia, M. Terrazas and R. Eritja, *Molecules*, 2015, **20**, 7602-7619.
- 15 A. T. Phan, M. Guéron and J.-L. Leroy, *J. Mol. Biol.*, 2000, **299**, 123-144.
- 16 G. Manzini, N. Yathindra and L. E. Xodo, *Nucleic Acids Res.*, 1994, **22**, 22, 4634-4640.
- 17 J. Kypr, I. Kejnovská, D. Renčiuk and M. Vorlíčková, *Nucleic Acids Res.* 2009, **37**, 6, 1713-1725.

RSC Advances Accepted Manuscript