

# RSC Advances



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. This *Accepted Manuscript* will be replaced by the edited, formatted and paginated article as soon as this is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name

ARTICLE

## A mechanistic study of carbonyl activation under solvent-free conditions: Evidence drawn from the synthesis of imidazoles

Kiran Pradhan<sup>a</sup>, Bipranch Kumar Tiwary<sup>b</sup>, Mussaraff Hossain<sup>a</sup>, Ranadhir Chakraborty<sup>b</sup>,  
Ashis Kumar Nanda<sup>a\*</sup>

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Syntheses of various imidazole and its derivatives, imidazole N-oxides and 1-hydroxyimidazole 3-oxides from sterically different dicarbonyl moieties provided insights into the Self-Catalytic effect of condensed phase reactions of carbonyl compounds. The self-catalytic activity in solvent-free multi-component syntheses was investigated using a combination of methods viz., reactivity, spectroscopy and theory. While IR spectroscopic studies revealed that reacting molecules were polarised in bulk, quantum mechanical calculations of associated HCHO monomers suggest an increase in the average dipole moment of each monomer and provide evidence for the presence of cooperative effects. A comparative study of the kinetics of un-catalysed and catalysed reactions with the help of HPLC provided insights to the mechanism.

### 1 Introduction

The application of a solvent-free synthetic methodology for organic compounds that were conventionally synthesized in a solvent medium is becoming more common in recent years.<sup>1</sup> Thus, well-known organic reactions such as the aldol,<sup>2</sup> Claisen,<sup>3</sup> Stobbe<sup>4</sup> and Knoevenagel condensations,<sup>5</sup> the Thorpe,<sup>6</sup> Tischchenko,<sup>7</sup> Reformatsky and Luche reactions,<sup>8</sup> the Baeyer-Villiger oxidation,<sup>9</sup> the pinacol rearrangement<sup>10</sup> and the oxidative couplings of phenols,<sup>11</sup> just to name a few, have been found not only to occur under solvent-free conditions but are more efficient than solution reactions. In addition to simplicity and cleanness of the procedure, the absence of any media has been found to lead to uncommon reactivities. Apart from these efforts, over the past two decades organic synthesis has been devoted to extending the scope of solvent free reactions, notably for newer multi-component reaction (MCR) strategies.<sup>12</sup> Solvent-free protocols have already been developed for almost all the classical multi-component reactions namely the Strecker,<sup>13</sup> Hantzsch,<sup>14</sup> Biginelli,<sup>15</sup> Mannich,<sup>16</sup> Passerini,<sup>17</sup> Ugi,<sup>18</sup> Gewald,<sup>19</sup> Petasis,<sup>20</sup> Radziwinski synthesis,<sup>21</sup> and so on. And to impart greater reactivity to the substrates and the reagents, these synthetic protocols have constructively exploited the use of transition metal catalysts and the very recent organocatalysts.

Lewis acids and organocatalysts have been shown to be

excellent catalysts for C-C bond forming reactions, and through carbonyl activation provide effective avenues for upgrading classical MCRs. In particular, close attention has to be paid to condensation reaction mechanisms that involve the activation of carbonyl-containing electrophiles. The vast number of investigations related to chemical transformation mediated by organocatalysts<sup>22</sup> or by Lewis acids in anhydrous organic media<sup>23</sup>, aqueous media<sup>24</sup> and in the absence of any media<sup>25</sup> has provided a basic understanding of the intricate dynamics of adduct formation between the acid and the carbonyl moieties and has shed light on the role that these adducts have in accelerating catalytic transformations. A noteworthy example is the dramatic increase in reaction rates for the Radziwinski synthesis attained by the activation of a carbonyl group by a Lewis acid center<sup>25</sup> or by a Lewis base center<sup>26</sup> through the formation of adducts.

In the present study we have attempted to explore in some details the mechanistic insight for better reaction under solvent-free condition compared to reactions in solution. Here we have chosen a well-studied method for imidazole synthesis.

### 2 Experimental

The compounds 1a-1i, 2a-2l, 3a-3g, 4a-4g and 5a-5e were all prepared under solvent-free conditions. All other reagents are commercially available and were used as purchased from the supplier. Grinding experiments were performed with an agate mortar and pestle, which was acetone rinsed and dried prior to use. The yield of the reaction was determined by the product peak area count in HPLC analysis with respect to the purified product. NMR spectra were recorded on Bruker Avance 300 spectrometer. Chemical shifts ( $\delta$  in ppm) were referenced to external SiMe<sub>4</sub>. IR spectra were recorded on a FTIR – 8300 SHIMADZU spectrophotometer. Analytical thin-layer

<sup>a</sup> Department of Chemistry, University of North Bengal, Darjeeling, India, 734013.  
E-mail- ashis\_nanda@hotmail.com, Mobile no- +91-9474329191

<sup>b</sup> Department of Biotechnology, University of North Bengal, Darjeeling, India, 734013..

† Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

chromatography was performed using silica gel aluminium sheets (Merck, TLC silica gel 60 F254). HPLC was performed on a Waters – 2487 Dual Lambda absorber with a RP-18 (Symmetry Shield) column. The solvent used was methanol with a flow rate of 0.5 ml/min. In all experiments the same column and same flow rate was maintained. The electrospray mass spectra were recorded on a MICROMASS QUATTRO II triple quadrupole mass spectrometer. The ESI capillary was set at 3.5 kV and cone voltage was 40 V. X-Ray diffraction data was collected on a Bruker APEX CCD-II diffractometer at 293 K. The structure was solved by a direct method using the program SHELXS-97.27

#### Computational details

The optimized geometry and the energy of each monomer were calculated using the CBS-QB3 model chemistry. The CBS-QB3 model chemistry<sup>28</sup> was employed for the calculation of the molecular structures since this method is known to predict thermochemical parameters with high accuracy.<sup>29</sup> Molecular geometry was fully optimized and harmonic vibrational frequencies were calculated using the analytic second derivatives to confirm the convergence to minima on the potential surface. All the calculations were carried out using the Gaussian 98 suite of programs.<sup>30</sup>

#### 2.1 General procedure for the synthesis of 2, 4, 5-trisubstituted imidazoles (1a-1i)

210 mg (1 mmole) of benzil, 1 mmole of the corresponding aldehyde and 770 mg (10 mmole) of ammonium acetate was taken in an agate mortar and thoroughly ground. The contents were transferred to a test tube and heated to 150-160°C for 4 minutes. The contents were cooled and water was added to the test tube and filtered. The product was recrystallised from ethanol. Completion of the reaction was checked by TLC.

#### 2.2 General procedure for the synthesis of 1, 2, 4, 5-tetrasubstituted imidazoles (2a-2l)

210 mg (1 mmole) of benzil, 1 mmole of aldehyde, 1 mmole of primary amine and 385 mg (5 mmole) of ammonium acetate was taken in an agate mortar and thoroughly ground. The contents were transferred to a test tube and heated to 150-160°C for 4 minutes. The contents were cooled and water was added to the test tube and filtered. The product was recrystallised from ethanol. Completion of the reaction was checked by TLC.

#### 2.3 General procedure for the synthesis of imidazole N-oxide (3a-3g)

1 mmole of the monoxime, 1 mmole of the aldehyde and 385 mg (5 mmole) of ammonium acetate was ground into an intimate mixture in an agate mortar and pestle. The mixture was then heated to 115-120°C in an oil bath with constant shaking. A black solution resulted which was cooled when a black sticky precipitate formed. To the black precipitate was then added a small volume of diethyl ether when a brown precipitate separated. The precipitate was then thoroughly washed with ethyl acetate, dissolved in ethanol and crystallized by addition of water to yield pure products.

#### 2.4 General procedure for the synthesis of N-substituted imidazole-1-oxide (4a-4g)


1 mmole of the monoxime, 1 mmole of the aldehyde and 1.5 mmole of the amine was ground for 2 minutes and subsequently heated in an oil bath at 115-120°C, when a melt was formed. After a further 8 minutes of heating, the completion of the reaction was indicated by TLC. On cooling the melt slowly solidified and to the product so formed was added a little amount of ether whereby a precipitate was obtained. The precipitate was further washed with hot ethyl acetate. Recrystallization from ethanol gave products with the same melting points.

#### 2.5 General procedure for the synthesis of 1-hydroxy imidazole-3-oxide (5a-5e)

2 mmole each of the monoxime and the aldehyde was thoroughly ground with 695 mg (10 mmole) of hydroxylamine hydrochloride in an agate mortar and pestle for a period of ca. 3 minutes during which it melts and then gets hardened slowly. The mixture was then transferred to a test tube and heated in an oil bath maintained at 110-120°C when it started to melt. Constant shaking for another 7 minutes gave the product which remained in the melt form even at room temperature. On completion of the reaction, checked by TLC, addition of 5 ml of diethyl ether or 5 ml of ethyl acetate precipitated the product. The water insoluble products were then washed with water and ethyl acetate to get the pure products.

### 3 Results and discussion

The Radziwinski Imidazole (R-I) synthesis is a very useful preparative method for imidazole and its derivatives. The manifold utilizes a diketone, an aldehyde and ammonium acetate as the ammonia source (Scheme 1). The pure products have been synthesized in quantitative yields in 4 minutes at the given temperature by heating in an oil bath under solvent-free conditions. The temperature conditions and product yield were optimised by HPLC studies (ESI).



Scheme 1 R-I synthesis of tri-substituted imidazoles (1a-1i)

Besides usual spectroscopic methods for determining the structure, we have used single crystal X-ray diffraction (Sc-XRD) data of one of the representative compounds, 4, 5-diphenyl-1H-imidazole, 1i, to confirm the structure. The single crystals of the compound suitable for Sc-XRD were obtained by the slow evaporation from methanol/hexane mixture. The compound crystallizes in a monoclinic crystal system with the space group P2<sub>1</sub>/c (Hall group -P<sub>2ybc</sub>); a = 11.0471(4) Å, b = 9.2483(3) Å, c = 11.5780(4) Å, α = 90°, β = 93.921(3)°, γ = 90°, Z

$\mu = 4$ ,  $\mu = 0.577 \text{ mm}^{-1}$ ,  $F_{000} = 468.0$  and  $K\alpha = 1.54184 \text{ \AA}$ . The ORTEP diagram is presented in Fig.1. The heterocyclic ring is planar. One interesting point in the diagram is the attachment of one proton in each nitrogen atom which reflects overall an excess of one proton. The  $N_7-C_{11}$  and  $N_9-C_{10}$  bond distances are identical (1.380 Å and 1.3799 Å respectively), similarly  $N_7-C_8$  and  $N_9-C_8$  bond distances are also close (1.3157 Å and 1.3462 Å respectively); corresponding N7 and N9 centered bond angles are also close ( $107.83^\circ$  and  $105.91^\circ$ ).

Intermolecular extra proton transfer rate in imidazole is reported in the order of  $0.3 \times 10^{-12} \text{ second}^{31}$  but there is no such data for intramolecular N to N proton transfer due to the shift of N–C double bond in imidazole; intuitively it is difficult to say that the process could be faster than X-Ray diffraction time ( $10^{-18} \text{ second}$ ). However, the bond lengths are in between C–N single bond (1.47 Å) and double bond (1.25 Å). Since both the nitrogen atoms are identical, and on probability basis H atoms might have added to both the atoms during computation. The extra proton does not originate from the imidazolium salt because in such case the counter ion should be in the unit cell; and non equivalence of the nitrogen atoms would have been observed.






Fig.1 ORTEP diagram of 1i derived from single crystal data.

However, these reactions are not very efficient in solution, and the yield of product is not very high. Existing literature reveals that the Debus-Radziszewski Imidazole synthesis in solution state takes around 24 hrs to achieve moderate to good yield.<sup>25</sup> By contrast, solvent-free reactions proceed efficiently under much simpler conditions to give products in quantitative yield (Scheme 2).



Scheme 2 R-I synthesis of tetra-substituted imidazoles (2a-2l)

Interestingly, very similar solvent-free reactions of various ammonia derivatives and alkyl/aryl monoximes at 115–120°C for 10 mins also gave the corresponding imidazole N-oxides (3a–3g), N-substituted imidazole N-oxides (4a–4g) and 1-hydroxy imidazole 3-oxide derivatives (5a–5e) in quantitative yields (Scheme 3). This is the first report of a solvent-free condensed phase protocol for the oxide and hydroxyl oxide derivatives of imidazoles. Apart from the usual characterizations of the compounds, further spectroscopic investigations, vibrational assignments, HOMO-LUMO, NBO and MEP analysis of two of the synthesized compounds 3a<sup>32</sup> and 3c<sup>33</sup> have already been done and reported. The syntheses also proves that acetic acid, generated in situ in the reaction, is not the only catalysing factor for enhancing the reaction rates under solvent-free conditions.



Scheme 3 Analogous R-I synthesis of imidazole N-Oxides (3a-3g), N-substituted imidazole N-Oxides (4a-4g) and 1-hydroxy imidazole 3-Oxides (5a-5e).

In order to understand, at least partially, why these reactions proceed so efficiently under solvent-free conditions, a combination of methods viz., reactivity, spectroscopy and theory have been used.

#### Kinetic Studies

Firstly, the reaction kinetics of imidazoles 1(b) and 1(c) were studied through HPLC monitoring of benzil and aldehyde (reactants) consumption along with product (imidazoles) formation. A good linearity was observed in each case with the plots of natural logarithm values (-ve for the reactants and +ve for the products) of peak area against time. From the slope of these curves the first order rate constants and half lives ( $t_{1/2}$ ) were determined. The observed dependence of reactants concentrations (logarithm) and product formation with the variation of time (at reaction temperature 125°C, temperature

optimised for maximum conversion) is shown in Fig. 2. The product peak in HPLC was not much prominent within the first 10 minutes. This information helps to understand more about the sequences of reactions which manifest together as a multi-component reaction.





Fig. 2 (–ve)/(+ve)  $\ln(I)$  vs time(mins) of [i] benzil during formation of imidazoles (1b) and (1c); [ii] rate curve of the product imidazoles (1b) and (1c) formation; I = Peak A.

The catalytic effect of some metal salts (5 mole %) at the same reaction temperature (125°C) was also compared. The corresponding rate constant and half life values are shown in Table 1.

Table 1 Rate constants and half lives of reactant consumption and product formation at 125°C under solvent-free conditions.

Catalyst	Rate of benzil consumption ( $t_{1/2}$ ) <sup>[a]</sup>	Rate of aldehyde consumption ( $t_{1/2}$ )	Rate of product formation ( $t_{1/2}$ )
Solvent-free(no catalyst)	0.011 (63.01)	0.008 (86.64)	0.026 (26.66)
Sm(NO <sub>3</sub> ) <sub>3</sub> ·6H <sub>2</sub> O	0.012 (57.76)	0.008 (86.64)	0.023 (30.13)
Yb(SO <sub>3</sub> CF <sub>3</sub> ) <sub>3</sub>	0.033 (21.00)	0.009 (77.02)	0.115 (6.03)
ZrO(NO <sub>3</sub> ) <sub>2</sub>	0.058 (11.95)	0.017 (40.77)	0.034 (20.38)
(NH <sub>4</sub> ) <sub>2</sub> Ce(NO <sub>3</sub> ) <sub>6</sub>	0.035 (19.80)	0.024 (28.88)	0.112 (6.19)
NiCl <sub>2</sub> ·6H <sub>2</sub> O	0.032 (21.66)	0.021 (33.00)	0.041 (16.90)

<sup>[a]</sup> Note: Half-life ( $t_{1/2}$ ) in minutes



Scheme 4 Mechanism of the R-I synthesis

The general catalytic process at the molecular level maybe represented as in Scheme 4.

The compiled results indicate that the catalysts acted at different stages of the reaction sequences of the MCR reaction. To cite a representative case of Ytterbium triflate, a threefold increase in the rate of benzil consumption and fivefold increase in imidazole formation as against the catalyst free reaction was observed. Considering that in a reaction, catalysts in general, are used in very small proportions. Therefore only a small mole-fraction of the substrate would have the chance to be associated with the catalyst and thus the probability of the substrate taking part in adduct formation would be further reduced. Hence with only traces of a catalyst (5 mole % in the present case) added to a reaction, its rate should have hardly been affected as only a few molecules would engage in the activated complex in comparison to the vast number of un-associated molecules in the reaction media. We also studied the reactions at different elevated temperatures and found that at the temperature range 150–160 °C, almost quantitative products formed in a very short time of 4 minutes; and that too without using any catalyst. This is a landmark record in imidazole synthesis (ESI).

#### Infrared studies

In order to contemplate the cause behind the efficacy in solvent-free procedure; we studied the IR-spectra of pure benzil in thin film as well as that with catalyst. The carbonyl region of a benzil thin film and that with 5 mmole % of ytterbium triflate has been presented in Fig. 3. It was observed that the free carbonyl band (with catalyst) at 1676 cm<sup>-1</sup> has apparently increased in intensity with concomitant shift to lower frequency compared to the corresponding band for benzil. It was earlier reported<sup>34</sup> that C=O stretching band of benzil appears at 1676 cm<sup>-1</sup> in the crystalline state and at 1685 cm<sup>-1</sup> in solution.




Fig. 3 IR spectra of benzil thin film, with and without a catalyst.

As the catalyst caused a red-shift and brought about enhancement in intensity, we have considered this phenomenon as a marker of catalytic effect. Similar shift of carbonyl stretching frequency for benzophenone in TiO<sub>2</sub> surface has also been reported.<sup>35</sup> The difference in stretching frequency (9 cm<sup>-1</sup> red-shift) indicates a greater degree of single bond character in the C=O bond (polarisation enhancement) in the solid state compared to that in the solution state for benzil. The red-shift of carbonyl stretching provides an

indication of bulk polarization of benzil in the solid state and similar effect is also observed with catalysts in the solid state. The extent of the shift caused by the presence of trace amounts of zirconyl nitrate and ytterbium triflate on benzil carbonyl stretching frequency in hexane solution was also studied. The IR spectra in the carbonyl range have been presented in Fig. 4.




Fig. 4 Solution IR spectra in the carbonyl range of benzil in hexane.

It was found that in the solution spectra, the free carbonyl peak at  $1685\text{ cm}^{-1}$  and all other associated peaks showed a little amount of red-shift in the presence of a catalyst. In addition to this, there was concomitant diminishing of the intensity of the free carbonyl peak with increasing intensities of some other peaks at lower frequencies (particularly the peak at  $1680\text{ cm}^{-1}$ ) in the presence of traces of catalyst. It is also apparent that the catalyst, zirconyl nitrate, caused a little bit more polarization than that with ytterbium triflate. Notably we have found that the catalytic effect of zirconyl nitrate is a bit greater for benzil consumption rate as well (Table 1). The conspicuous enhancement of the band at  $1680\text{ cm}^{-1}$  in the presence of catalyst in both solution state and in solid state is observed. The peak at  $1680\text{ cm}^{-1}$  has been described as another C=O stretching band associated with the different symmetry of the molecule.<sup>36</sup> Enhancement of the band near  $1680\text{ cm}^{-1}$  in presence of catalyst suggests that the catalysts not only polarize the carbonyl but also influence the conformation of benzil. The IR spectroscopic investigation thus strongly provides evidences in favour of bulk polarization in the condensed phase which consequently leads to the enhancement of the electrophilicity of the carbonyl carbon. The above observation suggests that catalysts bind to the carbonyl oxygen and the weak interaction activates the carbonyl group with the enhancement of polarization.

### Computational Studies

As further proof of concept, the CBS-QB3 model chemistry calculations of a HCHO monomer and trimer were performed. Geometry optimization for a linear arrangement of the trimer was done as shown in Fig. 5. In molecular association a number of minimum energy conformations with different geometries are feasible, we searched for minimum energy in linear arrangement and our calculation terminated on convergence to such minima.




Fig. 5 Minimum energy models with partial charges.

The partial charges on the carbon and the oxygen atom and the dipole moment of the associated monomer are found to be enhanced as we move from the monomer to the trimer (Table 2).


Table 2 Mulliken atomic charges, dipole moment and C-O bond distances in monomers and trimer from the CBS-QB3 model

Parameter	Monomer	Trimer
Mulliken charge at C (au) <sup>[a]</sup>	0.316740	0.372316
Mulliken charge at O (au) <sup>[a]</sup>	-0.316740	-0.372316
Dipole moment (Debye)	2.8542	3.2133 <sup>[b]</sup>
C – O bond distance ( $^{\circ}$ A)	1.20001	1.20259
Symmetry	C2v	C1

Note: <sup>[a]</sup> Atomic charges with Hydrogens summed into heavy atoms; <sup>[b]</sup> Average value.

It is evident that the atomic charges in the associated monomers are different to those in the isolated free monomer, indicating a sizeable charge reorganisation due to the association. These may be interpreted as atomic expressions of sizeable cooperative effects. This charge reorganization in the carbonyls, as a consequence, brings about enhanced polarization in the carbonyl groups in their environment as the partial charges increase in each atom. A seminal report has already established the possibility of a dipole-dipole type of intermolecular interaction and the occurrence of little or no H-bond in formaldehyde itself.<sup>37</sup> The theoretical ab initio calculation of conformationally similar formaldehyde dimer was done by Smith et al,<sup>38</sup> whereby they have shown that an excess electron can attach to the system forming a dipole bound anion. This work indirectly supports our proposition as well, to the cause of enhancement in electrophilic behavior of the carbonyl group under solvent-free condition. The theoretical results also suggest that, as a result of the cooperative effect of very weak forces a bulk amount of carbonyl groups get activated. Therefore, the activation can be attributed to a unique spatial organization of the carbonyl moieties. This would be best understood if we consider an arbitrary molecular assembly where polarization transfer

through a non-covalent bond (weak bond interactions) could well be conceived of as shown in Scheme 5.



Scheme 5 Polarization transfer through a non-covalent bond without and with a catalyst

This pseudo-conjugated pi-system in the proposed supramolecular assembly makes the system more chemically soft (more polarizable) compared to a free carbonyl which is also responsible for better reactivity in solvent free reaction media. Hence, considerable catalytic action became viable in condensed phase. Since, in such an assembly a set of reacting molecules participates in a cycle, the process undoubtedly becomes faster. In addition to this self-activating effect, if some oxophilic substance were present in catalytic amount it bonded to the terminal carbonyl oxygen affecting further activation of the trail of carbonyls and thereby resulting in further enhancement of the catalytic effect (Scheme 5). The present work emphasizes on meeting this challenge of using weak forces as a design tool for imparting new properties and performance in molecules and materials.

The present work incorporates the concept of Constitutional Dynamic Chemistry to resolve the existing paradox in the process of catalysis (involving carbonyl activation). According to Prof. Lehn, supramolecular entities are assembled entities of discrete number of molecular sub-units held together reversibly through weak interactions (non-covalent interactions).<sup>39</sup> When extended to the carbonyl system, one could conceive the carbonyl molecular sub-units to be held together through similar type of weak interactions, by virtue of it being polar. In our opinion this hypothesis unequivocally embraces all the catalytic effects on carbonyls in solution state as well as in solvent-free molten state. As a proof of concept, we have reported herein the un-catalyzed Radziwinski synthesis of imidazole and its derivatives under solvent-free conditions employing a multicomponent protocol.

## Conclusions

The polarizability of organized carbonyl functionalities in condensed phase contributes for the observed self-catalysis.

High yields of many different imidazoles were obtained from the simply mechanical grinding and heating of MCR starting materials, even in the absence of Lewis acid catalysts. The very weak dipole of carbonyls can induce polarization in bulk because the carbonyl bonds are very much polarisable and the net result is the enhancement of electrophilicity of carbonyls. In polar solvents, the weak but favorable conformation of the carbonyl cluster expectedly breaks due to stronger solute-solvent interactions. Thus solvents act adversely to the self-catalytic effect. This phenomenon can be well utilized to generate a self-catalytic effect without using any catalytic substance.

## Acknowledgements


The author gratefully acknowledges IISER, Kolkata for the Single crystal X-ray studies and CDRI, Lucknow for elemental analysis and mass spectra. The authors are also thankful to Dr. Anirban Misra, Professor and Mr. Tamal Goswami, Research Associate, Department of Chemistry, N.B.U for the computational calculations.

## References

- (a) K. Tanaka, F. Toda, *Chem. Rev.*, 2000, 100, 1025; F. Toda, *Cryst. Eng. Comm.*, 2002, 4, 215; G. W. V. Cave, C. L. Raston, J. L. Scott, *Chem. Commun.*, 2001, 2159; G. Rothenberg, A. P. Downie, C. L. Raston, J. L. Scott, *J. Am. Chem. Soc.*, 2001, 123, 8701; G. W. V. Cave, C. L. Raston, *Chem. Commun.*, 2000, 2199; V. P. Balema, J. W. Wiench, M. Pruski, V. K. Pecharsky, *Chem. Commun.*, 2002, 724.
- F. Toda, K. Tanaka and K. Hamai, *J. Chem. Soc., Perkin Trans.*, 1990, 3207.
- K. Yoshizawa, S. Toyota and F. Toda, *Tetrahedron Lett.*, 2001, 42, 7983.
- K. Tanaka, T. Sugino and F. Toda, *Green Chem.*, 2000, 2, 303.
- G. Kaupp, M. R. Naimi-Jamal and J. Schmeyers, *Tetrahedron*, 2003, 59, 375.
- K. Yoshizawa, S. Toyota and F. Toda, *Green Chem.*, 2002, 4, 68.
- D. C. Waddell and J. Mack, *Green Chem.*, 2009, 11, 79.
- K. Tanaka, S. Kishigami and F. Toda, *J. Org. Chem.*, 1991, 56, 4333.
- F. Toda, M. Yagi and K. Kiyoshige, *J. Chem. Soc., Chem. Commun.*, 1988, 958.
- F. Toda and T. Shigemasa, *J. Chem. Soc. Perkin Trans.1*, 1989, 209.
- F. Toda, K. Tanaka, and S. Iwata, *J. Org. Chem.*, 1989, 54, 13.
- N. Elders, D. van der Born, L. J. D. Hendrickx, B. J. J. Timmer, A. Krause, E. Janssen, F. J. J. de Kanter, E. Ruijter and R. V. A. Orru, *Angew. Chem., Int. Ed.* 2009, 48, 5856.
- (a) B. Karimi and D. Zareyee, *J. Mater. Chem.*, 2009, 19, 8665; H. Wang, X. Zhao, Y. Li and L. Lu, *Org. Lett.*, 2006, 8 (7), 1379; P. Galletti, M. Pori and D. Giacomini, *Eur. J. Org. Chem.*, 2011, 20, 3896.
- M. A. Zolfigol, E. Kolvari, A. Abdoli and M. Shiri, *Mol. Divers.*, 2010, 14(4), 809; V. Sivamurugan, R. S. Kumar, M. Palanichamy, V. Murugesan, *Journal of Heterocyclic Chemistry*, 2005, 42 (5), 969; G. V. M. Sharma, K. L. Reddy, P. S. Lakshmi and P. R. Krishna, *Synthesis*, 2006, 55.
- F. Bigi, S. Carloni, B. Frullanti, R. Maggi and G. Sartori, *Tetrahedron Lett.*, 1999, 40 (17), 3465; J. Peng and Y. Deng, *Tetrahedron Lett.*, 2001, 42 (34), 5917; R. Wang and Z. Liu, *J. Org. Chem.*, 2012, 77 (8), 3952.

- 16 (a) L. el Kaim, L. Gautier, L. Grimaud, L. M. Harwood and V. Michaut, *Green Chem.*, 2003, 5, 477; (b) Y. Hayashi, T. Urushima, S. Aratake, T. Okano and K. Obi, *Org. Lett.*, 2008, 10 (1), 21.
- 17 D. Koszelewski, W. Szymanski, J. Krysiak and R. Ostaszewski, *Synthetic Commun.*, 2008, 38 (7), 1120; T. Bousquet, M. Jida, M. Soueidan, R. Deprez- Poulain, F. Agbossou-Niedercorn L. Pelinski, *Tetrahedron Lett.*, 2012, 53 (3), 306.
- 18 (a) N. Liu, S. Cao, J. Wu, J. Yu, L. Shen, X. Feng and X. Qian, *Tetrahedron*, 2008, 64, 3966; L. El Kaim, L. Grimaud and S. Hadrot, *Tetrahedron Lett.*, 2006, 47 (23), 3945; M. Jida, S. Malaquin, R. Deprez-Poulain, G. Laconde and B. Deprez, *Tetrahedron Lett.*, 2010, 51 (39), 5109.
- 19 J. S. B. Forero, E. M. de Carvalho, J. J. Junior and F. M. da Silva, *Heterocyclic Lett.*, 2011, 1 (1), 61; K. Wang, D. Kim, and A. Domling, *J. Comb. Chem.*, 2010, 12, 111.
- 20 P. Nun, J. Martinez and F. Lamaty, *Synthesis*, 2010, 12, 2063.
- 21 B. Jiang, X. Wang, F. Shi, Shu-Jiang Tu, T. Ai, A. Ballew, G. Li, *J. Org. Chem.*, 2009, 74, 9486; G. Bratulescu, *Synthesis*, 2009, 2319; X. Diao, Y. Wang, Y. Jiang, D. Ma, *J. Org. Chem.*, 2009, 74, 7974; T. Horneff, S. Chuprakov, N. Chernyak, V. Gevorgyan, V. V. Fokin, *J. Am. Chem. Soc.*, 2008, 130, 14972; P. Saha, T. Ramana, N. Purkait, M. A. Ali, R. Paul, T. Punniyamurthy, *J. Org. Chem.*, 2009, 74, 8719; M. M. Heravia, M. Zakeria, N. Karimia, M. Saeedia, H. A. Oskooiea, N. T. Hosienib, *Synthetic Communications* 2010, 40, 1998; K. Hirano, S. Urban, C. Wang, F. Glorius, *Org. Lett.*, 2009, 11, 1019.
- 22 P.I. Dalko, L. Moisan, *Angew. Chem., Int. Ed.*, 2004, 43, 5138; D. Basavaiah, A. J. Rao, T. Satyanarayana, *Chem Rev.*, 2003, 103, 811; P. Langer, *Angew. Chem. Int. Ed.* 2000, 39, 3049-3052; H. Groger, J. Wilken, *Angew. Chem. Int. Ed. Engl.*, 2001, 40, 529; J.W.J. Kennedy, D.G. Hall, *Angew. Chem., Int. Ed.*, 2003, 42, 4732; B. List, *Tetrahedron*, 2002, 58, 5573.
- 23 Marc-Olivier Simon and Chao-Jun Li, *Chem Soc Rev*, 2011.
- 24 Chao-Jun Li, *Chem. Rev.* 2005, 105, 3095; Y. R. Leshkov and M. E. Davis, *ACS Catal.* 2011, 1, 1566; F. Cruz-Acosta, A. Santos-Exposito, P. de Armas and F. Garcia-Tellado, *Chem. Commun.*, 2009, 6839; R. N. Butler and A. G. Coyne, *Chem. Rev.* 2010, 110, 6302.
- 25 S. D. Sharma, P. Hazarika and D. Konwar, *Tetrahedron Lett.*, 2008, 49, 2216.
- 26 S. Samai, G. C. Nandi, P. Singh and M. S. Singh, *Tetrahedron*, 2009, 65, 10155.
- 27 G. M. Sheldrick, SHELXS-97, University of Gottingen, 1997
- 28 J. A. Montgomery, M. J. Frisch, J.W. Ochterski, G. A. Petersson Jr., *J. Chem. Phys.*, 2000, 112, 6532; J. A. Montgomery, M. J. Frisch, J. W. Ochterski, G. A. Petersson Jr., *J. Chem. Phys.*, 1999, 110, 2822.
- 29 L. A. Curtiss, K. Raghavachari, G. W. Trucks, J. A. Pople, *J. Chem. Phys.*, 1991, 94, 7221; L. A. Curtiss, K. Raghavachari, P. C. Redfern, V. Rassolov, J. A. Pople, *J. Chem. Phys.* 1998, 109, 7764; J. W. Ochterski, G. A. Petersson, J. A. Montgomery, *J. Chem. Phys.* 1996, 104, 2598; G. A. Petersson, D. K. Malick, W. G. Wilson, J. W. Ochterski, J. A. Montgomery, M. J. Frisch, *J. Chem. Phys.*, 1998, 109, 10 570.
- 30 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, et al., Gaussian 09, Revision A.11.4, Gaussian, Inc., Pittsburg PA, 2002.
- 31 W. Munch et al, *Solid State Ionics*, 2001, 145, 437
- 32 K.B. Benzon, H. T. Varghese, C. Y. Panicker, K. Pradhan, B. K. Tiwary, A. K. Nanda, C. V. Alsenoy, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 2015, 151, 965.
- 33 K. B. Benzon, H. T. Varghese, C. Y. Panicker, K. Pradhan, B. K. Tiwary, A. K. Nanda, C. V. Alsenoy, *Spectrochimica Acta Part A: Molecular and Bimolecular Spectroscopy*, 2015, 146, 307.
- 34 E. C. Lee, D. Kim, P. Jurecka, P. Tarakeshwar, P. Hobza and K. S. Kim, *J. Phys. Chem. A*, 2007, 111, 3446
- 35 L. Colombo, D. Kirin, V. Volovsek, N. E. Lindsay, J. F. Sullivan and J. R. Durig, *J. Phys. Chem.* 1989, 93, 6290.
- 36 L. M. Babkov, T. V. Bezrodnaya, G. A. Puchkovskaya, K. E. Uspenskii and V. V. Shimanovskaya, *Journal of Structural Chemistry* 2008, 49, 517.
- 37 T. Bercovici, J. King and R. S. Becker, *J. Chem. Phys.* 1972, 56, 3956; J. E. D. Bene, *J. Chem. Phys.* 1971, 60, 3812.
- 38 M. A. Dayle, J. S. Smith, Y. Elkadi and L. Adanokicz, *Chem. Phys. Lett.*, 1999, 305, 169.
- 39 J. M. Lehn, *Angew. Chem. Int. Ed.*, 1990, 29, 1304.





A supramolecular assembly of reacting molecules (carbonyls) held together by weak-bond interactions (as per the principles of Constitutional Dynamic Chemistry); high viability for reactions carried out in melt. The bulk polarised molecules participate in the reactions in a cycle rather than as individual moieties leading to greatly enhanced reaction rates; explains faster reaction rates of some solvent-free reactions as against reactions carried out in solvent media.