

# Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## ARTICLE

# A Supramolecular Strategy for Tuning the Energy Level of Naphthalenediimide: Promoted Formation of Radical Anions with Extraordinary Stability

Cite this: DOI: 10.1039/x0xx00000x

Qiao Song, Fei Li, Zhiqiang Wang and Xi Zhang\*

Received 00th January 2012,  
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

[www.rsc.org/](http://www.rsc.org/)

We reported a supramolecular strategy to promote and stabilize the formation of naphthalenediimide (NDI) radical anions. The LUMO and HOMO energy of NDI were lowered significantly by introducing cucurbit[7]uril (CB[7]) to each side of a designed NDI molecule through supramolecular complexation. This efficiently promoted the photo-induced electron transfer process between NDI and bromide anions in aqueous solution. The resulting NDI supramolecular radical anions are of outstanding stability. They are even stable in aqueous solution at higher temperatures of 40 °C and 60 °C. It is anticipated that this supramolecular strategy may provide a facile method for stabilizing radicals towards the development of novel materials with spin-based properties and optical properties in the visible and near-infrared regions.


## Introduction

Organic radicals are of prime interest because of their various spin-based applications and near-infrared optical applications.<sup>[1]</sup> Most organic radicals are highly reactive due to their unpaired electrons, which are often unstable under ambient conditions; only a few families of stable radicals at ambient conditions have been reported for many decades.<sup>[2]</sup> The 1,4,5,8-naphthalenediimides (NDIs) are  $\pi$ -electron-deficient compounds. NDI radical anions are one of the most widely studied organic radicals recently. As reported, there are two pathways to generate NDI radical anions, electron transfer (ET) or photo-induced electron transfer (PET). To display anion-involved ET and PET process, the NDIs must possess low LUMO energy levels. Electron-withdrawing N-substituents have been introduced to NDIs to lower the LUMO energy and promote the formation of NDI radical anions.<sup>[3]</sup> For example, NDI derivatives with strong electron-withdrawing phosphonium groups displayed LUMO energy as low as -4.90 eV. The acquired NDI radical anions are of extraordinary stability to air, light and chromatography, which can even be isolated as a single crystal.<sup>[4]</sup>

In addition to the above covalent modification, we wondered if it is possible to realize the stabilization of radicals through noncovalent synthesis, and the complicated covalent synthesis could be avoided to some extent. Recently, there exist examples to stabilize radicals through supramolecular interactions. Kim et al. reported several cases of the stabilization of radicals in host-guest complexes.<sup>[5]</sup> Methylviologen cation radicals and

tetrathiafulvalene cation radicals were incorporated into the hydrophobic cavity of the cucurbit[8]uril (CB[8]) in the molar ratio of 2:1, forming stable radical dimers, respectively. The stability of the radical dimers within CB[8] may be attributed to the hydrophobic cavity with a proper size. In other words, the CB[8] is used to protect the radical dimers through a direct encapsulation strategy.


In this paper, we aim to establish a general and facile strategy to stabilize radicals by tuning the LUMO and HOMO energy level of organic molecules through noncovalent interactions. To achieve this goal, we designed and synthesized a water-soluble NDI derivative with one quaternary ammonium alkyl chain as the N-substituent on each end, as shown in Scheme 1. Cucurbit[7]uril (CB[7]) is a macrocyclic host that contains seven electron-withdrawing carbonyl groups on each side, which was used to combine with NDI in aqueous solutions through noncovalent host-guest interactions.<sup>[6]</sup> Although the NDI core was not encapsulated into the CB[7] hydrophobic cavity, the LUMO and HOMO energy of NDI would be significantly lowered because of the electron-withdrawing effect of the carbonyl groups of CB[7]. This may greatly promote the PET process between the supramolecular complex of NDI and CB[7] (NDI/CB[7]) and bromide anion to form stable NDI radical anions. It is anticipated that this strategy of fabricating supramolecular radicals could provide a facile method for stabilizing radicals towards the development of novel materials with spin-based properties and optical properties in the visible and near-infrared regions.



**Scheme 1.** The formation of NDI radical anions is promoted and stabilized efficiently by introducing CB[7] to each side of NDI through supramolecular complexation.

## Results and discussion


The supramolecular complex between NDI derivative and CB[7] was prepared simply by mixing NDI and CB[7] in a molar ratio of 1:2 in aqueous solution. As shown in the  $^1\text{H}$  NMR spectra (Figure 1), peaks ascribed to the alkyl chains of NDI exhibited an upfield shift, indicating that CB[7] was introduced to each side of the NDI derivative. Furthermore, the ITC data indicated that complexation induced a significant change in the titration curve when CB[7] was added into the solution of NDI in a molar ratio of 2:1. One set of binding site model was utilized to fit the isothermal titration curve, which gave a binding constant of  $2.4 \times 10^4 \text{ M}^{-1}$  (see Figure S1 in the Supporting Information). Therefore, noncovalent host-guest interactions between the alkyl chains in the NDI derivative and CB[7] are responsible for the formation of the NDI/CB[7] supramolecular complex with well-defined composition.



**Figure 1.**  $^1\text{H}$  NMR spectra of NDI and NDI/CB[7] supramolecular complex (400 MHz,  $\text{D}_2\text{O}$ ,  $25^\circ\text{C}$ ).

To understand if the supramolecular complexation could promote the PET process, we did a comparative study by irradiating the aqueous solutions of NDI/CB[7] and NDI under UV light. The NDI/CB[7] solution turned dark red after 5 s of irradiation while NDI solution remained almost colourless at the same condition. As monitored by UV/vis spectra (Figure 2a), the absorption bands peaked at 363 nm and 383 nm ascribed to NDI vanished after UV irradiation. Concomitantly, several new absorption bands at long wavelengths appeared that are characteristic of NDI radical anions.<sup>[7]</sup> Moreover, the resulting NDI radical anions could be verified through electron paramagnetic resonance (EPR), which showed characteristic EPR signals after UV irradiation (see Figure S2 in the Supporting Information).  $^1\text{H}$  NMR and fluorescent spectra also support the formation of NDI radical anions for the NDI/CB[7] solution (see Figures S3 and S4 in the Supporting Information). We studied the reaction rates of both NDI and NDI/CB[7] by monitoring the absorption bands at 363 nm and 383 nm at different irradiation times (Figure 2b). Simulated according to the principle of first-order reaction, the reaction rate constant of NDI/CB[7] is  $0.50 \pm 0.01 \text{ s}^{-1}$ , which is around 10 times larger than that of NDI ( $0.051 \pm 0.001 \text{ s}^{-1}$ ). Therefore, the formation of NDI radical anions under UV irradiation is significantly promoted because of the supramolecular complexation.


The fluorescent lifetime of both NDI and NDI/CB[7] supramolecular complex was measured through time-resolved fluorescence measurements. As shown in Figure 2c, the fluorescent lifetime of NDI is 165 ps, while the fluorescent lifetime of NDI/CB[7] supramolecular complex is 100 ps. The shortened fluorescent lifetime further confirms that the PET process of NDI is promoted by the supramolecular complexation.




**Figure 2.** (a) UV/vis spectra of NDI/CB[7] before and after UV irradiation. (b) Reaction dynamics of NDI and NDI/CB[7] under UV irradiation. (c) Time-resolved fluorescence measurements of NDI and NDI/CB[7].

To investigate whether the LUMO and HOMO energy level of NDI was influenced by the supramolecular complexation, we evaluated the reduction potentials of NDI and NDI/CB[7] supramolecular complex through cyclic voltammetry (CV).<sup>[8]</sup> As demonstrated in Figure 3, they both underwent two reversible one-electron reductions, forming NDI<sup>-</sup> and NDI<sup>2-</sup> species, respectively. NDI/CB[7] showed two reduction peaks as 0.138 eV and -0.557 eV while NDI as -0.331 eV and -0.670 eV. To calculate the LUMO and HOMO energy of these two species, the UV/vis absorption spectra were utilized to estimate the HOMO-LUMO energy gap ( $\Delta E$ ) (Figure 4). Both

NDI and NDI/CB[7] gave identical  $\Delta E$  values of 3.15 eV. From the calculated LUMO and HOMO energy shown in Figure 4, the LUMO energy of NDI/CB[7] is 0.48 eV lower than NDI itself, and therefore the LUMO energy level is as low as -4.89 eV. This lowered LUMO energy level accounts for the promoted formation of NDI supramolecular radical anions under UV irradiation.



**Figure 3.** Cyclic voltammograms of NDI and NDI/CB[7].




**Figure 4.** Calculation of energy gap between HOMO and LUMO of NDI and NDI/CB[7] (left); LUMO and HOMO energy of NDI and NDI/CB[7] (right).

We wondered whether the lowered LUMO energy was due to the supramolecular complexation or simply because of physical mixing. To answer this question, a control experiment was carried out. Here, 1-adamantanamine hydrochloride (ADA), which has a much stronger binding strength with CB[7] than NDI,<sup>[9]</sup> was added to the solution of NDI/CB[7] supramolecular complex to destroy the complexation, which was verified by <sup>1</sup>H NMR spectra (see Figure S5 in the Supporting Information). The UV/vis spectra showed that the formation of NDI radical anions was greatly suppressed, which was similar to the NDI solution (see Figure S6 in the Supporting Information). This control experiment

reveals that the CB[7] should combine with the alkyl chains of the NDI derivative, and it is in this way that the carbonyl groups can stay close enough to affect the energy level of NDI.


The supramolecular radical anions formed in this way are of outstanding stability. The solution of NDI radical anions became colourless within a few seconds upon shaking against ambient atmosphere, suggesting that the NDI radical anions are very unstable. However, the NDI supramolecular radical anions changed very slowly, suggesting the stability of NDI radical anions has been significantly enhanced. For more quantitative data, we studied the stability of radical anions of NDI and NDI/CB[7] supramolecular complex in aqueous solutions via UV/vis spectroscopy. We monitored the absorption bands peaked at 363 nm and 383 nm at different times after UV irradiation. This gave the retention rate of the NDI radical anions as a function of time (Figure 5). At room temperature (25 °C), the NDI supramolecular radical anions decreased only by 14% after 6 h, while the radical anions in NDI vanished completely in 3 h. Furthermore, the supramolecular radical anions are comparably stable at higher temperature with retention rates of 80% and 70% after 6 h at 40 °C and 60 °C, respectively. Therefore, not only can we promote the formation of NDI radical anions, but also we can stabilize the NDI radical anions efficiently in a supramolecular way.



**Figure 5.** Stability of NDI and NDI/CB[7] monitored by UV/vis spectra.


The formed NDI radical anions can be utilized as reducing agent, which allows for the function of NDI/CB[7] supramolecular complex as supramolecular photo-reducing agent with high efficiency. Here, cytochrome *c* (cyt *c*) was selected as a model compound. Cyt *c* is a small redox protein in organism, participating in the mitochondrial electron transfer chain. It contains a heme group whose central iron atom can be reversibly reduced from the Fe(III) to the Fe(II)

state.<sup>[10]</sup> It was reported that NDI derivatives could be used to reduce cyt *c*.<sup>[11]</sup> The reduction kinetics of cyt *c* were investigated upon monitoring the absorption band peaked at 549 nm under UV irradiation at the existence of NDI or NDI/CB[7] supramolecular complex. As shown in Figure 6, the cyt *c* was fully reduced in 140 s in the presence of NDI while the reduction time was shortened to only 20 s with the aid of NDI/CB[7] supramolecular complex. Benefiting from the promoted PET of the NDI/CB[7] supramolecular complex, the reduction process of cyt *c* is significantly accelerated under UV irradiation, which provides a facile way to realize photo-reduction with promoted efficiency.



**Figure 6.** Reduction kinetics of cyt *c* at the existence of NDI or NDI/CB[7] supramolecular complex.

Importantly, this supramolecular strategy works for a wide range of NDI molecules. Taking NDI derivative (NDI-nap) bearing one naphthalene group on each end for example (Figure 7a), NDI-nap could combine with CB[7] in a molar ratio of 1:2,<sup>[12]</sup> which was proved by <sup>1</sup>H NMR spectrum and ITC (see Figures S7 and S8 in the Supporting Information). There was no change in the UV/vis spectra for NDI-nap alone, but the NDI-nap/CB[7] supramolecular complex presented characteristic absorption bands of NDI radical anions after UV irradiation (Figure 7b and 7c). Once again, we evaluated the LUMO and HOMO energy of the NDI-nap and NDI-nap/CB[7] based on CV and UV/vis spectra (see Figures S9 and S10 in the Supporting Information). The results show that the LUMO energy of NDI-nap/CB[7] is 0.34 eV lower than NDI-nap, which proves furthermore the feasibility of this supramolecular strategy.



**Figure 7.** (a) Molecular structure of NDI-nap. (b) UV/vis spectra of NDI-nap before and after UV irradiation. (c) UV/vis spectra of NDI-nap/CB[7] before and after UV irradiation.

## Conclusions

In conclusion, we have developed a supramolecular strategy to promote and stabilize the formation of NDI radical anions through the supramolecular complexation of cucurbit[7]uril and NDI derivatives. This strategy significantly lowers the LUMO and HOMO energy of NDI, favouring the formation of NDI supramolecular radical anions with extraordinary stability. It is highly anticipated that this line of research may provide a general method to stabilize radicals through a facile supramolecular route for the development of materials with spin-based properties and optical properties in the visible and near-infrared regions.

## Acknowledgements

This work was financially supported by the National Basic Research Program (2013CB834502) and the Foundation for Innovative Research Groups of NSFC (211210004). We thank Mr. Wei Cao and Ms. Qinqin Zhou in Tsinghua University with the assistance of cyclic voltammograms. We appreciate

the help of Prof. Anchi Yu and Ms. Huiyu Zhang from Renmin University of China with the help of time-resolved fluorescence measurements. We thank Ms. Meng Zhou in Tsinghua University with the measurement of COSY. We also thank Dr. Liulin Yang and Dr. Jiangfei Xu in Tsinghua University for their helpful discussions.

## Notes and references

The Key Lab of Organic Optoelectronics & Molecular Engineering, Department of Chemistry, Tsinghua University, Beijing 100084, P. R. China.

E-mail: [xi@mail.tsinghua.edu.cn](mailto:xi@mail.tsinghua.edu.cn)

Electronic Supplementary Information (ESI) available. See DOI: 10.1039/c000000x/

- (a) R. Haddon, *Nature*, 1975, **256**, 394; (b) F. Würthner, *Angew. Chem., Int. Ed.*, 2001, **40**, 1037; (c) M. Mas-Torrent, N. Crivillers, C. Rovira and J. Veciana, *Chem. Rev.*, 2012, **112**, 2506; (d) S. Fukuzumi, K. Ohkubo, F. D'Souza and J. Sessler, *Chem. Commun.*, 2012, **48**, 9801; (e) J. Barnes, A. Fahrenbach, D. Cao, S. Dyar, M. Frascioni, M. Giesener, D. Benitez, E. Tkatchouk, O. Chernyashkevskyy, W. Shin, H. Li, S. Sampath, C. Stern, A. Sarjeant, K. Hartlieb, Z. Liu, R. Carmieli, Y. Botros, J. Choi, A. Slawin, J. Ketterson, M. Wasielewski, W. Goddard and J. Stoddart, *Science*, 2013, **339**, 429; (f) A. Trabolsi, N. Khashab, A. Fahrenbach, D. Friedman, M. Colvin, K. Coti, D. Benitez, E. Tkatchouk, J. Olsen, M. Belowich, R. Carmieli, H. Khatib, W. Goddard, M. Wasielewski and J. Stoddart, *Nat. Chem.*, 2010, **2**, 42.
- (a) M. Gomberg, *J. Am. Chem. Soc.*, 1900, **22**, 757; (b) D. Griller and K. Ingold, *Acc. Chem. Res.*, 1976, **9**, 13; (c) P. Power, *Chem. Rev.*, 2003, **103**, 789; (d) C. Martin, M. Soleilhavou and G. Bertrand, *Chem. Sci.*, 2013, **4**, 3020; (e) R. Hicks, *Org. Biomol. Chem.*, 2007, **5**, 1321; (f) O. Mallow, M. Khanfar, M. Malischewski, P. Finke, M. Hesse, E. Lork, T. Augenstein, F. Breher, J. Harmer, N. Vasilieva, A. Zibarev, A. Bogomyakov, K. Seppelt and J. Beckmann, *Chem. Sci.*, 2015, **6**, 497.
- (a) S. Nelsen, *J. Am. Chem. Soc.*, 1967, **89**, 5925; (b) S. Guha, F. Goodson, L. Corson and S. Saha, *J. Am. Chem. Soc.*, 2012, **134**, 13679; (c) S. Guha, F. Goodson, S. Roy, L. Corson, C. Gravenmier and S. Saha, *J. Am. Chem. Soc.*, 2011, **133**, 15256; (d) S. Guha and S. Saha, *J. Am. Chem. Soc.*, 2010, **132**, 17674; (e) J. Chang, Q. Ye, K. Huang, J. Zhang, Z. Chen, J. Wu and C. Chi, *Org. Lett.*, 2012, **14**, 2964; (f) I. Campos, I. Nantes, M. Politi and S. Brochsztain, *Photochem. Photobiol.*, 2004, **80**, 518; (g) S. V. Bhosale, C. H. Jani and S. J. Langford, *Chem. Soc. Rev.*, 2008, **37**, 331; (h) D. Schmidt, D. Bialas and F. Würthner, *Angew. Chem., Int. Ed.*, 2014, **54**, 3611; (i) S. Lee, Y. Zu, A. Herrmann, Y. Geerts, K. Mullen and A. Bard, *J. Am. Chem. Soc.*, 1999, **121**, 3513; (j) M. Ajayakumar, D. Asthana and P. Mukhopadhyay, *Org. Lett.*, 2012, **14**, 4822; (k) C. Leong, B. Chan, T. Faust, P. Turner and D. D'Alessandro, *Inorg. Chem.*, 2013, **52**, 14246; (l) S. Seifert, D. Schmidt and F. Würthner, *Chem. Sci.*, 2015, **6**, 1663; (m) E. Iengo, G. Pantos, J. Sanders, M. Orlandi, C. Chiorboli, S. Fracasso and F. Scandola, *Chem. Sci.*, 2011, **2**, 676; (n) K.-R. Wang, D.-S. Guo, B.-P. Jiang and Y. Liu, *Chem. Commun.*, 2012, **48**, 3644; (o) B.-P. Jiang, D.-S. Guo and Y. Liu, *J. Org. Chem.*, 2011, **76**, 6101.
- S. Kumar, M. Ajayakumar, G. Hundal and P. Mukhopadhyay, *J. Am. Chem. Soc.*, 2014, **136**, 12004.
- (a) W. Jeon, H. Kim, C. Lee and K. Kim, *Chem. Commun.*, 2002, 1828; (b) A. Ziganshina, Y. Ko, W. Jeon and K. Kim, *Chem. Commun.*, 2004, 806.
- (a) W. Freeman, W. Mock and N. Shih, *J. Am. Chem. Soc.*, 1981, **103**, 7367; (b) J. W. Lee, S. Samal, N. Selvapalam, H. J. Kim and K. Kim,

- Acc. Chem. Res.*, 2003, **36**, 621; (c) K. Kim, N. Selvapalam, Y. Ko, K. Park, D. Kim and J. Kim, *Chem. Soc. Rev.*, 2007, **36**, 267; (d) F. Biedermann, V. Uzunova, O. Scherman, W. Nau and A. De Simone, *J. Am. Chem. Soc.*, 2012, **134**, 15318; (e) L. Cera and C. Schalley, *Chem. Sci.*, 2014, **5**, 2560; (f) F. Biedermann, W. Nau and H. Schneider, *Angew. Chem., Int. Ed.*, 2014, **53**, 11158; (g) H. Isobe, N. Tomita, J. Lee, H. Kim, K. Kim and E. Nakamura, *Angew. Chem., Int. Ed.*, 2000, **39**, 4257.
- 7 (a) S. Langford, M. Latter and C. Woodward, *Photochem. and Photobiol.*, 2006, **82**, 1530; (b) G. Andric, J. Boas, A. Bond, G. Fallon, K. Ghiggino, C. Hogan, J. Hutchison, M. Lee, S. Langford, J. Pilbrow, G. Troup and C. Woodward, *Aust. J. Chem.*, 2004, **57**, 1011.
- 8 C. Cardona, W. Li, A. Kaifer, D. Stockdale and G. Bazan, *Adv. Mater.*, 2011, **23**, 2367.
- 9 S. Liu, C. Ruspic, P. Mukhopadhyay, S. Chakrabarti, P. Zavalij and L. Isaacs, *J. Am. Chem. Soc.*, 2005, **127**, 15959.
- 10 (a) X. Liu, C. Kim, J. Yang, R. Jemmerson and X. Wang, *Cell*, 1996, **86**, 147; (b) J. Yang, X. Liu, K. Bhalla, C. Kim, A. Ibrado, J. Cai, T. Peng, D. Jones and X. Wang, *Science*, 1997, **275**, 1129; (c) R. Kluck, E. BossyWetzel, D. Green and D. Newmeyer, *Science*, 1997, **275**, 1132; (d) C. Bortolotti, L. Paltrinieri, S. Monari, A. Ranieri, M. Borsari, G. Battistuzzi and M. Sola, *Chem. Sci.*, 2012, **3**, 807; (e) S. Jiang, J. Cai, D. Wallace and D. Jones, *J. Biol. Chem.*, 1999, **274**, 29905.
- 11 (a) I. Campos, I. Nantes, M. Politi and S. Brochsztain, *Photochem. Photobiol.*, 2004, **80**, 518; (b) K. Tsukahara, R. Ueda and M. Goda, *Bull. Chem. Soc. Jpn.*, 2001, **74**, 1303.
- 12 (a) Y. Zeng, Y. Li, M. Li, G. Yang and Y. Li, *J. Am. Chem. Soc.*, 2009, **131**, 9100; (b) K. Liu, Y. Liu, Y. Yao, H. Yuan, S. Wang, Z. Wang and X. Zhang, *Angew. Chem., Int. Ed.*, 2013, **52**, 8285; (c) K. Liu, Y. Yao, Y. Kang, Y. Liu, Y. Han, Y. Wang, Z. Li and X. Zhang, *Sci. Rep.*, 2013, **3**, 2372.