

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemicalscience

Detection of Bioorthogonal Groups by Correlative Light and Electron Microscopy Allows Imaging of Degraded Bacteria in Phagocytes

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Daphne M. van Elstrand^a, Erik Bos^b, Wouter de Boer^a, Herman S. Overkleeft^a, Abraham J. Koster^{b*}, Sander I. van Kasteren^{a*}

The interaction between parasites and phagocytic immune cells is a key inter-species interaction in biology. Normally, phagocytosis results in the killing of invaders, but obligate intracellular parasites hijack the pathway to ensure their survival and replication. The *in situ* study of these parasites in the phagocytic pathway is very difficult, as genetic modification is often complicated and, if successful, only allows the tracking of pathogen phagocytosis up until the degradation of the engineered reporter constructs. Here we combine bioorthogonal chemistry with correlative light-electron microscopy (CLEM) to follow bacterial processing in the phagolysosomal system. Labelled bacteria are produced using bioorthogonal non-canonical amino tagging (BONCAT), precluding the need for any genetic modification. The bacterial proteome – even during degradation – was then visualised using a novel CLEM-based approach. This allowed us to obtain high resolution information about the subcellular localisation of the degrading bacteria, even after the proteolytic degradation of reporter constructs. To further explore the potential of CLEM-based imaging of bioorthogonal functionalities, azide-labelled glycans were imaged by this same approach, as well as active-subpopulations of enzymes using a 2-step activity-based protein profiling strategy.

Introduction

Phagocytic degradation is a question of great biological relevance, as it is one of the key mechanisms by which the immune system keeps pathogens at bay. As a consequence, subversion of the phagolysosomal pathway is a survival strategy employed by a wide range of parasites, which collectively are responsible for a great amount of human morbidity and mortality¹.

The interaction between immune cells and pathogenic bacteria is very difficult to study², as intracellular pathogens can be non-trivial to grow *ex vivo*³ and very difficult to genetically alter. Even in (rare) cases where these bacteria can be genetically modified⁴ the imaging of their encounters with host phagocytes is limited to encounters where successful infection is established. Encounters whereby the pathogens are killed and degraded are difficult to image as the proteolysis




Figure 1 - Overview: (a) phagocytosed azido-*E. coli* can be fluorescently visualised in an ultrathin cryosection using a cHc-reaction with a fluorophore; (b) overlay of this image on an electron micrograph provides an ultrastructural context for the signal with nanometre-resolution. As the bioorthogonal handle is stable to proteolysis, degraded bacteria can be visualized in this manner.

that is a hallmark of successful phagocytic maturation⁵ results in the degradation of reporter proteins and epitopes⁶. Bioorthogonal chemistry is a powerful tool for labelling of (sub)-populations of biomolecules in complex biological systems⁷ and could be employed to circumvent these problems. The approach relies on the introduction of a small, physiologically inert chemical group into a biomolecule of interest that can subsequently be visualised using a selective reaction⁸. The small size, biological stability of the chemical group, and the wide range of biomolecules that can be labelled with this approach makes this method a valuable part of the biochemist's toolkit^{9,10}.

^a Division of Bio-organic Synthesis, Leiden Institute of Chemistry, Gorlaeus Laboratories, Leiden University, The Netherlands and Institute for Chemical Immunology, Leiden, The Netherlands.

^b Department of Molecular Cell Biology, Section Electron Microscopy, Leiden University Medical Center, Leiden, The Netherlands.

* Correspondence should be addressed to: s.i.van.kasteren@chem.leidenuniv.nl or a.j.koster@lumc.nl

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

Bolstered by the recent successful imaging of a pathogen inside a host phagocyte through the use of a bioorthogonally modified cell wall component, *D*-alanine^{11–13}, we envisaged that bioorthogonal bacteria could also be used to image degradation events in host phagocytes. Bioorthogonal non-canonical amino acid tagging (BONCAT)^{14, 15} for pan-proteomic incorporation of bioorthogonal groups^{16, 17} would allow the labelling of a wide range of bacterial species without the need for genetic modification¹⁸. Furthermore, unlike reporter proteins, bioorthogonal groups, such as azides^{19, 20} have been shown to be stable in the harsh chemical environments of the phagolysosomal system and should therefore be detectable when extensive proteolysis has occurred.




Figure 2 - Optimizing CLEM imaging of azido-*E. coli*: (a) azido (i) and unlabelled wild-type *E. coli* were mixed in non-equal ratio (1:25) (ii). After Tokuyasu sample preparation and cryosectioning into 75 nm sections (iii), a cChC-reaction with AlexaFluor-488 was performed (iv). (b) CLEM-image of the above experiment. Green: AlexaFluor-488 label. Dotted circles: signal from the 100 nm TetraSpeck beads. Scale bar 1 μ m.

Information about subcellular localization is of key importance when studying parasite-phagocyte-interactions as movement between organelles may be key to the life cycle of certain parasites^{1, 21}. Only transmission electron microscopy (TEM)-based techniques allows the study of these pathogens in their subcellular context, as it provides substructural information on the position of any label/antigen within the cell²². However, in contrast to superresolution imaging^{23, 24}, no methods have been reported that allow the visualization of bioorthogonal groups using EM-based approaches²⁵.

Here we describe the development of a correlative light-electron microscopy (CLEM)-imaging-based visualisation of bioorthogonal groups that allows imaging of BONCAT-labelled bacteria inside phagocytes (Fig. 1); even as they are being degraded. This approach combines the benefits of confocal microscopy – which allows wide-field navigation to areas of interest²⁶ – with those of electron microscopy (EM) – which

provides narrow field high resolution information about the interior of the cell²². All approaches described here on the model organism *E. coli* are amenable to application to pathogens, which would open new avenues for studying the events leading to bacterial clearance and/or establishment of intracellular residence by intracellular pathogens.

Results and discussion

Optimizing the detection of bioorthogonal groups using CLEM

To develop the CLEM-based imaging of bioorthogonal groups, we initially focussed on azides as our bioorthogonal group of choice, as these are the most versatile of the groups in terms of both incorporation methods and available detection reactions^{27, 28}. We used the Met-auxotrophic strain B834(DE3) of *E. coli* as our model bacterium, as it can be readily labelled with azides by metabolic replacement of methionine with azidohomoalanine (Aha)^{16, 29, 30}. However, these labelling strategies can be translated to many other non-auxotrophic bacteria and eukaryotes very effectively¹⁸.

We first optimised Aha incorporation with respect to cell viability, and incorporation levels into the proteome (Fig. S1). It was observed that extended incubation times resulted in reduced viability (Fig. S2) and the formation of inclusion bodies (Fig. S3). This suggests detrimental effects on protein expression and folding of prolonged exposure to Aha^{18, 31}. Labelling for 1h in presence of 4 mM Aha gave robust signal and showed minimal inhibition of viability (Fig. S2). These conditions were used for all further imaging studies.

We focussed our initial development of the on-section labelling of CLEM-samples on cryosections prepared in accordance with Tokuyasu³², as this technique uses mild fixation and sample preparation techniques compatible with azide-chemistry. We prepared on sections of mixed azido-*E. coli* and unlabelled wt-*E. coli* controls at a non-equal ratio for use in optimising bioorthogonal on-section labelling (Fig. 2, S4).

Of the three available azide-specific bioorthogonal reactions available we focussed on optimising the copper-catalysed Huisgen cycloaddition (cChC)-reaction^{8, 33, 34} as it has the lowest background reactivity compared to the strain-promoted cycloaddition reactions, and Staudinger-Bertozzi ligation^{35, 36}. Downsides are largely associated with the toxicity of the catalyst²⁸. However, this is not of relevance here as the labelling is performed after aldehyde-fixation and cryosectioning of the samples. We found that glutaraldehyde-free fixation, followed by a blocking step before the cChC-reaction, combined with ligand-stabilisation of the Cu(I) catalytic species gave the best signal-to-noise levels. The addition of aminoguanidine during the cChC reduced the detrimental effects of the ascorbate by-products on the DAPI co-staining³⁷.

It was also found that care had to also be taken when using copper mounting grids on which the ultrathin sections were placed for CLEM-imaging.





Figure 3 - Confocal microscopy of (a) azido-*E. coli* or (b) GFP-*E. coli* after phagocytosis. BM-DCs were pulsed with either azido-*E. coli* or GFP-expressing *E. coli* (45 min pulse). Cells were fixed after a 2 hour chase and stained with DAPI (Blue), phalloidin-647 (Red) and, in case of azido-*E. coli*, AlexaFluor-488-alkyne (green = either GFP or AlexaFluor-488). i) DAPI/488nm overlay; ii) DAPI only; iii) all fluorescent channels overlay. Yellow arrows indicate a 488-single positive focus, white arrows a DAPI/488nm double positive focus.

If high concentrations of ascorbate and prolonged reaction times were used for the ccHc, we found that the grids dissolved. This could be prevented by shortening reaction times and keeping the ascorbate concentrations low (<10 mM).

After ccHc-labelling, the sections were first imaged with the confocal microscope (Fig. S4a) before embedding in methyl cellulose with uranyl acetate. The sections were EM imaged (Fig. S4b) and correlation of the confocal and EM images was performed using electron-dense fluorescent electron-dense beads³⁸ as fiducials (Fig. 2b and Fig. S4c-e).

Comparison of GFP-*E. coli* and azido-*E. coli* for imaging phagolysosomal degradation

Most CLEM studies employ the fusion of fluorescent proteins to the protein of interest or antibody-based approaches to allow their identification and localisation²². These labelling approaches have shown to be of great value for the imaging of specific proteins in their cellular context, but only in the cases where genetic modification of the organism has been possible and where the attachment of the fluorescent proteins does not affect protein function³⁹.



(Previous Page) Figure 4 - CLEM imaging of phagocytosed azido-*E. coli*: BM-DCs were pulsed with Azido-*E. coli* (45 min pulse). Cells were washed with PBS to remove unbound/non-internalized *E. coli*. Samples were fixed immediately after pulsing (a-d) or after a 3 h chase (e-h). Cells were subjected to Tokuyasu sample preparation and cryosectioned into 150 nm sections. Sections were reacted with AlexaFluor 488 alkyne using cHc-conditions (green), anti-LAMP1 (red) and DAPI (blue). DAPI staining and blue fiducials (indicated with circles in b and f) were used for correlation purposes. (a/e) confocal microscopy images; (b/f) CLEM image obtained from overlay LM and EM pictures; (c,d,g,h) CLEM details from b/f, showing LAMP-1 and 488nm channels (c/g) or 488 nm alone (d/h). Scale bar 500 nm.

Immunofluorescence has also been used, but combined with CLEM it either compromises ultrastructure (by virtue of the need of fixation and permeabilisation prior to CLEM-sample preparation)⁴⁰, or suffers from a notoriously low success rate due to compromised epitope availability in samples prepared for TEM⁴¹.

As our main envisaged application of bioorthogonal CLEM imaging was in detecting phagocytosed bacteria during degradation (Fig. 1), we first determined whether our BONCAT-based approach had advantages over genetic methods for the study of these events. The fate of azido-*E. coli* was compared to that of GFP-expressing *E. coli* (Fig. 3, S5, and S6). We incubated mouse bonemarrow-derived dendritic cells (BM-DCs)^{42, 43} with azido-*E. coli* or GFP-*E. coli* for 45 minutes. After washing, the cells were chased for 1h, 2h or 3h prior to fixation, bioorthogonal modification of the azides (where present), and confocal imaging (Fig. 3, S5) – time points in which maturation of a phagosome to a phagolysosome are known to take place in these cells⁴⁴.

To assess whether the fluorescent signal originated from an intact or (partially) degraded bacteria, we made use of extra-nuclear DAPI staining: colocalisation of the fluorescent signal with the extra-nuclear DAPI indicates the intactness of the bacterial DNA, which in turn indicates the intactness of the bacterium⁴⁵. Absence of this colocalisation (i.e. 488nm single positive foci) indicated the degradation of the bacterial genome and thus death.

The azide-based signal persisted significantly more than the GFP-signal after the killing of the bacterium; as indicated by the significantly larger number of DAPI-negative/azide-positive foci at all time points of the chase period compared to DAPI-negative/GFP-positive foci (Fig. S6). Many of the azide-positive foci were smaller than intact DAPI/azide double positive foci, indicating these signals to originate from partially degraded bacteria. Control samples of wt-*E. coli* were fully negative at all time points (Fig. S5).

CLEM-imaging of azido-*E. coli* after uptake by BM-DCs

We obtained ultrastructural information about the location of these smaller, DAPI-negative foci by performing CLEM analysis on azido-*E. coli*-treated BM-DCs samples at all four time points (Fig. 4 and S7). Co-staining with the lysosomal marker LAMP-1 revealed that these degraded fragments only partially resided in LAMP-positive late endosomes/lysosomes. This ties in with previous studies showing the existence of a second population of phagosomes⁴⁶ in DCs, which do not acidify and never

become LAMP-1 positive⁴⁶. This set of phagosomes has been implicated in DC-specific functions such as cross-presentation^{47, 48}.

Morphological information obtained from TEM showed that the azide-positive/DAPI-positive foci were intact bacteria, whereas the DAPI-negative foci showed no identifiable bacterial morphology, indicating that this technique allows the imaging of partially degraded bacteria inside mammalian phagocytes.

Application of CLEM to other bioorthogonal labelling strategies

The above approach highlights the potential of CLEM-imaging in detecting BONCAT-labelled bacteria inside phagocytes. However, bioorthogonal chemistry offers a much broader arena of applications to interrogate specific physiological phenomena²⁷. Two stalwart applications are the labelling of glycans in mammalian cells⁴⁹ and activity-based protein profiling⁵⁰. To firmly embed our technique within the bioorthogonal chemistry paradigm, we also applied CLEM-imaging in these settings.




Figure 5 - CLEM imaging of bioorthogonal tagged sialylated glycans in Jurkat cells. Green: AlexaFluor 488 alkyne; Blue: DAPI. DAPI staining was used for correlation purposes. (a) confocal microscopy images; (b) EM image; (c) CLEM image obtained from overlay LM and EM pictures; (d) CLEM detail from c of intracellular region; (e) CLEM detail from c of plasma membrane. Scale bar 2 μ m

We first recapitulated the archetypal imaging of sialic acid-containing glycans on mammalian cells⁴⁹, by imaging Jurkat cells incubated with Ac₄-N-azidoacetylmannosamine (Ac₄ManNAz; Fig. S8), an approach that results in cells with a bioorthogonally labelled sialoglycome⁵¹. We subjected these cells to the bioorthogonal labelling and imaging conditions as above and found that LM imaging revealed cell surface and intracellular staining as previously observed by Baskin *et al.*⁵² (Fig. 5a, S9). Subsequent correlation with EM-imaging of the same section of the cell (Fig. 5b-e) revealed the labelled sialic

acids to reside in the Golgi apparatus (Fig. 5d) and the plasma membrane (Fig. 5e). Incubation with unlabelled control (Ac₄ManNAc) gave no fluorescent labelling (Fig. 9a).

We next combined CLEM with two-step activity-based protein profiling^{53,54}. This technique has proven powerful for detecting active sub-populations of e.g. serine hydrolases⁵⁰, cysteine proteases^{55,56} and proteasomes⁵⁷ in a fashion whereby in the first step enzymes are covalently and irreversibly inhibited, after which the reporter moiety is ligated in a second step through bioorthogonal chemistry. One example that was of particular interest to us^{58,59} were the endo-lysosomal cysteine protease family: the cathepsins^{55,60} as these enzymes are not only covalently labelled by a series of analogues of the natural product E64⁶¹ and their activity is also restricted to the endo-lysosomal pathway⁶² (although their extracellular activity has also been reported⁶³). If their intracellular activity could be pinpointed to LAMP-1 negative vesicles in DCs, this would provide further visual evidence for the degradation of bacteria in LAMP-1-negative phagosomes observed earlier.

As such, their imaging using an azide-modified variant of E64 will offer an 'internal standard' of the selectivity of this approach: the signal of the enzyme is expected to be contained only within membrane-limited vesicles. Furthermore, it should in part overlap with the lysosomal marker LAMP-1.

We incubated BM-DCs with azide-modified E64 using conditions described⁶⁴ and after extensive washing (the 2-step probe is cell permeable and unbound ABP can therefore be removed prior to bioorthogonal labelling and imaging), the cells were fixed, sectioned and subjected to bioorthogonal labelling. The resulting fluorescent signal was found to be fully contained in membrane-limited structures (Fig. 6, S10) and indeed co-staining with the anti-LAMP-1 antibody revealed a population of vesicles negative for LAMP but with robust cathepsin activity.

These two examples show that CLEM-imaging of bioorthogonal handles is a versatile technique that can be used over a wide range of applications. In future, it will be interesting to see how the structural information will impact the application of activity-based probes for other enzymes and enzyme families, such as the aforementioned serine hydrolases⁶⁵, and the recently reported protease-specific probes⁶⁶.

Conclusions

By combining BONCAT of bacteria with CLEM-imaging, we have established a new approach that allowed us to visualise bioorthogonally modified bacteria in an ultrastructural cellular context, even during late stages of bacterial degradation. This approach is of great interest for the study of obligate intracellular parasites that are very hard to study by any other means. As the application of bioorthogonal chemistry is ever expanding, the CLEM-imaging method of bioorthogonal groups described here could also be of great benefit to the study of

labelled biomolecules in other fields in which bioorthogonal imaging has proven its value²⁵.




Figure 6 - CLEM imaging of active populations of cyteine proteases in BM-DCs. BM-DCs were incubated for 2h with 10 μ M DCG-04-azide (a-h). Cells were fixed in 2% PFA, subjected to Tokuyasu sample preparation and cryosectioned into 150 nm sections. Sections were reacted with AlexaFluor 488 alkyne using cChc- conditions (green), anti-LAMP1 (red) and DAPI (blue). DAPI staining and blue fiducials were used for correlation purposes. (a) confocal microscopy images; (b) CLEM image obtained from overlay LM and EM pictures; (c,d,) CLEM details from b, showing LAMP1, AlexaFluor 488 and DAPI (c) or only AlexaFluor 488 (d). Scale bar 1 μ m.

Here we demonstrate this value by imaging glycans on the surface and in the Golgi of mammalian cells and by imaging active sub-populations of cathepsins inside the endo-lysosomal pathway. Application of this approach to other bioorthogonal assays (for instance, lipid imaging⁶⁷ and the imaging of newly synthesized proteins⁶⁸), and perhaps using some of the more recently developed bioorthogonal chemistries⁶⁹ will allow the provision of additional structural information to the current imaging methods available for these types of groups.

Acknowledgements

SvK was funded by a Starting Grant from the European Research Council (ERC-Grant number 639005). We thank Ramon A. van der Valk and Remus T. Dame for providing the GFP-plasmid.

Notes and references

- D. M. Walker, S. Oghumu, G. Gupta, B. S. McGwire, M. E. Drew and A. R. Satoskar, *Cellular and molecular life sciences : CMLS*, 2014, 71, 1245-1263.
- M. B. Johnson and A. K. Criss, *Journal of Visualized Experiments : JoVE*, 2013, DOI: 10.3791/50729, 50729.
- R. W. Truman and J. L. Krahenbuhl, *Int J Lepr Other Mycobact Dis*, 2001, 69, 1-12.
- O. Lamrabet and M. Drancourt, *Tuberculosis (Edinb)*, 2012, 92, 365-376.
- O. V. Vieira, R. J. Botelho and S. Grinstein, *Biochem J*, 2002, 366, 689-704.
- H. Katayama, A. Yamamoto, N. Mizushima, T. Yoshimori and A. Miyawaki, *Cell Structure and Function*, 2008, 33, 1-12.
- R. K. Lim and Q. Lin, *Chem. Commun.*, 2010, 46, 1589-1600.
- H. C. Kolb, M. G. Finn and K. B. Sharpless, *Angew. Chem. Int. Ed. Engl.*, 2001, 40, 2005-2021.
- E. M. Sletten and C. R. Bertozzi, *Angew Chem Int Ed Engl*, 2009, 48, 6974-6998.
- C. P. Ramil and Q. Lin, *Chemical communications (Cambridge, England)*, 2013, 49, 11007-11022.
- G. W. Liechti, E. Kuru, E. Hall, A. Kalinda, Y. V. Brun, M. VanNieuwenhze and A. T. Maurelli, *Nature*, 2014, 506, 507-510.
- P. Shieh, M. S. Siegrist, A. J. Cullen and C. R. Bertozzi, *Proceedings of the National Academy of Sciences*, 2014, 111, 5456-5461.
- M. S. Siegrist, S. Whiteside, J. C. Jewett, A. Aditham, F. Cava and C. R. Bertozzi, *ACS Chemical Biology*, 2012, 8, 500-505.
- D. C. Dieterich, A. J. Link, J. Graumann, D. A. Tirrell and E. M. Schuman, *Proc. Natl. Acad. Sci. U.S.A.*, 2006, 103, 9482-9487.
- P. Landgraf, E. R. Antileo, E. M. Schuman and D. C. Dieterich, *Methods Mol Biol*, 2015, 1266, 199-215.
- K. L. Kiick, E. Saxon, D. A. Tirrell and C. R. Bertozzi, *Proc. Natl. Acad. Sci. U.S.A.*, 2002, 99, 19-24.
- K. L. Kiick and D. A. Tirrell, *Tetrahedron*, 2000, 56, 9487-9493.
- R. Hatzenpichler, S. Scheller, P. L. Tavormina, B. M. Babin, D. A. Tirrell and V. J. Orphan, *Environmental microbiology*, 2014, 16, 2568-2590.
- J. B. Pawlak, G. P. P. Gentil, T. J. Ruckwardt, J. S. Bremmers, N. J. Meeuwenoord, F. A. Ossendorp, H. S. Overkleeft, D. V. Filippov and S. I. van Kasteren, *Angew. Chem. Int. Ed. Engl.*, 2015, 54, 5628-5631.
- S. P. Ouellette, F. C. Dorsey, S. Moshich, J. L. Cleveland and R. A. Carabeo, *PLoS One*, 2011, 6, e16783.
- N. van der Wel, D. Hava, D. Houben, D. Fluitsma, M. van Zon, J. Pierson, M. Brenner and P. J. Peters, *Cell*, 2007, 129, 1287-1298.
- P. de Boer, J. P. Hoogenboom and B. N. G. Giepmans, *Nat Meth*, 2015, 12, 503-513.
- I. Nikić, T. Plass, O. Schraidt, J. Szymański, J. A. G. Briggs, C. Schultz and E. A. Lemke, *Angew. Chem. Int. Ed. Engl.*, 2014, 53, 2245-2249.
- R. S. Erdmann, H. Takakura, A. D. Thompson, F. Rivera-Molina, E. S. Allgeyer, J. Bewersdorf, D. Toomre and A. Schepartz, *Angew Chem Int Ed Engl*, 2014, 53, 10242-10246.
- D. M. van Elsland, E. Bos, H. S. Overkleeft, A. J. Koster and S. I. van Kasteren, *J Chem Biol*, 2015, DOI: 10.1007/s12154-015-0134-4, 1-5.
- B. N. G. Giepmans, *Histochem. Cell Biol.*, 2008, 130, 211-217.
- E. M. Sletten and C. R. Bertozzi, *Accounts Chem. Res.*, 2011, 44, 666-676.
- C. P. Ramil and Q. Lin, *Chem Commun*, 2013, 49, 11007-11022.
- D. J. Leahy, W. A. Hendrickson, I. Aukhil and H. P. Erickson, *Science*, 1992, 258, 987-991.
- W. B. Wood, *J Mol Biol*, 1966, 16, 118-133.
- A. K. Upadhyay, A. Murmu, A. Singh and A. K. Panda, *PLoS One*, 2012, 7, e33951.
- K. T. Tokuyasu, *J Cell Biol*, 1973, 57, 551-565.
- C. W. Tornøe, C. Christensen and M. Meldal, *J. Org. Chem.*, 2002, 67, 3057-3064.
- V. V. Rostovtsev, L. G. Green, V. V. Fokin and K. B. Sharpless, *Angew Chem Int Ed Engl*, 2002, 41, 2596-2599.
- M. Lo Conte, S. Staderini, A. Marra, M. Sanchez-Navarro, B. G. Davis and A. Dondoni, *Chem Commun (Camb)*, 2011, 47, 11086-11088.
- L. I. Willems, N. Li, B. I. Florea, M. Ruben, G. A. van der Marel and H. S. Overkleeft, *Angewandte Chemie-International Edition*, 2012, 51, 4431-4434.
- V. Hong, N. F. Steinmetz, M. Manchester and M. G. Finn, *Bioconj. Chem.*, 2010, 21, 1912-1916.
- W. Kukulski, M. Schorb, S. Welsch, A. Picco, M. Kaksonen and J. A. Briggs, *J Cell Biol*, 2011, 192, 111-119.
- C. L. Thomas and A. J. Maule, *J Gen Virol*, 2000, 81, 1851-1855.
- U. Schnell, F. Dijk, K. A. Sjollem and B. N. G. Giepmans, *Nature Methods*, 2012, 9, 152-158.
- G. Griffiths and J. M. Lucocq, *Histochem Cell Biol*, 2014, 142, 347-360.
- M. B. Lutz, N. Kukutsch, A. L. J. Ogilvie, S. Rößner, F. Koch, N. Romani and G. Schuler, *J. Immunol. Meth.*, 1999, 223, 77-92.
- M. A. West, R. P. A. Wallin, S. P. Matthews, H. G. Svensson, R. Zaru, H. G. Ljunggren, A. R. Prescott and C. Watts, *Science*, 2004, 305, 1153-1157.

44. L.-P. Erwig, K. A. McPhillips, M. W. Wynes, A. Ivetic, A. J. Ridley and P. M. Henson, *Proc. Natl. Acad. Sci. U.S.A.*, 2006, 103, 12825-12830.
45. E.-Y. Seo, T.-S. Ahn and Y.-G. Zo, *Applied and Environmental Microbiology*, 2010, 76, 1981-1991.
46. A. Savina, C. Jancic, S. Hugues, P. Guermonprez, P. Vargas, I. C. Moura, A.-M. Lennon-Duménil, M. C. Seabra, G. Raposo and S. Amigorena, *Cell*, 2006, 126, 205-218.
47. I. Cebrian, G. Visentin, N. Blanchard, M. Jouve, A. Bobard, C. Moita, J. Enninga, L. F. Moita, S. Amigorena and A. Savina, *Cell*, 2011, 147, 1355-1368.
48. O. P. Joffre, E. Segura, A. Savina and S. Amigorena, *Nature Reviews Immunology*, 2012, 12, 557-569.
49. E. Saxon and C. R. Bertozzi, *Science*, 2000, 287, 2007-2010.
50. A. E. Speers, G. C. Adam and B. F. Cravatt, *J. Am. Chem. Soc.*, 2003, 125, 4686-4687.
51. R. M. Werner, L. M. Williams and J. T. Davis, *Tetrahedron Lett.*, 1998, 39, 9135-9138.
52. J. M. Baskin, J. A. Prescher, S. T. Laughlin, N. J. Agard, P. V. Chang, I. A. Miller, A. Lo, J. A. Codelli and C. R. Bertozzi, *Proc. Natl. Acad. Sci. U.S.A.*, 2007, 104, 16793-16797.
53. B. F. Cravatt, A. T. Wright and J. W. Kozarich, *Annu Rev Biochem*, 2008, 77, 383-414.
54. L. I. Willems, H. S. Overkleeft and S. I. van Kasteren, *Bioconj. Chem.*, 2014, 25, 1181-1191.
55. D. Greenbaum, K. F. Medzhradszky, A. Burlingame and M. Bogoy, *Chem Biol*, 2000, 7, 569-581.
56. L. E. Sanman and M. Bogoy, *Annu Rev Biochem*, 2014, 83, 249-273.
57. H. Ovaa, P. F. van Swieten, B. M. Kessler, M. A. Leeuwenburgh, E. Fiebiger, A. M. van den Nieuwendijk, P. J. Galardy, G. A. van der Marel, H. L. Ploegh and H. S. Overkleeft, *Angew Chem Int Ed Engl*, 2003, 42, 3626-3629.
58. S. I. van Kasteren, I. Berlin, J. D. Colbert, D. Keane, H. Ovaa and C. Watts, *ACS chemical biology*, 2011, 6, 1198-1204.
59. S. I. van Kasteren and H. S. Overkleeft, *Curr. Opin. Chem. Biol.*, 2014, 23, 8-15.
60. V. Turk, V. Stoka, O. Vasiljeva, M. Renko, T. Sun, B. Turk and D. Turk, *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics*, 2012, 1824, 68-88.
61. A. J. Barrett, A. A. Kembhavi, M. A. Brown, H. Kirschke, C. G. Knight, M. Tamai and K. Hanada, *Biochem J*, 1982, 201, 189-198.
62. C. Watts, *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics*, 2012, 1824, 14-21.
63. J. S. Mort, A. D. Recklies and A. R. Poole, *Arthritis & Rheumatism*, 1984, 27, 509-515.
64. S. Hoogendoorn, G. H. van Puijvelde, J. Kuiper, G. A. van der Marel and H. S. Overkleeft, *Angew Chem Int Ed Engl*, 2014, 53, 10975-10978.
65. Y. Liu, M. P. Patricelli and B. F. Cravatt, *Proc Natl Acad Sci U S A*, 1999, 96, 14694-14699.
66. L. E. Edgington, M. Verdoes and M. Bogoy, *Curr. Opin. Chem. Biol.*, 2011, 15, 798-805.
67. H. C. Hang, J. P. Wilson and G. Charron, *Acc Chem Res*, 2011, 44, 699-708.
68. K. E. Beatty, J. C. Liu, F. Xie, D. C. Dieterich, E. M. Schuman, Q. Wang and D. A. Tirrell, *Angew. Chem. Int. Ed. Engl.*, 2006, 45, 7364-7367.
69. K. Lang and J. W. Chin, *ACS Chem Biol*, 2014, 9, 16-20.