

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Observation of Electrochemically Generated Nitrenium Ions by Desorption Electrospray Ionization Mass Spectrometry

Timothy A. Brown^a, Niloufar Hosseinei Nassab^a, Hao Chen^{*b}, and Richard N. Zare^{*a}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

We report the observation of the electrochemically generated nitrenium ions of 4,4'-dimethoxydiphenylamine and di-p-tolylamine in solution by mass spectrometry. This setup takes inspiration from desorption electrospray ionization mass spectrometry to sample directly from the surface of a rotating waterwheel working electrode for mass spectrometric analysis. Detection of the 4,4'-dimethoxydiphenylamine nitrenium ion was expected based upon para-methoxy resonance stabilization, whereas observation of the di-p-tolylamine nitrenium ion might be unexpected because resonance stabilization from the para-substituted position is unavailable. However, the short timescale analysis of the setup allows for the isolation of the di-p-tolylamine nitrenium ion, which is electrogenerated in solution and detected mass spectrometrically.

Introduction

Nitrenium ions are reactive cation intermediates that are isoelectronic to carbenes.^{1–3} Understanding nitrenium ions impacts not only organic synthesis but also has biological significance due to their reactivities toward nucleobases.^{4–7} Nitrenium ions can be formed in a number of ways including chemically,^{4,8} photochemically,^{9–11} and electrochemically.^{12–15} Arylamines have been studied as model systems for formation of the fleeting nitrenium species in solution.^{8–14,16} Mass spectrometry (MS) has been utilized to observe nitrenium ions before via ion/molecule reactions,^{4,8,17,18} however the mass spectrometric characterization of these arylamine nitrenium intermediates from electrochemical generation has been scarce outside of the observation of the nitrenium ion of clozapine.^{7,15,19} Herein we present the first application of desorption electrospray ionization (DESI)^{20,21} MS in detecting nitrenium ions generated electrochemically from arylamines.

Experimental

The experimental design in Figure 1 greatly resembles the waterwheel setup previously reported,^{22,23} which employs a round rotating platinum working electrode immersed in an




Figure 1. Top: The setup used for detection of electrochemical reaction intermediates, employing a rotating waterwheel electrode. Wheel rotation is indicated by the curved arrow.

acetonitrile solution containing 1 mM lithium triflate as the electrolyte. The distance between the MS inlet and the working electrode surface is approximately 2 mm. As the working electrode rotates, a thin layer of liquid film develops on the electrode surface (approximately 1 mm in thickness). A plain carbon cloth counter electrode and an Ag/AgCl reference electrode are immersed in the reservoir of electrolyte solution. A metal contact (not shown) rests against the platinum working electrode to complete the three-electrode system. A potentiostat (WaveNow, Pine Research Instrumentation, Durham, NC) is used to apply a potential across the three electrodes.

Above the rotating waterwheel system a custom spray probe directs a stream of sample microdroplets to the surface of the working electrode. Studies have shown that electrospray ionization and DESI sources can function as an electrochemical cell^{24–27} and so no high voltage is applied to the sample spray to minimize in-source oxidation. The spray droplets hit the surface of the thin film of electrolyte solution, now thinner than 1 mm due to the N₂ nebulizing gas pressure,

^a Department of Chemistry, Stanford University, Stanford, CA 94305-5080, USA
Email: zare@stanford.edu

^b Center for Intelligent Chemical Instrumentation, Department of Chemistry and Biochemistry and Edison Biotechnology Institute, Ohio University, Athens, OH 45701-2979, USA. Email: chen2@ohio.edu

^c Electronic Supplementary Information (ESI) available: MS/MS data for structural confirmation and experimental procedure details. See DOI: 10.1039/x0xx00000x

on the surface of the working electrode. Much like in DESI-MS and easy ambient supersonic spray^{28–30} MS, tinier secondary microdroplets are directed into the mass spectrometer^{31,32} and are analyzed with an LTQ Orbitrap XL hybrid mass spectrometer (Thermo Fisher Scientific, San Jose, CA) where the m/z ratio is determined utilizing the high mass accuracy and high resolving power of the Orbitrap mass analyzer.³³

Results and discussion

Electrogeneration of DMDPA nitrenium ion

4,4'-dimethoxydiphenylamine (DMDPA) was chosen as a model system because nitrenium formation from DMDPA has been extensively studied.^{14,16} DMDPA is proposed to be electrochemically oxidized to the nitrenium ion by losses of two electrons and one proton, passing through a radical cation intermediate (Figure 2). In this experiment, 100 μM of DMDPA was prepared in 1 mM solution of lithium triflate (LiOTf) in acetonitrile. When the analyte solution is sprayed at an injection flow rate of 10 $\mu\text{L}/\text{min}$ onto the working electrode rotating at 1.0 rev/s, m/z 230.1175 is observed, ascribed to the protonated DMDPA cation (Figure 3a, theoretical m/z 230.1176, error -0.2 ppm) as well as m/z 229.1098, attributed to the DMDPA radical cation (Figure 3a, theoretical m/z 229.1097, error 0.0 ppm). When an oxidation potential of 1.5 V is applied across the rotating working electrode the intensity of 229.1097 increases almost ten-fold (Figure 3b), suggesting that the DMDPA radical cation is formed through electrochemical oxidation of DMDPA. A peak at m/z 228.1020 is also observed, which is ascribed to the DMDPA nitrenium ion (Figure 3b, theoretical m/z 228.1019, error $+0.5$ ppm). Upon CID, the DMDPA nitrenium ion gives rise to fragment ions of m/z 213 and 197, corresponding to the losses of CH_3 and CH_3O radicals, respectively, consistent with its structure (Figure S3, Supporting information). The MS signals of the DMDPA radical cation and DMDPA nitrenium ion increase greatly when an oxidizing potential is applied to the working electrode, as is observed in the extracted ion chromatograms (EIC) at specific m/z values (Figure 3c, 3d). This indicates that these species are formed via electrochemical oxidation of DMDPA and is the first electrogenerated nitrenium ion observed with DESI-MS.

Electrogeneration of DPTA nitrenium ion

The electrochemical oxidation of di-*p*-tolylamine (DPTA) was also studied. DPTA is proposed to undergo nitrenium formation by the same pathway as DMDPA (Figure 2), but each system lacks the para-methoxy groups that can stabilize both the radical cation and nitrenium species through resonance. Because of this the half-life time of these species in solution is estimated to be shorter compared to corresponding DMDPA intermediates.

100 μM of DPTA was prepared in 1 mM solution of LiOTf in acetonitrile. When the analyte solution is sprayed at an injection flow rate of 10 $\mu\text{L}/\text{min}$ onto the working electrode rotating at 1.0 rev/s, a mass peak at m/z 198.1275 is observed,




Figure 2. The general scheme for electrochemical generation of nitrenium ions from secondary arylamines.

ascribed to the protonated DMDPA cation (Figure 4a, theoretical m/z 198.1277, error -0.9 ppm). When an oxidation potential of 3.0 V is applied across the rotating working electrode the emergence of m/z 197.1199, corresponding to the DPTA radical cation (Figure 4b, theoretical m/z 197.1199 error 0.0 ppm), is suggested to arise from electrochemical oxidation of DPTA. The increase in the signal intensity of 196.1121, believed to be the DPTA nitrenium ion (Figure 4b, theoretical m/z 196.1121, error 0.0 ppm) is also inferred to




Figure 3. Positive-ion mode mass spectra of DMDPA. a) 0.0 V applied to the working electrode; b) 1.5 V applied to the working electrode; c) EIC for the 229.1097 peak as a function of the applied potential; and d) EIC for the 228.1020 peak as a function of the applied potential.

stem from the electrochemical oxidation of DPTA. For DPTA, the para-methyl groups provide hydrogens that are available for elimination, leading to the formation of a constitutional isomer of the desired amine. Performing the analysis on the analogous d-14 labelled DPTA determined that both the nitrenium ion and the elimination product are formed as a function of an oxidative potential and is discussed further in the Electronic Supplementary Information.




Figure 4. Positive-ion mode mass spectra of DPTA. a) 0.0 V applied to the working electrode; b) 3.0 V applied to the working electrode; c) EIC for the 197.1199 peak as a function of the applied potential; and d) EIC for the 196.1121 peak as a function of the applied potential.

Upon CID, the DPTA nitrenium ion gives rise to a fragment ion of m/z 181 by loss of a CH_3 radical, consistent with its structure. The MS signals of the DPTA radical cation and DPTA nitrenium ion increase greatly when an oxidizing potential is applied to the working electrode, as is observed in the extracted ion chromatograms (Figure 4c, 4d). This observation indicates that these species are formed via electrochemical oxidation of DPTA.

Conclusions

The results presented here show evidence for the electrochemical generation of nitrenium ions through the proposed radical cation intermediate. The short timescale of electrogeneration and isolation allows for the observation of the reactive intermediates. This study emphasizes the power of DESI-MS as an analytical tool in identifying reactive intermediates in electrochemical processes.

Acknowledgements

T. A. Brown thanks the Center for Molecular Analysis and Design (CMAD) for a graduate fellowship. H. Chen is grateful to the support from NSF Career Award (CHE-1149367), NSF (CHE-1455554) for his visit to the Zare laboratory. This work is supported by a subcontract with the University of Utah (Agreement # 10029173-S2) for which the Air Force Office of Scientific Research (Grant FA9550-12-1-0481) is the prime sponsor.

Notes and references

- 1 D. E. Falvey, in *Reactive Intermediate Chemistry*, eds. R. A. Moss, M. S. Platz and M. J. Jr, John Wiley & Sons, Inc., 2003, pp. 593–650.
- 2 P. G. Gassman, *Acc. Chem. Res.*, 1970, **3**, 26–33.
- 3 G. I. Borodkin and V. G. Shubin, *Russ. Chem. Rev.*, 2008, **77**, 395.
- 4 Z. C. Liu and J. P. Uetrecht, *J. Pharmacol. Exp. Ther.*, 1995, **275**, 1476–1483.
- 5 J. L. Maggs, D. Williams, M. Pirmohamed and B. K. Park, *J. Pharmacol. Exp. Ther.*, 1995, **275**, 1463–1475.
- 6 D. P. Williams, M. Pirmohamed, D. J. Naisbitt, J. L. Maggs and B. K. Park, *J. Pharmacol. Exp. Ther.*, 1997, **283**, 1375–1382.
- 7 W. Lohmann, H. Hayen and U. Karst, *Anal. Chem.*, 2008, **80**, 9714–9719.
- 8 H. Chen, H. Chen, R. G. Cooks and H. Bagheri, *J. Am. Soc. Mass Spectrom.*, 2004, **15**, 1675–1688.
- 9 D. Chiapperino, S. Mcllroy and D. E. Falvey, *J. Am. Chem. Soc.*, 2002, **124**, 3567–3577.
- 10 R. J. Moran, C. Cramer and D. E. Falvey, *J. Org. Chem.*, 1997, **62**, 2742–2751.
- 11 S. I. Thomas and D. E. Falvey, *J. Org. Chem.*, 2007, **72**, 4626–4634.
- 12 A. Rieker and B. Speiser, *J. Org. Chem.*, 1991, **56**, 4664–4671.
- 13 E. M. Genies and M. Lapkowski, *J. Electroanal. Chem. Interfacial Electrochem.*, 1987, **236**, 189–197.
- 14 D. Serve, *J. Am. Chem. Soc.*, 1975, **97**, 432–434.
- 15 S. M. van Leeuwen, B. Blankert, J.-M. Kauffmann and U. Karst, *Anal. Bioanal. Chem.*, 2005, **382**, 742–750.
- 16 U. Svanholm and V. D. Parker, *J. Am. Chem. Soc.*, 1974, **96**, 1234–1236.
- 17 M. J. Yurkovich, P. Duan, R. C. Shea, M. A. Watkins, S. M. Mandell, E. M. Tippmann, S. Jason Li, M. S. Platz and H. I. Kenttämä, *Int. J. Mass Spectrom.*, 2009, **287**, 16–20.
- 18 H. Chen, X. Zheng, P. Yang and R. G. Cooks, *Chem. Commun.*, 2004, 688–689.

- 19 F. T. G. van den Brink, L. Büter, M. Odijk, W. Olthuis, U. Karst and A. van den Berg, *Anal. Chem.*, 2015, **87**, 1527–1535.
- 20 Z. Takáts, J. M. Wiseman, B. Gologan and R. G. Cooks, *Science*, 2004, **306**, 471–473.
- 21 Z. Takáts, J. M. Wiseman and R. G. Cooks, *J. Mass Spectrom.*, 2005, **40**, 1261–1275.
- 22 T. A. Brown, H. Chen and R. N. Zare, *J. Am. Chem. Soc.*, 2015, **137**, 7274–7277.
- 23 T. A. Brown, H. Chen and R. N. Zare, *Angew. Chem. Int. Ed.*, 2015, n/a–n/a.
- 24 M. Abonnenc, L. Qiao, B. Liu and H. H. Girault, *Annu. Rev. Anal. Chem.*, 2010, **3**, 231–254.
- 25 M. Prudent, M. Méndez, C. Roussel, B. Su, N. Lion, J. S. Rossier and H. H. Girault, *Chim. Int. J. Chem.*, 2009, **63**, 283–287.
- 26 G. J. Van Berkel and V. Kertesz, in *Electrospray and MALDI Mass Spectrometry*, ed. R. B. Cole, John Wiley & Sons, Inc., 2010, pp. 75–122.
- 27 M. Benassi, C. Wu, M. Nefliu, D. R. Ifa, M. Volný and R. G. Cooks, *Int. J. Mass Spectrom.*, 2009, **280**, 235–240.
- 28 R. Haddad, R. Sparrapan, T. Kotiaho and M. N. Eberlin, *Anal. Chem.*, 2008, **80**, 898–903.
- 29 R. Haddad, H. M. S. Milagre, R. R. Catharino and M. N. Eberlin, *Anal. Chem.*, 2008, **80**, 2744–2750.
- 30 A. M. A. P. Fernandes, G. D. Fernandes, D. Barrera-Arellano, G. F. de Sá, R. D. Lins, M. N. Eberlin and R. M. Alberici, *J. Mass Spectrom.*, 2014, **49**, 335–341.
- 31 A. B. Costa and R. Graham Cooks, *Chem. Phys. Lett.*, 2008, **464**, 1–8.
- 32 A. B. Costa and R. G. Cooks, *Chem. Commun.*, 2007, 3915–3917.
- 33 R. H. Perry, R. G. Cooks and R. J. Noll, *Mass Spectrom. Rev.*, 2008, **27**, 661–699.

