

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/xxxxxxxxxx

How π back-donation quantitatively controls the CO stretching response in classical and non-classical metal carbonyl complexes[†]

Giovanni Bistoni,^{*ab} Sergio Rampino,^{*b} Nicola Scafuri,^c Gianluca Ciancaleoni,^b Daniele Zuccaccia,^d Leonardo Belpassi,^b and Francesco Tarantelli^{*ab}

Received Date

Accepted Date

DOI: 10.1039/xxxxxxxxxx

www.rsc.org/journalname

The CO stretching response upon coordination to a metal M to form $[(L)_nM(CO)]^m$ complexes (L is an auxiliary ligand) is investigated in relation to the σ donation and π back-donation components of the M–CO bond and to the electrostatic effect exerted by the ligand-metal fragment. Our analysis encompasses over 30 carbonyls, in which the relative importance of donation, back-donation and electrostatics are varied either through the ligand, in a series of $[(L)Au(CO)]^{0/+}$ gold(I) complexes, or through the metal in a series of anionic, neutral and cationic homoleptic carbonyls. Charge-displacement analysis is used to obtain well-defined, consistent measures of σ donation and π back-donation charges, as well as of the σ and π components of CO polarization. It is found that all complexes feature a comparable charge flow of σ symmetry (both in the M–CO bonding region and in the CO fragment itself), which is therefore largely uncorrelated to CO response. By contrast, π back-donation is exceptionally variable and is found to correlate tightly with the change in CO bond distance, with the shift in CO stretching frequency, and with the extent and direction (C→O or C←O) of the CO π polarization. As a result, we conclusively show that π back-donation can be an important bond component also in "non-classical" carbonyls and we actually provide the framework in which the spectroscopic data on coordinated CO can be used to extract quantitative information on the π donor properties of metal-ligand moieties.

1 Introduction

The high affinity of carbon monoxide (CO) towards metals (M) is known since the end of the nineteenth century¹ and its relevance has kept growing thereafter, both in pure^{2,3} and applied chemistry^{4,5}. This has led many chemists to study in detail the coordination bond between M and CO in metal carbonyl complexes, which is commonly described in terms of the Dewar-Chatt-Duncanson (DCD) model^{6–8}. According to this scheme, the interaction between M and CO involves the donation of electron

charge from carbon's lone pair to empty M orbitals of σ symmetry (M←CO σ donation), and a back-donation from filled M to empty CO orbitals of π symmetry (M→CO π back-donation). The effectiveness of this model for the description of the M–CO bond has been consolidated over the years by a large number of theoretical studies based on a variety of techniques, among which energy^{9,10} and charge¹¹ decomposition schemes, Natural Bond Orbitals (NBO) analysis¹², Electron Localization Function (ELF) approaches¹³ and the Quantum Theory of Atoms in Molecules (QTAIM)^{14–17}.

On the experimental side, discussions on the nature of the M–CO bond are mostly based on the analysis of the variation in the CO stretching frequency ν_{CO} (via IR spectroscopy) and bond distance r_{CO} (via X-ray crystallography) with respect to free CO ($\nu_{free-CO} = 2143 \text{ cm}^{-1}$, $r_{free-CO} = 1.12822 \text{ \AA}$). In most metal carbonyl complexes the CO bond appears weakened, i.e., the stretching frequency decreases ($\Delta\nu_{CO} = \nu_{CO} - \nu_{free-CO} < 0$) and the bond distance increases ($\Delta r_{CO} = r_{CO} - r_{free-CO} > 0$), but in a minority of (mainly late-metal cationic) complexes, which are sometimes termed "non-classical",¹⁸ the CO bond appears strengthened ($\Delta\nu_{CO} > 0$ and $\Delta r_{CO} < 0$). These differences in the CO

^a Dipartimento di Chimica, Biologia e Biotecnologie, Università di Perugia, Via Elce di Sotto 8, 06123 Perugia, Italy

^b Istituto di Scienze e Tecnologie Molecolari del CNR, Via Elce di Sotto 8, 06123 Perugia, Italy E-mail: giovanni@thch.unipg.it; srampino@thch.unipg.it; francesco.tarantelli@unipg.it


^c Institut Charles Gerhardt, Université Moltpellier 2, ENSCM 5253, cc 1501, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France

^d Dipartimento di Chimica, Fisica e Ambiente, Via del Cotonificio 108, 33100 Udine, Italy

[†] Electronic Supplementary Information (ESI) available: Additional figures and DFT-optimized XYZ geometries for all the complexes studied. See DOI: 10.1039/b000000x/

stretching response to the M–CO bond formation in metal carbonyl complexes are commonly explained in terms of the relative importance of the DCD constituents of the M–CO bond. In particular, M→CO π back-donation is represented as exerting a bond-weakening effect on CO, while M←CO σ donation is thought to act in the opposite way^{19,20}. This view relies on a molecular-orbital picture in which both the π acceptor and σ donor CO orbitals have a C–O anti-bonding character. However, while there is general agreement on the effect thus played by π back-donation, the role of σ donation has been brought into question in the last fifteen years^{21–23}. In particular, these studies suggest that the σ CO donor orbital has rather a weak bonding character and that the CO bond strengthening in “non-classical” complexes is an electrostatic effect due to the (positively charged) ligand-metal moiety, whereby the CO bonding orbitals of both σ and π symmetry are polarized in the C←O direction, thus enhancing the covalency of the CO bond.

One way to schematically depict M(CO) bonding resorts to a simple Valence Bond (VB) picture. Focusing on the M(CO) moiety of a generic $[(L)_nM(CO)]^m$ complex, three VB structures differing for the extent of π back-donation can be written:



In going from structure a) to structure b) and c), where one has zero, one and two π^* orbitals of CO engaged in back-bonding, the CO bond multiplicity goes from three to two to one. The relative weight of each structure will of course depend on the π donor properties of the specific $[(L)_nM]^m$ fragment. At the same time, the electronic structure of CO is also affected by the electric field generated by this fragment, especially in those cases when $m \neq 0$. For CO in the presence of an electric field generated, for instance, by a positively charged metal fragment (exemplified here with the symbol \oplus), three analogue VB structures can be written:



The presence of such electric field would in this case favour the triple bonded structure d) over structures e) and f) featuring a double and single bond, respectively. (An opposite effect, of course, is expected to occur when the electric field is generated by an anionic ligand-metal fragment). The DCD bonding structure and the electrostatic polarization effect may thus a priori act in different directions with different weight, so that their interplay in driving CO stretching response may be difficult to disentangle.

Still, however, carbonyl complexes showing blue shift ($\Delta\nu_{CO} > 0$) of the CO stretching frequency are most often assumed to lack back-donation from the metallic fragment^{18,24,25}. Exemplary in this respect is the set of complexes $[(L)Au(CO)]^{0/+}$ of gold(I) that have been experimentally characterized.^{24,26–31} Until last year, to our knowledge, nine gold(I) carbonyl complexes had been spectroscopically characterized: the ligand free $[Au(CO)]^+$ (observed in neon matrix²⁶) and its derivatives with ligands Cl^- ²⁷, Br^- ²⁷, CF_3^- ²⁴, CO ²⁸, Mes_3P ²⁹, $SI\text{dipp}$ ³⁰, $Idipp$ ³⁰ and $[HB(3,5-(CF_3)_2Pz)_3]^-$ ³¹, where Mes stands for 2,4,6- $Me_3C_6H_2$, $SI\text{dipp}$ for 1,3-bis(2,6-diisopropylphenyl)imidazolin-2-ylidene, $Idipp$ for 1,3-bis(2,6-diisopropylphenyl)imidazol-2-ylidene and $[HB(3,5-(CF_3)_2Pz)_3]^-$ is a fluorinated tris(pyrazol)borate ligand. All of




Fig. 1 The experimentally characterized gold(I) carbonyl complexes discussed in this work, with the observed carbonyl stretching frequency shifts, $\Delta\nu_{CO}$ (in cm^{-1}), and literature references.

them exhibit blue shift of the CO frequency and therefore are classified as “non classical”. This has been taken by some authors as a proof that the gold fragment gives no or poor back-donation^{32,33}. However, in apparent contradiction, both theoretical and experimental studies have shown that the π donor character of gold is usually far from being negligible^{34–36} (especially toward carbon monoxide³⁷) with important effects in catalysis^{38,39}. Recently, furthermore, a gold(I) complex showing $\Delta\nu_{CO} < 0$ has been fully characterized,⁴⁰ bearing a neutral *o*-carborane diphosphine (DPCb) as ancillary ligand. Such an “exception”, which is even more singular when considering that the formal positive charge should strengthen the CO bond, made the authors speak of “enhanced π back-donation” from the $[(DPCb)Au]^+$ fragment. For the reader’s convenience, an overview of the experimentally characterized systems, with the reported $\Delta\nu_{CO}$ values and reference to the original papers, is displayed in Figure 1. An additional gold(I) system, $[\{MeB[3-(Mes)Pz]_3\}Au(CO)]$, has been preliminarily reported as red-shifted in Ref.⁴¹.

The relationship of the DCD constituents of coordination bonds, determined unambiguously via charge-displacement (CD) analysis^{35,42}, with spectroscopic observables has been the subject of some of our recent work^{36,37,43}, and in the present work we have used this analysis to study systematically an extensive series of carbonyl compounds. The unique power of CD analysis lies in the fact that it provides a complete picture, across the entire molecular space, of the charge flow of σ and π character accompanying the formation of a coordination bond, and it permits a well-defined, consistent measure of the charge transfer (CT) as-

sociated both with the DCD components of the M–CO bond and with the σ and π components of the polarization⁴⁴ taking place at the CO ligand itself. As a result, as we hope we will have convinced the reader by the end of the paper, this work provides a definitive and quantitative account of the role and interplay of the DCD components of the M–CO bond and of CO polarization in driving CO stretching response to coordination.

We thus investigate the relation between Δv_{CO} and Δr_{CO} and the charge displacements of σ and π symmetry along the M–C–O axis in response to the M–CO bond formation in metal carbonyl complexes. We carry out our analysis first on an exhaustive series of 23 gold(I) carbonyls of formula $[(\text{L})\text{Au}(\text{CO})]^{0/+}$, where L is a varying auxiliary ligand (including none), which includes 8 of the experimentally characterized complexes and which is evenly partitioned between charged and neutral complexes, as well as between "classical" (CO bond elongated and frequency red-shifted) and "non-classical" (CO bond shortened and frequency blue-shifted). The choice of binary gold complexes seems to be particularly simple and useful, as it permits to isolate and study systematically the effect of the trans ligand across a wide variety of metal binding properties and electronic effects. We begin our analysis (Sec. 3.1) by studying in greater detail the two "extreme" cases of the "naked" Au^+ , $[\text{Au}(\text{CO})]^+$, which displays the experimentally largest blue-shift, and of $[(\text{DPCb})\text{Au}(\text{CO})]^+$, which is the only known case of a positively charged but significantly red-shifted gold(I) complex. Having thus highlighted the main findings, we then thoroughly confirm them by extending the study to the whole series of complexes (Sec. 3.2). To complete the work we then also investigate the role of the metal itself in driving CO response to coordination, by studying a series of homoleptic $[(\text{CO})_n\text{M}(\text{CO})]^m$ complexes, with M including Hg, Ir, Ni, Fe, Cr, Mo, Co, Ru (Section 3.3). Finally, concludes the work an ad hoc study of CO in a uniform axial electric field (Section 3.4) in order to isolate the impact of CO polarization, and of its σ and π components, on CO stretching response.

2 Methodology and computational details

In the Charge-Displacement (CD) analysis framework, a chemical bond A–B is analyzed in terms of the difference $\Delta\rho(x,y,z)$ between the electron density of the adduct AB and that of the two non-interacting fragments A and B frozen at their in-adduct geometries. A partial progressive integration of $\Delta\rho(x,y,z)$ along a suitably chosen "bond axis" z yields the so called charge-displacement function (CDF)⁴²

$$\Delta q(z) = \int_{-\infty}^z dz' \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Delta\rho(x,y,z') dx dy. \quad (1)$$

The CDF defines, at each point z , the exact amount of electron charge displaced from right to left (the direction of decreasing z) upon bond formation through a plane perpendicular to the z axis through the point z (negative CDF values indicate a charge flow in the opposite direction). If both the adduct and its constituting fragments have proper symmetry, $\Delta\rho(x,y,z)$ can be decomposed into additive components of σ and π symmetry with respect to the bond axis z (see Ref.³⁵ for further details).

All of the complexes studied in this work have general formula

$[(\text{L})_n\text{M}(\text{CO})]^m$. Since the M–CO bond is under investigation, the appropriate fragments are the ligand-metal moiety $[(\text{L})_n\text{M}]^m$ and carbon monoxide CO, and the z reference axis joins the M and C centres. For the purpose of separating the σ and π components of $\Delta\rho(x,y,z)$, we group the orbitals of adduct and fragments according to the irreducible representations of the complex (and fragments) symmetry groups, which in the present cases are either the C_{2v} group (where the A_1 representation corresponds to σ , while B_1 and B_2 correspond to π) or the C_{3v} group (A_1 corresponding to σ donation, and E_1 and E_2 to π back-donation). No CO orbital is of A_2 symmetry, therefore this representation is not relevant for the DCD analysis of the M–CO bond and is found to represent only a (minor) rearrangement internal to the ligand-metal fragment. Among the gold(I) complexes considered, $[(\text{PF}_3)_3\text{Au}(\text{CO})]^+$, $[(\text{PH}_3)_3\text{Au}(\text{CO})]^+$, $[(\text{P}(\text{CH}_3)_3)_3\text{Au}(\text{CO})]^+$, $[(\text{CF}_3)_3\text{Au}(\text{CO})]$ and $[(\text{CH}_3)_3\text{Au}(\text{CO})]$ belong to the C_{3v} point group. All others have C_{2v} symmetry. The symmetry point groups of the homoleptic complexes considered in Section 3.3 are listed in Table 2. The reduced symmetries C_{2v} and C_{3v} have been used to separate the σ and π components of the electron density difference also for these complexes.

The CDFs of the σ and π components of $\Delta\rho(x,y,z)$ provide a thorough, spatially detailed picture of the DCD donation and back-donation charge flows³⁵. Well-defined measures of the net charge transfer and of its donation and back-donation contributions (hereafter CT_{net} , $\text{CT}_{\text{don}}^{\sigma}$ and $\text{CT}_{\text{back}}^{\pi}$, respectively) can be obtained by taking the CDFs values at a plausible inter-fragment boundary, which we take to be the z point where equal-valued isodensity surfaces of the fragments become tangent^{35,36}.

As mentioned in the Introduction, in the present context the CDF also provides precious additional information concerning CO polarization. Since the C–O bond is collinear with the M–C z axis of integration, the CDF in the C–O bond region represents the electron displacement within CO with respect to free CO in response to the M–CO bond formation. The amount of charge flowing across a plane normal to the CO bond through its mid-point (i.e., the CDF value at $z = r_{\text{CO}}/2$) can be usefully taken as a quantitative estimate of such "polarization", and the total value can again itself be decomposed in σ and π components. We shall refer to these values as to $\text{CT}_{r_{\text{CO}}/2}^{\sigma}$, $\text{CT}_{r_{\text{CO}}/2}^{\pi}$ and $\text{CT}_{r_{\text{CO}}/2}^{\text{net}}$, respectively.

Geometry optimizations and the calculation of harmonic frequencies and electron densities were carried out by means of Density Functional Theory (DFT) with the ADF package^{45–47}. Becke's exchange functional⁴⁸ in combination with the Lee-Yang-Parr correlation functional⁴⁹ (BLYP) was adopted. We used an all electron triple-zeta basis set with two polarization functions (TZ2P) and a small frozen core for all atoms. Relativistic effects were included via the zeroth-order regular approximation (ZORA) Hamiltonian^{50–52}. An assessment of the effect of the exchange-correlation functional and of the basis set on the CDF is given in the supplementary information, where a comparison is also made with results from fully relativistic calculations carried out with a recently implemented parallel version of the Dirac-Kohn-Sham program BERTHA^{53–55}.

The purely electrostatic effect on the CO charge rearrangement was investigated using a uniform axial electric field (see also

Refs.^{56–58}) orientated along the C–O bond axis z (more details are given in Section 3.4). The density difference $\Delta\rho(x,y,z)$ in this case was formulated as the electron density of CO in the presence of the electric field at the actual minimum energy configuration minus that of free CO at the same geometry.

3 Results and Discussion

As mentioned in the Introduction, we first describe here a detailed investigation of the M–CO bond in $[\text{Au}(\text{CO})]^+$ and $[(\text{DPCb})\text{Au}(\text{CO})]^+$ (Section 3.1). We then extend the analysis to a whole series of 21 $[(\text{L})\text{Au}(\text{CO})]^{0/+}$ complexes (Section 3.2) and, finally, to a series of nine homoleptic complexes of general formula $[(\text{CO})_n\text{M}(\text{CO})]^m$ (Section 3.3). The full list of complexes considered is in Tables 1 and 2. The purely electrostatic effect is investigated in a last section (3.4) where an analysis of CO in a uniform axial electric field is carried out.

Three of the experimentally characterized gold complexes, with ligands DPCb, $[\text{HB}(3,5\text{-}(\text{CF}_3)_2\text{Pz})_3]^-$ and Mes_3P do not satisfy the symmetry requirements discussed in Section 2. $[(\text{DPCb})\text{Au}(\text{CO})]^+$, however, is only slightly asymmetric in its minimum configuration and has been here constrained to C_{2v} symmetry (the difference in energy with respect to the unconstrained optimized configuration is as small as 1 kcal/mol). The other two have been excluded from our analysis because they are much more asymmetric and to constrain them to C_{3v} symmetry would probably alter their properties significantly.

As Table 1 shows, the experimental CO stretching frequency for the three complexes $[(\text{CF}_3)\text{Au}(\text{CO})]$, $[(\text{Cl})\text{Au}(\text{CO})]$ and $[(\text{Br})\text{Au}(\text{CO})]$ (the first of which is measured in the solid state and the others in solution) is actually blue-shifted rather than red-shifted as the calculations consistently suggest for all the neutral systems (the computed $\nu_{\text{free-CO}}$ is 2143 cm^{-1}). Regarding this apparent inconsistency, Frenking *et al.* recently found that the experimental blue shift is actually due to the presence of intermolecular interactions and not to the properties of the single molecule⁵⁹. They proved this by computing the CO frequency of small aggregates of $[(\text{CF}_3)\text{Au}(\text{CO})]$ and of $[(\text{Cl})\text{Au}(\text{CO})]$ and finding that the frequency increases from smaller to larger values than that of free CO. Indeed, Au–Au interactions have been experimentally observed for these two complexes in the solid state^{24,60} and are likely to occur also in solution, especially for ligands with little steric hindrance. For this reason, and since experimental data are available only for a small subset of the complexes considered here, we shall base our discussion on the DFT values of $\Delta\nu_{\text{CO}}$ and Δr_{CO} (computed $r_{\text{free-CO}} = 1.137$). In fact, we shall most often refer to the latter parameter only, because the non-uniform influence of vibrational mode coupling, and the more complicated CO vibration modes in the homoleptic carbonyls, make $\Delta\nu_{\text{CO}}$ a less reliable parameter than Δr_{CO} for a quantitative analysis of its relation with the M–CO bond characteristics.

3.1 $[\text{Au}(\text{CO})]^+$ and $[(\text{DPCb})\text{Au}(\text{CO})]^+$

We start our analysis with an in-depth investigation of the gold carbon coordination bond in $[\text{Au}(\text{CO})]^+$ and $[(\text{DPCb})\text{Au}(\text{CO})]^+$. As mentioned in the Introduction, among the experimentally




Fig. 2 Total CDF and its symmetry (C_{2v}) components for the Au–CO bond in the complex $[\text{Au}(\text{CO})]^+$. Black dots indicate the z position of the atomic nuclei. A solid vertical line marks the boundary between the Au^+ and the CO fragments (see Section 2 for its definition). A dashed vertical line indicates the midpoint of the C–O bond ($z = r_{\text{CO}}/2$). At the top: isodensity surfaces ($\pm 0.0025\text{ e/a.u.}^3$) for the A_1 , B_1 and B_2 components of $\Delta\rho(x,y,z)$. Red surfaces identify charge depletion areas, blue surfaces identify charge accumulation areas.

characterized gold carbonyl complexes, these two systems display the most different spectroscopic properties. $[\text{Au}(\text{CO})]^+$ (observed in neon matrix²⁶) shows a CO stretching frequency much higher than that of free CO (experimental $\Delta\nu_{\text{CO}} = 94\text{ cm}^{-1}$) while $[(\text{DPCb})\text{Au}(\text{CO})]^+$ ⁴⁰ represents a unique case of cationic complex with red-shifted CO stretching frequency (experimental $\Delta\nu_{\text{CO}} = -30\text{ cm}^{-1}$). The computed values ($\Delta\nu_{\text{CO}} = 75\text{ cm}^{-1}$ and -76 cm^{-1} , respectively, see Table 1) reflect this opposite behavior.

We focus first on $[\text{Au}(\text{CO})]^+$, showing in Figure 2 the CDFs for the overall density difference and its symmetry-separated components. We recall here that, at a given point z , a positive CDF value corresponds to a charge flow from right to left (ie., in the $\text{Au}^+ \leftarrow \text{CO}$ direction) while a negative value corresponds to a charge flow in the opposite ($\text{Au}^+ \rightarrow \text{CO}$) direction. The total CDF is positive over both the Au–C and C–O bond regions and also at the oxygen far side of CO, indicating a continuous flow of electrons in the direction from CO towards gold. The negative values of the curve on the left side of Au^+ indicate a rearrangement in the opposite direction, which was shown in Ref.⁴² to be due to gold sd hybridization. The total CDF results from an A_1 component which is large and positive in the Au–carbon region (identifying σ donation) and a $B_1 + B_2$ component which is negative in the same zone (identifying π back-donation) plus a negligible A_2 component. These components are easily recognized in the isodensity plots of the respective density difference shown on top of the Figure.

The net charge transfer CT_{net} from CO to Au^+ (the CDF value at the boundary solid vertical line) amounts to 0.16 e resulting

Table 1 Computed $\Delta\nu_{\text{CO}}$ (cm^{-1}) and Δr_{CO} (\AA) and charge-transfer results (e) obtained from the CD analysis for the considered series of $[(\text{L})\text{Au}(\text{CO})]^{0/+}$ complexes. In boldface are data for the experimentally observed complexes. Reference values are $\nu_{\text{free-CO}} = 2106 \text{ cm}^{-1}$ (experimental: 2143 cm^{-1}), $r_{\text{free-CO}} = 1.137 \text{ \AA}$. ^a The vibrational coupling between the CO and the ligand has been eliminated through isotopic substitution ($[(\text{C}^{28}\text{N})\text{Au}(\text{CO})]$ and $[(\text{C}^3\text{H})\text{Au}(\text{CO})]$).

	$\Delta\nu_{\text{CO}}$ (exp. $\Delta\nu_{\text{CO}}$)	Δr_{CO}	CT_{net}	$\text{CT}_{\text{don}}^{\sigma}$	$\text{CT}_{\text{back}}^{\pi}$	$\text{CT}_{r_{\text{CO}/2}}$	$\text{CT}_{r_{\text{CO}/2}}^{\sigma}$	$\text{CT}_{r_{\text{CO}/2}}^{\pi}$
"Non classical" behavior								
$[(\text{CO})\text{Au}(\text{CO})]^{+}$	72 (74²⁸)	-0.010	0.08	0.21	-0.13	0.15	0.06	0.09
$[(\text{PF}_3)\text{Au}(\text{CO})]^{+}$	80	-0.009	0.09	0.22	-0.13	0.15	0.06	0.09
$[\text{Au}(\text{CO})]^{+}$	75 (94²⁶)	-0.008	0.16	0.34	-0.18	0.16	0.07	0.09
$[(\text{Ne})\text{Au}(\text{CO})]^{+}$	76	-0.008	0.15	0.33	-0.18	0.16	0.07	0.09
$[(\text{C}_2\text{H}_4)\text{Au}(\text{CO})]^{+}$	63	-0.007	0.08	0.24	-0.16	0.13	0.06	0.07
$[(\text{PH}_3)\text{Au}(\text{CO})]^{+}$	60	-0.007	0.06	0.22	-0.16	0.12	0.05	0.07
$[(\text{C}_2\text{H}_2)\text{Au}(\text{CO})]^{+}$	63	-0.007	0.06	0.23	-0.17	0.13	0.06	0.07
$[(\text{Xe})\text{Au}(\text{CO})]^{+}$	62	-0.006	0.08	0.27	-0.19	0.13	0.06	0.07
$[(\text{P}(\text{CH}_3)_3)\text{Au}(\text{CO})]^{+}$	38	-0.004	0.05	0.22	-0.17	0.10	0.05	0.05
$[(\text{NHC})\text{Au}(\text{CO})]^{+}$	39	-0.004	0.00	0.20	-0.20	0.10	0.05	0.05
$[(\text{C}_5\text{H}_5\text{N})\text{Au}(\text{CO})]^{+}$	46	-0.003	0.01	0.23	-0.22	0.10	0.05	0.05
$[(\text{Sidipp})\text{Au}(\text{CO})]^{+}$	23 (54³⁰)	-0.002	-0.01	0.20	-0.21	0.08	0.05	0.03
$[(\text{Idipp})\text{Au}(\text{CO})]^{+}$	17 (49³⁰)	-0.001	-0.02	0.20	-0.22	0.08	0.05	0.02
"Classical" behavior								
$[(\text{CF}_3)\text{Au}(\text{CO})]$	-2 (51²⁴)	0.002	-0.02	0.22	-0.24	0.05	0.05	0.00
$[(\text{CN})\text{Au}(\text{CO})]$	-6 ^a	0.003	-0.06	0.20	-0.26	0.05	0.05	0.00
$[(\text{H})\text{Au}(\text{CO})]$	-23 ^a	0.004	-0.07	0.19	-0.26	0.02	0.04	-0.02
$[(\text{CH}_3)\text{Au}(\text{CO})]$	-42	0.007	-0.07	0.21	-0.28	0.01	0.04	-0.03
$[(\text{C}_6\text{H}_5)\text{Au}(\text{CO})]$	-45	0.007	-0.06	0.22	-0.28	0.02	0.04	-0.02
$[(\text{I})\text{Au}(\text{CO})]$	-45	0.008	-0.08	0.24	-0.32	0.02	0.05	-0.03
$[(\text{Cl})\text{Au}(\text{CO})]$	-29 (13²⁷)	0.008	-0.11	0.23	-0.33	0.02	0.05	-0.03
$[(\text{Br})\text{Au}(\text{CO})]$	-40 (10²⁷)	0.008	-0.09	0.24	-0.33	0.02	0.05	-0.03
$[(\text{F})\text{Au}(\text{CO})]$	-23	0.009	-0.13	0.22	-0.35	0.02	0.05	-0.04
$[(\text{DPCb})\text{Au}(\text{CO})]^{+}$	-76 (-30⁴⁰)	0.010	-0.06	0.26	-0.32	0.03	0.05	-0.02

from a donation component $\text{CT}_{\text{don}}^{\sigma}$ of 0.34 e and a back-donation component $\text{CT}_{\text{back}}^{\pi}$ of 0.18 e. The first important comment here is therefore that, in a system like this showing a large blue-shift of the CO stretching frequency, back-donation is actually a significant component of the interaction, estimated to be more than half as large as the donation.

An analogous significant contribution from the electron charge rearrangement of π symmetry was also recently highlighted in Ref. ²⁹ through a Natural Orbitals for Chemical Valence-Extended Transition State (NOCV-ETS) ⁶¹ energy decomposition analysis. In particular, the π contribution to the overall orbital interaction energy ΔE_{orb} was found to be surprisingly large (32.5% of the overall ΔE_{orb}). The authors were cautious, however, in attributing such contribution exclusively to π back-donation, as ΔE_{orb} not only accounts for genuine inter-fragment orbital interactions but also for the polarization of the orbitals within each fragment.

This uncertainty may be dissolved here, because, as discussed in Section 2, the interfragment charge transfer and its components are automatically separated from the corresponding components of CO polarization in the CDF picture. Inspection of Figure 2 is in fact particularly revealing in this respect. Focusing on the CDFs in the carbonyl region, we notice immediately that the positive value of the total function indicates that the CO bond is on the whole polarized in the C←O direction. Remarkably, furthermore, this polarization results from the concordant positive contributions of *both* the σ and π components. We see indeed that, while the σ CDF keeps its (positive) sign on the right hand side of C and even beyond the oxygen site, an inversion (from negative to positive) is seen to occur for the π component pre-

cisely at the carbon site, leading to a maximum located about the mid-point of the C–O bond. In both cases, therefore, there is a displacement of electrons from oxygen towards carbon, which is due to the presence of the positively charged metal fragment. As discussed in Section 2, we can quantify the extent of CO polarization by taking the CDFs values at the mid-point of the CO bond (dashed vertical line in Figure 2). For the case in exam, the C←O polarization amounts to $\text{CT}_{r_{\text{CO}/2}} = 0.16$ e, resulting from a σ contribution $\text{CT}_{r_{\text{CO}/2}}^{\sigma}$ of 0.07 e and a π contribution $\text{CT}_{r_{\text{CO}/2}}^{\pi}$ of 0.09.

We now turn to $[(\text{DPCb})\text{Au}(\text{CO})]^{+}$, with its CDFs reported in Figure 3. This is analogous to Figure 2 except that here the B_1 (dashed blue line) and B_2 (dotted-dashed line) components are not identical and are shown separately in the plot. We notice an immediate striking contrast with the previous $[\text{Au}(\text{CO})]^{+}$ case, in that the back-donation components globally dominate over σ donation in the coordination bond region, so that the total CDF is everywhere negative, indicating a continuous, though modest, flow of electrons from $[(\text{DPCb})\text{Au}]^{+}$ to CO. This confirms the already cited findings of Ref. ⁴⁰. We note that π back-donation is in turn largely dominated by the B_2 component. The net charge transfer at the inter-fragment boundary is -0.06 e, resulting from a σ donation component of 0.26 e (A_1) and a π back-donation component of -0.32 e (-0.07 due to the B_1 component and -0.25 due to the B_2 component).

The polarization of the electron cloud in the carbonyl region also differs from that in $[\text{Au}(\text{CO})]^{+}$ in a remarkable way. In analogy with $[\text{Au}(\text{CO})]^{+}$, the σ CDF remains positive in the CO region and the B_1 component turns positive at the C site, reflecting the




Fig. 3 Total CDF and its symmetry (C_{2v}) components for the Au–CO bond in the complex $[(\text{DPCb})\text{Au}(\text{CO})]^+$. Black dots indicate the z position of the atomic nuclei. A solid vertical line marks the boundary between the $[(\text{DPCb})\text{Au}]^+$ and the CO fragments (see Section 2 for its definition). A dashed vertical line indicates the midpoint of the C–O bond ($z = r_{\text{CO}}/2$). At the top: isodensity surfaces ($\pm 0.0025 \text{ e/a.u.}^3$) for the A_1 , B_1 and B_2 components of $\Delta\rho(x, y, z)$. Red surfaces identify charge depletion areas, blue surfaces identify charge accumulation areas.

polarization of the CO bonding orbitals due to the electrostatic effect of the metal fragment. However, by contrast, the B_2 component maintains its negative sign also in the CO region, i.e. the back-donation it represents is so pronounced to penetrate the CO region and extend even beyond oxygen. As a consequence, the CO bond is on the whole slightly polarized in the $\text{C} \leftarrow \text{O}$ ($\text{CT}_{r_{\text{CO}}/2}^\sigma = 0.03 \text{ e}$), but this results from a σ polarization in the same direction ($\text{CT}_{r_{\text{CO}}/2}^\sigma = 0.05 \text{ e}$) and a π polarization in the opposite $\text{C} \rightarrow \text{O}$ direction ($\text{CT}_{r_{\text{CO}}/2}^\pi = -0.02 \text{ e}$).

It is worth deepening the comparison between the two complexes examined so far. In both, the metallic fragment bears a formal positive charge. However, $[\text{Au}(\text{CO})]^+$ behaves “non-classically” (blue-shifted $\Delta\nu_{\text{CO}}$), while $[(\text{DPCb})\text{Au}(\text{CO})]^+$ behaves “classically” (red-shifted $\Delta\nu_{\text{CO}}$). The CD analysis reveals that the σ donation component of the metal–CO bond is roughly comparable in the two cases ($\text{CT}_{\text{don}}^\sigma$ 0.34 vs 0.26 e), while π back-donation is almost twice as large in $[(\text{DPCb})\text{Au}(\text{CO})]^+$ ($\text{CT}_{\text{back}}^\pi$ 0.32 vs 0.18 e) and its extent acts to substantially reduce the $\text{C} \leftarrow \text{O}$ polarization of the CO bond. The polarization of the CO σ bonding orbitals is comparable in the two complexes ($\text{CT}_{r_{\text{CO}}/2}^\sigma$ 0.07 vs 0.05 e), but that of the π bonding orbitals is not ($\text{CT}_{r_{\text{CO}}/2}^\pi$ 0.09 vs -0.02 e). These findings suggest that π electron displacement upon coordination is the main factor driving CO bond response. In particular, if the presence of the metal fragment, especially if positively charged, is capable of polarizing the π CO bonding orbitals, even in the presence of a significant back-donation, the CO bond is strengthened; if, on the other hand, π back-donation is strong and extended enough to contrast CO polarization, even in

the presence of an equally cationic metal fragment, the CO bond results weakened.

3.2 The complete $[(\text{L})\text{Au}(\text{CO})]^{0/+}$ series

We now need of course to verify if the above preliminary surmise stands the test of a wider series of carbonyl compounds. To this end, we have extended the analysis to all of the 23 $[(\text{L})\text{Au}(\text{CO})]^{0/+}$ complexes listed in Table 1, which collects the spectroscopic data $\Delta\nu_{\text{CO}}$ and Δr_{CO} as well as the various computed CT figures. The complexes are listed in order of increasing Δr_{CO} and the experimentally characterized compounds are those shown in boldface. As briefly discussed at the beginning of Section 3, it is seen that, according to our computed shifts, the neutral complexes plus $[(\text{DPCb})\text{Au}(\text{CO})]^+$ behave “classically”, the remaining cationic complexes behave “non-classically”. The σ donation and π back-donation CDFs for these complexes are collected, respectively, in the top and bottom panel of Figure 4. Red lines are for the complexes showing red shift of ν_{CO} , blue lines are for those showing blue shift.

Two eye-catching features emerge upon inspection of Figure 4. The first is that all systems exhibit a surprisingly similar σ charge rearrangement (top panel) in the CO fragment region, in contrast with a much wider variability on the metal fragment side and despite the fact that some of the complexes are neutral and some cationic. In fact, as Table 1 shows, if one excludes the special cases of the naked Au^+ , of the “inert” ligands Ne and Xe, and of the “anomalous” $[(\text{DPCb})\text{Au}(\text{CO})]^+$, even the net ligand-to-metal σ donation, $\text{CT}_{\text{don}}^\sigma$, varies by only 0.05 e across the whole series of ligands. On the contrary, the π CDF (bottom panel of Figure 4) appears to be strongly influenced by the nature of the ligand over the whole molecular region, and $\text{CT}_{\text{back}}^\pi$ varies by 0.22 e over the ligand series. The second important observation is that, in the CO region, the complexes showing a blue-shifted ν_{CO} (blue lines) all invariably exhibit a flow of π electrons in the $\text{C} \leftarrow \text{O}$ direction ($\text{CT}_{r_{\text{CO}}/2}^\pi > 0$), due to the positively charged metallic fragment, while the complexes showing red-shifted ν_{CO} (red lines) exhibit a negative $\text{CT}_{r_{\text{CO}}/2}^\pi$, i.e., charge flows in the opposite $\text{C} \rightarrow \text{O}$ direction (with the exception of two complexes for which $\text{CT}_{r_{\text{CO}}/2}^\pi$ is essentially vanishing and red-shift is also negligibly small).

It thus appears quite clearly that in the series of gold(I) carbonyls: (i) σ donation is much less tunable than π back-donation, being very little dependent on the nature and the charge of the ligand; and (ii) whereas the net CO bond polarization turns out to be invariably oriented in the $\text{C} \leftarrow \text{O}$ direction ($\text{CT}_{r_{\text{CO}}/2}^\pi > 0$), the direction of its π density component can vary and appears to be tightly connected with the direction of the CO stretching shift and bond-length change. These findings are given a definitive illustration in Figures 5 and 6 where the correlation of Δr_{CO} with CT_{net} , $\text{CT}_{\text{don}}^\sigma$ and $\text{CT}_{\text{back}}^\pi$ and with $\text{CT}_{r_{\text{CO}}/2}$, $\text{CT}_{r_{\text{CO}}/2}^\sigma$ and $\text{CT}_{r_{\text{CO}}/2}^\pi$, respectively, is reported. In both Figures, black triangles are used for the overall CT, red squares for its σ component and blue circles for its π component. Empty symbols are for the neutral species, filled ones are for the cationic species. Focusing first on Figure 5, no correlation is found, as expected, between Δr_{CO} and $\text{CT}_{\text{don}}^\sigma$, while a good inverse correlation ($R^2 = 0.945$) can be seen between




Fig. 4 σ donation (top panel) and π back-donation (bottom panel) CDFs for the Au–CO bond in the series of $[(L)Au(CO)]^{0/+}$ complexes of Table 1. Red lines (blue lines) are used for complexes showing red-shift (blue-shift) in the computed $\Delta\nu_{CO}$. The z origin is placed at the C atom for all complexes and black dots indicate the position of C and O (the latter varying negligibly, less than 0.02 Å, among the complexes). A dashed vertical line marks the C–O midpoint. The position of the Au atom across the series varies more significantly and its range is marked by a rectangle. Similarly, a gray vertical band indicates the range of the interfragment boundary.




Fig. 5 Correlation between the computed Δr_{CO} in the considered series of $[(L)Au(CO)]^{0/+}$ complexes and the CT_{net} (black triangles), CT_{don}^{σ} (red squares) and CT_{back}^{π} (blue circles). Empty symbols are for the neutral species, filled symbols for the cationic species.




Fig. 6 Correlation between the computed Δr_{CO} in the considered series of $[(L)Au(CO)]^{0/+}$ complexes and $CT_{rCO/2}$ (black triangles), $CT_{rCO/2}^{\sigma}$ (red squares) and $CT_{rCO/2}^{\pi}$ (blue circles). Empty symbols are for the neutral species, filled symbols for the cationic species.

Δr_{CO} and CT_{back}^{π} , a trace of which remains in the plot of Δr_{CO} vs CT_{net} . The same bond weakening effect of π back-donation is also evident in the plot of $\Delta\nu_{CO}$ vs CT_{back}^{π} (see supplementary information), though correlation, as earlier warned, is made worse by mode coupling ($R^2 = 0.849$). Figure 6 shows the correlation of Δr_{CO} with $CT_{rCO/2}$ and its components $CT_{rCO/2}^{\sigma}$ and $CT_{rCO/2}^{\pi}$. Not surprisingly, as these quantities are all directly related to the charge rearrangement of the CO bond itself, correlations are here quantitatively better (R^2 is 0.970 for that with $CT_{rCO/2}^{\pi}$). Obviously, as Δr_{CO} correlates well with both π back-donation and CO π electron polarization, the latter two quantities are also in mutual correlation.

3.3 Homoleptic complexes: the $[(CO)_nM(CO)]^m$ series

In the previous Sections we considered a series of gold(I) complexes where the donor/acceptor properties of the M–CO bond were varied through the ligand L. We now extend the analysis to a series of homoleptic carbonyls of formula $[(CO)_nM(CO)]^m$, where the relative extent of the DCD constituents of the M–CO bond and CO polarization are varied essentially by changing the metal. The full list of the considered homoleptic complexes is in Table 2, reporting their spectroscopic shifts and CD decomposition results. Complexes are listed in order of increasing value of Δr_{CO} . We omit for brevity a presentation of the complete CDFs. The computed structures for these systems are in agreement with experimental X-ray data where available^{62–66}. $Hg(CO)_2^{2+}$ and $Ir(CO)_6^{3+}$, both cationic, behave "non classically"; with experimental blue-shifted ν_{CO} at 2279.5 cm^{-1} for the former and at 2254, 2276 and 2298 cm^{-1} for the latter.^{65,66} On the opposite side, the anionic complexes show exceptionally low CO stretching frequency, the most red-shifted being that of $Fe(CO)_4^{2-}$ at 1730 cm^{-1} (this is the first anionic carbonyl complex spectroscopically observed^{67,68}). In between are $Mo(CO)_6$, $Fe(CO)_5$ (for which both the axial and equatorial M–CO bonds have been investigated),⁶⁹ $Ni(CO)_4$ and $Cr(CO)_6$. The complexes present therefore a wide range of ν_{CO} variation but $\Delta\nu_{CO}$ turns out not to be

a good parameter for analyzing correlations with the CD data because normal-mode coupling varies significantly with the different structure of the complexes. We therefore base our discussion, as already done also for the gold(I) complexes, on the computed Δr_{CO} . This varies in a range of 0.087 Å over the series, from -0.018 to 0.069 Å (Table 2).

The Table shows that also in this series of compounds the range of variation in π back-donation (0.69 e) is much larger than that of σ donation (0.15 e). In particular, almost no back-donation is found for $\text{Hg}(\text{CO})_2^{2+}$ while $\text{CT}_{\text{back}}^{\pi}$ for $[\text{Fe}(\text{CO})_4]^{2-}$ is as high as 0.71 e. This picture is completely consistent with the simple VB view discussed in the Introduction, in that we go from a purely σ M–CO bond (structure a) for $\text{Hg}(\text{CO})_2^{2+}$ to a situation in which all π^* CO orbitals are engaged in back-bonding (structure c) for $[\text{Fe}(\text{CO})_4]^{2-}$. Also the charge rearrangement (polarization) in the carbonyl region is seen to follow a similar trend, with a much narrower range of $\text{CT}_{r_{\text{CO}/2}}^{\sigma}$ values (between 0.02 and 0.10 e) than that of $\text{CT}_{r_{\text{CO}/2}}^{\pi}$ (from 0.21 to -0.30 e). As before, no clear correlation can be discerned between Δr_{CO} and the σ CT data, while $\text{CT}_{\text{back}}^{\pi}$ and $\text{CT}_{r_{\text{CO}/2}}^{\pi}$ values are seen to decrease almost monotonically as Δr_{CO} increases.

A plot of Δr_{CO} vs either $\text{CT}_{\text{back}}^{\pi}$ or $\text{CT}_{r_{\text{CO}/2}}^{\pi}$ for the whole set of complexes studied, including the present homoleptic carbonyls in addition to the gold(I) series, appears in fact to suggest, because the range of variation is now significantly enlarged, that a quadratic fit, rather than a linear one, may better represent the actual correlation. (An evident non-linear relationship has already been found between the electric field strength and Δr_{CO} .⁵⁶) Figure 7 shows very clearly this to be indeed the case, with the accuracy of all fits improved with respect to the sole subset of gold complexes. Once again, in the homoleptic series, the carbonyl complexes featuring CO bond strengthening (blue-shifted $\Delta \nu_{\text{CO}}$ and negative Δr_{CO}), i.e. the cationic $\text{Hg}(\text{CO})_2^{2+}$ and $\text{Ir}(\text{CO})_6^{3+}$, show a flow of π electrons in the C←O direction. All other complexes, where the CO bond weakens (red-shifted $\Delta \nu_{\text{CO}}$ and positive Δr_{CO}) show opposite-direction flows.

3.4 CO in a uniform axial electric field

The observation that the CO bond is lengthened or shortened upon formation of the M–CO bond according to whether the CO bonding orbitals of π symmetry are polarized in the C→O or C←O direction, respectively, is certainly remarkable. To verify that this is a general fact, actually independent of CO coordination, we discuss in this last Section an ad hoc study of the electron cloud rearrangement and stretching response of CO in an external uniform axial electric field oriented along the C–O bond axis.

In Figure 8, we show the computed CO stretching Δr_{CO} reported versus the π and σ components of $\text{CT}_{r_{\text{CO}/2}}$ while the latter vary as a result of the applied field. In the same figure also the points representing the computed Δr_{CO} and π and σ components of $\text{CT}_{r_{\text{CO}/2}}$ are reported for the whole series of carbonyl complexes studied in this work. Let us focus first on the stretching response to the electric field. When the field is absent, the system corresponds to free CO and Δr_{CO} , $\text{CT}_{r_{\text{CO}/2}}^{\sigma}$ and $\text{CT}_{r_{\text{CO}/2}}^{\pi}$ are all zero. As the field increases on the left, in the direction that induces (lin-




Fig. 7 Correlation between CO bond-length change upon coordination, Δr_{CO} , and (a) M→CO π back-donation $\text{CT}_{\text{back}}^{\pi}$ (upper panel) (b) CO π electron polarization $\text{CT}_{r_{\text{CO}/2}}^{\pi}$ (lower panel), for the whole set of complexes studied. Black-colored points refer to the gold(I) complexes, blue-colored points to the homoleptic carbonyls (filled circles are for positively charged complexes, empty circles for neutral or negatively charged ones). Both the linear fit ($R^2=0.959$ for the upper panel, $R^2=0.945$ for the lower panel) and the quadratic fit ($R^2=0.984$ for the upper panel and $R^2=0.996$ for the lower panel) curves are shown.




Fig. 8 The dashed lines show Δr_{CO} versus the σ (red color) and π (blue) CO polarization $\text{CT}_{r_{\text{CO}/2}}$ for a CO molecule placed in a uniform axial electric field of magnitude ranging from -0.11 to 0.07 a.u. in steps of 0.02 a.u. (colored square points). For comparison, also shown is the correlation between Δr_{CO} and σ and π CO polarization (red and blue circles, respectively) for the whole series of Au and homoleptic complexes studied. The empty circles are for neutral or negatively charged complexes, the filled circles for cationic ones.

Table 2 Symmetry group, experimental $\Delta\nu_{\text{CO}}$ (and related symmetry mode), computed Δr_{CO} and results obtained from the CD analysis for the considered series of homoleptic carbonyl complexes $[(\text{CO})_n\text{M}(\text{CO})]^m$. Distances in Å, charge transfers in e.

	sym.	exp. $\Delta\nu_{\text{CO}}$	Δr_{CO}	CT_{net}	$\text{CT}_{\text{don}}^{\sigma}$	$\text{CT}_{\text{back}}^{\pi}$	$\text{CT}_{r_{\text{CO}}/2}$	$\text{CT}_{r_{\text{CO}}/2}^{\sigma}$	$\text{CT}_{r_{\text{CO}}/2}^{\pi}$
"Non classical" behavior									
Hg(CO) ₂ ²⁺	C _{∞v}	136 ⁶⁶ (A ₁)	-0.018	0.29	0.31	-0.02	0.13	0.08	0.21
Ir(CO) ₆ ³⁺	O _h	155 ⁶⁵ (A _{1g})	-0.015	0.17	0.31	-0.13	0.08	0.10	0.18
		133 ⁶⁵ (E _g)							
		111 ⁶⁵ (T _{1u})							
"Classical" behavior									
Ni(CO) ₄	T _d	-23 ⁷⁰ (A ₁)	0.012	-0.16	0.16	-0.32	-0.01	0.04	-0.05
Fe(CO) ₅ (ax.)	D _{3h}	-22 ⁶⁹ (A ₁)	0.014	-0.18	0.23	-0.41	-0.02	0.06	-0.08
		-101 ⁶⁹ (A ₁ ')							
		-109 ⁶⁹ (A ₂ ')							
Cr(CO) ₆	O _h	-24 ⁷¹ (A _{1g})	0.016	-0.21	0.17	-0.37	-0.02	0.05	-0.07
		-116 ⁷¹ (E _g)							
		-143 ⁷¹ (T _{1u})							
Mo(CO) ₆	O _h	-22 ⁷¹ (A _{1g})	0.016	-0.23	0.14	-0.37	-0.02	0.04	-0.06
		-118 ⁷¹ (E _g)							
		-140 ⁷¹ (T _{1u})							
Fe(CO) ₅ (eq.)	D _{3h}	(cfr. ax.)	0.018	-0.20	0.20	-0.40	-0.03	0.05	-0.08
Co(CO) ₄ ⁻	T _d	-255 ⁷² (A ₁)	0.038	-0.36	0.16	-0.52	-0.15	0.03	-0.18
Ir(CO) ₄ ⁻	T _d	-248 ⁷³ (A ₁)	0.039	-0.28	0.30	-0.58	-0.14	0.04	-0.18
Ru(CO) ₄ ²⁻	T _d	-407 ⁷⁴ (A ₁)	0.066	-0.42	0.25	-0.67	-0.24	0.03	-0.27
Fe(CO) ₄ ²⁻	T _d	-413 ⁷⁵ (A ₁)	0.069	-0.55	0.16	-0.71	-0.28	0.02	-0.30

early) C→O (negative) polarization, C–O bond length increases quadratically and π polarization is seen to increase much more rapidly than σ polarization. Conversely, as the field increases on the right, inducing C←O polarization, the C–O bond shortens (much less rapidly).

When we now compare these curves with the relation observed between Δr_{CO} and the σ and π components of CO polarization induced by metal coordination, rather than by an applied field (disconnected circles in the Figure), we notice immediately that the π circles follow quite closely the correlation between field-induced polarization and stretching, while, in striking contrast, the σ circles deviate from the field-induced line (a clear indication of a much more pronounced "chemical" signature) and, moreover, span a very narrow range of (positive) polarization, essentially without any correlation with the widely varying Δr_{CO} . This is indeed a very strong confirmation that the CO stretching response to any solicitation causing electron charge rearrangement, be it the formation of a M–CO coordination bond or the effect of an external electric field, is driven essentially exclusively by the charge rearrangement of π symmetry: whether induced by an external electric field or by metal coordination, C→O (C←O) polarization of the π bond orbitals invariably and tightly correlates with bond lengthening (shortening).

4 Conclusions

In this work we have carried out an in-depth analysis of the M–CO bond in $[(\text{L})_n\text{M}(\text{CO})]^m$ metal carbonyl complexes, with the aim of elucidating on quantitative grounds the σ donation and π back-donation effects on the CO stretching response, in particular the change in bond length Δr_{CO} , to the M–CO bond formation. The analysis was carried out for a large variety of car-

bonyls, in which the relative extent of the DCD constituents were varied both through L in a series of $[(\text{L})\text{Au}(\text{CO})]^{0/+}$ gold(I) carbonyl complexes and through M in a series of anionic, neutral and cationic $[(\text{CO})_n\text{M}(\text{CO})]^m$ homoleptic carbonyls. Crucially for the purpose of this investigation, reliable and consistent measures, not only of σ donation and π back-donation charges, but also of the σ and π components of CO polarization were obtained by the well-established charge-displacement analysis of electron-density differences, as resulting from accurate DFT calculations. The nature of the M–CO bond in the considered complexes was found to range smoothly between the two extreme cases of an almost purely σ bonded complex (Hg(CO)₂²⁺, $\text{CT}_{\text{back}}^{\pi}=0.02$ e) and of a strongly back-bonded complex ($[\text{Fe}(\text{CO})_4]^{2-}$, $\text{CT}_{\text{back}}^{\pi}=0.71$ e). Conversely, all complexes were found to feature a narrowly comparable σ donation component, with $\text{CT}_{\text{don}}^{\sigma}$ values ranging from 0.14 to 0.34 e. The same picture holds accurately for the electron cloud rearrangement over the carbonyl region: all considered complexes feature a comparable σ polarization of CO and a much more variable π polarization. Quite remarkably, no correlation is found between Δr_{CO} and the σ displacements, while Δr_{CO} , π back-donation and CO π polarization all correlate tightly with one another.

These results show that the driving force of the CO stretching response to the M–CO bond formation is provided exclusively by the changes taking place in the π electron density. In the complexes studied, such π charge rearrangement is found to result from the interplay between π back-donation (structures a-c of the Introduction) and the electrostatic effect (structures d-f) exerted by the metal ligand-fragment. In particular, cationic metal-ligand fragments polarize the π CO bonding orbitals in the C←O direction, thus shortening the bond and enhancing the covalency, as

highlighted in Ref.²³. This effect, on the other hand, is contrasted by π back-donation shifting charge in the opposite direction. The net direction $C\leftarrow O$ or $C\rightarrow O$ of the polarization of π CO bonding orbitals is found to invariably determine whether the CO bond is strengthened or weakened, respectively. This is most evident in the $[(DPCb)Au(CO)]^+$ complex, where π back-donation is so strong as to invert the polarization of the π CO bonding orbitals in the $C\rightarrow O$ direction despite the formal positive charge on the ligand-metal fragment, making it the only example of a cationic gold(I) carbonyl complex with “classical” behavior ($\Delta r_{CO} > 0$). An ad hoc study of CO in a uniform axial electric field demonstrates that it is indeed the polarization of the π CO bonding orbitals, no matter how induced (whether by the coordination bond to M or by an electric field), that drives direction and magnitude of the CO stretching response to the M–CO bond formation.

Regarding the fundamental question of what can be inferred on the nature of the M–CO bond from the analysis of Δr_{CO} (and less reliably, due to mode coupling, $\Delta \nu_{CO}$) in metal carbonyl complexes, we conclude that the value of Δr_{CO} quantifies to an excellent extent the π backdonation component of the M–CO bond, since such component directly correlates with the π polarization. In particular, where $CT_{r_{CO}/2}^\pi$ changes its sign (ie. the polarization of π CO bonding orbitals changes direction determining whether the CO bond is weakened or strengthened), CT_{back}^π is approximately as high as the average extent of σ donation among the complexes herein considered. This indicates that π back-donation is an important component also in the class of “non-classical” complexes, as those of gold(I) considered in this work.

5 Acknowledgements

This work has been supported by Ministero dell'Istruzione, Università e Ricerca (MIUR) through the FIRB *Futuro in Ricerca* project n. RBFR1022UQ.

References

- 1 A. Roberts, *Chem. News*, 1898, **78**, 260.
- 2 F. Calderazzo, *Angew. Chem. Int. Edit.*, 1977, **16**, 299–311.
- 3 J.-K. Shen, Y.-C. Gao, Q.-Z. Shi and F. Basolo, *Coordin. Chem. Rev.*, 1993, **128**, 69–88.
- 4 F. Fischer and H. Tropsch, *Brennst. Chem.*, 1923, **4**, 276–285.
- 5 J. Fierro, *catlet*, 1993, **22**, 67–91.
- 6 M. J. S. Dewar, *Bull. Soc. Chim. Fr.*, 1951, **18**, C71–C79.
- 7 J. Chatt and L. A. Duncanson, *J. Chem. Soc.*, 1953, 2939–2947.
- 8 *Modern Coordination Chemistry: The Legacy of Joseph Chatt*, ed. N. Winterton and J. Leigh, The Royal Society of Chemistry, 2002, pp. X001–X004.
- 9 T. Ziegler and A. Rauk, *Inorg. Chem.*, 1979, **18**, 1755–1759.
- 10 K. Nakashima, X. Zhang, M. Xiang, Y. Lin, M. Lin and Y. Mo, *J. Theor. Comput. Chem.*, 2008, **7**, 639–654.
- 11 S. Dapprich and G. Frenking, *J. Phys. Chem.*, 1995, **99**, 9352–9362.
- 12 F. Weinhold and C. R. Landis, *Valency and bonding: a natural bond orbital donor-acceptor perspective*, Cambridge University Press, 2005.
- 13 E. Matito and M. Solà, *Coordin. Chem. Rev.*, 2009, **253**, 647–665.
- 14 P. Macchi and A. Sironi, *Coordin. Chem. Rev.*, 2003, **238**, 383–412.
- 15 J. Pilme, B. Silvi and M. E. Alikhani, *J. Phys. Chem. A*, 2003, **107**, 4506–4514.
- 16 F. Cortés-Guzmán and R. F. Bader, *Coordin. Chem. Rev.*, 2005, **249**, 633–662.
- 17 D. Tiana, E. Francisco, M. Blanco, P. Macchi, A. Sironi and A. Martín Pendás, *J. Chem. Theory Comput.*, 2010, **6**, 1064–1074.
- 18 A. J. Lupinetti, S. H. Strauss and G. Frenking, *Progress in Inorganic Chemistry*, 2001, **49**, 1–112.
- 19 P. K. Hurlburt, J. J. Rack, J. S. Luck, S. F. Dec, J. D. Webb, O. P. Anderson and S. H. Strauss, *J. Am. Chem. Soc.*, 1994, **116**, 10003–10014.
- 20 H. Willner and F. Aubke, *Angew. Chem. Int. Edit.*, 1997, **36**, 2402–2425.
- 21 A. S. Goldman and K. Krogh-Jespersen, *J. Am. Chem. Soc.*, 1996, **118**, 12159–12166.
- 22 A. J. Lupinetti, S. Fau, G. Frenking and S. H. Strauss, *J. Phys. Chem. A*, 1997, **101**, 9551–9559.
- 23 G. Frenking, C. Loschen, A. Krapp, S. Fau and S. H. Strauss, *J. Comput. Chem.*, 2007, **28**, 117–126.
- 24 S. Martínez-Salvador, J. Forníes, A. Martín and B. Menjón, *Angew. Chem. Int. Edit.*, 2011, **50**, 6571–6574.
- 25 R. H. Crabtree, *The organometallic chemistry of the transition metals*, Wiley, 2005.
- 26 B. Liang and L. Andrews, *J. Phys. Chem. A*, 2000, **104**, 9156–9164.
- 27 D. B. Dell'Amico, F. Calderazzo, P. Robino and A. Segre, *J. Chem. Soc., Dalton Trans.*, 1991, **0**, 3017–3020.
- 28 H. Willner, J. Schaebs, G. Hwang, F. Mistry, R. Jones, J. Trotter and F. Aubke, *J. Am. Chem. Soc.*, 1992, **114**, 8972–8980.
- 29 H. V. R. Dias, C. Dash, M. Yousufuddin, M. A. Celik and G. Frenking, *Inorg. Chem.*, 2011, **50**, 4253–4255.
- 30 C. Dash, P. Kroll, M. Yousufuddin and H. V. R. Dias, *Chem. Commun.*, 2011, **47**, 4478–4480.
- 31 H. V. R. Dias and W. Jin, *Inorg. Chem.*, 1996, **35**, 3687–3694.
- 32 D. J. Gorin and F. D. Toste, *Nature*, 2007, **446**, 395–403.
- 33 I. Krossing, *Angew. Chem. Int. Edit.*, 2011, **50**, 11576–11578.
- 34 M. Alcarazo, T. Stork, A. Anoop, W. Thiel and A. Fürstner, *Angew. Chem. Int. Edit.*, 2010, **49**, 2542–2546.
- 35 N. Salvi, L. Belpassi and F. Tarantelli, *Chem. Eur. J.*, 2010, **16**, 7231–7240.
- 36 G. Bistoni, L. Belpassi and F. Tarantelli, *Angew. Chem. Int. Edit.*, 2013, **52**, 11599–11602.
- 37 G. Ciancaleoni, N. Scafuri, G. Bistoni, A. Macchioni, F. Tarantelli, D. Zuccaccia and L. Belpassi, *Inorg. Chem.*, 2014, **53**, 9907–9916.
- 38 A. S. K. Hashmi, *Angew. Chem. Int. Edit.*, 2008, **47**, 6754–6756.
- 39 A. M. Echavarren, *Nature Chem.*, 2009, **1**, 431–433.

- 40 M. Joost, L. Estévez, S. Mallet-Ladeira, K. Miqueu, A. Amgoune and D. Bourissou, *Angew. Chem. Int. Edit.*, 2014, **53**, 14512–14516.
- 41 H. V. R. Dias and M. Fianchini, *Angew. Chem. Int. Edit.*, 2007, **46**, 2188–2191.
- 42 L. Belpassi, I. Infante, F. Tarantelli and L. Visscher, *J. Am. Chem. Soc.*, 2008, **10**, 1048–1060.
- 43 G. Ciancaleoni, L. Biasiolo, G. Bistoni, A. Macchioni, F. Tarantelli, D. Zuccaccia and L. Belpassi, *Chem. Eur. J.*, 2015, **21**, 2467–2473.
- 44 D. Cappelletti, E. Ronca, L. Belpassi, F. Tarantelli and F. Pirani, *Accounts Chem. Res.*, 2012, **45**, 1571–1580.
- 45 SCM, Theoretical Chemistry, Vrije Universiteit, Amsterdam, The Netherlands, *ADF User's Guide. Release 2012.01*, 2012.
- 46 C. Fonseca Guerra, J. G. Snijders, G. te Velde and E. J. Baerends, *Theor. Chem. Acc.*, 1998, **99**, 391–403.
- 47 G. te Velde, F. M. Bickelhaupt, E. J. Baerends, C. Fonseca Guerra, S. J. A. van Gisbergen, J. G. Snijders and T. Ziegler, *J. Comput. Chem.*, 2001, **22**, 931–967.
- 48 A. D. Becke, *Phys. Rev. A*, 1988, **38**, 3098–3100.
- 49 C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B*, 1988, **37**, 785–789.
- 50 E. van Lenthe, E. J. Baerends and J. G. Snijders, *J. Chem. Phys.*, 1993, **99**, 4597–4610.
- 51 E. van Lenthe, E. J. Baerends and J. G. Snijders, *J. Chem. Phys.*, 1994, **101**, 9783–9792.
- 52 E. van Lenthe, A. Ehlers and E.-J. Baerends, *J. Chem. Phys.*, 1999, **110**, 8943–8953.
- 53 S. Rampino, L. Belpassi, F. Tarantelli and L. Storchi, *J. Chem. Theory Comput.*, 2014, **10**, 3766–3776.
- 54 L. Storchi, S. Rampino, L. Belpassi, F. Tarantelli and H. M. Quiney, *J. Chem. Theory Comput.*, 2013, **9**, 5356–5364.
- 55 L. Belpassi, L. Storchi, H. M. Quiney and F. Tarantelli, *Phys. Chem. Chem. Phys.*, 2011, **13**, 12368–12394.
- 56 N. Hush and M. Williams, *J. Mol. Spectrosc.*, 1974, **50**, 349–368.
- 57 M. Head-Gordon and J. C. Tully, *Chem. Phys.*, 1993, **175**, 37–51.
- 58 P. Bagus and G. Pacchioni, *Electrochim. Acta*, 1991, **36**, 1669–1675.
- 59 M. A. Celik, C. Dash, V. A. Adiraju, A. Das, M. Yousufuddin, G. Frenking and H. V. R. Dias, *Inorg. Chem.*, 2012, **52**, 729–742.
- 60 P. G. Jones, *Z. Naturforsch. B*, 1982, **37**, 823–824.
- 61 M. P. Mitoraj, A. Michalak and T. Ziegler, *J. Chem. Theory Comput.*, 2009, **5**, 962–975.
- 62 J. E. Ellis, *Organometallics*, 2003, **22**, 3322–3338.
- 63 J. E. Ellis, *Inorg. Chem.*, 2006, **45**, 3167–3186.
- 64 L. J. Farrugia and C. Evans, *J. Phys. Chem. A*, 2005, **109**, 8834–8848.
- 65 C. Bach, H. Willner, F. Aubke, C. Wang, S. J. Rettig and J. Trotter, *Angew. Chem. Int. Edit.*, 1996, **35**, 1974–1976.
- 66 M. Bodenbinder, G. Balzer-Jöllenebeck, H. Willner, R. Batchelor, F. Einstein, C. Wang and F. Aubke, *Inorg. Chem.*, 1996, **35**, 82–92.
- 67 W. Heiber and F. Leutert, *Berichte der deutschen chemischen Gesellschaft (A and B Series)*, 1931, **64**, 2832–2839.
- 68 W. Hieber, J. Sedlmeier and R. Werner, *Chem. Ber.*, 1957, **90**, 278–286.
- 69 L. H. Jones, R. S. McDowell, M. Goldblatt and B. I. Swanson, *J. Chem. Phys.*, 1972, **57**, 2050.
- 70 L. Perrin, E. Clot, O. Eisenstein, J. Loch and R. H. Crabtree, *Inorg. Chem.*, 2001, **40**, 5806–5811.
- 71 L. H. Jones, R. S. McDowell and M. Goldblatt, *inch*, 1969, **8**, 2349–2363.
- 72 W. Beck and R. E. Nitzschmann, *Z. Naturforsch.*, 1962, **17B**, 577–581.
- 73 L. Garlaschelli, R. Della Pergola, S. Martinengo and J. E. Ellis, in *Inorganic Syntheses*, John Wiley & Sons, Inc., 2007, pp. 211–215.
- 74 J. D. Cotton, M. I. Bruce and F. G. A. Stone, *J. Chem. Soc. A*, 1968, 2162–2165.
- 75 W. F. Edgell, M. T. Yang, B. J. Bulkin, R. Bayer and N. Koizumi, *J. Am. Chem. Soc.*, 1965, **87**, 3080–3088.