


# Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



## Rapid Oxidative Hydrogen Evolution from a Family of Square-Planar Nickel Hydride Complexes

Received 00th January 20xx,  
Accepted 00th January 20xx

S. Ramakrishnan,<sup>a†</sup> S. Chakraborty,<sup>b†</sup> W. W. Brennessel,<sup>b</sup> C. E. D. Chidsey<sup>a\*</sup> and W. D. Jones<sup>b\*</sup>

DOI: 10.1039/x0xx00000x

www.rsc.org/

A series of square-planar nickel hydride complexes supported by bis(phosphinite) pincer ligands with varying substituents (–OMe, –Me, and –Bu<sup>t</sup>) on the pincer backbone have been synthesized and completely characterized by NMR spectroscopy, IR spectroscopy, elemental analysis, and X-ray crystallography. Their cyclic voltammograms show irreversible oxidation peaks (peak potentials from 101 to 316 mV vs. Fc<sup>+</sup>/Fc) with peak currents consistent with overall one-electron oxidations. Chemical oxidation by the one-electron oxidant Ce(NBu<sub>4</sub>)<sub>2</sub>(NO<sub>3</sub>)<sub>6</sub> was studied by NMR spectroscopy, which provided quantitative evidence for post-oxidative H<sub>2</sub> evolution leading to a solvent-coordinated nickel(II) species with the pincer backbone intact. Bulk electrolysis of the unsubstituted nickel hydride (**3a**) showed an overall one-electron stoichiometry and gas chromatographic analysis of the headspace gas after electrolysis further confirmed stoichiometric production of dihydrogen. Due to the extremely high rate of the post-oxidative chemical process, electrochemical simulations have been used to establish a lower limit of the bimolecular rate constant ( $k_f > 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ) for the H<sub>2</sub> evolution step. To the best of our knowledge, this is the fastest known oxidative H<sub>2</sub> evolution process observed in transition metal hydrides. Quantum chemical calculations based on DFT indicate that the one-electron oxidation of the nickel hydride complex provides a strong chemical driving force (–90.3 kcal/mol) for the production of H<sub>2</sub> at highly oxidizing potentials.

### Introduction


Virtual H<sub>2</sub> carriers such as alcohols and saturated *N*-heterocycles that are liquids at room temperature constitute important alternatives to the use of H<sub>2</sub> in commercial fuel cells.<sup>1–3</sup> They are easy to store compared to gaseous dihydrogen and in principle, could be part of a reversible fuel cell with the dehydrogenated products also being liquids at room temperature. Developing catalysts containing earth-abundant elements to dehydrogenate these fuels to produce protons and electrons ergoneutrally, is therefore relevant for fuel cell applications.<sup>4–6</sup>

Scheme 1 shows a conceptual pathway for the dehydrogenation of organic fuels involving homogenous molecular catalysts.<sup>4</sup> Dehydrogenation of the fuel in the presence of a base typically leads to transition metal hydride intermediates, ubiquitous in transfer hydrogenation processes<sup>7–9</sup>, which shuttle H<sub>2</sub> equivalents from a sacrificial donor (for example, an alcohol) to an acceptor molecule such as a ketone. Subsequent electro-oxidation and deprotonation

of the metal hydride, a two-electron one-proton process, results in a solvent-coordinated species, as shown in Scheme 1. To design an energy-efficient electrocatalytic cycle for this transformation, it is important that both of the steps shown in Scheme 1 are close to ergoneutrality. In this work, we focused on the latter step, i.e. the electrochemical oxidation of a series of metal hydride complexes.

The electro-oxidation of metal hydrides has been a subject of interest due to its predominantly irreversible nature which often suggests a subsequent fast chemical step leading to undesired products.<sup>10–13</sup> Tilset and coworkers have reported that electron removal from the metal center in Group VI hydrides enhances the acidity of the hydride by approximately 20 pKa units leading to deprotonation, forming a reduced

**Scheme 1** Electrocatalytic Cycle for Electro-Dehydrogenation of H-containing Liquid Fuels (L<sub>n</sub>=Ligand, S=Coordinating Solvent, M=Transition metal, n=integer, B=Base).



<sup>a</sup> Department of Chemistry, Stanford University, Stanford, California 94305, USA. chidsey@stanford.edu


<sup>b</sup> Department of Chemistry, University of Rochester, Rochester, New York 14627, USA. jones@chem.rochester.edu

† These authors contributed equally. Electronic Supplementary Information (ESI) available: supplementary electrochemical data, electrochemical simulation data, Cartesian coordinates and energies of DFT optimized structures, sample input files and X-Ray Analyses. See DOI: 10.1039/x0xx00000x


## ARTICLE

## Chemical Science

metal species that would spontaneously lose another electron at the operating potential, thereby constituting an overall two electron process.<sup>10,14</sup> Depending on the pKa of the oxidized hydride, a suitable base can act as the proton acceptor and a solvent molecule such as acetonitrile as a ligand that stabilizes the coordinatively unsaturated metal species resulting from proton loss, as shown in equation 1 for a generic neutral hydride complex.<sup>10,14,15</sup>



In the absence of a suitable base the stoichiometry of the electrochemical oxidation of the metal hydride has been found to be an overall one-electron process in a few systems rather than the two-electron process discussed above.<sup>10,13</sup> One of the observed outcomes of this one-electron process has been H<sub>2</sub> evolution<sup>13,16,17</sup> with a net oxidation of one electron per metal hydride, as shown in equation 2.






One mechanism of H<sub>2</sub> evolution has been suggested by Collman et al.<sup>13</sup> in the case of a porphyrin-based Ru-hydride system. They hypothesize that two molecules of the oxidized Ru-hydride converge in an associative fashion to form a bridging [Ru(III)-H...H-Ru(III)] complex which subsequently loses H<sub>2</sub> by solvent coordination to yield two equivalents of the Ru(II)-solvento species, the overall transformation shown in equation 2. The other mechanism that has been evoked<sup>10</sup> is deprotonation-oxidation, wherein a neutral metal hydride accepts a proton from the cation radical to make an H<sub>2</sub> complex. A solvent molecule then displaces H<sub>2</sub> from the latter, and the Ru(I) species loses an electron to the electrode and coordinates a solvent, leading to the same overall stoichiometry shown in equation 2.

In this work, we have studied the oxidation of a new family of d<sup>8</sup> square-planar nickel hydride complexes supported by bis(phosphinite) pincer ligands (POCOP ligands).<sup>18-21</sup> This work expands the understanding of metal hydride oxidation. These nickel pincer complexes were synthesized and completely characterized by <sup>1</sup>H, <sup>31</sup>P, <sup>13</sup>C NMR spectroscopy, solid-state IR spectroscopy, elemental analysis, and X-ray crystallography. One-electron oxidation of all these complexes was found to be irreversible, and the nature of the post-oxidative chemical transformations leading to facile H<sub>2</sub> evolution were explored by a combination of chemical and electrochemical studies, density functional theory (DFT) based calculations and electrochemical simulations. The bimolecular rate constant for H<sub>2</sub> evolution was found to be at least 10<sup>7</sup> M<sup>-1</sup> s<sup>-1</sup> which is, to the best of our knowledge, the fastest known oxidative H<sub>2</sub> evolution reaction observed in transition metal hydride complexes.<sup>13,22</sup> This reactivity highlights the challenges of developing earth-abundant metal hydride complexes as catalytic intermediates in the ergoneutral oxidation of liquid fuels.

## Results and Discussion

## Synthesis and Characterization of Nickel Pincer Complexes

The desired nickel pincer hydrides **3a-d** were successfully synthesized in three consecutive steps following a procedure similar to the one reported by Guan et al.<sup>23</sup> Related bis(phosphinite) pincer ligands (**1a-d**, POCOP ligands) were prepared by Brookhart's method involving the treatment of substituted resorcinol derivatives with two equivalents of P(<sup>t</sup>Bu)<sub>2</sub>Cl in the presence of NaH.<sup>24,25</sup> These ligands require handling under inert atmosphere to avoid hydrolysis, which takes place in ambient atmosphere.<sup>26</sup> Nickel pincer chlorides **2a** and **2d** were prepared through cyclometallation of **1a** and **1d** with anhydrous NiCl<sub>2</sub>.<sup>23,25</sup> Nickel chlorides **2b** and **2c** were synthesized similarly (78-84% isolated yields) and characterized by <sup>1</sup>H, <sup>31</sup>P{<sup>1</sup>H}, <sup>13</sup>C{<sup>1</sup>H} NMR spectroscopy, and elemental analysis (Scheme 2). Solid-state structures of **2b** and **2c** were also determined by X-ray crystallography (see supplementary information). Both of the complexes adopt a square planar geometry around the Ni(II) center, a typical structure for this type of pincer complexes reported by Zargarian<sup>26</sup> and Guan.<sup>23</sup> Alternatively, **2a-d** could also be formed by refluxing a mixture of a resorcinol derivative, P(<sup>t</sup>Bu)<sub>2</sub>Cl, dimethylaminopyridine (DMAP), NiCl<sub>2</sub>, and 1,4-dioxane (solvent) in a single reaction vessel for several days

Scheme 2 Synthesis of the <sup>t</sup>Bu[POCOP]NiCl Complexes.Scheme 3 Single-Pot Synthesis of the <sup>t</sup>Bu[POCOP]NiCl Complexes.Scheme 4 Synthesis of the <sup>t</sup>Bu[POCOP]NiH Complexes.

(Scheme 3). Although this method avoids the separation of the air-sensitive POCOP ligands, the corresponding isolated yields of **2a-d** were much lower (19-37%).

Reactions of **2a-d** with excess  $\text{LiAlH}_4$  (20 equivalents) at room temperature generated the desired nickel pincer hydride complexes **3a-d** as light yellow solids in good isolated yields (65-78%, Scheme 4). Complexes **3b-d** are not reported in the literature and therefore were completely characterized by  $^1\text{H}$ ,  $^{31}\text{P}\{^1\text{H}\}$ ,  $^{13}\text{C}\{^1\text{H}\}$  NMR spectroscopy, IR spectroscopy, and elemental analysis. The  $^1\text{H}$  NMR spectra of **3b-d** recorded in  $\text{C}_6\text{D}_6$  revealed the characteristic Ni-H resonances as triplets at  $\delta -8.06$  ( $^2J_{\text{H-P}} = 52\text{Hz}$ ),  $-7.98$  ppm ( $^2J_{\text{H-P}} = 52\text{Hz}$ ), and  $-7.90$  ( $^2J_{\text{H-P}} = 52\text{Hz}$ ), respectively. The singlet resonance observed in the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra of all of the nickel hydride species is consistent with a  $\text{C}_2$ -symmetric structure. Solid-state IR spectroscopy showed distinct, moderately strong Ni-H stretching bands near  $1791\text{ cm}^{-1}$  (for **3b**),  $1768\text{ cm}^{-1}$  (for **3c**), and  $1810\text{ cm}^{-1}$  (for **3d**). Compounds **3b-d** have been further characterized by X-ray crystallography (Figure 1, see supporting information). The hydride ligands in all of these nickel hydride complexes were located and refined. Crystal




Fig. 1 X-ray Crystal Structure of **3b** (50% probability level). Selected bond lengths ( $\text{\AA}$ ) and angles (deg): Ni(1)–H(1) 1.47(2), Ni(1)–C(1) 1.8876(14), Ni(1)–P(1) 2.1155(6), Ni(1)–P(2) 2.1161(6), P(1)–Ni(1)–P(2) 165.288(17), C(1)–Ni(1)–H(1) 177.9(8), P(1)–Ni(1)–H(1) 95.7(8), P(2)–Ni(1)–H(1) 99.0(8).

data and refinement parameters for the nickel hydride complexes are given in Table 1.

#### Synthesis of the Nickel Pincer Acetonitrile Complex

The related nickel pincer acetonitrile complex (**4a**) was independently synthesized from two different routes (Scheme 5). In one of the methods, the chloride ligand in **2a** was abstracted by an  $\text{Ag(I)}$  salt in acetonitrile to form the desired complex **4a**. Tetrafluoroborate ( $\text{BF}_4^-$ ), remains as a counter-anion. In the second method, the corresponding nickel hydride complex **3a** was protonated by  $\text{HBF}_4 \cdot \text{Et}_2\text{O}$  to afford complex **4a**. In this reaction,  $\text{H}_2$  is released as the only other byproduct. Complex **4a** has been characterized by  $^1\text{H}$ ,  $^{31}\text{P}\{^1\text{H}\}$ ,  $^{13}\text{C}\{^1\text{H}\}$  NMR spectroscopy, IR spectroscopy, and elemental analysis. In the  $^1\text{H}$  NMR spectrum, the resonance corresponding to the Ni– $\text{NCCH}_3$  appeared at  $\delta 2.63$  as a broad triplet. A strong  $\text{C}\equiv\text{N}$  stretch for the acetonitrile moiety present in complex **4a** was observed at  $2290\text{ cm}^{-1}$  in the IR spectrum. A similar nickel

Table 1 Crystal Data and Refinement Parameters for **3b-d**

	<b>3b</b>	<b>3c</b>	<b>3d</b>
empirical formula	$\text{C}_{23}\text{H}_{42}\text{O}_3\text{P}_2\text{Ni}$	$\text{C}_{23}\text{H}_{42}\text{O}_2\text{P}_2\text{Ni}$	$\text{C}_{30}\text{H}_{56}\text{O}_2\text{P}_2\text{Ni}$
crystal system	monoclinic	monoclinic	monoclinic
space group	$P2_1/n$	$P2_1/n$	$P2_1/n$
a, $\text{\AA}$	9.756(3)	9.528(3)	15.131(3)
b, $\text{\AA}$	19.408(6)	20.201(7)	12.589(2)
c, $\text{\AA}$	14.026(4)	13.4341(5)	17.529(3)
$\alpha$ , deg	90	90	90
$\beta$ , deg	104.528(6)	100.908(7)	106.717(4)
$\gamma$ , deg	90	90	90
Volume, $\text{\AA}^3$	2570.9(14)	2539.1(16)	3197.8(11)
Z	4	4	4
no. of data	64153	80520	66747
no. of unique data,	14241, 0.0784	13995, 0.0707	14087, 0.0980
R1, wR2 ( $I > 2\sigma(I)$ )	0.0493, 0.0990	0.0571, 0.1174	0.0527, 0.1088
R1, wR2 (all data)	0.0949, 0.1152	0.0868, 0.1264	0.0988, 0.1271


acetonitrile complex was reported by Crabtree and Hazari during their investigation related to the electrocatalytic production of hydrogen.<sup>27</sup>


#### Electrochemistry of the Nickel Hydride Complexes

The cyclic voltammograms of the hydride complexes **3a-d** are shown in Figure 2A. A mixture of THF–MeCN 1:4 by volume respectively, was employed as the solvent mixture for all electrochemistry experiments as **3a-d** are only partially soluble in MeCN. All of these complexes show an irreversible oxidation peak, assigned to a metal-centered oxidation, at slightly different peak potentials depending on the substituent on the aromatic ring (peak potentials ranging from 101 to 316 mV vs  $\text{Fc}^+/\text{Fc}$ ). The complete irreversibility in the CV of these hydride complexes suggests a fast chemical step involving a hydride, proton or H-atom transfer following initial electron transfer from the molecule to the electrode.<sup>28</sup> The peak areas in these CVs are comparable to the peak area observed in an equimolar solution of ferrocene or the corresponding chloride complex **2a** (Figure 2B, S4) under similar conditions. This provides evidence that the peak is due to a one-electron oxidation and argues against the deprotonation pathway in equation 1, which would result in a two-electron oxidation.

On scanning further positive from the peak potential of the initial Ni(III)H/Ni(II)H oxidation in the CV, another wave was observed ca. 0.5 V positive of the initial wave (Figure 2C). This second partially reversible wave probably corresponds to the

Scheme 5 Synthesis of the Nickel Pincer Acetonitrile Complexes.





**Fig. 2** (A) CV of 1 mM solutions of **3a-d** in 1:4 by volume THF-MeCN (0.1 M TBABF<sub>4</sub>, TBA = Tetra-*N*-butyl ammonium) at 250 mV/s. (B) Comparison of CVs of 4 mM solutions of **2a**, **3a** and Ferrocene in 1:4 by volume THF-MeCN (0.1 M TBABF<sub>4</sub>) measured at a scan rate of 250 mV/s. (C) CV of **3a** beyond the first one-electron oxidation in THF (0.1 M TBABF<sub>4</sub>) for different scan rates.

Ni(III/II) couple of a new species generated during the first oxidation, with a ligand being less donating compared to the hydride ligand in **3a**, as evidenced by the more positive  $E_{1/2}$  of this couple. The magnitude of this second oxidation peak relative to the first decreases with decreasing scan rate suggesting that this second Ni(II) species is partially converting to another species on the seconds time scale.

To probe the nature of the products formed upon oxidation, a one-electron chemical oxidant, Ce(NBu<sub>4</sub>)<sub>2</sub>(NO<sub>3</sub>)<sub>6</sub>,<sup>29</sup> was added in 1/3 equivalent increments to **3a** dissolved in a 50:50 mixture of CD<sub>3</sub>CN and C<sub>6</sub>D<sub>6</sub> in a J. Young NMR tube and the resultant changes monitored by <sup>1</sup>H and <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy

(Figure 3). The triplet centered at  $\delta$  -8.39 ( $J=52$  Hz) corresponds to the hydride resonance in **3a**. With incremental addition of the Ce(IV) oxidant to **3a**, we observed a gradual disappearance of the hydride resonance of **3a** from the <sup>1</sup>H NMR spectrum. In the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum, the intensity of the peak at  $\delta$  221.42 ppm corresponding to **3a** also decreased with the addition of the chemical oxidant. After the addition of 1 equivalent of the Ce(IV) salt, the hydride resonance was completely replaced with the new resonance at  $\delta$  199.57 ppm corresponding to the nickel-solvento species **4a** (Figure 3). After the initial addition of the Ce(IV) salt, bubbles appeared in the NMR tube as well as a singlet peak in the <sup>1</sup>H NMR spectrum ( $\delta$  4.42 ppm, Figure 3 inset) corresponding to H<sub>2</sub> production.<sup>30</sup> To quantify the amount of H<sub>2</sub> produced, **3a** was taken in two NMR tubes (5 mg each, 0.011 mmol) each sealed with a septum and dissolved in 0.2 ml d<sup>8</sup>-THF. Sub-stoichiometric amounts of 2,6-dimethylbenzoic acid and Ce(NBu<sub>4</sub>)<sub>2</sub>(NO<sub>3</sub>)<sub>6</sub>, each in 0.8 ml of CD<sub>3</sub>CN were added to the first and second NMR tubes respectively. The integration of the H<sub>2</sub> peak in the <sup>1</sup>H NMR produced ( $\delta$  4.55 ppm in this solvent mixture) was quantified using acid addition to the second test tube as a calibrant (Figure S1). To be noted is the fact that the calibration sample contains twice the amount of H<sub>2</sub> per Ni-H as we are using an acid as the proton source. By this method, H<sub>2</sub> production on one electron oxidation by Ce(IV) amounted to 95%, which corresponds to the stoichiometry in equation 2. These observations provide strong evidence for the oxidative H<sub>2</sub> evolution pathway (equation 2).

To further quantify the electron stoichiometry and the amount of dihydrogen produced, we performed a constant potential electrolysis of 3 mL of a 1.23 mM solution of **3a** in THF:MeCN (1:4 by volume), maintaining a potential *ca.* 200 mV positive of the Ni(III)H/Ni(II)H peak potential. For this amount of **3a** and a net one-electron oxidation, the total charge to be passed for 100% Faradaic efficiency would be 356 mC. Integrating the current versus time plot for the electrolysis (Figure 4A) gave 354 mC, confirming a net one-electron stoichiometry with respect to **3a** for the oxidation. After the electrolysis was complete, a 3 mL aliquot of the headspace above the solution was analyzed by gas chromatography. The results shown in Figure 4B clearly indicate hydrogen production based on the retention time of a standard 1% H<sub>2</sub> gas sample. Comparison of the areas of the sample versus the reference gave an H<sub>2</sub> partial pressure of at least 0.007 atm in the 4.5 mL of headspace which corresponds to  $1.3 \times 10^{-3}$  mmol H<sub>2</sub> and a yield of at least 70% based on equation 2. The yield is a lower bound given the solubility of H<sub>2</sub> in the solvent mixture as well as possible crossover between the two chambers of the H-cell through diffusion. The new peak in the post-electrolysis cyclic voltammogram shown in Figure 4A inset (blue) is tentatively assigned to the Ni-solvento species (**4a**) formed during the electrolysis (Figure S7).

#### Mechanistic Insights from DFT-based Calculations

In order to better understand the mechanism of the oxidative H<sub>2</sub> evolution process, we calculated the relative free energies




Fig. 3  $^1\text{H}$  and  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra for the oxidation of **3a** with ca. (A) 0 equiv. (B) 0.33 equiv. (C) 0.66 equiv. and (D) 1 equiv.  $\text{Ce}(\text{NBU}_4)_2(\text{NO}_3)_6$ .


of plausible intermediates along two different reaction coordinates corresponding to the disproportionation and deprotonation-oxidation mechanisms (Figure 5). The relative free energies of all intermediates are referenced to the  $\mathbf{3a}^+/\mathbf{3a}$  couple. As a result, note that the free energies of  $\mathbf{3a}^+$  and  $\mathbf{3a}$  are the same in Figure 5.<sup>31</sup>

**Mechanism 1: Bimolecular Disproportionation ( $E,E,C_i$ ).** Two Ni(III)-H cations are generated from the oxidation of two molecules of **3a** at the  $\mathbf{3a}^+/\mathbf{3a}$  potential which is taken as the reference for comparing the relative free energies of all other intermediates. From two molecules of Ni(III)-H, the formation of an  $\text{H}_2$ -bridged structure (see supporting information) is energetically downhill by -34.8 kcal/mol which is the highest driving force in the overall transformation.  $\text{H}_2$  loss from this

structure to generate two cationic coordinatively unsaturated Ni(II) species is again downhill by about -11.5 kcal/mol. The final step involving ligation of these species by MeCN is a very energetically favorable process leading to the products.

**Mechanism 2: Deprotonation-Oxidation ( $E,C_iE_r'$ ).** As seen in Figure 5 the deprotonation of the *protic* H from the newly generated Ni(III)-H cation by the *hydridic* H of a Ni(II)-H in the depletion layer of the electrode could also result in  $\text{H}_2$  evolution. However, formation of the bridging structure (see supporting information) in this case is downhill only by ca. -3 kcal/mol.  $\text{H}_2$  dissociation from this bimolecular intermediate leads to a coordinatively unsaturated neutral Ni(I) species and the cationic Ni(II) species. A second electron-transfer step oxidizing the Ni(I) to Ni(II), which is energetically favorable at

Fig. 4 (A) Current vs. time plot for the bulk electrolysis of **3a** in a 1:4 THF-MeCN solvent mixture in presence of 0.1 M TBABF<sub>4</sub>. Constant potential electrolysis was performed at 500 mV vs.  $\text{Fc}^+/\text{Fc}$  which is ca. 200 mV positive of the peak oxidation potential of **3a**. Inset shows the CVs of the sample taken before (red) and after (blue) the electrolysis experiment at 1 V/s, which shows the formation of **4a**. (B) Gas chromatogram for the headspace of the solution after bulk electrolysis. Inset shows the magnified image of the  $\text{H}_2$  peak superimposed on the 1%  $\text{H}_2$  standard.



**Fig. 5** Reaction coordinate diagram with mechanisms a) E,E,C<sub>i</sub> (in blue) and b) E,C<sub>i</sub>E<sub>i</sub>' (in red) leading to H<sub>2</sub> evolution. Relative free energies given in eV and kcal/mol. Level of theory: Gaussian09<sup>TM</sup>/M06/LANL2DZ(Ni)/6-311G(d) (C,H,O,N,P)/SMD(MeCN). 'E' and 'C' denote electrochemical and chemical steps respectively, and 'r' and 'i' denote reversible and irreversible steps respectively.

the (Ni(III)H/Ni(II)H) electrode potential, followed by MeCN coordination yields the products. The last step is the same as in mechanism 1.

From the electrode potential of the Ni(III)H/Ni(II)H couple, all intermediates in either pathway are energetically downhill. However, the association step to form the Ni(III)-Ni(III) H<sub>2</sub> bridging structure has a much higher driving force (-34.8 kcal/mol) compared to the analogous step in mechanism 2 (-3.2 kcal/mol).

Changes in the pK<sub>a</sub> and BDFE of **3a** on one-electron oxidation were computed at the same level of theory as above using isodesmic reaction schemes with reference compounds (see computational details). The results shown in Tables S1 and S2 indicate substantial thermodynamic activation towards deprotonation as well as H-atom abstraction (See supplementary information) which is in line with the high exergonicity of the whole process at the (Ni(III)H/Ni(II)H) potential leading to H<sub>2</sub> evolution. Our calculations predict a decrease in pK<sub>a</sub> of about 46 units on going from Ni(II)H to the Ni(III)H cation (Table S2). This result was cross-verified with five other density functionals ranging from local GGA to hybrid-GGA<sup>32,33</sup> and was found to be consistent (Figure S8). This difference in pK<sub>a</sub> is much higher than the values reported in the literature for octahedral Ru hydrides.<sup>10</sup> This result is consistent with the high oxidation potential of the parent hydride **3a** as compared to those reported for the Ru hydrides.



Given the calculated pK<sub>a</sub> values of the Ni(III)H cation we expect deprotonation of the metal hydride by the solvent. However, our observation of quantitative H<sub>2</sub> production (vide

supra) from the Ni complexes provides evidence against any *net* contribution from deprotonation of the Ni(III)H cation by the solvent. The likely pathway involves half the nickel hydrides deprotonating after oxidation at the electrode surface, with the released protons reacting with Ni(II) hydride diffusing towards the electrode surface in the depletion layer to make H<sub>2</sub>, followed by the oxidation of the deprotonated Ni at the electrode to a Ni(II) species (E<sub>r</sub>C<sub>i</sub>E<sub>i</sub>').

#### Electrochemical Simulation to Obtain Rate Constant for the Following Chemical Reaction.

Fast scan-rate CV experiments using ultramicroelectrodes were performed in an attempt to observe a return wave after oxidative of **3a** (Figure S2), which would then enable a reliable estimate of the rate of the following chemical reaction. However, the CV's were irreversible up to a scan-rate of 2000 V/s. Concentration dependent CVs down to 0.125 mM **3a** also did not show a return reduction wave (Figure S3). Even higher scan rates or lower concentrations did not yield anything substantive due to the ratio of faradaic to capacitive current becoming too low for meaningful interpretations. We therefore simulated our electrochemical data in DigiElch 7<sup>TM</sup> to estimate the lower limit of the rate of the following chemical reaction. CVs were simulated from two different regimes, viz. scan rate of 4 V/s using a 3 mm diameter disk electrode, and scan rate of 400 V/s using an 11 μm diameter disk electrode were chosen (see supplementary information for complete details of the simulations). 400 V/s was the highest scan rate at

Scheme 6 Mechanisms for hydrogen production.

**Mechanism 1: E<sub>r</sub>E<sub>r</sub>C<sub>i</sub>****Mechanism 2: E<sub>r</sub>C<sub>i</sub>E<sub>r</sub>'**

which the faradaic and capacitive components of the current could be reasonably separated.

To estimate the standard electron-transfer rate constant  $k_s$ , experimentally measured CVs of the corresponding chloride complex **2a** were simulated with a planar 1D semi-infinite diffusion model as implemented in DigiElch 7™, shown in Figure S4. Estimates of the lower limit of the uncompensated resistance of the cells,  $R_u$ , were made based on the cell geometry and published electrolyte conductivities.<sup>34</sup> A sensitivity analysis was then performed to estimate the range


**Table 2** Simulated lower limits of  $E^\circ$  and  $k_f$  for E<sub>r</sub>E<sub>r</sub>C<sub>i</sub> and E<sub>r</sub>C<sub>i</sub>E<sub>r</sub>' mechanisms in the two limits:

At 4V/s, Limit I:  $k_s = 0.1$  cm/s &  $R_u = 650 \Omega$ . Limit II:  $k_s = 0.003$  cm/s &  $R_u = 253 \Omega$ .  
At 400 V/s, Limit I:  $k_s = 5$  cm/s &  $R_u = 9 \times 10^5 \Omega$ . Limit II:  $k_s = 0.2$  cm/s &  $R_u = 5 \times 10^5 \Omega$ .

Mechanism	Scan Rate (V/s)	$E^\circ_{(\text{min})}$ (V) vs. $\text{Fc}^+/\text{Fc}$		$k_{f(\text{min})}$ ( $\text{M}^{-1}\text{s}^{-1}$ )	
		I	II	I	II
E <sub>r</sub> E <sub>r</sub> C <sub>i</sub>	400	0.33	0.33	$1.0 \times 10^7$	$1.0 \times 10^7$
	4	0.30	0.30	$1.0 \times 10^6$	$1.0 \times 10^6$
E <sub>r</sub> C <sub>i</sub> E <sub>r</sub> '	400	0.35	0.35	$5.0 \times 10^6$	$5.0 \times 10^6$
	4	0.32	0.32	$7.0 \times 10^4$	$2.5 \times 10^4$

of  $R_u$  and  $k_s$  that fit this data (see supplementary information for details).

Using this information, CVs of **3a** were simulated according to both of the mechanisms (Scheme 6) viz. bimolecular disproportionation (E<sub>r</sub>E<sub>r</sub>C<sub>i</sub>) and deprotonation-oxidation (E<sub>r</sub>C<sub>i</sub>E<sub>r</sub>') and the results are summarized in Figure 6 and Table 2. The reported  $k_f$  and  $E^\circ$  correspond to approximately the lowest values above which there is no return wave in the CVs, which was verified by sensitivity analysis of the simulation (Figure S5b).



**Fig. 6** Experiments and simulations of CVs of **3a** at (A) and (B) 4 V/s with a 3 mm diameter working electrode; (C) and (D) 400 V/s with a 11  $\mu\text{m}$  diameter working electrode. See supplementary information for details.



At 400 V/s, the predicted lower limit for  $k_f$  is on the order of  $10^7 \text{ M}^{-1} \text{ s}^{-1}$  for  $E_r E_c C_i$  and  $5 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$  for  $E_r C_i E_r'$ , both mechanisms therefore being extremely facile. Higher scan-rate simulations with smaller electrodes provide a higher and therefore better estimate for  $k_f$  and  $E^\circ$  than their lower scan-rate counterparts. Consistent with these expectations, the simulations at 4 V/s gave slightly lower values for  $k_{f(\text{min})}$  and  $E^\circ_{(\text{min})}$ , shown in Table 2. It is interesting to note the difference in the bimolecular rates of these two mechanisms predicted by the simulations, with  $k_f$  ( $E_r E_c C_i$ ) being slightly larger than  $k_f$  ( $E_r C_i E_r'$ ) in either frequency regime by a factor of two, although both mechanisms fit the experimental data equally well.

## Conclusions

If earth-abundant metal hydrides are to be employed as electrocatalysts in fuel cells (Scheme 1), it is important to fully understand their reactivity on electron transfer out of the metal-hydrogen bond. The nickel (II) hydrides synthesized in this work have proven to be highly stable targets for such electrochemical studies. We have shown in this work that the one-electron oxidation in these  $d^8$  square-planar nickel hydride complexes leads to highly exergonic and extremely rapid post-oxidative  $\text{H}_2$  evolution instead of the production of protons and electrons as was originally desired. This work therefore highlights an important challenge in using transition metal hydrides to electro-oxidize hydrogen containing liquid fuels in a fuel cell. It is possible that site isolation of these hydrides on a solid support would suppress this bimolecular  $\text{H}_2$  evolution pathway on oxidation, and lead to protons and electrons instead. However the high positive potentials required for the first oxidation of these hydrides render them unlikely targets for ergoneutral electrocatalysts.

## Experimental Details

### Cyclic Voltammetry

Cyclic voltammetry experiments were performed with a SP-200 potentiostat from Bio-Logic Co. with a glassy carbon 3 mm diameter disk electrode (MF-2012) or a carbon fibre tip 11  $\mu\text{m}$  diameter ultramicroelectrode (MF-2007) from BASi Inc. as the working electrode, a  $\text{Ag}/\text{AgNO}_3$  (10 mM), MeCN electrode as the non-aqueous reference electrode, and a Pt wire as the counter electrode in a typical three-electrode cell. TBABF<sub>4</sub> (0.1 M) was used as the supporting electrolyte. All experiments were done under  $\text{N}_2$  in a glove-box.

Bulk electrolysis was performed in an 'H-cell' with two cylindrical glass chambers of 2 cm diameter connected by a glass joint fitted with a sintered glass frit. A 2 cm x 3 cm rectangular piece of carbon cloth suspended into the analyte solution with a copper wire was used as the working electrode. A finely polished and air-dried silver wire (0.032" diameter) was used as the pseudo reference in the same chamber. A polished Pt wire (0.032" diameter) was used as the counter electrode in the other compartment of the cell. Both compartments were sealed with rubber septa to prevent the

escape of any gas generated. After bulk electrolysis, percent  $\text{H}_2$  was determined by headspace analysis using a Shimadzu Gas Chromatograph equipped with a 100  $\mu\text{L}$  sample loop, a TCD, and a CarbonPLOT column with  $\text{N}_2$  carrier gas. Percent  $\text{H}_2$  was quantified by a one point calibration with a 1%  $\text{H}_2$  standard.

### General Experimental Information

Unless otherwise noted, all organometallic compounds were prepared and handled under a nitrogen atmosphere using standard Schlenk and glovebox techniques. Dry and oxygen-free solvents (benzene, THF, pentane, and ether) were collected from an Innovative Technology PS-MD-6 solvent purification system and used throughout the experiments. Toluene and  $\text{C}_6\text{D}_6$  were distilled from a deep purple solution of Na and benzophenone under a nitrogen atmosphere.  $\text{CH}_3\text{CN}$  was dried using  $\text{CaH}_2$  and stored over 4 $\text{\AA}$  molecular sieves. Other solvents such as  $\text{CD}_2\text{Cl}_2$  and  $\text{CDCl}_3$  were used as received from commercial sources.  $^1\text{H}$ ,  $^{13}\text{C}$ , and  $^{31}\text{P}$  NMR spectra were recorded on a Bruker Avance-400 MHz spectrometer. Chemical shift values in  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were referenced internally to the residual solvent resonances, whereas  $^{31}\text{P}$  NMR spectra were referenced externally to 85%  $\text{H}_3\text{PO}_4$  (0 ppm). Infrared spectra were recorded in the solid state on a Thermo Scientific Nicolet 4700 FT-IR spectrometer equipped with smart orbit diamond attenuated total reflectance (ATR) accessory. Elemental analysis was performed by the CENTC Elemental Analysis Facility at the University of Rochester Using a Perkin Elmer 2400 Series II elemental analyzer in CHN mode. Preparation of 1,3-( $^t\text{Bu}_2\text{PO}$ )<sub>2</sub> $\text{C}_6\text{H}_4$  (**1a**),<sup>24</sup> 5-OCH<sub>3</sub>-1,3-( $^t\text{Bu}_2\text{PO}$ )<sub>2</sub> $\text{C}_6\text{H}_3$  (**1b**),<sup>24</sup> 5-CH<sub>3</sub>-1,3-( $^t\text{Bu}_2\text{PO}$ )<sub>2</sub> $\text{C}_6\text{H}_3$  (**1c**),<sup>24</sup> 4,6-( $^t\text{Bu}$ )<sub>2</sub>-1,3-( $^t\text{Bu}_2\text{PO}$ )<sub>2</sub> $\text{C}_6\text{H}_2$  (**1d**),<sup>25</sup> **2a**,<sup>23</sup> **2d**,<sup>25</sup> and **3a**<sup>23</sup> have been previously reported in the literature.

### Synthesis of [5-OCH<sub>3</sub>-1,3-( $^t\text{Bu}_2\text{PO}$ )<sub>2</sub> $\text{C}_6\text{H}_2$ ]NiCl (**2b**)

Under a nitrogen atmosphere, 50 mL of toluene was added to a mixture of **1b** (0.86 g, 2.0 mmol) and anhydrous  $\text{NiCl}_2$  (285 mg, 2.2 mmol), giving an orange suspension. While the solution was being refluxed for 24 h, an orange solution resulted with some undissolved residues. After the mixture was cooled to room temperature, the residues were removed by gravity filtration and a clear orange filtrate was obtained. The volume of the orange filtrate was reduced to ca. 5 mL, and then  $\text{Et}_2\text{O}$  was added to cause precipitation of a yellow solid. The product was collected by filtration, washed with  $\text{Et}_2\text{O}$ , and dried under vacuum to afford a yellow powder of **2b** (875 mg, 84% yield). X-ray-quality crystals were grown by allowing a layer of pentane to slowly diffuse into a saturated  $\text{CH}_2\text{Cl}_2$  solution of the complex at 0  $^\circ\text{C}$ .  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ,  $\delta$ ): 1.42 (virtual triplet,  $\text{PC}(\text{CH}_3)_3$ ,  $J_{\text{P-H}} = 8 \text{ Hz}$ , 36H), 3.65 (s,  $\text{OCH}_3$ , 3H), 5.98 (s, *ArH*, 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ ,  $\delta$ ): 28.13 (s,  $\text{PC}(\text{CH}_3)_3$ ), 39.32 (t,  $J_{\text{P-C}} = 7.1 \text{ Hz}$ ,  $\text{PC}(\text{CH}_3)_3$ ), 55.47 (s,  $\text{OCH}_3$ ), 91.98 (t,  $J_{\text{P-C}} = 6.1 \text{ Hz}$ , *ArC*), 114.24 (t,  $J_{\text{P-C}} = 22.2 \text{ Hz}$ , *ArC*), 161.38 (s, *ArC*), 169.68 (t,  $J_{\text{P-C}} = 10.1 \text{ Hz}$ , *ArC*).  $^{31}\text{P}\{^1\text{H}\}$  NMR (162 MHz,  $\text{CDCl}_3$ ,  $\delta$ ): 185.66 (s). Anal. Calcd for  $\text{C}_{23}\text{H}_{41}\text{ClO}_3\text{P}_2\text{Ni}$ : C, 52.96; H, 7.92. Found: C, 52.93; H, 7.89.

**Synthesis of [5-CH<sub>3</sub>-1,3-(<sup>t</sup>Bu<sub>2</sub>PO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>]NiCl (2c)**

This compound was prepared from **1c** (1.66 g, 4.02 mmol) and anhydrous NiCl<sub>2</sub> (573 mg, 4.42 mmol) in 78% yield (1.59 g) by a procedure similar to that used for **2b**. X-ray-quality crystals were grown by allowing a layer of pentane to slowly diffuse into a saturated CH<sub>2</sub>Cl<sub>2</sub> solution of the complex at 0 °C. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, δ): 1.41 (virtual triplet, PC(CH<sub>3</sub>)<sub>3</sub>, J<sub>p-H</sub> = 8 Hz, 36H), 2.13 (s, CH<sub>3</sub>, 3H), 5.18 (s, ArH, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>, δ): 21.55 (s, CH<sub>3</sub>), 28.13 (s, PC(CH<sub>3</sub>)<sub>3</sub>), 39.27 (t, J<sub>p-C</sub> = 7.1 Hz, PC(CH<sub>3</sub>)<sub>3</sub>), 105.92 (t, J<sub>p-C</sub> = 6.1 Hz, ArC), 120.56 (t, J<sub>p-C</sub> = 21.2 Hz, ArC), 139.02 (s, ArC), 169.44 (t, J<sub>p-C</sub> = 10.1 Hz, ArC). <sup>31</sup>P{<sup>1</sup>H} NMR (162 MHz, CDCl<sub>3</sub>, δ): 184.04 (s). Anal. Calcd for C<sub>23</sub>H<sub>41</sub>ClO<sub>2</sub>P<sub>2</sub>Ni: C, 54.63; H, 8.17. Found: C, 54.75; H, 8.07.

**Synthesis of [5-OCH<sub>3</sub>-1,3-(<sup>t</sup>Bu<sub>2</sub>PO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>]NiH (3b)**

Under a nitrogen atmosphere a suspension of LiAlH<sub>4</sub> (729 mg, 19.2 mmol) and **2b** (500 mg, 0.96 mmol) in 40 mL of benzene was stirred at room temperature for 24 h. The resulting mixture was filtered through a short plug of Celite to afford a yellow solution. After the solvent was evaporated under vacuum, the desired nickel hydride **3b** was isolated as a light yellow powder (322 mg, 69% yield). X-ray-quality crystals were grown by layering CH<sub>3</sub>OH on a saturated THF solution of the hydride at 0 °C and allowing it to slowly diffuse. <sup>1</sup>H NMR (400 MHz, C<sub>6</sub>D<sub>6</sub>, δ): -8.06 (t, J<sub>p-H</sub> = 52 Hz, NiH, 1H), 1.32 (virtual triplet, PC(CH<sub>3</sub>)<sub>3</sub>, J<sub>p-H</sub> = 8 Hz, 36H), 3.36 (s, OCH<sub>3</sub>, 3H), 6.56 (s, ArH, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, C<sub>6</sub>D<sub>6</sub>, δ): 28.56 (t, J<sub>p-C</sub> = 3.0 Hz, PC(CH<sub>3</sub>)<sub>3</sub>), 37.88 (t, J<sub>p-C</sub> = 9.1 Hz, PC(CH<sub>3</sub>)<sub>3</sub>), 55.06 (s, OCH<sub>3</sub>), 92.12 (t, J<sub>p-C</sub> = 6.1 Hz, ArC), 133.36 (t, J<sub>p-C</sub> = 15.2 Hz, ArC), 162.50 (s, ArC), 169.09 (t, J<sub>p-C</sub> = 10.1 Hz, ArC). <sup>31</sup>P{<sup>1</sup>H} NMR (162 MHz, C<sub>6</sub>D<sub>6</sub>, δ): 218.20 (s). IR (ATR, cm<sup>-1</sup>): ν<sub>Ni-H</sub> = 1790. Anal. Calcd for C<sub>23</sub>H<sub>42</sub>O<sub>3</sub>P<sub>2</sub>Ni: C, 56.70; H, 8.69. Found: C, 56.28; H, 8.63.

**Synthesis of [5-CH<sub>3</sub>-1,3-(<sup>t</sup>Bu<sub>2</sub>PO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>]NiH (3c)**

This compound was prepared from **2c** (500 mg, 0.99 mmol) and LiAlH<sub>4</sub> (751 mg, 19.8 mmol) in 65% yield (294 mg) by a procedure similar to that used for **3b**. X-ray-quality crystals were grown by layering CH<sub>3</sub>OH on a saturated hexane solution of the nickel hydride at 0 °C and allowing it to slowly diffuse. <sup>1</sup>H NMR (400 MHz, C<sub>6</sub>D<sub>6</sub>, δ): -7.98 (t, J<sub>p-H</sub> = 52 Hz, NiH, 1H), 1.33 (virtual triplet, PC(CH<sub>3</sub>)<sub>3</sub>, J<sub>p-H</sub> = 8 Hz, 36H), 2.19 (s, CH<sub>3</sub>, 3H), 6.67 (s, ArH, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, C<sub>6</sub>D<sub>6</sub>, δ): 21.94 (s, CH<sub>3</sub>), 28.57 (s, PC(CH<sub>3</sub>)<sub>3</sub>), 37.85 (t, J<sub>p-C</sub> = 8.1 Hz, PC(CH<sub>3</sub>)<sub>3</sub>), 105.78 (s, ArC), 138.72 (s, ArC), 139.10 (t, J<sub>p-C</sub> = 16.2 Hz, ArC), 168.94 (t, J<sub>p-C</sub> = 9.1 Hz, ArC). <sup>31</sup>P{<sup>1</sup>H} NMR (162 MHz, C<sub>6</sub>D<sub>6</sub>, δ): 216.22 (s). IR (ATR, cm<sup>-1</sup>): ν<sub>Ni-H</sub> = 1768. Anal. Calcd for C<sub>23</sub>H<sub>42</sub>O<sub>2</sub>P<sub>2</sub>Ni: C, 58.62; H, 8.98. Found: C, 58.78; H, 8.92.

**Synthesis of [4,6-(<sup>t</sup>Bu)<sub>2</sub>-1,3-(<sup>t</sup>Bu<sub>2</sub>PO)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>]NiH (3d)**

This compound was prepared from **2d** (500 mg, 0.83 mmol) and LiAlH<sub>4</sub> (630 mg, 16.6 mmol) in 78% yield (379 mg) by a procedure similar to that used for **3b**. X-ray-quality crystals were grown by layering CH<sub>3</sub>OH on a saturated hexane solution of the nickel hydride at 0 °C and allowing it to slowly diffuse.

<sup>1</sup>H NMR (400 MHz, C<sub>6</sub>D<sub>6</sub>, δ): -7.90 (t, J<sub>p-H</sub> = 56 Hz, NiH, 1H), 1.36 (virtual triplet, PC(CH<sub>3</sub>)<sub>3</sub>, J<sub>p-H</sub> = 8 Hz, 36H), 1.60 (s, C(CH<sub>3</sub>)<sub>3</sub>, 18H), 7.32 (s, ArH, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, C<sub>6</sub>D<sub>6</sub>, δ): 28.53 (s, PC(CH<sub>3</sub>)<sub>3</sub>), 30.48 (s, C(CH<sub>3</sub>)<sub>3</sub>), 34.38 (s, C(CH<sub>3</sub>)<sub>3</sub>), 37.64 (t, J<sub>p-C</sub> = 9.1 Hz, PC(CH<sub>3</sub>)<sub>3</sub>), 123.09 (s, ArC), 125.83 (t, J<sub>p-C</sub> = 5.1 Hz, ArC), 146.07 (t, J<sub>p-C</sub> = 16.2 Hz, ArC), 163.92 (t, J<sub>p-C</sub> = 9.1 Hz, ArC). <sup>31</sup>P{<sup>1</sup>H} NMR (162 MHz, C<sub>6</sub>D<sub>6</sub>, δ): 215.04 (s). IR (ATR, cm<sup>-1</sup>): ν<sub>Ni-H</sub> = 1810. Anal. Calcd for C<sub>30</sub>H<sub>56</sub>O<sub>2</sub>P<sub>2</sub>Ni: C, 63.28; H, 9.91. Found: C, 62.95; H, 9.75.

**Synthesis of [(1,3-(<sup>t</sup>Bu<sub>2</sub>PO)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>)Ni(NCCH<sub>3</sub>)]BF<sub>4</sub> (4a)**

*Method A (through halide substitution):* Under a nitrogen atmosphere, a flame-dried 50 mL Schlenk flask was charged with **2a** (300 mg, 0.61 mmol), AgBF<sub>4</sub> (142 mg, 0.73 mmol), and 15 mL CH<sub>3</sub>CN. The resulting mixture was stirred at room temperature for four hours to obtain a light yellow solution with some insoluble white precipitate. The solution was filtered through a short plug of Celite, washed with THF (2x10 mL), and the resulting filtrate was evaporated under vacuum to afford the desired nickel-acetonitrile complex **4a** as a light yellow powder (271 mg, 76% yield). *Method B (through protonation):* Under a nitrogen atmosphere, a flame-dried 50 mL Schlenk flask was loaded with **3a** (100 mg, 0.22 mmol), HBF<sub>4</sub>•Et<sub>2</sub>O (33 μL, 0.24 mmol), and 10 mL CH<sub>3</sub>CN. The resulting mixture was stirred at room temperature for one hour and the resulting solution was evaporated under vacuum to afford the desired product (89 mg, 69% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, δ): 1.46 (virtual triplet, PC(CH<sub>3</sub>)<sub>3</sub>, J<sub>p-H</sub> = 7.2 Hz, 36H), 2.28 (s, free CH<sub>3</sub>CN, 3H), 2.63 (s, Ni-NCCH<sub>3</sub>, 3H), 6.49 (d, J<sub>H-H</sub> = 8.0 Hz, ArH, 2H), 7.04 (t, J<sub>H-H</sub> = 7.6 Hz, ArH, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>, δ): 4.17 (s, Ni-NCCH<sub>3</sub>), 27.89 (s, PC(CH<sub>3</sub>)<sub>3</sub>), 40.04 (t, J<sub>p-C</sub> = 7.9 Hz, PC(CH<sub>3</sub>)<sub>3</sub>), 106.17 (t, J<sub>p-C</sub> = 5.8 Hz, ArC), 122.66 (t, J<sub>p-C</sub> = 18.4 Hz, Ni-NCCH<sub>3</sub>), 131.01 (s, ArC), 136.06 (s, ArC), 169.78 (t, J<sub>p-C</sub> = 8.1 Hz, ArC), resonances arising from free CH<sub>3</sub>CN are also present in the spectrum. <sup>31</sup>P{<sup>1</sup>H} NMR (162 MHz, CDCl<sub>3</sub>, δ): 197.60 (s). <sup>31</sup>P{<sup>1</sup>H} NMR (162 MHz, CD<sub>3</sub>CN, δ): 199.55 (s). IR (ATR, cm<sup>-1</sup>): ν<sub>C=N</sub> = 2290.

**Computational Details**



All quantum chemical calculations based on density functional theory (DFT)<sup>35-37</sup> were performed using the Gaussian09 software package.<sup>38</sup> Intermediates were optimized in the gas phase using the M06 functional<sup>39</sup> to obtain the lowest energy structures along with their electronic energies. A 6-311g\* basis set<sup>40-43</sup> was employed for all the s- and p-block elements and the LANL2DZ basis set<sup>44,45</sup> with effective core potentials for Ni. Subsequent harmonic analyses were performed on all the optimized structures to obtain bond frequencies and thermal corrections to the free energy,<sup>46,47</sup> which were added to the previously calculated electronic energies to obtain the absolute free energies in the gas phase. Solvation energies for the optimized structures were obtained at the same level of theory using the SMD (Solvation Model Density) model<sup>48</sup> for acetonitrile, which is a polarizable continuum solvent model.

The standard reaction free energy for each set of reactants and products was then calculated according to equation 3.



Standard state corrections were made to the free energies to account for the change in going from 1 mol per 24.46 L (gas phase) to 1 M (solution phase), which amounts to  $RT \ln(24.46)$  or 1.89 kcal/mol.

$$\Delta G_{\text{reaction}}^{\circ \text{SMD(MeCN)}} = \Delta G_{\text{products}}^{\circ \text{SMD(MeCN)}} - \Delta G_{\text{reactants}}^{\circ \text{SMD(MeCN)}} \quad (3)$$

To compute the bond dissociation free energy (BDFE) and pKa for **3a** in MeCN, an isodesmic scheme, involving the nickel hydride complexes **3a** and **3a<sup>+</sup>** and a set of reference molecules, was employed as shown in equations 4-10.



$$\text{BDFE}(\text{M-H}) = \Delta G_1^{\circ} + \text{BDFE}(\text{R-H}) \quad (6)$$



$$\text{pKa}(\text{M-H}) = -\log_{10}[\exp(-\Delta G_2^{\circ}/RT)] + \text{pKa}(\text{B-H}) \quad (9)$$

$$\begin{aligned} \Delta \text{pKa} &= \text{pKa}(\text{M}^{\text{III}}\text{-H}) - \text{pKa}(\text{M}^{\text{I}}\text{-H}) \\ &= (\Delta G_2^{\circ}(\text{3a}^{\dagger}) - \Delta G_2^{\circ}(\text{3a})) / (2.303RT) \end{aligned} \quad (10)$$

Reported values of BDFE(R-H) and pKa(B-H) in MeCN solvent for a set of reference molecules were taken from literature;<sup>49,50</sup>  $\Delta G_1^{\circ}$  and  $\Delta G_2^{\circ}$  were computed by our DFT method discussed above, employing the SMD model for MeCN.<sup>48</sup> For pKa calculations, additional benchmarking was performed using other density functionals viz. M06L,<sup>51</sup> B3LYP,<sup>52-54</sup> B3P86,<sup>53,55</sup> BP86<sup>52,55,56</sup> and  $\omega$ B97xD.<sup>57</sup> All electrochemical simulations were performed using DigiElch 7<sup>TM</sup>.<sup>58</sup>

## Acknowledgements

This work was funded by the Center for Electrocatalysis, Transport Phenomena, and Materials (CETM) for Innovative Energy Storage, an Energy Frontier Research Center funded by the U.S. Department of Energy, and the ESD NYSTAR program. S.R would like to thank Andrew Ingram and Kate Waldie for the GC experiment.

## Notes and references

- 1 R. H. Crabtree, *Energy Environ. Sci.*, 2008, **1**, 134.

- 2 C. M. Araujo, D. L. Simone, S. J. Konezny, A. Shim, R. H. Crabtree, G. L. Soloveichik and V. S. Batista, *Energy Environ. Sci.*, 2012, **5**, 9534.
- 3 S. P. Annen, V. Bambagioni, M. Bevilacqua, J. Filippi, A. Marchionni, W. Oberhauser, H. Schönberg, F. Vizza, C. Bianchini and H. Grützmacher, *Angew. Chem. Int. Ed. Engl.*, 2010, **49**, 7229–33.
- 4 P. Driscoll, E. Deunf, L. Rubin, O. Luca, R. Crabtree, C. Chidsey, J. Arnold and J. Kerr, in *ECS Transactions*, The Electrochemical Society, 2011, vol. 35, pp. 3–17.
- 5 P. F. Driscoll, E. Deunf, L. Rubin, J. Arnold and J. B. Kerr, *J. Electrochem. Soc.*, 2013, **160**, G3152–G3158.
- 6 M. Bevilacqua, C. Bianchini, A. Marchionni, J. Filippi, A. Lavacchi, H. Miller, W. Oberhauser, F. Vizza, G. Granozzi, L. Artiglia, S. P. Annen, F. Krumeich and H. Grützmacher, *Energy Environ. Sci.*, 2012, **5**, 8608.
- 7 W. Baratta, G. Chelucci, S. Gladiali, K. Siega, M. Toniutti, M. Zanette, E. Zangrando and P. Rigo, *Angew. Chem. Int. Ed. Engl.*, 2005, **44**, 6214–9.
- 8 A. Fujii, S. Hashiguchi, N. Uematsu, T. Ikariya and R. Noyori, *J. Am. Chem. Soc.*, 1996, **118**, 2521–2522.
- 9 M. Yamakawa, H. Ito and R. Noyori, *J. Am. Chem. Soc.*, 2000, **122**, 1466–1478.
- 10 K. T. Smith, C. Roemming and M. Tilset, *J. Am. Chem. Soc.*, 1993, **115**, 8681–8689.
- 11 O. B. Ryan, M. Tilset and V. D. Parker, *J. Am. Chem. Soc.*, 1990, **112**, 2618–2626.
- 12 O. B. Ryan, K. Smith and M. Tilset, *J. Organomet. Chem.*, 1991, **421**, 315–322.
- 13 J. P. Collman, P. S. Wagenknecht and N. S. Lewis, *J. Am. Chem. Soc.*, 1992, **114**, 5665–5673.
- 14 V. Skagestad and M. Tilset, *J. Am. Chem. Soc.*, 1993, **115**, 5077–5083.
- 15 C. Bianchini, M. Peuzzini, A. Ceccanti, F. Laschi and P. Zanello, *Inorganica Chim. Acta*, 1997, **259**, 61–70.
- 16 T.-Y. Cheng, D. J. Szalda, J. Zhang and R. M. Bullock, *Inorg. Chem.*, 2006, **45**, 4712–4720.
- 17 M. J. Krafft, M. Bubrin, A. Paretzki, F. Lissner, J. Fiedler, S. Zálíš and W. Kaim, *Angew. Chem. Int. Ed. Engl.*, 2013, **52**, 6781–4.
- 18 O. V. Ozerov, C. Guo, L. Fan and B. M. Foxman, *Organometallics*, 2004, **23**, 5573–5580.
- 19 L.-C. Liang, P.-S. Chien and P.-Y. Lee, *Organometallics*, 2008, **27**, 3082–3093.
- 20 R. B. Lansing, K. I. Goldberg and R. A. Kemp, *Dalton Trans.*, 2011, **40**, 8950–8.
- 21 N. Grüger, L.-I. Rodríguez, H. Wadehoff and L. H. Gade, *Inorg. Chem.*, 2013, **52**, 2050–2059.
- 22 C. Rettenmeier, H. Wadehoff and L. H. Gade, *Chem. - A Eur. J.*, 2014, **20**, 9657–9665.

- 23 S. Chakraborty, J. A. Krause and H. Guan, *Organometallics*, 2009, **28**, 582–586.
- 24 I. Göttker-Schnetmann, P. White and M. Brookhart, *J. Am. Chem. Soc.*, 2004, **126**, 1804–11.
- 25 T. Chen, L. Yang, L. Li and K.-W. Huang, *Tetrahedron*, 2012, **68**, 6152–6157.
- 26 V. Pandarus and D. Zargarian, *Organometallics*, 2007, **26**, 4321–4334.
- 27 O. R. Luca, J. D. Blakemore, S. J. Konezny, J. M. Praetorius, T. J. Schmeier, G. B. Hunsinger, V. S. Batista, G. W. Brudvig, N. Hazari and R. H. Crabtree, *Inorg. Chem.*, 2012, **51**, 8704–9.
- 28 A. J. Bard and L. R. Faulkner, *Electrochemical Methods: Fundamentals and Applications*, Wiley, New York, 2nd edn., 2001.
- 29 H. A. Muathen, *Indian. J. Chem.*, 1991, **30B**, 522–524.
- 30 G. R. Fulmer, A. J. M. Miller, N. H. Sherden, H. E. Gottlieb, A. Nudelman, B. M. Stoltz, J. E. Bercaw and K. I. Goldberg, *Organometallics*, 2010, **29**, 2176–2179.
- 31 B. H. Solis and S. Hammes-Schiffer, *Inorg. Chem.*, 2011, **50**, 11252–62.
- 32 S. F. Sousa, P. A. Fernandes and M. J. Ramos, *J. Phys. Chem. A*, 2007, **111**, 10439–10452.
- 33 T. Sperger, I. a. Sanhueza, I. Kalvet and F. Schoenebeck, *Chem. Rev.*, 2015, **115**, 9532–9586.
- 34 K. M. Kadish, J. Q. Ding and T. Malinski, *Anal. Chem.*, 1984, **56**, 1741–1744.
- 35 W. Kohn and L. J. Sham, *Phys. Rev.*, 1965, **140**, A1133–A1138.
- 36 P. Hohenberg, *Phys. Rev.*, 1964, **136**, B864–B871.
- 37 R. G. Parr and W. Yang, *Density Functional Theory of Atoms and Molecules*, Oxford University Press, New York, 1989.
- 38 M. J. Frisch et al., Wallingford CT 2010.
- 39 Y. Zhao and D. G. Truhlar, *Theor. Chem. Acc.*, 2007, **120**, 215–241.
- 40 T. Clark, J. Chandrasekhar, G. W. Spitznagel and P. V. R. Schleyer, *J. Comput. Chem.*, 1983, **4**, 294–301.
- 41 M. J. Frisch, J. A. Pople and J. S. Binkley, *J. Chem. Phys.*, 1984, **80**, 3265.
- 42 R. Krishnan, J. S. Binkley, R. Seeger and J. A. Pople, *J. Chem. Phys.*, 1980, **72**, 650.
- 43 A. V. Mitin, J. Baker and P. Pulay, *J. Chem. Phys.*, 2003, **118**, 7775.
- 44 P. J. Hay and W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 270.
- 45 W. R. Wadt and P. J. Hay, *J. Chem. Phys.*, 1985, **82**, 284.
- 46 R. F. Ribeiro, A. V. Marenich, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. B*, 2011, **115**, 14556–62.
- 47 J. Ho, A. Klamt and M. L. Coote, *J. Phys. Chem. A*, 2010, **114**, 13442–4.
- 48 A. V. Marenich, C. J. Cramer and D. G. Truhlar, *J. Phys. Chem. B*, 2009, **113**, 6378–96.
- 49 J. J. Warren, T. A. Tronic and J. M. Mayer, *Chem. Rev.*, 2010, **110**, 6961–7001.
- 50 R. H. Morris, *J. Am. Chem. Soc.*, 2014, **136**, 1948–59.
- 51 Y. Zhao and D. G. Truhlar, *J. Chem. Phys.*, 2006, **125**, 194101.
- 52 A. D. Becke, *Phys. Rev. A*, 1988, **38**, 3098–3100.
- 53 A. D. Becke, *J. Chem. Phys.*, 1993, **98**, 5648.
- 54 C. Lee, W. Yang and R. G. Parr, *Phys. Rev. B*, 1988, **37**, 785–789.
- 55 J. P. Perdew, *Phys. Rev. B*, 1986, **33**, 8822–8824.
- 56 J. P. Perdew and W. Yue, *Phys. Rev. B*, 1986, **33**, 8800–8802.
- 57 J.-D. Chai and M. Head-Gordon, *J. Chem. Phys.*, 2009, **131**, 174105.
- 58 M. Rudolph, *J. Electroanal. Chem.*, 2003, **543**, 23–39.