

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).


Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemicalscience

Manuscript ID: SC-EDG-08-2015-003259

Table of contents entry



The sydnone-dibenzocyclooctyne and norbornene-tetrazine cycloadditions are both bioorthogonal and mutually orthogonal, used for simultaneous labeling of two targets.

EDGE ARTICLE

Discovery of new mutually orthogonal bioorthogonal cycloaddition pairs through computational screening†

Cite this: DOI: 10.1039/x0xx00000x

Maruthi Kumar Narayanam,^a Yong Liang,^b K. N. Houk^{*,b,c} and Jennifer M. Murphy^{*,a}Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/


Density functional theory (DFT) calculations and experiments in tandem led to discoveries of new reactivities and selectivities involving bioorthogonal sydnone cycloadditions. Dibenzocyclooctyne derivatives (DIBAC and BARAC) were identified to be especially reactive dipolarophiles, which undergo the (3+2) cycloadditions with *N*-phenyl sydnone with the rate constant of up to $1.46 \text{ M}^{-1} \text{ s}^{-1}$. Most significantly, the sydnone-dibenzocyclooctyne and norbornene-tetrazine cycloadditions were predicted to be mutually orthogonal. This was validated experimentally and used for highly selective fluorescence labeling of two proteins simultaneously.

Introduction

The interrogation of the molecular details of biological processes in living systems demands exquisite selectivity combined with high reactivity. Despite many challenges, the design and tuning of reaction partners has provided a myriad of bioorthogonal chemical reactions, which have had a major impact on the study of many biological processes *in vivo*.¹ The discovery of useful bioorthogonal reactions is generally empirical. The modified Staudinger reaction, reported by Bertozzi in 2000,² involves organic azides and triaryl phosphines, but is somewhat slow (typical second-order rate constant of $2 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1}$), necessitating the use of high concentrations of phosphine reagents. A remarkable advance in this field was Bertozzi et al.'s azide-cyclooctyne (3+2) cycloaddition that has enabled cellular component labeling in living systems.³ Stimulated by this distortion-accelerated process, nearly all recent additions to the bioorthogonal reaction repertoire are cycloadditions,⁴ especially 1,3-dipolar⁵ and Diels-Alder⁶ reactions. A great deal of progress has been made towards developing new reactions for labeling biomolecules with extraordinarily high rate constants and monitoring multicomponent processes in complex environments.⁷

First discovered in 1935, sydnones are remarkably stable mesoionic heterocycles that contain an azomethine imine moiety.⁸ In the early 1960s, Huisgen reported that, upon treatment with alkynes at $170 \text{ }^\circ\text{C}$, sydnones undergo (3+2) cycloadditions to form pyrazoles, via a bicyclic intermediate which eliminates carbon dioxide.⁹ Although sydnones have been used for the synthesis of biologically interesting molecules,¹⁰ the potential of sydnones as bioorthogonal reaction partners was overlooked until recently. Using a high-throughput immunoassay screening, Taran and co-workers discovered that

sydnones are biocompatible and undergo 1,3-dipolar cycloadditions with terminal alkynes in the presence of a copper phenanthroline complex to produce 1,4-pyrazoles in up to 99% yield (Scheme 1a).¹¹ Chin developed a Cu-free strain-promoted (3+2) cycloaddition between *N*-phenyl sydnone and bicyclo-[6.1.0]-nonyne (BCN) at ambient temperature to afford the corresponding pyrazole in 30 minutes, with an isolated yield of 99% (Scheme 1b).¹² The rate constant in MeOH-H₂O (55:45) was determined to be $0.054 \text{ M}^{-1} \text{ s}^{-1}$ at $21 \text{ }^\circ\text{C}$.¹³



Scheme 1 Reactivity of the (3+2) cycloadditions involving *N*-phenyl sydnone (Only one regioisomer is depicted).

These results inspired our investigation into the reactivity of sydnone as 1,3-dipoles for bioorthogonal cycloadditions. Since many strained alkenes and alkynes have been well established as bioorthogonal 2π cycloaddends, it is now efficient to evaluate reactivities towards *N*-phenyl sydnone through computational screening. Here we report the computation-guided discovery of two new bioorthogonal (3+2) cycloaddition reactions of *N*-phenyl sydnone with biarylazacyclooctynone (BARAC) and dibenzoazacyclooctyne (DIBAC), which demonstrate significant rate enhancement over the previously reported sydnone-BCN cycloaddition.¹² In addition, with the aid of DFT calculations, we design and identify two mutually orthogonal reaction pairs, which enable highly selective fluorescence labeling of two proteins simultaneously.

Results and discussion

To better understand the sydnone-BCN cycloaddition and to accelerate the discovery of new bioorthogonal chemistry involving *N*-phenyl sydnone, DFT calculations were performed at the M06-2X/6-311+G(d,p)//M06-2X/6-31G(d) level of theory.^{14,15} We have previously established the reliability of prediction of rates in aqueous solution with this method.¹⁶ As shown in Figure 1, the (3+2) cycloaddition of sydnone **1** with BCN **4** via transition state **TS1** requires an activation free energy of 22.2 kcal/mol in water, and the formation of cycloadduct **8** is exergonic by 22.6 kcal/mol. The conversion of intermediate **8** to final pyrazole product **5** by the release of CO₂ via transition state **TS2** has almost no barrier. The overall reaction is strongly exergonic by 111.1 kcal/mol. This indicates that, after the 1,3-dipolar cycloaddition, the subsequent CO₂ release occurs spontaneously, and that the reaction rate is entirely controlled by the (3+2) cycloaddition step.




Fig. 1 Energetics and transition states for the (3+2) cycloaddition of *N*-phenyl sydnone (**1**) with BCN **4** and subsequent CO₂ release.

As shown in Figure 2a, we located transition-state structures **TS3-10** for eight (3+2) cycloadditions of *N*-phenyl sydnone with strained alkenes and alkynes previously used as bioorthogonal reagents. On the basis of the DFT-computed activation free energies (**TS1** and **TS3-10**, Figures 1 and 2a) and the experimental rate constant of the sydnone-BCN cycloaddition ($k_2 = 5 \times 10^{-2} \text{ M}^{-1} \text{ s}^{-1}$, Scheme 1b), we predicted a series of rate constants (k_2) of unexplored bioorthogonal sydnone cycloadditions. It was found that the range of reactivities is enormous, with activation free energies varying from 30 to 19 kcal/mol, which correspond to a 10^8 range in rate constants at room temperature. For cyclooctyne, 1,3-disubstituted cyclopropene (Cp(1,3)), difluorocyclooctyne (DIFO), and *trans*-cyclooctene (TCO) (**TS5-8**), the predicted rate constants range from 10^{-3} to $10^{-1} \text{ M}^{-1} \text{ s}^{-1}$ (shown in yellow, Figure 2a). These data are very close to the reported rate constant for the sydnone-BCN cycloaddition. This suggests that these reactions may be employed for biomolecular labeling but may not be suitable for *in vivo* applications due to the moderate rate constants. To our delight, calculations predicted that when




Fig. 2 (a) DFT-computed activation free energies for the (3+2) cycloadditions of sydnone **1** with eight strained alkenes and alkynes at the CPCM(water)-M06-2X/6-311+G(d,p)//M06-2X/6-31G(d) level of theory and the predicted rate constants in water at 25 °C. (b) DFT-computed activation free energies and predicted rate constants for the (4+2) cycloadditions of tetrazine **9** with norbornene, DIBAC, and BARAC.

N-phenyl sydnone reacts with DIBAC and BARAC (**TS9-10**), the rate constants will be 1 and $10 \text{ M}^{-1} \text{ s}^{-1}$, respectively (shown in green, Figure 2a). By contrast, sydnone **1** is predicted to be inert to norbornene and 3,3-disubstituted cyclopropene (Cp(3,3)) (**TS3-4**) under physiological conditions, according to the expected rate constants of $10^{-7} \text{ M}^{-1} \text{ s}^{-1}$ (shown in red, Figure 2a). This dramatic reactivity difference between dibenzocyclooctyne derivatives (DIBAC and BARAC) and norbornene in the sydnone cycloaddition was then exploited for design of mutually orthogonal bioorthogonal reaction pairs.


Figure 2b shows that diaryltetrazine **9** reacts smoothly with norbornene (**TS11**). However, the Diels-Alder reactions of DIBAC and BARAC (**TS12-13**) with diaryltetrazine **9** will not occur at room temperature, according to the predicted rate constants of $2 \times 10^{-6} \text{ M}^{-1} \text{ s}^{-1}$ (shown in red, Figure 2b). As a result, the sydnone-dibenzocyclooctyne and norbornene-tetrazine cycloadditions are predicted to be mutually orthogonal.

Our previous computational studies showed that the distortion energy of substrates is crucial to the reactivity of bioorthogonal cycloadditions.^{16,17} Usually, the more pre-distorted substrate is more reactive as it takes less energy to reach its transition-state geometry. Dibenzocyclooctyne derivatives (DIBAC and BARAC) have much smaller alkyne bond angles than normal alkynes (ca. 153° versus 180°),^{17a} while norbornene is just pre-distorted by ca. 8° as compared to normal alkenes.^{17b} Therefore, in sydnone cycloadditions, dibenzocyclooctyne derivatives (DIBAC and BARAC) are significantly more reactive than norbornene. However, DIBAC and BARAC exhibit steric hindrance due to the two aryl hydrogen atoms ortho to the alkyne,¹⁶ so their Diels-Alder reactions with bulky disubstituted tetrazines are dramatically slower than that of norbornene.

To test these predictions, we first measured the experimental rate constant for the 1,3-dipolar cycloaddition of *N*-phenyl sydnone (**1**) with BARAC **6** (for details, see the ESI†). As expected, a significantly enhanced second-order rate constant of $1.46 \text{ M}^{-1} \text{ s}^{-1}$ was obtained for the (3+2) cycloaddition in MeCN-

H_2O (1:1) at 23°C (Scheme 1c). This is about 30 times larger than that for the sydnone-BCN cycloaddition.¹²

Scheme 2 summarizes experiments to test the predicted mutual orthogonality between sydnone-DIBAC and norbornene-tetrazine cycloadditions. The second-order rate constant for the (3+2) cycloaddition of DIBAC **10** with sydnone **1** in MeOH- H_2O (55:45) was determined to be $0.902 \text{ M}^{-1} \text{ s}^{-1}$ (for details, see the ESI†). The Diels-Alder reaction of diaryltetrazine **13** with 5-norbornene-2-acetic acid (**12**) was found to have a k_2 of $1.05 \text{ M}^{-1} \text{ s}^{-1}$ (for details, see the ESI†) in MeOH- H_2O (55:45), which is comparable to rate constants reported in the literature for similar systems.¹⁸ To test potential cross-reactivity, sydnone **1** and norbornene **12** or diaryltetrazine **13** and DIBAC **10** were incubated in a deuterated methanol and monitored via NMR spectroscopy. As predicted, no reaction was observed between either reaction pair at room temperature after 24 hours.



Scheme 2 Mutual orthogonality between sydnone-DIBAC and norbornene-tetrazine cycloadditions ($\text{R}=\text{CH}_2\text{CH}_2\text{NH}_2$, $\text{R}'=\text{NH}(\text{CH}_2)_3\text{NHBoc}$; only one regioisomer is depicted).




Fig. 3 Modifications of protein surfaces via bioorthogonal cycloadditions. (a) Dibenzocyclooctyne (DIBAC) and 5-norbornene-2-acetic acid (Nor) were appended to BSA and OVA (20 mg/mL in PBS) respectively, via NHS ester-amine coupling conditions (20 mM labeling reagent). The labeled proteins **BSA-DIBAC** and **OVA-Nor** (2 mg/mL) simultaneously react with sydnone-BODIPY (**Syd-630**) and tetrazine-BODIPY (**Tz-504**). (b) Gel analysis of BSA functionalized with DIBAC and OVA functionalized with Nor incubated for 1 h with either **Syd-630**, **Tz-504**, both reagents simultaneously, or no reagent (-). (c) Gel analysis of DIBAC-modified BSA and Nor-modified OVA with both **Syd-630** and **Tz-504** simultaneously for 1-60 min or no reagent (-). The protein loading on the gels was assessed with Coomassie stain.

Probing multiple biomolecules, within their native cellular habitat, requires the selective modification of each system with chemical moieties that are compatible with each other. A remarkable expansion of the bioorthogonal toolbox has equipped chemical biologists with a range of selective reactions.^{4,7} However, the highly complex nature of biological processes often demands concurrent modification and tracking of multiple targets within the same system. Clever manipulation of these various reactions has enabled the use of sequential click chemistry reactions to tag multiple targets in a single system without perturbations to the system.¹⁹ Hilderbrand and others have demonstrated the orthogonality between the Diels-Alder tetrazine ligation and the 1,3-dipolar azide cycloaddition to facilitate multicomponent labeling.²⁰

Leveraging the excellent orthogonality between sydnone-DIBAC and norbornene-tetrazine cycloadditions, we examined the application of this chemistry towards dual fluorescence protein labeling. The DIBAC and norbornene moieties were appended to the surface of two ubiquitous proteins, bovine serum albumin (**BSA**) and ovalbumin (**OVA**) using standard coupling conditions (Figure 3a). The modified proteins (**BSA-DIBAC** and **OVA-Nor**) were treated with either red fluorescent sydnone-BODIPY630 (**Syd-630**) or green fluorescent tetrazine-BODIPY504 (**Tz-504**) or both and analyzed via in gel fluorescence imaging. When the modified proteins were jointly treated with **Syd-630**, only **BSA-DIBAC** was fluorescently labeled (lane 7, Figure 3b). Likewise, in the presence of **Tz-504**, **OVA-Nor** showed selective labeling without any detection of **BSA-DIBAC** fluorescence labeling (lane 8, Figure 3b). Co-administration of both fluorophore conjugates in “one pot” resulted in the concurrent labeling of both proteins (lane 9, Figure 3b). Unmodified proteins **BSA** and **OVA** showed no fluorescence labeling after exposure to **Syd-630** and **Tz-504** (lanes 4 and 5, Figure 3b), indicating the absence of nonspecific reactivity. In a time-dependent experiment, the protein **BSA-DIBAC** underwent labeling and exhibited fluorescence within 1 min (lane 7, Figure 3c), whereas the protein **OVA-Nor** exhibited fluorescence labeling after 15 min (lane 8, Figure 3c). These experiments demonstrate the exquisite specificity of the sydnone and tetrazine imaging agents for DIBAC- and norbornene-modified proteins, respectively, as well as their mutual compatibility within a single environment. Moreover, for the new sydnone-DIBAC cycloaddition, the ability to undergo fluorescence labeling within 1 min using as little as 12 nM protein (with 500 μ M dye) is very promising for *in vivo* imaging.

Conclusions

With the aid of computational screening, we have discovered two rapid cycloadditions between *N*-phenyl sydnone and dibenzocyclooctyne derivatives (DIBAC and BARAC) that proceed in aqueous media, at physiological temperature without the use of a catalyst. The predicted mutually orthogonal sydnone-DIBAC and norbornene-tetrazine cycloaddition pairs

have been successfully applied to fluorescence labeling of two proteins simultaneously. DFT calculations shown in Figure 2a also predicted that sydnone is inert to less distorted 3,3-disubstituted cyclopropene. Our previous study has proven that 3,3-disubstituted cyclopropenes react with nitrile imines via 1,3-dipolar cycloaddition but not with tetrazines via Diels-Alder reaction.^{19d} This suggests that precise control over cycloaddition selectivity (dibenzocyclooctynes with sydnones, norbornenes with tetrazines, and 3,3-disubstituted cyclopropenes with nitrile imines) can enable labeling of three targets in a biological environment. We anticipate that these results will complement the expanding toolkit of bioorthogonal reactions and advance the efforts to simultaneously examine multiple processes within a biological system.

Acknowledgements

This work was supported by the UCLA Center for In Vivo Imaging in Cancer Biology (NIH P50 CA086306), the National Institutes of Health (R01GM109078), and the National Science Foundation (CHE-1361104). Calculations were performed on the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by the NSF (OCI-1053575). We thank Prof. James Wohlschlegel and Dr. Ajay Vashisht for helpful discussions and assistance with protein labeling experiments. We thank Prof. Jennifer Prescher at UCI for helpful comments.

Notes and references

^aCrump Institute for Molecular Imaging, David Geffen School of Medicine, University of California, Los Angeles, California 90095, USA. E-mail: jmmurphy@mednet.ucla.edu

^bDepartment of Chemistry and Biochemistry, University of California, Los Angeles, California 90095, USA. E-mail: houk@chem.ucla.edu

^cDepartment of Chemical and Biomolecular Engineering, University of California, Los Angeles, California 90095, USA.

[†]Electronic Supplementary Information (ESI) available: experimental and computational details. See DOI: 10.1039/b000000x/

1 (a) E. M. Sletten and C. R. Bertozzi, *Angew. Chem., Int. Ed.*, 2009, **48**, 6974-6998; (b) M. Grammel and H. C. Hang, *Nat. Chem. Biol.*, 2013, **9**, 475-484; (c) C. S. McKay and M. Finn, *Chem. Biol.*, 2014, **21**, 1075-1101; (d) A. Borrmann and J. C. van Hest, *Chem. Sci.*, 2014, **5**, 2123-2134.

2 E. Saxon and C. R. Bertozzi, *Science*, 2000, **287**, 2007-2010.

3 (a) N. J. Agard, J. A. Prescher and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2004, **126**, 15046-15047; (b) E. M. Sletten and C. R. Bertozzi, *Acc. Chem. Res.*, 2011, **44**, 666-676.

4 (a) J. C. Jewett and C. R. Bertozzi, *Chem. Soc. Rev.*, 2010, **39**, 1272-1279; (b) D. M. Patterson, L. A. Nazarova and J. A. Prescher, *ACS Chem. Biol.*, 2014, **9**, 592-605.

5 (a) J. M. Baskin, J. A. Prescher, S. T. Laughlin, N. J. Agard, P. V. Chang, I. A. Miller, A. Lo, J. A. Codelli and C. R. Bertozzi, *Proc. Natl. Acad. Sci. U.S.A.*, 2007, **104**, 16793-16797; (b) J. A. Codelli, J. M. Baskin, N. J. Agard and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2008, **130**, 11486-11493; (c) X. Ning, J. Guo, M. A. Wolfert and G. J. Boons, *Angew.*

- Chem., Int. Ed.*, 2008, **47**, 2253-2255; (d) M. F. Debets, S. S. van Berkel, S. Schoffelen, F. P. J. T. Rutjes, J. C. M. van Hest and F. L. van Delft, *Chem. Commun.*, 2010, **46**, 97-99; (e) J. C. Jewett, E. M. Sletten and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2010, **132**, 3688-3690; (f) J. Dommerholt, S. Schmidt, R. Temming, L. J. Hendriks, F. P. Rutjes, J. van Hest, D. J. Lefeber, P. Friedl and F. L. van Delft, *Angew. Chem., Int. Ed.*, 2010, **49**, 9422-9425; (g) T. Plass, S. Milles, C. Koehler, C. Schultz and E. A. Lemke, *Angew. Chem., Int. Ed.*, 2011, **50**, 3878-3881; (h) M. F. Debets, S. S. Van Berkel, J. Dommerholt, A. J. Dirks, F. P. Rutjes and F. L. Van Delft, *Acc. Chem. Res.*, 2011, **44**, 805-815; (i) R. K. Lim and Q. Lin, *Acc. Chem. Res.*, 2011, **44**, 828-839; (j) Z. Yu, Y. Pan, Z. Wang, J. Wang and Q. Lin, *Angew. Chem., Int. Ed.*, 2012, **51**, 10600-10604; (k) Z. Yu, T. Y. Ohulchanskyy, P. An, P. N. Prasad and Q. Lin, *J. Am. Chem. Soc.*, 2013, **135**, 16766-16769; (l) Z. Yu and Q. Lin, *J. Am. Chem. Soc.*, 2014, **136**, 4153-4156.
- 6 (a) M. L. Blackman, M. Royzen and J. M. Fox, *J. Am. Chem. Soc.*, 2008, **130**, 13518-13519; (b) N. K. Devaraj, R. Weissleder and S. A. Hilderbrand, *Bioconjugate Chem.*, 2008, **19**, 2297-2299; (c) N. K. Devaraj and R. Weissleder, *Acc. Chem. Res.*, 2011, **44**, 816-827; (d) D. S. Liu, A. Tangpeerachaiikul, R. Selvaraj, M. T. Taylor, J. M. Fox and A. Y. Ting, *J. Am. Chem. Soc.*, 2012, **134**, 792-795; (e) J. L. Seitchik, J. C. Peeler, M. T. Taylor, M. L. Blackman, T. W. Rhoads, R. B. Cooley, C. Refakis, J. M. Fox and R. A. Mehl, *J. Am. Chem. Soc.*, 2012, **134**, 2898-2901; (f) K. Lang, L. Davis, S. Wallace, M. Mahesh, D. J. Cox, M. L. Blackman, J. M. Fox and J. W. Chin, *J. Am. Chem. Soc.*, 2012, **134**, 10317-10320; (g) K. Lang, L. Davis, J. Torres-Kolbus, C. Chou, A. Deiters and J. W. Chin, *Nature Chem.*, 2012, **4**, 298-304; (h) W. Chen, D. Wang, C. Dai, D. Hamelberg and B. Wang, *Chem. Commun.*, 2012, **48**, 1736-1738; (i) J. Yang, J. Šečkutě, C. M. Cole and N. K. Devaraj, *Angew. Chem., Int. Ed.*, 2012, **51**, 7476-7479; (j) J. Šečkutě, J. Yang and N. K. Devaraj, *Nucleic Acids Res.*, 2013, **41**, e148; (k) Q. Li, T. Dong, X. Liu and X. Lei, *J. Am. Chem. Soc.*, 2013, **135**, 4996-4999; (l) J. Yang, Y. Liang, J. Šečkutě, K. N. Houk and N. K. Devaraj, *Chem Eur J*, 2014, **20**, 3365-3375; (m) H. Wu, J. Yang, J. Šečkutě and N. K. Devaraj, *Angew. Chem., Int. Ed.*, 2014, **53**, 5805-5809; (n) H. Wu, B. T. Cisneros, C. M. Cole and N. K. Devaraj, *J. Am. Chem. Soc.*, 2014, **136**, 17942-17945; (o) A. Darko, S. Wallace, O. Dmitrenko, M. M. Machovina, R. A. Mehl, J. W. Chin and J. M. Fox, *Chem. Sci.*, 2014, **5**, 3770-3776; (p) S. B. Engelsma, L. I. Willems, C. E. van Paaschen, S. I. van Kasteren, G. A. van der Marel, H. S. Overkleeft and D. V. Filippov, *Org. Lett.*, 2014, **16**, 2744-2747; (q) D. N. Kamber, Y. Liang, R. J. Blizzard, F. Liu, R. A. Mehl, K. N. Houk and J. A. Prescher, *J. Am. Chem. Soc.*, 2015, **137**, 8388-8391.
- 7 (a) M. F. Debets, J. C. van Hest and F. P. Rutjes, *Org. Biomol. Chem.*, 2013, **11**, 6439-6455; (b) K. Lang and J. W. Chin, *ACS Chem. Biol.*, 2014, **9**, 16-20; (c) H.-W. Shih, D. N. Kamber and J. A. Prescher, *Curr. Opin. Chem. Biol.*, 2014, **21**, 103-111.
- 8 J. C. Earl and A. W. Mackney, *J. Chem. Soc.*, 1935, 899-900.
- 9 R. Huisgen, R. Grashey, H. Gotthardt and R. Schmidt, *Angew. Chem., Int. Ed. Engl.*, 1962, **1**, 48-49.
- 10 D. L. Browne and J. P. Harrity, *Tetrahedron*, 2010, **66**, 553-568.
- 11 S. Kolodych, E. Rasolofonjatovo, M. Chaumontet, M. C. Nevers, C. Créminon and F. Taran, *Angew. Chem., Int. Ed.*, 2013, **52**, 12056-12060.
- 12 S. Wallace and J. W. Chin, *Chem. Sci.*, 2014, **5**, 1742-1744.
- 13 Shortly after Chin's report (ref 12), 4-chloro-sydnones were reported to react much faster with BCN than non-chlorinated analogues: L. Plougastel, O. Koniev, S. Specklin, E. Decuypere, C. Créminon, D.-A. Buisson, A. Wagner, S. Kolodych and F. Taran, *Chem. Commun.*, 2014, **50**, 9376-9378. For example, the rate constant for the (3+2) cycloaddition between *N*-phenyl 4-chloro-sydnone and BCN in PBS/DMSO was reported to be 0.872 M⁻¹ s⁻¹ at 25 °C.
- 14 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, *Gaussian 09*, Revision D.01, Gaussian Inc., Wallingford CT, 2013.
- 15 (a) Y. Zhao and D. G. Truhlar, *Theor. Chem. Acc.*, 2008, **120**, 215-241; (b) Y. Zhao and D. G. Truhlar, *Acc. Chem. Res.*, 2008, **41**, 157-167.
- 16 Y. Liang, J. L. Mackey, S. A. Lopez, F. Liu and K. N. Houk, *J. Am. Chem. Soc.*, 2012, **134**, 17904-17907.
- 17 (a) C. G. Gordon, J. L. Mackey, J. C. Jewett, E. M. Sletten, K. N. Houk and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2012, **134**, 9199-9208; (b) S. A. Lopez and K. N. Houk, *J. Org. Chem.*, 2013, **78**, 1778-1783; (c) F. Liu, Y. Liang and K. N. Houk, *J. Am. Chem. Soc.*, 2014, **136**, 11483-11493.
- 18 M. R. Karver, R. Weissleder and S. A. Hilderbrand, *Bioconjugate Chem.*, 2011, **22**, 2263-2270.
- 19 (a) B. C. Sanders, F. Friscourt, P. A. Ledin, N. E. Mbuja, S. Arumugam, J. Guo, T. J. Boltje, V. V. Popik and G.-J. Boons, *J. Am. Chem. Soc.*, 2010, **133**, 949-957; (b) H. Stöckmann, A. A. Neves, H. A. Day, S. Stairs, K. M. Brindle and F. J. Leeper, *Chem. Commun.*, 2011, **47**, 7203-7205; (c) T. Plass, S. Milles, C. Koehler, J. Szymański, R. Mueller, M. Wießler, C. Schultz and E. A. Lemke, *Angew. Chem., Int. Ed.*, 2012, **51**, 4166-4170; (d) D. N. Kamber, L. A. Nazarova, Y. Liang, S. A. Lopez, D. M. Patterson, H.-W. Shih, K. N. Houk and J. A. Prescher, *J. Am. Chem. Soc.*, 2013, **135**, 13680-13683.
- 20 (a) M. R. Karver, R. Weissleder and S. A. Hilderbrand, *Angew. Chem., Int. Ed.*, 2012, **51**, 920-922; (b) L. I. Willems, N. Li, B. I. Florea, M. Ruben, G. A. van der Marel and H. S. Overkleeft, *Angew. Chem., Int. Ed.*, 2012, **51**, 4431-4434; (c) D. M. Patterson, L. A. Nazarova, B. Xie, D. N. Kamber and J. A. Prescher, *J. Am. Chem. Soc.*, 2012, **134**, 18638-18643; (d) C. M. Cole, J. Yang, J. Šečkutě and N. K. Devaraj, *ChemBioChem*, 2013, **14**, 205-208; (e) D. M. Patterson, K. A. Jones and J. A. Prescher, *Mol. BioSyst.*, 2014, **10**, 1693-1697; (f) I. Nikić, T. Plass, O. Schraidt, J. Szymański, J. A. G. Briggs, C. Schultz and E. A. Lemke, *Angew. Chem., Int. Ed.*, 2014, **53**, 2245-2249; (g) A. Sachdeva, K. Wang, T. Elliott and J. W. Chin, *J. Am. Chem. Soc.*, 2014, **136**, 7785-7788.