



**Nickel–Catalyzed Arylation of Heteroaryl-containing
Diarylmethanes: Exceptional Reactivity of the
Ni(NIXANTPHOS)-based Catalyst**

Journal:	<i>Chemical Science</i>
Manuscript ID	SC-EDG-09-2015-003704
Article Type:	Edge Article
Date Submitted by the Author:	30-Sep-2015
Complete List of Authors:	Walsh, Patrick; UPenn, chem Cao, Xinyu; UPenn, Chem Sha, Sheng-Chun; UPenn, Chem Li, Minyan; University of Pennsylvania, Department of Chemistry Kim, Byeong-Seon; University of Pennsylvania, Department of Chemistry Morgan, Catherine; UPenn, Chem Huang, Rudan; Beijing Institute of Technology, Chem



Journal Name

ARTICLE

Nickel-Catalyzed Arylation of Heteroaryl-containing Diarylmethanes: Exceptional Reactivity of the Ni(NIXANTPHOS)-based Catalyst


Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Xinyu Cao^{a, b}, Sheng-Chun Sha^b, Minyan Li^b, Byeong-Seon Kim^b, Catherine Morgan^b, Rudan Huang^a, Xiaodong Yang^c, and Patrick J. Walsh^{*, b}

Nickel(0)-catalyzed cross-coupling of heteroaryl-containing diarylmethanes with both aryl bromides and chlorides has been achieved. The success of this reaction relies on the introduction of a unique nickel/NIXANTPHOS-based catalyst system, which provides a direct route to triarylmethanes from heteroaryl-containing diarylmethanes. Reactivity studies indicate the Ni(NIXANTPHOS)-based catalyst exhibits enhanced reactivity over XANTPHOS derivatives and other Ni(phosphine)-based catalysts in the reactions examined.



1 Introduction


Triarylmethanes have attracted significant attention due to their diverse applications as leuco dye precursors,¹ fluorescent probes,² and photochromic agents.³ They also find uses in medicinal chemistry.⁴ Given the utility of triarylmethanes, it is not surprising that considerable effort has been invested in their synthesis.^{5,6} Recently, triarylmethanes have been prepared using transition metal catalyzed cross-coupling reactions, usually employing palladium-based catalysts.⁷ To decrease costs and increase sustainability of these processes, much effort has focused on employing earth abundant first row transition metals in reactions traditionally catalyzed by palladium.^{8–11} Given nickel's position directly above palladium, and its high reactivity with aryl halides,¹² nickel stands out as a clear starting point for the development of first-row transition metal catalyzed processes. Indeed, it has been used successfully in many cross-coupling reactions.^{12–14}

The synthesis of triarylmethanes via nickel-catalyzed reactions has been achieved through traditional Kumada and Suzuki–Miyaura coupling approaches.^{8–11} In 2012 and 2013, the Jarvo group demonstrated the synthesis of triarylmethanes


with high ee by coupling of enantioenriched benzylic ethers with aryl Grignard reagents^{8a} (Scheme 1A) and arylboronic esters (Scheme 1B).⁹ Related strategies were reported by the

Previous work


A. Jarvo (2012)




B. Jarvo (2013)




C. Watson (2013)



D. Yang (2015)




E. Jarvo (2015)



This work

F. C-H Functionalization



Scheme 1 Synthetic Approaches to Triarylmethanes.

Watson¹⁰ and Yang¹¹ groups with enantioenriched benzylic pivalates and aryl boronic acid derivatives. More recently, the

^aKey Laboratory of Cluster Science of Ministry of Education, School of Chemistry, Beijing Institute of Technology, Beijing 100081, PR China

^bDepartment of Chemistry, University of Pennsylvania, 231 S. 34th St., Philadelphia, PA 19104, USA.

^cKey Laboratory of Medicinal Chemistry for Natural Resource, School of Chemical Science and Technology, Yunnan University, Kunming, 650091, PR China

E-mail: pwalsh@sas.upenn.edu;


Web: <https://sites.google.com/site/titaniumupenn/>; Fax: +1 215-573-6743

† Electronic supplementary information (ESI) available: Data for new compounds, experimental procedures, 1H and 13C NMR spectra See DOI: 10.1039/x0xx00000x

Jarvo group reported Suzuki reaction as a complementary method to the previously reported nickel-catalyzed Kumada coupling.^{8b}

Our approach to triarylmethanes differs from these important contributions in that it involves the catalytic functionalization of weakly acidic sp³-hybridized C–H bonds via a deprotonative–cross–coupling process (DCCP).^{15–16} In our prior studies, diarylmethanes were reversibly deprotonated in the presence of a Pd(NIXANTPHOS) catalyst and the resulting carbanions coupled with aryl bromides^{17a,c} and chlorides^{17b} in high yields. It is noteworthy that early work on the deprotonation/cross coupling of both aromatic C(sp²)–H's and C(sp³)–H's of fluorinated hydrocarbons were reported by Daugulis employing base metal catalysts (Cu and Fe).^{17d–g}

One of the most significant findings of our early investigations is that under the basic reaction conditions, van Leeuwen's wide bite-angle ligand,¹⁸ NIXANTPHOS' N–H (pK_a ~22)¹⁹ is deprotonated under the reaction conditions and the resulting heterobimetallic catalyst displays exceptional reactivity when compared with other bidentate phosphine-based palladium catalysts.¹⁷ The most striking comparison is between the NIXANTPHOS- and XANTPHOS-based palladium catalysts. Despite the outward similarity of these ligand scaffolds, the NIXANTPHOS-based catalyst exhibited over 90% assay yield (AY, determined by ¹H NMR) in the coupling of 1-chloro-4-*tert*-butylbenzene with diphenylmethane to afford the triarylmethane product, while the parent XANTPHOS-based catalyst, under identical conditions, exhibited only 1% conversion (Equation 1). Furthermore, other ligands known to participate in cross-coupling reactions with aryl chlorides, including the Buchwald family of ligands, were examined in this reaction and found to give less than 2% product. These results attest to the difficulty of this reaction, particularly with aryl chlorides.



Ligand	HPLC AY (%)
NIXANTPHOS	91
N-Bn-NIXANTPHOS	1
XANTPHOS	0
DPEPhos	0
dipp	0
XPhos	1
SPhos	1
PCy ₃	0

Z = N-H, NIXANTPHOS
 Z = N-CH₂Ph, N-Bn-NIXANTPHOS
 Z = C(CH₃)₂, XANTPHOS
 Z = H, H, DPEPhos

In the present study, our goals were to 1) develop a nickel catalyst to promote the DCCP of heteroaryl-containing diarylmethane derivatives and 2) determine if the high reactivity imparted to palladium when chelated by a deprotonated NIXANTPHOS ligand translates to nickel catalysts. Herein, we describe the synthesis of triarylmethanes

by DCCP of heteroaryl-containing diarylmethanes and aryl bromides and chlorides catalyzed by a Ni(NIXANTPHOS)-based system. To determine if the exceptional reactivity of the Ni(NIXANTPHOS)-based catalyst translates to additional reactions, we also report preliminary results on the previously unknown arylation of a 2-pyridylmethyl amine. Compared with 37 mono- and bidentate phosphine ligands, the Ni(NIXANTPHOS)-based catalyst outperformed the other nickel catalysts in the arylation of 2-pyridylmethyl amine under the conditions examined.

2 Results and Discussion


2.1 Catalyst Identification

We evaluated nickel precursors in the presence of phosphine ligands by screening 38 electronically diverse mono- and bidentate phosphines using Ni(COD)₂ as the nickel source. Coupling of 2-benzylpyridine **1a** (1.2 equiv) with 1-bromo-4-*tert*-butylbenzene **2a** (1 equiv) in the presence of NaN(SiMe₃)₂ (3 equiv), catalytic Ni(COD)₂ (10 mol %) and ligand (20 mol % for monodentate phosphine ligands and 10 mol % for bidentate ligands) were performed in cyclopentyl methyl ether (CPME) on microscale (0.01 mmol, 0.1 M) at 110 °C. After heating for 16 h, the reactions were worked up and analyzed by HPLC (retention times, UV/vis characteristics and integration against an internal standard, see Supporting Information for details). Of the 38 different phosphine ligands tested, NIXANTPHOS was the *only promising hit*. Interestingly, as discussed below, the XANTPHOS-based catalyst was much less active in this transformation. Translation of the microscale results with commercially available NIXANTPHOS to laboratory scale (0.1 mmol) under the same reaction conditions resulted in 78% assay yield (AY, as determined by ¹H NMR in all cases) (Table 1, entry 1). Examination of different nickel to ligand ratios indicated that a 1:1 ratio at 10 mol % loading provided the best yields (entries 2–3). Decreasing the equivalents of NaN(SiMe₃)₂ from 3 to 2 resulted in an increase in the yield of coupling product **3aa** to 83% (entry 4). Reducing the concentration to 0.05 M resulted in generation of product **3aa** in 85% (entry 5).

Optimization was continued by adjusting other reaction parameters. Changing the ratio of 2-benzyl pyridine to aryl bromide and base to 1.0 : 1.5 : 2.0 rendered product **3aa** in 93% AY (entry 6). Changing the reaction times from 16 to 8 or 24 h led to lower yields (entries 7–8). Lowering the temperature from 110 °C to 50 °C and room temperature increased the AY to 99% at both the lower temperatures (entries 9–10). Decreasing the catalyst loading to 2.5 mol % at room temperature also resulted in 99% AY (entry 11). Further decreasing the catalyst loading to 1 mol % at RT led to 72% AY (entry 12). Under the optimized conditions of entry 11, the triarylmethane product was isolated in 99% yield. Substitution of XANTPHOS for NIXANTPHOS at twice the catalyst loading (5 mol %), under otherwise identical conditions afforded only

30% AY. This result prompted a series of experiments probing the difference in reactivity between these two structurally similar ligands (see later sections).

Table 1 Optimization of Ni–NIXANTPHOS Catalyzed DCCP of **1a** with **2a**^a



Entry	1a : 2a : base	Ni/L (mol %)	T (°C)	t (h)	Concn.	Yield ^b (%)
1	1.2 : 1 : 3	10/10	110	16	0.1	78
2	1.2 : 1 : 3	10/15	110	16	0.1	71
3	1.2 : 1 : 3	10/20	110	16	0.1	62
4	1.2 : 1 : 2	10/10	110	16	0.1	83
5	1.2 : 1 : 2	10/10	110	16	0.05	85
6	1 : 1.5 : 2	10/10	110	16	0.05	93
7	1 : 1.5 : 2	10/10	110	8	0.05	77
8	1 : 1.5 : 2	10/10	110	24	0.05	81
9	1 : 1.5 : 2	10/10	50	16	0.05	99
10	1 : 1.5 : 2	10/10	rt	16	0.05	99
11 ^c	1 : 1.5 : 2	2.5/2.5	rt	16	0.05	99(99)
12	1 : 1.5 : 2	1/1	rt	16	0.05	72

^a Reactions conducted on a 0.10 mmol scale. ^b Yield determined by ¹H NMR spectroscopy of the crude reaction mixtures. ^c Isolated yield after chromatographic purification.


2.2 Scope of the Diphenylmethane Derivatives

Under the optimized conditions of entry 12 in Table 1, the scope of heteroaryl-containing diarylmethanes with 4-*tert*-butyl bromobenzene (**2a**) was examined (Scheme 2). The isomeric 2-, 3- and 4-benzylpyridine substrates have pK_a values of 28.2, 30.1, and 26.7 in DMSO, respectively. It is not entirely surprising that their reactions required different conditions to achieve high yields (Scheme 2, see Supporting Information for details). Nonetheless, coupling of 2-, 3- and 4-benzylpyridine furnished products **3aa**, **3ba** and **3ca** in 99, 67 and 96% isolated yields, respectively. 3-Benzylpyridine was the most challenging of the benzyl pyridines, because it is the most difficult to deprotonate. Nonetheless, the more acidic dipyridin-3-ylmethane (**1d**) underwent coupling with aryl bromide **2a** and 2 equiv of LiN(SiMe₃)₂ in 75% isolated yield. Unlike analogous palladium-catalyzed reactions with less acidic diarylmethanes,⁷ such as diphenylmethane (pK_a = 32 in DMSO)²⁰ the Ni(NIXANTPHOS)-based catalyst failed to give product with substrates with benzylic C–H pK_a's values greater than about 30. This result is not totally surprising. Based on the assumption that the reaction pathways are similar,²¹ we hypothesize that the decreased electronegativity of nickel (1.91) relative to palladium (2.20, Pauling scale) renders the Ni catalyst less electrophilic.²² This, combined with the smaller size of Ni compared to Pd²³ and the small value of the equilibrium for deprotonation (pK_a of HN(SiMe₃)₂ in THF =

26),²⁴ likely results in very little or no transmetalation with the anion derived from diphenylmethane.

In addition to benzylpyridines, other diphenylmethane derivatives underwent nickel-catalyzed coupling reactions. These include fluorene (**1e**) and xanthene (**1f**, pK_a in DMSO = 30.0). The triarylmethane derivatives **3ea** and **3fa** were isolated in 99 and 81% yield, respectively. Notably, reactions with the more acidic fluorene required only 1 mol % catalyst loading. Related triarylmethanes with these core structures have been found to have various applications.^{25–26}

Scheme 2 Scope of Heteroaryl-containing Diarylmethanes in Ni–NIXANTPHOS Catalyzed DCCP^{a,b}




^a Reactions conducted on a 0.1 mmol scale at 0.05 M. ^b Isolated yields after chromatographic purification. ^c 2.5 mol % Ni(COD)₂/2.5 mol % NIXANTPHOS. ^d 10 mol % Ni(COD)₂/10 mol % NIXANTPHOS. ^e 1 mol % Ni(COD)₂/1 mol % NIXANTPHOS. ^f 2 equiv of NaN(SiMe₃)₂. ^g 4 equiv of NaN(SiMe₃)₂. ^h 2.5 equiv of LiN(SiMe₃)₂. ⁱ 2 equiv of LiN(SiMe₃)₂. ^j 1.5 equiv of NaN(SiMe₃)₂. ^k room temperature. ^l 50 °C. ^m 110 °C.

2.3 Scope of the Aryl Halide

To probe the ability of the Ni(NIXANTPHOS)-based catalyst to promote reactions with different electrophiles, 2-benzylpyridine was coupled with various aryl bromides (Scheme 3). Bromobenzene (**2b**) gave coupling product **3ab** in 93% yield with 5 mol % catalyst loading at room temperature. Under the same conditions, electron rich 4-bromoanisole (**2c**) and 4-bromo-*N,N*-dimethylaniline (**2d**) furnished products **3ac** and **3ad** in 91 and 83% yields, respectively. Aryl bromide with a 4-fluoro- group afforded triarylmethane **3ae** in 83% isolated yield, (5 mol % catalyst loading at 50 °C). Keto-containing 4-bromobenzophenone delivered **3af** in 74% yield with 10 mol % catalyst at room temperature. Aryl bromide bearing a 3-CF₃ substituent furnished the product **3ag** in 86% yield with 5 mol % catalyst at 50 °C.



Scheme 3 Scope of Aryl Bromides in Ni–NIXANTPHOS Catalyzed DCCP with **1a**^{a,b}



^a Reactions conducted on a 0.1 mmol scale at 0.05 M. ^b Isolated yields after chromatographic purification. ^c 5 mol % Ni(COD)₂/5 mol % NIXANTPHOS. ^d 10 mol % Ni(COD)₂/10 mol % NIXANTPHOS. ^e room temperature. ^f 50 °C.

We next examined our Ni(NIXANTPHOS)–based catalyst with more challenging aryl chlorides (Scheme 4). We were pleased to find that the Ni(NIXANTPHOS)–based catalyst gave good to excellent yields in the coupling of aryl chlorides with 2-benzylpyridine using catalyst loadings as low as 2.5 mol %. Coupling chlorobenzene, 1-*tert*-butyl-4-chlorobenzene, and 4-chloroanisole furnished products in 99 (**3aa**), 92 (**3ab**) and 99% (**3ac**) yields with 2.5–5 mol % catalyst loadings at room temperature. 4-Chloro-1-fluorobenzene (**4e**) led to triarylmethane products **3ae** in 90% yield with 10 mol % catalyst loading at room temperature. 4-Chlorobenzonitrile (**4h**) delivered **3ah** in 79% yield with 10 mol % catalyst loading at 50 °C, despite the known reaction of nitriles with silyl amide bases.²⁷ Chlorobenzophenone afforded **3af** in 92% yield (5 mol % catalyst loading at 50 °C) while 3-chlorobenzotrifluoride (**4g**) furnished **3ag** in 90% yield with 5 mol % catalyst loading at 50 °C. Heteroaromatic 6-chloroquinoline **4i** successfully underwent reaction in 73% yield with 5 mol % catalyst loading at room temperature. Overall, good scope of the nickel-catalyzed DCCP was demonstrated in Schemes 2–4, and aryl chlorides were found to be better substrates than aryl bromides (based on yields).


Scheme 4 Scope of Aryl Chlorides in Ni–NIXANTPHOS Catalyzed DCCP with **1a**^{a,b}



^a Reactions conducted on a 0.1 mmol scale at 0.05 M. ^b Isolated yield after chromatographic purification. ^c 2.5 mol % Ni(COD)₂/2.5 mol % NIXANTPHOS. ^d 5 mol % Ni(COD)₂/5 mol % NIXANTPHOS. ^e 10 mol % Ni(COD)₂/10 mol % NIXANTPHOS. ^f room temperature. ^g 50 °C.

To demonstrate the potential utility of our method, we conducted a gram scale reaction of chlorobenzophenone (**4f**, 5 mmol, 1.08 g) and the 2-benzylpyridine (**1a**, 10 mmol, 1.61 mL) in 85% yield (Scheme 5), suggesting the reaction is scalable.

Scheme 5 Gram scale reaction via nickel-catalyzed DCCP reaction



2.4 Comparison of NIXANTPHOS and XANTPHOS DERIVATIVES

We previously demonstrated that oxidative addition of aryl chlorides to a Pd⁰(NIXANTPHOS) catalyst took place at room temperature in the presence of base.^{17b} This result was surprising because the low energy pathway to oxidative addition of unactivated aryl chlorides to Pd(0) has been shown to proceed through a Pd–L intermediate, where L is a monodentate phosphine.²⁸ Palladium complexes of bulky trialkyl phosphines and certain Buchwald ligands will oxidatively add aryl chlorides at room temperature or below.²⁹ In contrast, Pd(0) complexes of bidentate phosphines can oxidatively add aryl chlorides via a higher energy PdL₂ pathway.³⁰ Such reactions usually take place around 90 °C, rather than at room temperature with our Pd⁰(NIXANTPHOS)–based catalyst.³¹ We also demonstrated that the NIXANTPHOS N–H must be deprotonated to perform the oxidative addition of aryl chlorides at room temperature, and that the actual catalyst is a heterobimetallic Pd/main group metal complex (main group = K, Na, Li). Although the precise mechanism of the oxidative addition in this system is currently unknown, we

speculate that cooperativity between the metals plays an important role. As a point of reference, the structurally related Pd⁰(XANTPHOS) exhibited no reactivity with aryl chlorides under the same reaction conditions.^{17b}

With this backdrop, we were interested to determine if the exceptional reactivity imparted to palladium when bound to NIXANTPHOS would translate to an increase in reactivity of analogous nickel complexes. Nickel-based phosphine catalysts are more reactive toward oxidative addition than most palladium complexes.^{12,32} Unfortunately, the nickel-based catalysts proved more challenging to study than their palladium counterparts, due to precipitate formation. Nonetheless, some experiments are presented that highlight the significant differences in reactivity between the nickel NIXANTPHOS and XANTPHOS systems.

As discussed earlier, our initial ligand screen indicated that the NIXANTPHOS-based nickel catalyst outperformed the XANTPHOS-based nickel catalyst by a significant margin. Additionally, the NIXANTPHOS ligand will be deprotonated under the nickel-catalyzed reaction conditions outlined in these arylation reactions.

To probe this difference in reactivity, we compared conversions of NIXANTPHOS- and XANTPHOS-based nickel catalysts in the arylation of 2-benzylpyridine with bromobenzene. As shown in Scheme 6, the Ni(NIXANTPHOS)-based catalyst exhibited higher conversion (64%) after 4 h at room temperature than the XANTPHOS analogue (7% conversion by ¹H NMR). The presence of a significant quantity of precipitate in the reaction with XANTPHOS was cause for concern, however. We isolated the precipitate by filtration, but attempts to redissolve it in common organic solvents for the purpose of characterization were thwarted by the insolubility. It has been reported that Pd(XANTPHOS)₂ also exhibits very low solubility.³³


We were worried that precipitate formation would decrease the amount of active catalyst in the solution. We therefore synthesized the structurally similar 4,7-di-*tert*-butyl-XANTPHOS (*t*-Bu-XANTPHOS, Scheme 6, see Supporting Information for details). Palladium complexes of *t*-Bu-XANTPHOS exhibit greater solubility than analogous XANTPHOS complexes in palladium catalyzed reactions.²⁸ When *t*-Bu-XANTPHOS was used in the nickel-catalyzed reactions, no solubility problems were encountered. However, we were surprised to find that under the same conditions the reaction with *t*-Bu-XANTPHOS exhibited *lower* conversions than both NIXANTPHOS- and XANTPHOS-based catalysts. It is possible that transmetalation is turnover limiting and the more hindered Ni(*t*-Bu-XANTPHOS)-based catalyst undergoes slower transmetalation.

When the reaction was conducted with the Ni(NIXANTPHOS)-based catalyst, 2-benzylpyridine and chlorobenzene, the conversion was 48% after 4 h. The similarity of conversion in the reactions with chloro and bromobenzene under otherwise identical conditions suggests that oxidative addition is not turnover limiting (Scheme 6).

At this point, we can conclude that the Ni(NIXANTPHOS)-based catalyst indeed exhibits much higher activity than the structurally related XANTPHOS derived catalysts. Recall also

that, as noted in the discussion of the initial screening, the Ni(NIXANTPHOS) catalyst outperformed the other 36 mono- and bidentate phosphines of the initial reactivity screen. Determination of the origin of the difference in reactivity between Ni(NIXANTPHOS)- and Ni(XANTPHOS)-based catalysts has proven challenging in the arylation of 2-benzylpyridine under our reaction conditions.

Scheme 6 Reactivity Comparisons with Ni-based Catalysts




2.5. Determination of the Reactivity of the Ni(NIXANTPHOS)-based Catalyst in the Arylation of 2-Pyridylmethyl amines.

As outlined in Equation 1, we previously demonstrated that the reactivity of the Pd(NIXANTPHOS)-based catalyst is remarkable compared to other palladium phosphine-based catalyst. Our hypothesis is this is due to the deprotonation of the palladium-ligated NIXANTPHOS ligand, which gives rise to a heterobimetallic catalyst that exhibits increased reactivity due to cooperativity between the palladium and main group metal. The significance of the nickel results above is that similar high reactivity is observed with the Ni(NIXANTPHOS)-based catalyst, suggesting that perhaps the exceptional reactivity imparted to palladium by NIXANTPHOS is likewise translatable to nickel.


To further explore this possibility, we compared the reactivity of the Ni(NIXANTPHOS)-based catalyst to 37 different Ni(phosphine)-based catalysts using microscale techniques in a novel deprotonative cross-coupling process of a 2-pyridylmethyl amine with bromobenzene and LiN(SiMe₃)₂ as base (see Supporting Information for a list of all ligands and details). This reaction and the results are illustrated in Scheme 7. The Ni(NIXANTPHOS)-based catalyst exhibited the highest product/internal standard ratio

(2.29), outperforming the other 37 ligands. The other ligands with the highest product/internal standard ratio, were ^tBu-MePhos (1.22), Di-^tBuXPhos (1.20), cataCXium PlntB (1.17), JohnPhos (1.13), Kwong's Indole ligand (1.12)³⁴ and CataXCium PtB (1.12).³⁵

Scheme 7 Reactivity Comparisons with Ni-based Catalysts and 2-Pyridylmethyl Amine **5a** with Bromobenzene.




Ligand (Loading %)	Product/Internal Standard Ratio
NIXANTPHOS (10%)	2.29
^t Bu-MePhos (20%)	1.22
Di- ^t BuXPhos (20%)	1.20
cataCXium PlntB (20%)	1.17
cataCXium PtB (20%)	1.13
Indole ligand (20%)	1.12
JohnPhos (20%)	1.12



The results with the Ni(NIXANTPHOS)-based catalyst was scaled to laboratory scale using 5 mol % Ni(COD)₂ and 7.5 mol % NIXANTPHOS. The isolated yield was 93% (Scheme 8). The XANTPHOS based catalyst, however, exhibited only 24% conversion under the same reaction conditions.

Scheme 8 Reactivity Comparisons with Ni(NIXANTPHOS) and Ni(XANTPHOS)-based Catalysts in the arylation of 2-Pyridylmethyl Amine **5a** with Bromobenzene



3 Conclusions

In summary, we have presented an efficient method to form triarylmethane derivatives from diarylmethanes and aryl bromides and chlorides employing a Ni(NIXANTPHOS)-based catalyst. The method enables functionalization of C(sp³)-H's and does not require prefunctionalized organometallic reagents. A reactivity comparison between NIXANTPHOS- and XANTPHOS-based catalysts points to enhanced reactivity of NIXANTPHOS-based nickel catalyst. To test the generality of the Ni(NIXANTPHOS)-based catalyst, a novel arylation of a 2-pyridylmethyl amine was examined. Out of the 38 mono- and bidentate cross-coupling ligands examined with Ni(COD)₂, NIXANTPHOS again showed the highest reactivity. These results are the first hint that the exceptional reactivity of NIXANTPHOS-based palladium catalysts may be translatable to other transition metal catalysts. This topic is currently under investigation in our research group.

Acknowledgements

We thank the National Science Foundation [CHE-1152488] and National Institutes of Health (NIGMS 104349) for financial support. This work is funded by International Graduate Exchange Program of Beijing Institute of Technology.

Notes and references

- (a) R. Muthyala, A. R. Katritzky and X. Lan, *Dyes Pigm.*, 1994, **25**, 303; (b) A. R. Katritzky, V. Gupta, C. Garot, C. V. Stevens and M. F. Gordeev, *Heterocycles*, 1994, **38**, 345; (c) D. F. Duxbury, *Chem. Rev.*, 1993, **93**, 381; (d) E. S. Lewis, J. M. Perry, and R. H. Grinstead, *J. Am. Chem. Soc.*, 1970, **92**, 899.
- (a) A. C. Bhasikuttan, J. Mohanty, W. M. Nau and H. Pal, *Angew. Chem., Int. Ed.*, 2007, **46**, 4120; (b) H. Abe, J. Wang, K. Furukawa, K. Oki, M. Uda, S. Tsuneda and Y. Ito, *Bioconjugate Chem.*, 2008, **19**, 1219; (c) H. N. Kim, M. H. Lee, H. J. Kim, J. S. Kim and J. Yoon, *Chem. Soc. Rev.*, 2008, **37**, 1465; (d) M. Beija, C. A. M. Afonso and J. M. G. Martinho, *Chem. Soc. Rev.*, 2009, **38**, 2410.
- M. Irie, *J. Am. Chem. Soc.*, 1983, **105**, 2078.
- (a) R. S. Dothager, K. S. Putt, B. J. Allen, B. J. Leslie, V. Nesterenko and P. J. Hergenrother, *J. Am. Chem. Soc.*, 2005, **127**, 8686; (b) S. K. Das, G. Panda, V. Chaturvedi, Y. S. Manju, A. K. Galkwad and S. Sinha, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 5586; (c) G. Panda, Shagufta; J. K. Mishra, V. Chaturvedi, A. K. Srivastava, R. Srivastava and B. S. Srivastava, *Bioorg. Med. Chem.*, 2004, **12**, 5269; (d) K. Risberg, I. J. Guldvik, R. Palchadhuri, Y. G. Xi, J. F. Ju, O. Fodstad, P. J. Hergenrother and Y. J. Andersson, *Immunother.*, 2011, **34**, 438.
- (a) R. M. Roberts, A. M. Elkhawaga, K. M. Sweeney and M. F. Elzohry, *J. Org. Chem.*, 1987, **52**, 1591; (b) C. Ramesh, J. Banerjee, R. Pal and B. Das, *Adv. Synth. Catal.*, 2003, **345**, 557; (c) J. S. Yadav, B. V. S. Reddy and S. Sunitha, *Adv. Synth. Catal.*, 2003, **345**, 349; (d) V. Nair, K. G. Abhilash and N. Vidya, *Org. Lett.*, 2005, **7**, 5857; (e) J. Esquivias, R. G. Arrayas and J. C. Carretero, *Angew. Chem., Int. Ed.*, 2006, **45**, 629; (f) V. Nair, N. Vidya and K. G. Abhilash, *Synthesis*, 2006, 3647; (g) S. H. Lin and X. Y. Lu, *J. Org. Chem.*, 2007, **72**, 9757; (h) S. Podder, J. Choudhury, U. K. Roy and S. Roy, *J. Org. Chem.*, 2007, **72**, 3100; (i) I. Alonso, J. Esquivias, R. Gomez-Arrayas and J. C. Carretero, *J. Org. Chem.*, 2008, **73**, 6401; (j) G. J. Li, E. J. Wang, H. Y. Chen, H. F. Li, Y. H. Liu and P. G. Wang, *Tetrahedron*, 2008, **64**, 9033; (k) Z. X. Li, Z. Duan, J. X. Kang, H. Q. Wang, L. J. Yu and Y. J. Wu, *Tetrahedron*, 2008, **64**, 1924; (l)

- C. R. Liu, M. B. Li, C. F. Yang, S. K. Tian, *Chem. Commun.*, 2008, 1249; (m) Z. Y. Wang, X. Y. Sun and J. Wu, *Tetrahedron*, 2008, **64**, 5013.
- 6 (a) Shi reported a Fe-catalyzed cross dehydrogenative coupling via a Friedel–Crafts-type mechanism with substituted anisoles: Y. Z. Li, B. J. Li, X. Y. Lu, S. Lin and Z. J. Shi, *Angew. Chem., Int. Ed.*, 2009, **48**, 3817; (b) Han reported a Pd-catalyzed arylation of diphenylmethyl acetate with heteroarenes: F. Q. Yuan, L. X. Gao and F. S. Han, *Chem. Commun.*, 2011, **47**, 5289.
- 7 (a) J. Y. Yu and R. Kuwano, *Org. Lett.*, 2008, **10**, 973; (b) T. Niwa, H. Yorimitsu and K. Oshima, *Org. Lett.*, 2007, **9**, 2373; (c) S. Tabuchi, K. Hirano, T. Satoh and M. Miura, *J. Org. Chem.*, 2014, **79**, 5401; (d) M. Nambo and G. M. Crudden, *ACS Catal.*, 2015, **5**, 4734; (e) A. López-Pérez, J. Adrio and J. C. Carretero, *Org. Lett.*, 2009, **11**, 5514; (f) Molander, G. A.; Elia, M. D. *J. Org. Chem.* 2006, **71**, 9198; (g) Yu, J.-Y.; Kuwano, R. *Org. Lett.* 2008, **10**, 973; (h) Nambo, M.; Crudden, C. M. *Angew. Chem., Int. Ed.* 2014, **53**, 742.
- 8 (a) B. L. H. Taylor, M. R. Harris and E. R. Jarvo, *Angew. Chem., Int. Ed.*, 2012, **51**, 7790; (b) A. G. Johnson, M. M. Tranquilli, M. R. Harris and E. R. Jarvo, *Tetrahedron Lett.*, 2015, **56**, 3486; (c) C. Gosmini, J. M. Begouin and A. Moncomble, *Chem. Commun.*, 2008, 3221; (d) B. D. Sherry and A. Fürstner, *Acc. Chem. Res.*, 2008, **41**, 1500; (e) W. M. Czaplik, M. Mayer, J. Cvengroš and A. J. von Wangelin, *ChemSusChem*, 2009, **2**, 396; (f) D. S. Surry and S. L. Buchwald, *Chem. Sci.*, 2010, **1**, 13; (g) G. Evano, C. Theunissen and A. Pradal, *Nat. Prod. Rep.*, 2013, **30**, 1467.
- 9 M. R. Harris, L. E. Hanna, M. A. Greene, C. E. Moore and E. R. Jarvo, *J. Am. Chem. Soc.*, 2013, **135**, 3303.
- 10 Q. Zhou, H. D. Srinivas, S. Dasgupta and M. P. Watson, *J. Am. Chem. Soc.*, 2013, **135**, 3307.
- 11 Q. Chen, X. H. Fan, L. P. Zhang and L. M. Yang, *RSC Adv.*, 2015, **5**, 15338.
- 12 (a) B. M. Rosen, K. W. Quasdorf, D. A. Wilson, N. Zhang, A. M. Resmerita, N. K. Garg and V. Percec, *Chem. Rev.*, 2011, **111**, 1346; (b) S. Z. Tasker, E. A. Standley and T. F. Jamison, *Nature*, 2014, **509**, 299; (c) W. M. Czaplik, M. Mayer, J. Cvengroš and A. J. von Wangelin, *ChemSusChem*, 2009, **2**, 396; (d) J. Cornella, C. Zarate and R. Martin, *Chem. Soc. Rev.*, 2014, **43**, 8081; (e) M. Henrion, V. Ritleng and M. J. Chetcuti, *ACS Catal.*, 2015, **5**, 1283; (f) F. Glorius, *Angew. Chem., Int. Ed.*, 2008, **47**, 8347.
- 13 For some recent examples of nickel-catalyzed coupling reactions see (a) J. A. Fernandez-Salas, E. Marelli and S. P. Nolan, *Chem. Sci.*, 2015, **6**, 4973; (b) A. R. Martin, D. J. Nelson, S. Meiries, A. M. Z. Slawin and S. P. Nolan, *Eur. J. Org. Chem.*, 2014, 3127; (c) S. Ge, R. A. Green and J. F. Hartwig, *J. Am. Chem. Soc.*, 2014, **136**, 1617; (d) S. Ge and J. F. Hartwig, *J. Am. Chem. Soc.*, 2011, **133**, 16330; (e) A. Borzenko, N. L. Rottaloria, P. M. MacQueen, C. M. Lavoie, R. McDonald and M. Stradiotto, *Angew. Chem., Int. Ed.*, 2015, **54**, 3773; (f) M. Grigalunas, T. Ankner, P.-O. Norrby, O. Wiest and P. Helquist, *J. Am. Chem. Soc.*, 2015, **137**, 7019; (g) A. M. Whittaker and V. M. Dong, *Angew. Chem., Int. Ed.*, 2015, **54**, 1312; (h) O. Gutierrez, J. C. Tellis, D. N. Primer, G. A. Molander, M. C. Kozlowski, *J. Am. Chem. Soc.*, 2015, **137**, 4896.
- 14 For examples of Ni-catalyzed C-H functionalization reactions see (a) H. Shiota, Y. Ano, Y. Aihara, Y. Fukumoto and N. Chatani, *J. Am. Chem. Soc.*, 2011, **133**, 14952; (b) L. Ackermann, B. Punji and W. Song, *Adv. Synth. Catal.*, 2011, **353**, 3325; (c) I. Hyodo, M. Tobisu and N. Chatani, *Chem. Asian J.*, 2012, **7**, 1357; (d) X. Wu, Y. Zhao and H. Ge, *Chem. Eur. J.*, 2014, **20**, 9530; (e) M. Li, J. Dong, X. Huang, K. Li, Q. Wu, F. Song and J. You, *Chem. Commun.*, 2014, **50**, 3944; (f) W. Song, S. Lackner and L. Ackermann, *Angew. Chem., Int. Ed.*, 2014, **53**, 2477; (g) X. Wu, Y. Zhao and H. Ge, *J. Am. Chem. Soc.*, 2014, **136**, 1789; (h) J. S. Bair, Y. Schramm, A. G. Sergeev, E. Clot, O. Eisenstein and J. F. Hartwig, *J. Am. Chem. Soc.*, 2014, **136**, 13098; (i) C. Lin, W. Yu, J. Yao, B. Wang, Z. Liu and Y. Zhang, *Org. Lett.*, 2015, **17**, 1340; (j) S.-Y. Yan, Y.-J. Liu, B. Liu, Y.-H. Liu and B.-F. Shi, *Chem. Commun.*, 2015, **51**, 4069.
- 15 For selective catalytic functionalization of weakly acidic sp³-hybridized C–H bonds, see (a) S. C. Sha, J. D. Zhang and P. J. Walsh, *Org. Lett.*, 2015, **17**, 410; (b) M. Y. Li, S. Berritt and P. J. Walsh, *Org. Lett.*, 2014, **16**, 4312; (c) M. Y. Li, B. Yucel, J. Adrio, A. Bellomo and P. J. Walsh, *Chem. Sci.*, 2014, **5**, 2383; (d) T. J. Jia, A. Bellomo, K. E. Baina, S. D. Dreher and P. J. Walsh, *J. Am. Chem. Soc.*, 2013, **135**, 3740.
- 16 For related approaches with azaaryl methane derivatives see (a) T. Niwa, H. Yorimitsu, K. Oshima, *Org. Lett.*, 2007, **9**, 2373; (b) G. Song, Y. Su, X. Gong, K. Han, X. Li, *Org. Lett.*, 2011, **13**, 1968; (c) G. I. McGrew, J. Temaismithi, P. J. Carroll, P. J. Walsh, *Angew. Chem., Int. Ed.*, 2010, **49**, 5541.
- 17 (a) J. Zhang, A. Bellomo, A. D. Creamer, S. D. Dreher and P. J. Walsh, *J. Am. Chem. Soc.*, 2012, **134**, 13765; (b) J. Zhang, A. Bellomo, N. Trongsiwat, T. Z. Jia, P. J. Carroll, S. D. Dreher, M. T. Tudge, H. L. Yin, J. R. Robinson, E. J. Schelter and P. J. Walsh, *J. Am. Chem. Soc.*, 2014, **136**, 6276; (c) A. Bellomo, J. Zhang, N. Trongsiwat and P. J. Walsh, *Chem. Sci.*, 2013, **4**, 849; (d) I. Popov, S. Lindeman, O. Daugulis, *J. Am. Chem. Soc.*, 2011, **133**, 9286; (e) H.-Q. Do, O. Daugulis, *J. Am. Chem. Soc.*, 2007, **129**, 12404; (f) H.-Q. Do, R. K. M. Khan, O. Daugulis, *J. Am. Chem. Soc.*, 2008, **130**, 15185; (g) L. D. Tran, O. Daugulis, *Org. Lett.*, 2010, **12**, 4277.
- 18 P. C. J. Kamer, P. W. N. M. van Leeuwen and J. N. H. Reek, *Acc. Chem. Res.*, 2001, **34**, 895.
- 19 F. G. Bordwell, *Acc. Chem. Res.*, 1988, **21**, 456.
- 20 F. G. Bordwell, W. S. Matthews, N. R. Vanier, *J. Am. Chem. Soc.*, 1975, **97**, 442.
- 21 While many catalytic reactions are known to proceed through a Ni(0)/Ni(II) redox couple, reactions with other oxidation states are also known. See: J. Cornella, E. Gómez-Bengoa, R. J. Martin, *Am. Chem. Soc.*, 2013, **135**, 1997.
- 22 For discussion of how the electronegativity of Ni impacts reactivity see: (a) S. Z. Tasker, E. A. Standley, T. F. Jamison, *Nature*, 2014, **509**, 299; (b) T. T. Tsou, J. K. Kochi, *J. Am. Chem. Soc.*, 1979, **101**, 6319. This article presents an excellent investigation into the mechanism of oxidative addition of Ni(0) into aryl halides and the factors determining rate, selectivity, and solvent effects of such reactions. (c) E. L. Lanni, A. J. McNeil, *J. Am. Chem. Soc.* 2009, **131**, 16573.
- 23 B. Cordero, V. Gómez, A. E. Platero-Prats, M. Revés, J. Echeverría, E. Cremades, F. Barragán, S. Alvarez, *Dalton Trans.* 2008, 2832.
- 24 R. R. Fraser, T. S. Mansour, S. Savard, *J. Org. Chem.* **1985**, **50**, 3232-3234.
- 25 (a) G. Pizzo, M. R. Piscopo, I. Pizzo and G. Giuliana, *Clin Oral Invest.*, 2007, **11**, 189; (b) J. L. Gabriel, T. F. J. Miller, M. R. Wolfson and T. H. Shaffer, *ASAIO Journal.*, 1996, **42**, 968.
- 26 J. Shi, X. Zhang and D. C. Neckers, *J. Org. Chem.*, 1992, **57**, 4418.
- 27 D. A. Culkun and J. F. Hartwig, *Acc. Chem. Res.*, 2003, **36**, 234.
- 28 (a) G. C. Fu, *Acc. Chem. Res.*, 2008, **41**, 1555; (b) R. Martin and S. L. Buchwald, *Acc. Chem. Res.*, 2008, **41**, 1461; (c) D. S. Surry and S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2008, **47**, 6338; (d) U. Christmann and R. Vilar, *Angew. Chem., Int. Ed.*, 2005, **44**, 366; (e) F. Barrios-Landeros and J. F. Hartwig, *J. Am. Chem. Soc.*, 2005, **127**, 6944; (f) F. Barrios-Landeros, B. P. Carrow and J. F. Hartwig, *J. Am. Chem. Soc.*, 2009, **131**, 8141; (g) A. K. de K. Lewis, S. Caddick, F. G. N. Cloke, N. C. Billingham, P. B. Hitchcock and J. Leonard, *J. Am. Chem. Soc.*, 2003, **125**, 10066; (h) E. Galardon, S. Ramdeehul, J. M.

- Brown, A. Cowley, K. K. Hii and A. Jutand, *Angew. Chem., Int. Ed.*, 2002, **41**, 1760; See references on computational studies of oxidative addition of aryl chlorides to Pd(0): (i) F. Schoenebeck and K. N. Houk, *J. Am. Chem. Soc.*, 2010, **132**, 2496; (j) H. M. Senn, and T. Ziegler, *Organometallics*, 2004, **23**, 2980; (g) S. Kozuch, C. Amatore, A. Jutand and S. Shaik, *Organometallics*, 2005, **24**, 2319; (k) M. Ahlquist and P.-O. Norrby, *Organometallics*, 2007, **26**, 550.
- 29 J. P. Wolfe, R. A. Singer, B. H. Yang and S. L. Buchwald, *J. Am. Chem. Soc.*, 1999, **121**, 9550; (b) T. Kinzel, Y. Zhang and S. L. Buchwald, *J. Am. Chem. Soc.*, 2010, **132**, 14073; (c) Y. Yang, N. J. Oldenhuis, S. L. Buchwald, *Angew. Chem., Int. Ed.*, 2013, **52**, 615.
- 30 (a) Y. Ben-David, M. Portnoy, D. Milstein, *J. Am. Chem. Soc.*, 1989, **111**, 8742; (b) Y. Ben-David, M. Portnoy and D. Milstein, *J. Chem. Soc. Chem. Commun.*, 1989, 1816.
- 31 M. Portnoy and D. Milstein, *Organometallics*, 1993, **12**, 1665. Note that the oxidative addition of 10-fold excess of PhCl to Pd(dipp)₂ required 90 °C to go to completion in 2 h in dioxane. The temperature range of the kinetic measurements was 30–70 °C.
- 32 T. T. Tsou and J. K. Kochi, *J. Am. Chem. Soc.*, 1979, **101**, 6319.
- 33 L. M. Klingensmith, E. R. Strieter, T. E. Barder and S. L. Buchwald, *Organometallics*, 2006, **25**, 82.
- 34 (a) H. W. Lee, F. L. Lam, C. M. So, C. P. Lau, A. S. C. Chan, F. Y. Kwong, *Angew. Chem.*, 2009, **121**, 7572; (b) C. M. So, C. P. Lau, F. Y. Kwong, *Org. Lett.*, 2007, **9**, 2795; (c) C. M. So, C. P. Lau, F. Y. Kwong, *Chem. Eur. J.*, 2011, **17**, 761; (d) S. M. Wong, C. M. So, K. H. Chung, C. P. Lau, F. Y. Kwong, *Eur. J. Org. Chem.*, 2012, **2012**, 4172; (e) P. Y. Yeung, K. H. Chung, F. Y. Kwong, *Org. Lett.*, 2011, **13**, 2912.
- 35 (a) D. S. Surry, S. L. Buchwald, *Chem. Sci.*, 2011, **2**, 27; (b) N. C. Bruno, M. T. Tudge, S. L. Buchwald, *Chem. Sci.*, 2013, **4**, 916.

TOC

