


# Chemical Science

Accepted Manuscript




This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## Table of Contents



Catalytic enantioselective method of consecutive Suzuki-Miyaura alkylations of *gem*-chloro(iodo)alkanes to form two C-C bonds in one pot transformation is described.



Journal Name

ARTICLE

## Enantioselective Assembly of Tertiary Stereocenters via Multicomponent Chemoselective Cross-Coupling of Geminal Chloro(iodo)alkanes†

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

X. Jiang,<sup>a</sup> K. Kulbitski,<sup>a</sup> G. Nisnevich<sup>a</sup> and M. Gandelman<sup>a</sup>

Tertiary stereocenters represent a ubiquitous and highly important motif in many pharmaceutically active compounds and natural products. Development of efficient and straightforward approaches to their enantioselective construction from readily available simple substrates is an important yet challenging goal for synthetic chemistry. Herein we describe an efficient, versatile and facile method for the highly enantioselective construction of tertiary stereocenters via unprecedented consecutive one-pot Suzuki reactions of non-activated racemic 1-chloro-1-iodoalkanes with alkylboranes. It represents the first cross-coupling approach which employs simple and readily available primary alkyl substrates for the direct multicomponent assembly of enantioenriched tertiary stereocenters. A simple and effective preparation of 1-chloro-1-iodoalkanes from ubiquitous  $\alpha$ -chloroalkanoic acids is also described. Collectively, the developed methods open a door to efficient catalytic enantioselective synthesis of alkanes bearing tertiary stereocenters from carboxylic acids just in few steps.

### Introduction

Metal catalyzed cross-couplings are extremely useful and efficient reactions for construction of carbon-carbon bonds.<sup>1</sup> Contrary to  $C_{sp^2}$ -based electrophiles, coupling of  $C_{sp^3}$ -based partners has proven a daunting challenge due to the relatively slow oxidative addition of alkyl halides and the undesirable  $\beta$ -hydride elimination that alkyl halides might undergo.<sup>2</sup> Development of efficient catalysts for alkyl-alkyl couplings of primary and secondary alkyl electrophiles has emerged as an important frontier with a potential to significantly extend the synthetic power of the cross-coupling transformations.<sup>3</sup> In this context, utilization of secondary alkyl electrophiles (or nucleophiles) in cross-couplings is especially attractive as it opens a door to the catalytic asymmetric synthesis of stereogenic tertiary centers.<sup>3d-e,4</sup> All-carbon tertiary stereocenters represent an ubiquitous and highly important motif in many pharmaceutically active compounds and natural products, and the development of efficient and straightforward approaches to their construction is an important goal for organic chemistry.<sup>5</sup>


There are two major approaches that utilize cross-coupling reactions of non-activated alkyl partners for the synthesis of enantioenriched products with tertiary stereocenters.<sup>4-6</sup> The first stereospecific approach depends on substrate control (electrophile or nucleophile) as the source of stereochemical information.<sup>4a</sup> This process is mediated by achiral catalysts, and requires use of the enantioenriched secondary electrophiles or nucleophiles in the stoichiometric amount. The second approach employs racemic secondary electrophiles (or nucleophiles) as starting materials, and the stereoselectivity of the product is controlled by substoichiometric amounts of the chiral catalyst (Scheme 1a).<sup>4a</sup> Recently, remarkable catalyst-controlled enantioselective cross-couplings have been developed for a number of families of non-activated secondary alkyl halides.<sup>7</sup> It was demonstrated that this process is stereoconvergent as both stereoisomers of the racemic alkyl halides were converted into a single stereoisomer of product as the reaction proceeds through a common achiral intermediate.<sup>7c</sup> Notably, all reported cross-coupling processes for the preparation of tertiary stereocenters utilize secondary alkyl partners as starting materials.<sup>4a</sup> However, synthesis of non-symmetrical secondary alkyl halides (or, alternatively, alkylmetal reagents) might be multistep and tedious, typically involving reaction of Grignard reagents (functional group sensitive) with carbonyls or epoxides. Therefore, alternative approaches which rely on new easily available substrates for stereoselective construction of tertiary centers via cross-coupling reactions could

<sup>a</sup> Dr. X. Jiang, K. Kulbitski, Dr. G. Nisnevich, Prof. Dr. M. Gandelman  
Schulich Faculty of Chemistry  
Technion-Israel Institute of Technology  
Technion City, Haifa32000, Israel  
E-mail: chmark@tx.technion.ac.il

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

significantly extend the potential of this method in terms of applicability and functional group compatibility.




**Scheme 1.** Cross-coupling approaches for stereoconvergent construction of tertiary stereocenters (a) established method employing secondary partners; (b) One-pot sequential cross-couplings of geminal 1-chloro-1-iodoalkanes with primary alkylboranes (this work).

## Results and Discussion

Herein we describe a novel multicomponent method for the facile and highly enantioselective construction of tertiary stereocenters via Suzuki reactions of non-activated racemic geminal dihaloalkanes with primary alkylboranes. This represents the first example of cross-coupling that employs simple primary alkyl electrophiles and nucleophiles for the assembly of tertiary centers (Scheme 1b). We demonstrate the unprecedented use of geminal chloro(iodo)alkanes in cross-coupling reactions, in which the chemoselective functionalization of the C-I bond (bond dissociation energy ( $BDE_1$ ) = 57 kcal/mol)<sup>8</sup> with the first alkylborane, followed by functionalization of the stronger C-Cl bond ( $BDE_2$  = 84 kcal/mol)<sup>8</sup> with a second alkylborane in a consecutive step resulting in a one-pot transformation. Importantly, *the same catalyst system* and reaction conditions are used for both steps of this stereoconvergent transformation. Moreover, we disclose a general and efficient approach to access 1-chloro-1-iodoalkanes. This chemistry relies on the facile iododecarboxylation of  $\alpha$ -chloroalkanoic acids, which, in turn, are easily prepared by  $\alpha$ -chlorination of the corresponding alkanolic acids.<sup>9</sup> Collectively these methods constitute a short, straightforward and efficient approach for the construction of highly enantioenriched alkanes bearing tertiary stereocenters (up to 99% *ee*) from abundant primary alkanolic acids.

1-chloro-1-iodoalkanes constitute an attractive class of polyfunctional compounds. However, the development of their chemistry has been an uneven process, most likely, due to the lack of selective, practical and high yielding methods of their synthesis.<sup>10</sup> Recently, we reported a robust, general and efficient method for synthesis of alkyl iodides from carboxylic acids through their simple treatment with commercially available 1,3-diiodo-5,5-dimethylhydantoin (DIH) under visible light (VIS) irradiation.<sup>11</sup> We now report that, contrary to other iododecarboxylation approaches,<sup>10d</sup> our method can be

uniquely extended to  $\alpha$ -chloroalkanoic acids. Thus, these compounds react smoothly with DIH under VIS-irradiation to generate the corresponding 1-chloro-1-iodoalkanes (Scheme 2). These products are isolated in an essentially pure form after a simple aqueous work-up of the reaction mixture (see supporting information). All 1-chloro-1-iodoalkanes presented in this paper (Tables 1, 3, 4) were prepared and isolated by this method, starting from the corresponding  $\alpha$ -chloroalkanoic acids, in a selective manner and high yields.




**Scheme 2.** Synthesis of geminal chloro(iodo)alkanes via iododecarboxylation of  $\alpha$ -chloroalkanoic acids.

Having an efficient synthetic access to these synthons in hand, we examined their applicability in the construction of tertiary stereocenters through a direct multicomponent reaction. Based on the substantial difference in the  $BDE$ 's of C-I and C-Cl bonds, we hypothesized that it might be possible to selectively activate the C-I bond in a cross-coupling reaction with alkylmetal  $R_1-M$ , following by functionalization of the C-Cl bond in a second cross-coupling with a different alkylmetal  $R_2-M$  (see Scheme 1b). Ideally, one catalyst system might be able to mediate both steps. Since the starting material, 1-chloro-1-iodoalkane, is racemic, a stereoconvergent synthesis of enantiomerically enriched products with tertiary chiral centers may be envisioned if an appropriate chiral catalyst could be identified.


Initiating an active program on studies of this general concept of multicomponent assembly of the tertiary chiral centers, we started from investigating homobenzylic geminal chloroiodides in the asymmetric Suzuki alkyl-alkyl coupling reactions. Recently, Fu *et al* reported the first asymmetric Suzuki cross-coupling of secondary homobenzylic bromides with alkyl-9-BBN.<sup>7a</sup> In this seminal work, it was demonstrated that  $ArCH_2$ -group in electrophile is necessary for ensuring good enantioselectivity, most likely, due to its weak secondary interaction with the catalyst. We firstly investigated the reaction of homobenzylic dihalide substrate **1a** in the reaction with different alkylborane agents in the presence of  $NiCl_2 \cdot glyme$  and diamine ligand **2** (Table 1). Gratifyingly, subjecting **1a** to the coupling conditions with tripropylborane, following by addition of  $Ph(CH_2)_3-9-BBN$  to the reaction mixture (after complete conversion of **1a**), resulted in formation of the tertiary asymmetric alkane **3a** which was isolated in 73% yield and 86% *ee* (Table 1, entry 1). Remarkably, the same catalyst system is operative in both steps of the transformation. The use of alkyl-9-BBN as the nucleophile source in both stages of the reaction proved to be superior to other tested boron reagents. The replacement of  $Pr_3B$  with  $Pr-9-BBN$  furnished the product in even higher yield

and enantioselectivity (84% yield, 96% *ee*; entry 2). Inversion of the addition sequence, i.e. first adding the bulkier Ph(CH<sub>2</sub>)<sub>3</sub>-9-BBN in the first step following by the less bulky Pr-9-BBN, proved to be less efficient (entry 3).

**Table 1:** Investigation of the double alkyl-alkyl Suzuki cross-coupling reaction.<sup>[a]</sup>



Entry	Borane A	Borane B	R <sub>1</sub>	R <sub>2</sub>	Yield <sup>[b]</sup> , <i>ee</i> (%)
1	Pr <sub>3</sub> B	Ph(CH <sub>2</sub> ) <sub>3</sub> -9-BBN	Pr	Ph(CH <sub>2</sub> ) <sub>3</sub>	73, 86
2	Pr-9-BBN	Ph(CH <sub>2</sub> ) <sub>3</sub> -9-BBN	Pr	Ph(CH <sub>2</sub> ) <sub>3</sub>	84, 96
3	Ph(CH <sub>2</sub> ) <sub>3</sub> -9-BBN	Pr-9-BBN	Ph(CH <sub>2</sub> ) <sub>3</sub>	Pr	70, 82



[a] Reactions were conducted with **1a** (0.2 mmol), NiCl<sub>2</sub>·glyme (0.02 mmol), ligand (0.024 mmol), borane A (0.26 mmol), KO<sup>t</sup>Bu (0.2 mmol), *i*-BuOH (0.26 mmol) in *i*-Pr<sub>2</sub>O (2 mL) at RT. After 24 h, borane B (0.4 mmol), KO<sup>t</sup>Bu (0.28 mmol), *i*-BuOH (0.4 mmol) were added and adjusted the total volume to 4 mL with *i*-Pr<sub>2</sub>O. [b] Isolated yields.


Any deviation from these reaction conditions resulted in inferior results (Table 2). We tested the chiral secondary diamine ligands **4-9** in the multicomponent cross-coupling of **1a** with Pr-9-BBN and Ph(CH<sub>2</sub>)<sub>3</sub>-9-BBN; however, none of these outperformed the original ligand **2** (compare entry 1 with entries 2-7). Reaction using ligand **6**, bearing electron rich aryl substituents on the ethylenediamine backbone, provided **3a** with enantioselectivity comparable to ligand **2** (94% *ee*, entry 4). However analogs of **6** with electron poor aryls or bulky naphthyl substituents gave the product in significantly lower *ee* (entries 2, 3 and 5).

Interestingly, bis-pyrrolidine ligand **8**, which was recently used by us for the efficient synthesis of chiral secondary fluoroalkanes via Suzuki cross-coupling of 1-bromo-1-fluoroalkanes,<sup>12</sup> also afforded a product with high enantioselectivity, albeit with considerably lower yield (entry 6). Utilizing other sources of Ni(II) or Ni(0) instead of NiCl<sub>2</sub>·glyme, as well as lower loadings of Ni salt and ligand, eroded the reaction efficiency (entries 8, 9 and 13). While dioxane is unsuitable as a solvent (entry 12), diethyl ether proved to be as efficient as diisopropyl ether (entry 11). We preferred to use the higher boiling diisopropyl ether in order to ensure a constant concentration of the reaction partners in the course of the process.


Having arrived at the optimal reaction conditions, we explored the substrates scope of this multicomponent coupling (Tables 3 and 4). The transformation is very versatile as the R, R<sub>1</sub> and R<sub>2</sub> fragments can be easily varied upon assembly of tertiary center; therefore, diverse combinations can be envisioned (see Fig. 1b). We found that the

introduction of a relatively compact alkyl group in the first step ensures overall high enantioselectivity (*vide supra*; see Table 1).

**Table 2:** Optimization of reaction conditions.<sup>[a]</sup>



Entry	Variation from the "standard" conditions	Yield <sup>[b]</sup> , <i>ee</i> (%)
1		84, 96
2	<b>4</b> as ligand instead of <b>2</b>	79, 86
3	<b>5</b> as ligand instead of <b>2</b>	71, 73
4	<b>6</b> as ligand instead of <b>2</b>	80, 94
5	<b>7</b> as ligand instead of <b>2</b>	82, 68
6	<b>8</b> as ligand instead of <b>2</b>	65, 90
7	<b>9</b> as ligand instead of <b>2</b>	67, 78
8	NiBr <sub>2</sub> ·diglyme as catalyst instead of NiCl <sub>2</sub> ·glyme	77, 89
9	Ni(cod) <sub>2</sub> as catalyst instead of NiCl <sub>2</sub> ·glyme	68, 87
10	Hexanol instead of <i>i</i> -BuOH	80, 92
11	Et <sub>2</sub> O as solvent instead of <i>i</i> -Pr <sub>2</sub> O	81, 96
12	Dioxane as solvent instead of <i>i</i> -Pr <sub>2</sub> O	75, 80
13	6% Cat. and 8% ligand instead of 10%/12%	61, 84



[a] Reactions were conducted with **1a** (0.2 mmol), Ni salt (0.02 mmol), ligand (0.024 mmol), Propyl-9-BBN (0.26 mmol), KO<sup>t</sup>Bu (0.2 mmol), *i*-BuOH or Hexanol (0.26 mmol) in *i*-Pr<sub>2</sub>O or dioxane (2 mL) RT. After 24 h, Ph(CH<sub>2</sub>)<sub>3</sub>-9-BBN (0.4 mmol), KO<sup>t</sup>Bu (0.28 mmol), *i*-BuOH or Hexanol (0.4 mmol) were added and adjusted the total volume to 4 mL with the corresponding solvent. [b] Isolated yield.

**Table 3:** Scope of R<sub>1</sub>-9-BBN.<sup>[a]</sup>

Entry	R <sub>1</sub>	R <sub>2</sub>	Product	Yield <sup>[b]</sup> , <i>ee</i> (%)
1	Et	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>3b</b>	79, 92
2	Pr	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>3a</b>	85, 96
3	Bu	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>3c</b>	80, 98
4	Pentyl	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>3d</b>	81, 80
5	Hexyl	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>3e</b>	83, 89
6	Ph(CH <sub>2</sub> ) <sub>3</sub>	4-MeOC <sub>6</sub> H <sub>4</sub> (CH <sub>2</sub> ) <sub>3</sub>	<b>3f</b>	68, 78


[a] Reactions were conducted with **1a** (0.5 mmol), Ni salt (0.05 mmol), ligand **2** (0.06 mmol), R<sub>1</sub>-9-BBN (0.65 mmol), KO<sup>t</sup>Bu (0.5 mmol), *i*-BuOH (0.65 mmol) in *i*-Pr<sub>2</sub>O 5 mL at RT. After 24 h, R<sub>2</sub>-9-BBN (1.0 mmol), KO<sup>t</sup>Bu (0.7 mmol), *i*-BuOH (1.0 mmol) were added and adjusted the total volume to 10 mL with *i*-Pr<sub>2</sub>O. [b] Isolated yields.

As such, several unbulky alkyl-9-BBN were initially examined as trans-metallation partners in the first step (Table 3). Although a very good to high enantioselectivity is obtained in each case, the variation of the chain length in R<sub>1</sub> group has a distinguishable influence. While the shorter ethyl, propyl and butyl motifs ensure highly enantioenriched products (entries 1-3), the longer pentyl, hexyl and 3-phenylpropyl units lead to slightly diminished *ee* (entries 4-6).

Selecting propyl-9-BBN as a representative nucleophile for the first cross-coupling event, we demonstrated the flexibility

of the reaction with respect to the alkylborane, utilized in the second step (R<sub>2</sub>-9-BBN), and 2-aryl-*gem*-chloro(iodo)ethanes (ArCH<sub>2</sub>CHCl; **1a-k**). These results are summarized in Table 4. A wide array of non-activated homoaryl geminal chloriodides was tolerated in the reaction. Both phenyl as well as *ortho*-, *meta*- and *para*-substituted phenyls with both electron donating and withdrawing groups participated in the reaction affording the products in high yields and enantioselectivities with variety of organoboron partners. Interestingly, having an electron donating methoxy-group in the *para*-position of the aromatic ring in homoaryl dihalide (i.e. compound **1b**) was beneficial providing **10e-10l** in enantioselectivities up to 99% *ee* (entries 6-13).

**Table 4:** Scope of 2-aryl-*gem*-chloro(iodo)ethanes and R<sub>2</sub>-9-BBN.<sup>[a]</sup>



Entry	Ar	Substrate	R	Product	Yield <sup>[b]</sup> , ee (%)
1	Ph	<b>1a</b>	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>3a</b>	85, 96
2	Ph	<b>1a</b>	4-FC <sub>6</sub> H <sub>4</sub> (CH <sub>2</sub> ) <sub>3</sub>	<b>10a</b>	81, 92
3	Ph	<b>1a</b>	4-MeC <sub>6</sub> H <sub>4</sub> (CH <sub>2</sub> ) <sub>3</sub>	<b>10b</b>	85, 87
4	Ph	<b>1a</b>	4-MeOC <sub>6</sub> H <sub>4</sub> (CH <sub>2</sub> ) <sub>3</sub>	<b>10c</b>	88, 82
5	Ph	<b>1a</b>	Ph(CH <sub>2</sub> ) <sub>4</sub>	<b>10d</b>	81, 78
6	4-MeOC <sub>6</sub> H <sub>4</sub>	<b>1b</b>	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>10e</b>	82, >99.5
7	4-MeOC <sub>6</sub> H <sub>4</sub>	<b>1b</b>	2-MeOC <sub>6</sub> H <sub>4</sub> (CH <sub>2</sub> ) <sub>3</sub>	<b>10f</b>	88, 99.5
8	4-MeOC <sub>6</sub> H <sub>4</sub>	<b>1b</b>	4-MeOC <sub>6</sub> H <sub>4</sub> (CH <sub>2</sub> ) <sub>3</sub>	<b>10g</b>	87, 94
9	4-MeOC <sub>6</sub> H <sub>4</sub>	<b>1b</b>	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub> (CH <sub>2</sub> ) <sub>3</sub>	<b>10h</b>	82, 95
10	4-MeOC <sub>6</sub> H <sub>4</sub>	<b>1b</b>	4-FC <sub>6</sub> H <sub>4</sub> (CH <sub>2</sub> ) <sub>3</sub>	<b>10i</b>	85, 94
11	4-MeOC <sub>6</sub> H <sub>4</sub>	<b>1b</b>	4-MeC <sub>6</sub> H <sub>4</sub> (CH <sub>2</sub> ) <sub>3</sub>	<b>10j</b>	86, 92
12	4-MeOC <sub>6</sub> H <sub>4</sub>	<b>1b</b>	TBSO(CH <sub>2</sub> ) <sub>3</sub>	<b>10k</b>	80, 81
13	4-MeOC <sub>6</sub> H <sub>4</sub>	<b>1b</b>	Cyclohexyl(CH <sub>2</sub> ) <sub>3</sub>	<b>10l</b>	84, 78
14	2-MeC <sub>6</sub> H <sub>4</sub>	<b>1c</b>	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>10m</b>	78, 87
15	3-MeC <sub>6</sub> H <sub>4</sub>	<b>1d</b>	TBSO(CH <sub>2</sub> ) <sub>3</sub>	<b>10n</b>	84, 80
16	4-MeC <sub>6</sub> H <sub>4</sub>	<b>1e</b>	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>10o</b>	86, 78
17	2-FOC <sub>6</sub> H <sub>4</sub>	<b>1f</b>	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>10p</b>	83, 72
18	3-FOC <sub>6</sub> H <sub>4</sub>	<b>1g</b>	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>10q</b>	80, 82
19	4-FOC <sub>6</sub> H <sub>4</sub>	<b>1h</b>	TBSO(CH <sub>2</sub> ) <sub>3</sub>	<b>10r</b>	79, 38
20	4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	<b>1i</b>	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>10s</b>	81, 89
21	2-Napthyl	<b>1j</b>	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>10t</b>	58, 71
22	3-MeOC <sub>6</sub> H <sub>4</sub>	<b>1k</b>	Ph(CH <sub>2</sub> ) <sub>3</sub>	<b>10u</b>	78, 88


[a] Reactions were conducted with **1** (0.5 mmol), Ni salt (0.05 mmol), ligand **2** (0.06 mmol), *n*-Propyl-9-BBN (0.65 mmol), KO<sup>t</sup>Bu (0.5 mmol), *i*-BuOH (0.65 mmol) in *i*-Pr<sub>2</sub>O (5 mL) at RT. After 24 h, R-9-BBN (1.0 mmol), KO<sup>t</sup>Bu (0.7 mmol), *i*-BuOH (1.0 mmol) were added and adjusted the total volume to 10 mL with *i*-Pr<sub>2</sub>O. [b] Isolated yields.

An increase of electron density in the aromatic ring possibly strengthens its secondary interaction with the catalyst in the enantiodiscriminative step. Indeed, 1-chloro-1-fluoro-2-(*p*-fluorophenyl)ethane (**1h**), bearing a strongly withdrawing fluoride substituent in *para*-position, reacts with considerably lower enantioselectivity (compare entries 6 vs. 20; 12 vs. 19). 1-chloro-1-iodo-2-(*meta*-methoxyphenyl)ethane (**1k**) leads to the tertiary alkane **10u** with lower optical purity (88% *ee*) than the analogous *para*-substituted substrate **1b** (compare entries 6 and 22). It is conceivable that the secondary interactions between the aromatic system and the catalyst are relatively weak in case of the substrate **1k** due to steric congestion

imposed by the *meta*-methoxy substituent. Noteworthy, the high yields and enantioselectivities are obtained in these reactions under the mild conditions and at room temperature.

When *gem*-dihaloalkane **1a** was cross-coupled with hexyl-9-BBN under the standard reaction conditions, the corresponding unsymmetrical secondary alkyl chloride **11** was isolated in 90% yield (Scheme 3a). However, the product was a racemic mixture of enantiomers. A complete absence of *ee* in **11** was surprising, since when such cross-coupling was performed with the corresponding fluoro(halo)alkanes as substrates the resulting secondary alkyl fluorides were isolated in highly enantioenriched form.<sup>12</sup> This difference can be explained by the assumption that the C-Cl bond in secondary alkyl chlorides (e.g., **11**) undergoes fast reversible oxidative addition/reductive elimination process under our cross-coupling conditions which leads to epimerization. This process is apparently impossible in case of stronger C-F bond. Since the first coupling step leads to racemic alkyl chlorides, we can conclude that the second cross-coupling event represents an enantiodiscriminative step under the given developed reaction conditions.<sup>13</sup>

It should be mentioned, that the developed multicomponent transformation is efficient and chemoselective *only* for geminal alkanes possessing a combination of *chloro- and iodo-substituents*. Geminal homo-dihaloalkanes are not suitable substrates for this method. It was previously reported, that CH<sub>2</sub>Cl<sub>2</sub> undergoes dialkylation with BuMgBr via Kumada cross-couplings to result in nonane even if a large excess of dichloromethane is used and reaction is carried out at -20 °C.<sup>14</sup> Similarly, reaction of 1,1-diiodo-2-phenylethane under our coupling conditions mainly resulted in double alkylation already in the first step.



**Scheme 3.** (a) Generation of the racemic intermediate **11** (b) Enantioselective alkyl-alkyl Suzuki cross-coupling of geminal bromo-iodoalkane **12**.

When 1-bromo-1-iodo-2-phenylethane **12** was tested in the representative multicomponent cross-coupling, the asymmetric alkane **3a** was obtained in 42% yield and 83% *ee* (Scheme 3b). However, the main side products were doubly alkylated compounds **13a** and **13b** even if this reaction is

performed at 10 °C. As such, the obtained exclusive chemoselectivity (in the first step) in our method is quite surprising, given that both coupling steps are mediated with the same catalyst under the same reaction conditions.

## Conclusions

In conclusion, we have developed an efficient, versatile and facile method for the highly enantioselective construction of tertiary stereocenters through an unprecedented Suzuki reactions of non-activated racemic geminal chloro(iodo)-alkanes with alkylboranes. We have established the first cross-coupling approach which employs simple and readily available primary alkyl substrates for the direct multicomponent assembly of enantioenriched tertiary stereocenters. The ability to selectively functionalize an alkyl C-I bond in the presence of geminal C-Cl bond is an important factor for the success of this one-pot two-step transformation. Importantly, the same catalyst system is used for both steps of this stereoconvergent transformation, and the selectivity is achieved under mild conditions and at room temperature. In addition, we have disclosed a general and efficient preparation of 1-chloro-1-iodoalkanes, the electrophiles used in the described cross-coupling transformation. These highly functional compounds have become readily available due to developed direct iododecarboxylation of the corresponding ubiquitous  $\alpha$ -chloroalkanoic acids. Collectively, the developed methods open a door to efficient catalytic enantioselective synthesis of alkanes bearing tertiary stereocenters from carboxylic acids just in few steps. Further studies on the reactivity of unactivated geminal dihaloalkanes are underway in our labs.

## Acknowledgements

We acknowledge the financial support from German-Israeli Project Cooperation (DIP) and KAMIN program of the Industry, Trade and Labour Ministry.

## Notes and references

- (a) *Metal-Catalyzed cross-coupling reactions*, (Eds.: A. De Meijere, F. Diederich), Wiley-VCH, 2004. (b) *Transition metals for organic synthesis*, (Eds.: M. Beller, C. Bolm), Wiley-VCH, 2004.
- For pioneering examples, see: (a) A. Devasagayaraj, T. Stüdemann, P. Knochel, *Angew. Chem.* 1995, **107**, 2952; *Angew. Chem. Int. Ed.* 1995, **34**, 2723. (b) T. Ishiyama, A. Shigeru, N. Miyaoura, A. Suzuki, *Chem. Lett.* 1992, **21**, 691. (c) J. K. Kochi, M. Tamura, *J. Am. Chem. Soc.* 1971, **93**, 1483.
- (a) A. C. Frisch, M. Beller, *Angew. Chem.* 2005, **117**, 680; *Angew. Chem. Int. Ed.* 2005, **44**, 674. (b) A. Fürstner, R. Martin, *Chem. Lett.* 2005, **34**, 624. (c) M. R. Netherton, G. C. Fu, *Adv. Synth. Catal.* 2004, **346**, 1525. (d) A. Rudolph, M. Lautens, *Angew. Chem.* 2009, **121**, 2694; *Angew. Chem. Int. Ed.* 2009, **48**, 2656. (e) R. Jana, T. P. Pathak, M. S. Sigman, *Chem. Rev.* 2011, **111**, 1471.
- (a) E. C. Swift, E. R. Jarvo, *Tetrahedron* 2013, **69**, 5799. (b) F. Glorius, *Angew. Chem.* 2008, **120**, 8474; *Angew. Chem. Int. Ed.* 2008, **49**, 8347.
- (a) *Comprehensive asymmetric catalysis*, (Eds.: E. N. Jacobsen, A. Pfaltz, H. Yamamoto), Springer-Verlag, 2004. (b) *Asymmetric catalysis on industrial scale: challenges, approaches and solutions*, (Eds.: H.-U. Blaser, E. Schmidt), Wiley-VCH, 2004.
- For construction of tertiary carbon stereocenters by methods alternative to cross-coupling reactions of non-activated alkyl partners, see: (a) B. M. Trost, M. L. Crawley, *Chem. Rev.* 2003, **103**, 2921. (b) D. Mc. Cartney, P. J. Guiry, *Chem. Soc. Rev.* 2011, **40**, 5122. (c) R. S. Harutyunyan, T. Den. Hartog, K. Geurts, A. J. Minnaard, B. L. Feringa, *Chem. Rev.* 2008, **108**, 2824.
- (a) B. Saito, G. C. Fu, *J. Am. Chem. Soc.* 2008, **130**, 6994. (b) N. A. Owston, G. C. Fu, *J. Am. Chem. Soc.* 2010, **132**, 11908. (c) Z. Lu, A. Wilsily, G. C. Fu, *J. Am. Chem. Soc.* 2011, **133**, 8154. (d) S. L. Zultanski, G. C. Fu, *J. Am. Chem. Soc.* 2011, **133**, 15362. (e) A. Wilsily, F. Tramutola, N. A. Owston, G. C. Fu, *J. Am. Chem. Soc.* 2012, **134**, 5794.
- S. J. Blanksby, G. B. Ellison, *Acc. Chem. Res.* 2003, **36**, 255.
- R. J. Crawford, *J. Org. Chem.* 1983, **48**, 1364.
- (a) R. W. Hoffmann, G. P. Nell, R. Leo, K. Harms, *Chem. Eur. J.* 2000, **6**, 3359. (b) C. De. Lima, M. Julia, J.-N. Verpeaux, *Synlett.* 1992, **2**, 133. (c) A. J. Fry, J. N. Cawse, *J. Org. Chem.* 1967, **32**, 1677. (d) C. Robert, Jr. Neuman, M. L. Rahm, *J. Org. Chem.* 1966, **31**, 1857.
- K. Kulbitski, G. Nisnevich, M. Gandelman, *Adv. Synth. Catal.* 2011, **353**, 1438.
- (a) X. Jiang, S. Sakthivel, K. Kulbitski, G. Nisnevich, M. Gandelman, *J. Am. Chem. Soc.* 2014, **136**, 9548. (b) X. Jiang, M. Gandelman, *J. Am. Chem. Soc.* 2015, **137**, 2542.
- A stereoconvergent mechanism of Ni-catalysed cross-couplings of secondary alkyl halides was previously suggested by Fu et al. For discussion of the origin of enantioselectivity in products, see ref. 7d, e. and 4a.
- Z. Csok, O. Vechorkin, S. B. Harkins, R. Scopelliti, X. Hu, *J. Am. Chem. Soc.* 2008, **130**, 8156.