

# Soft Matter

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

## ARTICLE

## Mechanical signatures of microbial biofilms in micropillar-embedded growth chambers

Cite this: DOI:

10.1039/x0xx00000x

S.C. Chew,<sup>ab</sup> B. Kundukad,<sup>c</sup> W.K. Teh<sup>a</sup>, P. Doyle,<sup>cd</sup> L. Yang<sup>ae</sup>, S. A. Rice<sup>ae</sup> and S. Kjelleberg<sup>ae\*</sup>Received 00th January 2012,  
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

[www.rsc.org/](http://www.rsc.org/)

Biofilms are surface-attached communities of microorganisms embedded in an extracellular matrix and are essential for the cycling of organic matter in natural and engineered environments. They are also the leading cause of many infections, for example those associated with chronic wounds and implanted medical devices. The extracellular matrix is a key biofilm component that determines its architecture and defines its physical properties. Here, we use growth chambers embedded with micropillars to study net mechanical forces (differential pressure) exerted during biofilm formation *in situ*. Pressure from the biofilm is transferred to the micropillars via the extracellular matrix, and reduction of major matrix components decreases the magnitude of micropillar deflections. The spatial arrangement of micropillar deflections caused by pressure differences in the different biofilm strains may potentially be used as mechanical signatures for biofilm characterization. Hence, we submit that micropillar-embedded growth chambers provide insights into the mechanical properties and dynamics of the biofilm and its matrix.

### Introduction

Microorganisms excrete and embed themselves in a matrix of extracellular polymeric substances (EPS)<sup>1</sup> to form surface-attached communities, or biofilms, as their predominant lifestyle in nature.<sup>2</sup> Biofilms play essential roles in sustaining natural environments and human society, including the remediation of natural habitats, treating wastewater and bioleaching.<sup>3</sup> Biofilms can also be problematic in industry, for example, biomass accumulation in pipes disrupts flow and leads to corrosion, contamination of production lines compromises food quality, and biofilm formation on reverse osmosis membranes limits the production of clean water and increases running costs.<sup>4</sup> Biofilms have a significant impact on healthcare, where the US National Institutes of Health (NIH) has estimated that more than 80% of chronic infections and 65% of microbial infections are biofilm-associated (Program Announcement Number PA-03-047, National Institutes of Health).

Biofilm formation is a highly regulated process whereby microorganisms employ physiological

cooperation and spatial organization to increase both their metabolic efficiency and adaptation to changes in their local environment.<sup>5</sup> The result of this process is a multicellular structure that is usually heterogeneous in architecture. Cell growth and death, EPS production and degradation are factors that generate mechanical forces that move biomass to expand or remodel the biofilm. However, research into measuring mechanical forces generated by the biofilm, specifically how they contribute to development of the biofilm structure, is still in its infancy for many biofilms. A recent study has found that *Bacillus subtilis* floating pellicles maintain a low internal stress that drives biofilm spreading after relaxation from confinement and recovery during biofilm ablation.<sup>6</sup> For *B. subtilis* grown at the air-solid surface interface, such as on agar media, biofilm spreading is mediated by the osmotic pressure resulting from the EPS absorbing water rather than the internal pressure caused by colony growth.<sup>7</sup> Lateral mechanical forces are spatially focused by localized cell death in *B. subtilis*, driving macroscopic movement and vertical buckling of the biofilm matrix. This results

in a wrinkled morphology that may increase resistance to liquid wetting and gas penetration.<sup>8,9</sup>

Moreover, the mechanical strength and viscoelasticity of the EPS is expected to impact on external and internal mechanical forces experienced by the biofilm, and thus influence any movement or rearrangement of biomass and affect biofilm structure and morphology. For example, the EPS of *P. aeruginosa* biofilms can shear under flow, creating string-like extensions that eventually result in the development of streamers.<sup>10</sup> *P. aeruginosa* biofilms are also enhanced in surface spreading and streamer formation when loose viscoelastic matrices are produced, and reduced in surface spreading and streamer formation when densely crosslinked elastic matrices are produced.<sup>11</sup> As such, it is important to study the mechanical forces or pressure that drive the spreading and self-organization of these microbial populations, which can also provide a conceptual framework for other microbial systems.

Micropillar arrays were developed for the study and measurement of cellular/subcellular traction forces that eukaryotic cells exert upon adhesion to substrates. In the standard micropillar setup and model (Fig. 1a), the cell is required to adhere to, and therefore leverage on, more than one micropillar for attachment and traction. The micropillar array has been extensively used for studies of cell spreading and single cell migration as well as movement of monolayers of mammalian cells. For eukaryotic cells, the cellular/subcellular traction forces are generated by the cytoskeletal motor protein myosin II, which causes rearrangement of actin filaments when coupled to adhesion sites of the cell.<sup>12,13</sup> In contrast, apart from the measurement of cooperative retraction forces by bundled type IV pili in *Neisseria gonorrhoeae*, for which large forces of 200 pN to 1 nN have been recorded,<sup>14</sup> micropillars have not been used in force studies of microbial cells. Unlike mammalian cells, microbial cells are considered rigid, with the Young's modulus (E) of live *Escherichia coli* being recorded as  $1.9 \pm 0.9$  to  $3.0 \pm 0.6$  MPa, and  $6.1 \pm 1.5$  MPa for dead *E. coli*.<sup>15</sup> As such, they have well-defined shapes that do not deform easily and are unlikely to generate high traction forces within the cell body. In addition, micropillar dimensions are usually in the order of 1-3  $\mu\text{m}$  in diameter, and require sufficient spacing intervals for deflections to be measured. Thus, most microbial cells are too small to attach to and interact with the multiple micropillars required to generate force measurements. However, within the biofilm, cells are collectively held together by an extracellular matrix, of which intercellular forces between cells have been reported to range from 6.5-6.8 nN based

upon atomic force microscopy.<sup>16</sup> The E for biofilms vary widely, with reported values of  $37.82 \pm 5.87$  kPa<sup>17</sup> and  $25.0 \pm 2.5$  kPa<sup>8</sup> for early stage laboratory biofilms grown on agar, and 200 to 9000 kPa<sup>18</sup> for an environmental biofilm.

The multicellular biofilm can spread over a large area and grow to hundreds of micrometers in thickness. The extracellular matrix secreted by the cells within the biofilm is expected to be the major component that binds to multiple micropillars while holding the cells together (Fig. 1b). We hypothesized that once the biofilm was formed, forces generated by multicellular microbial behaviour involving cell growth, death, motility and differentiation would create internal mechanical forces or pressure. While a uniform pressure generates equal lateral forces in all directions and does not deflect the micropillars, the heterogeneous growth and development of the biofilm may produce net mechanical forces or differential pressure sufficient to deflect the micropillars.

This work describes the application of micropillar-embedded growth chambers as a tool for the mechanical characterization of bacterial biofilms. We investigated biofilms formed by several medical and environmental model bacterial species, and mutants of these defective in the production of key matrix components. This includes *E. coli* strains causing urinary tract infections,<sup>19</sup> as well as non-virulent strains used in the industrial production of recombinant therapeutics. *P. aeruginosa* is a common opportunistic pathogen. The mucoid type, in which the bacterium overexpresses alginate along with exopolysaccharides Pel and Psl in its matrix, is often isolated from cystic fibrosis patients where it is a major biofilm forming bacterial species.<sup>20</sup> *Staphylococcus aureus* and *Staphylococcus epidermidis* often cause biofilm-associated wound infections and device-related infections.<sup>21, 22</sup> The environmental bacterium *Shewanella oneidensis* reduces heavy metals and many complex xenobiotics.<sup>23</sup> We show here that micropillars can measure forces at the substratum related to biofilm growth and matrix mechanical properties. The various biofilms generate deflection patterns in the micropillar array that are reflective of their growth dynamics and EPS mechanical properties.

## Experimental

### Biofilm cultivation

The bacterial strains used in this study include *E. coli* SAR18 F<sup>+</sup>; *P. aeruginosa* PAO1 $\Delta\text{muca}$ ,

PAO1 $\Delta$ *mucA* $\Delta$ *pslBCD* and PAO1 $\Delta$ *mucA* $\Delta$ *pelA*; *S. aureus* 15981; *S. epidermidis* 1457 and 1457 $\Delta$ *atlE*; *S. oneidensis* MR-1 (Table 1). Overnight cultures of bacterial strains were grown at their optimal temperatures and growth medium to an OD<sub>600</sub> = 2.0. For *E. coli* SAR18 F<sup>+</sup>, *P. aeruginosa*  $\Delta$ *mucA*, PAO1  $\Delta$ *mucA* $\Delta$ *pslBCD* and PAO1  $\Delta$ *mucA* $\Delta$ *pelA*, growth was at 37°C in Luria Broth (LB) medium. For *S. aureus* 15981, *S. epidermidis* 1457 and *S. epidermidis* 1457 $\Delta$ *atlE*, growth was at 37°C in Tryptic Soy Broth (TSB) medium. *S. oneidensis* MR-1 was grown at 30°C in LB medium. Subsequently for biofilm formation, 10  $\mu$ L of overnight cultures were added to the wells of the growth chambers with 190  $\mu$ L of medium to a final OD<sub>600</sub> = 0.1 and incubated for 24 hours under static conditions at the optimal temperature of 37°C or 30°C for each organism as described above. The growth curves of the bacterial strains using the above cultivation conditions are provided in Fig. S1.

Table 1. List of strains

Strain	Description	Ref.
SAR18 F <sup>+</sup>	<i>E. coli</i> CSH26 wild-type carrying transfer constitutive IncF plasmids that promotes cell-cell adhesion and biofilm formation mediated by F pili	24
PAO1 $\Delta$ <i>mucA</i>	Mucoid <i>P. aeruginosa</i> PAO1 expressing alginate, Pel and Psl exopolysaccharides	25
PAO1 $\Delta$ <i>mucA</i> $\Delta$ <i>pelA</i>	Psl mutant of mucoid PAO1, expressing alginate and Pel exopolysaccharides only	26
PAO1 $\Delta$ <i>mucA</i> $\Delta$ <i>pslBCD</i>	Pel mutant of mucoid PAO1, expressing alginate and Psl exopolysaccharides only	26
15981	<i>S. aureus</i> wild-type	26
1457	<i>S. epidermidis</i> wild-type	27
1457 $\Delta$ <i>atlE</i>	Autolysin ( <i>atlE</i> ) mutant of <i>S. epidermidis</i> 1457, unable to release extracellular DNA (eDNA) into matrix	27
MR-1	<i>S. oneidensis</i> wild-type	23

### Micropillar arrays

PDMS (polydimethylsiloxane) micropillar arrays were moulded from silicon masters made with conventional high-resolution photolithography and deep reactive ion-etching techniques.<sup>28</sup> The micropillars had a height (H) of 3  $\mu$ m, diameter (D)

of 1.5  $\mu$ m and center-to-center spacing of 4  $\mu$ m. The Young's modulus (E) of the micropillars was 1.0  $\pm$  0.3 MPa as verified using atomic force microscopy. The PDMS micropillar arrays were established at the bottom of growth chambers to form 300  $\mu$ L wells, 9.4 mm in width and 10.7 mm in length. The growth chambers were placed in 100% ethanol and sonicated to ensure micropillars remained in upright position and to avoid collapse of the micropillars during introduction of high surface tension fluids. The ethanol solution was replaced with sterile water, and then LB or TSB medium to a final volume of 190  $\mu$ L each well.

### Microscopy and imaging

For the imaging of micropillar deflections, bright-field Z-stack micrographs of the micropillars after allowing for 24 h of bacterial attachment and biofilm growth were captured (ZEISS Axio Imager M1). Biofilms were stained with Congo Red (Sigma-Aldrich, Singapore) to improve contrast between the biofilm and micropillars. The positions of the micropillar tops and bottoms were located by their centroid positions using a Gaussian or Mexican Hat filter (background subtraction) in IMARIS software (Bitplane, Zürich). The deflection vector  $r$  is given by the difference in positions:

$$r = \sqrt{[(x_t - x_b)^2 + (y_t - y_b)^2]} \quad (1)$$

where  $x_t$  and  $y_t$  are x and y positions of the centroid of the micropillar top respectively, and  $x_b$  and  $y_b$  are x and y positions of the centroid of the micropillar base respectively.

For the biomass quantification of biofilms formed by the respective strains after 24 h of growth, confocal images of the biofilm were captured (Zeiss LSM780 confocal scanning laser microscope) and analyzed by COMSTAT (www.comstat.dk).<sup>29, 30</sup> Biofilms were cultivated in chambers without micropillars. The Gram-negative *E. coli*, *P. aeruginosa* and *S. oneidensis* were tagged with the green fluorescent protein (Gfp) for visualization and quantification of live biomass, and stained with propidium iodide (PI) for visualization and quantification of dead biomass. The Gram-positive *S. aureus* and *S. epidermidis* were stained with SYTO9 and PI for visualization and quantification of live and dead biomass.

### Force calculations

In the case where the mammalian cell body is located on top of the micropillars, the micropillar is

modelled as a cantilevered beam deflected by a horizontal traction force applied on the micropillar top,  $f$  (see Fig. 1a). For small deformations, the force required to deflect the micropillar may be approximated by Hooke's law using the Euler-Bernoulli beam theory and the following equation:<sup>31</sup>

$$f = \frac{3EIr}{H^3} \quad (2)$$

where  $I$  = moment of inertia. The moment of inertia for a solid beam with a circular cross-section is given by:

$$I = \frac{\pi D^4}{64} \quad (3)$$

Substituting for  $I$  gives:

$$f = \frac{3\pi ED^4 r}{64H^3} \quad \text{or} \quad f = Kr \quad (4)$$

where  $K$  is the spring constant of the micropillar and is given by

$$K = \frac{3\pi ED^4}{64H^3} \quad (5)$$

In the case of the biofilm, the microbial cell does not extend over more than one micropillar. Instead, biofilms form around the micropillars (Fig. 1b-c). Lateral growth and motion from the biofilm in between the micropillars is expected to apply tension and compression along the entire height of the pillar. A previous study found that biofilms are most stiff and mechanically homogeneous within 10  $\mu\text{m}$  in the  $z$ -direction from the substratum.<sup>32</sup> In such cases, and for short micropillars such as those used here, the micropillar may be modelled as a cantilevered beam with the biofilm applying a uniform load across its height with total force,  $f$  (Fig. 1b). The total force applied to the micropillar is then related to the deflection at the micropillar top according to the equation:<sup>33</sup>

$$f = \frac{8EIr}{H^3} \quad (6)$$

The force is larger by a constant of 8/3, as more force is required to achieve the same amount of deflection when uniformly applied along the micropillar as opposed to being focused at the free end. The spring constant  $K$  for a uniform force applied along the micropillar height is then:

$$K = \frac{8EI}{H^3} \quad \text{or} \quad K = \frac{\pi ED^4}{8H^3} \quad (7)$$

The differential pressure ( $p$ ) within the biofilm is calculated by dividing the force by the entire cross-section of the micropillar.

$$p = \frac{f}{D \times H} \quad (8)$$




Fig. 1. Micropillar deflection caused by mammalian cell versus biofilm. (a) For mammalian cells, which attach to the top of the micropillar, traction forces are generated by actin re-arrangement that deflect the micropillar. Thus, micropillar deflection is modelled as a beam bending with a lateral force applied at one end. (b) The biofilm grows over and between the micropillars, applying pressure along the entire height of the micropillar. Micropillar deflection is thus modelled as beam bending with uniform load applied along its height. (c) Left: control micropillar array without biofilm. Right: micropillar array covered with biofilm. The area size is  $52.4 \times 52.4 \mu\text{m}$ .

## Results and Discussion

### Distribution of pressure differences in various biofilms

The bacterial strains attached to the micropillars at the substratum of the growth chambers and formed biofilms that deflected the micropillars over 24 h. The deflection magnitudes of 120 micropillars within a  $50 \times 50 \mu\text{m}$  area of substratum from at least three different experimental replicates for each strain were used to calculate the differential pressure applied to the micropillar over 24 h. A histogram was compiled to compare the distribution of pressure differences (Fig. 2), and the average, maximum and standard deviation of differential pressure were calculated for biofilms of each wild-type and mutant strain (Table 2). For mucoid  $P$ .

*aeruginosa* (PAO1 $\Delta$ *mucA*) and *S. epidermidis*, most of the differences in pressure after 24 h of biofilm growth fell within the 4-8 kPa range. This range accounted for  $54.8 \pm 9.2\%$  and  $39.0 \pm 7.4\%$ , of the pressure differences in *P. aeruginosa* and *S. epidermidis* respectively. For *E. coli* and *S. aureus* respectively, most of the pressure differences fell within the 8-12 kPa range, accounting for  $46.4 \pm 5.4\%$  and  $39.6 \pm 3.8\%$  respectively. For *S. oneidensis*, a majority of  $95 \pm 6\%$  of pressure differences were below 4 kPa. *E. coli* had the highest average differential pressure at  $8.5 \pm 0.1$  kPa. This was then followed by *S. aureus* at  $8.0 \pm 0.6$  kPa, *S. epidermidis* at  $7.9 \pm 0.6$  kPa, *P. aeruginosa* at  $7.2 \pm 0.2$  kPa and *S. oneidensis* at  $1.8 \pm 0.6$  kPa. The average differential pressure may be reflective of the heterogeneity within the biofilm of the particular species. In this study *E. coli*, *S. aureus* and *S. epidermidis* formed a heterogeneous biofilm with many microcolonies and channels. Mucooid *P. aeruginosa* also formed heterogeneous biofilms, but with smaller microcolonies and more undifferentiated areas. *S. oneidensis* did not form differentiated biofilms under our growth conditions.




Fig. 2. Distribution of differential pressures within biofilms formed by mucooid *P. aeruginosa*, and wild-type *E. coli*, *S. epidermidis* and *S. oneidensis* strains. Frequency of the differential pressure within 0-4 kPa, 4-8 kPa, 8-12 kPa, 12-16 kPa, 16-20 kPa and >20 kPa was indicated, respectively. At least three experimental replicates were used.

While *E. coli* had the highest average differential pressure, *S. aureus* and *S. epidermidis* were able to achieve differential pressures greater than 20 kPa. However, this accounted for less than 1% of the differential pressure. *S. epidermidis* had the highest maximum differential pressure and standard deviation, followed by *S. aureus*, *E. coli*, *P. aeruginosa* and *S. oneidensis* (Table 2).

Table 2. Average, maximum and standard deviation of differential pressure detected in wild-type and mutant strains.

Strain	Differential pressure (kPa)*		
	Average	Maximum	S.D.^
<i>E. coli</i> SAR18 F+	$8.5 \pm 0.1$	$16.1 \pm 2.7$	3.1
<i>P. aeruginosa</i> PAO1 $\Delta$ <i>mucA</i>	$7.2 \pm 0.2$	$12.0 \pm 1.0$	2.3
<i>P. aeruginosa</i> PAO1 $\Delta$ <i>mucA</i> $\Delta$ <i>pelA</i>	$3.1 \pm 0.7$	$6.5 \pm 1.0$	1.2
<i>P. aeruginosa</i> PAO1 $\Delta$ <i>mucA</i> $\Delta$ <i>pslBCD</i>	$6.2 \pm 0.3$	$10.3 \pm 1.5$	1.8
<i>S. aureus</i> 15981	$8.0 \pm 0.6$	$18.7 \pm 2.2$	3.2
<i>S. epidermidis</i> 1457	$7.9 \pm 0.6$	$20.0 \pm 3.1$	3.7
<i>S. epidermidis</i> 1457 $\Delta$ <i>atlE</i>	$5.3 \pm 0.9$	$20.5 \pm 5.0$	3.7
<i>S. oneidensis</i> MR-1	$1.8 \pm 0.6$	$5.3 \pm 1.6$	1.0

\*Calculated from 120 micropillars within a  $50 \times 50 \mu\text{m}$  area. At least three biological replicates were used.

^Standard deviation.




Fig. 3. Live and dead biomass of biofilms formed by various strains in 24 h static cultures.

Differential pressure arises from the pressure of a growing biofilm biomass exerting itself onto the micropillars. To examine the correlation between differential pressure and the total biofilm biomass generated by the strains, we measured the biomass of static biofilm cultures for each strain (Fig. 3). The live and dead measurements were combined to determine the total biomass. Among wild-type strains and mucooid *P. aeruginosa*, *S. aureus* and *S. epidermidis* biofilms had the highest total biomass with  $5.8 \pm 0.5 \mu\text{m}^3 \mu\text{m}^{-2}$  and  $5.4 \pm 0.1 \mu\text{m}^3 \mu\text{m}^{-2}$ , respectively. *E. coli* displayed the second highest total biomass at  $5.1 \pm 1 \mu\text{m}^3 \mu\text{m}^{-2}$ . Biofilms of

PAO1 $\Delta$ *mucA*, *S. aureus* and *S. epidermidis* had significant proportions of dead biomass at  $62 \pm 7\%$ ,  $49 \pm 2\%$  and  $43 \pm 8\%$  respectively. *E. coli* biofilms displayed a small fraction,  $10 \pm 5\%$ , of dead biomass. This suggests that cell death was not a major factor for its higher average differential pressure, as would be predicted based on previous data for *B. subtilis*.<sup>8</sup> PAO1 $\Delta$ *mucA* and *S. oneidensis* generated the second lowest and lowest total biomass at  $3.0 \pm 0.1 \mu\text{m}^3 \mu\text{m}^{-2}$  and  $1.7 \pm 0.3 \mu\text{m}^3 \mu\text{m}^{-2}$ , respectively, and may explain their comparatively lower average differential pressures. These results indicate that although the generation of total biomass could partly account for the differential pressure, other species-specific biofilm matrix factors might also be involved.

### Matrix components of the biofilm strongly influence differential pressure generated by the biofilm

EPS mechanical properties were hypothesized to influence the deflection of the micropillars. To examine whether micropillars can be used to investigate the impact of specific EPS components on differential pressure in the biofilm, mucoid *P. aeruginosa*, *S. epidermidis* strains, and their respective matrix mutant strains were assessed (Table 2 and Fig. 4). *P. aeruginosa*  $\Delta$ *mucA $\Delta$ *pelA* and *P. aeruginosa*  $\Delta$ *mucA $\Delta$ *pslBCD* mutants are unable to synthesize the major *P. aeruginosa* matrix components Pel and Psl exopolysaccharides, respectively.<sup>26</sup> Both *S. aureus* and *S. epidermidis* release a large amount of eDNA into their matrix, which contributes significantly to biofilm formation.<sup>27, 34</sup> *S. epidermidis*  $\Delta$ *atlE* mutants are unable to release eDNA into the biofilm matrix and hence are expected to produce weaker biofilms.<sup>27</sup>**

The average and maximum differential pressure measured for the *P. aeruginosa*  $\Delta$ *mucA $\Delta$ *pelA* biofilms were  $3.1 \pm 0.7$  and  $6.5 \pm 1.0$  kPa respectively (Table 2, Fig. 4). The average and maximum differential pressure measured for the *P. aeruginosa*  $\Delta$ *mucA $\Delta$ *pslBCD* biofilms were  $6.2 \pm 0.3$  and  $10.3 \pm 1.5$  kPa respectively (Table 2, Fig. 4). Thus, the loss of the exopolysaccharide Pel significantly impaired the ability of the biofilm to deflect the micropillars, whereas loss of the exopolysaccharide Psl reduced differential pressure to a lesser extent compared to the *P. aeruginosa*  $\Delta$ *mucA* parent strain. In the present study, the live and total biomasses of *P. aeruginosa*  $\Delta$ *mucA $\Delta$ *pslBCD*, at  $3.5 \pm 0.6 \mu\text{m}^3 \mu\text{m}^{-2}$  and  $6.0 \pm 0.1 \mu\text{m}^3 \mu\text{m}^{-2}$  respectively, was much greater than the *P. aeruginosa*  $\Delta$ *mucA $\Delta$ *pelA* strain, at  $0.8 \pm 0.1 \mu\text{m}^3 \mu\text{m}^{-2}$  and  $3.9 \pm 0.1 \mu\text{m}^3 \mu\text{m}^{-2}$ , respectively, and****

the *P. aeruginosa*  $\Delta$ *mucA* strain, at  $1.1 \pm 0.2 \mu\text{m}^3 \mu\text{m}^{-2}$  and  $3.0 \pm 0.1 \mu\text{m}^3 \mu\text{m}^{-2}$ , respectively (Fig. 3). This finding agrees with the report that the Pel polysaccharide is important for lateral growth and spreading,<sup>11</sup> which would increase the overall pressure in the biofilm. In contrast, the Psl exopolysaccharide increased the cross-linking density and elasticity of the matrix, resulting in increased mechanical stiffness that restricted lateral growth and spreading. Biofilms containing the Pel polysaccharide were also shown to vary in rheology in different locations and over time, whereas biofilms expressing only Psl were spatially and temporally homogeneous in rheology.<sup>11</sup> This may also explain the relative larger contribution of Pel to differential pressure compared to Psl. The expression of both Pel and Psl in *P. aeruginosa*  $\Delta$ *mucA* biofilms had a synergistic effect on increasing pressure differences (Fig. 4), which may act to improve overall biofilm dynamics and formation.

eDNA is another well known matrix component crucial for maintaining the biofilm mechanic properties. The *S. epidermidis* eDNA deficient  $\Delta$ *atlE* mutant biofilms showed a decreased ability to deflect the micropillars, with an average differential pressure of  $5.2 \pm 0.9$  kPa compared to  $7.9 \pm 0.6$  kPa for the parental strain. Despite this, a high differential pressure could still be achieved in the *S. epidermidis*  $\Delta$ *atlE* biofilm. Indeed, the maximum differential pressure was  $20.5 \pm 5.0$  kPa, which suggests that other major biofilm matrix components (e.g. polysaccharide intercellular adhesin) played a more important role here in generating pressure differences.




Fig. 4. Comparison of quartiles, maximum and minimum of differential pressure data for mucoid *P. aeruginosa*, wild-type *S. epidermidis* and their respective EPS mutant strains.




Fig. 5. The top image of each panel is a  $50 \times 50 \mu\text{m}$  2D sectional bright field image of the micropillars deflected by (a) no biofilm or biofilms formed by (b) *E. coli*  $F^+$  SAR18 (c) *P. aeruginosa* PAO1 $\Delta\text{mucA}$  (d) *S. aureus* 15981 (e) *S. epidermidis* 1457 and (f) *S. oneidensis* MR-1. The bottom image for each panel is the deflection vector map. The vectors are depicted by black arrows, which have been enlarged by a factor of 3.5 to increase clarity. The contour of the biofilm is drawn in red, and the blue lines approximate the slope of major edges of the biofilm contour. Images are representative of three experimental replicates.



## Characteristic deflection patterns of bacterial strains are caused by biofilm heterogeneity and microcolony formation

Deflections of 120 micropillars within a  $50 \times 50 \mu\text{m}$  area were plotted as vectors (black arrows) on a 2D Cartesian plane for the mucoid *P. aeruginosa* and wild-type strains (Fig. 5). Micropillars in the control chamber remained undeflected and generated null vectors (Fig. 5a). *E. coli* SAR18 F<sup>+</sup> (Fig. 5b), *P. aeruginosa* PAO1 $\Delta\text{mucA}$  (Fig. 5c), *S. aureus* 15891 (Fig. 5d) and *S. epidermidis* 1457 (Fig. 5e) developed heterogeneous biofilms that exhibited extensive cell clustering and microcolony formation. The formation of such microcolonies deflected the micropillars in various directions and disrupted the regular micropillar positioning. As a measure of regularity in the deflections and to examine how similar in direction the vectors were across the different biofilms, the cosine similarity of each vector to its right horizontal and lower vertical neighbour was calculated according to the equation:

$$\text{similarity} = \cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|} \quad (9)$$

where  $\theta$  of  $0^\circ$  gives similarity of 1,  $\theta$  of  $90^\circ$  give similarity of 0 and  $\theta$  of  $180^\circ$  gives similarity of -1.

*E. coli*, mucoid *P. aeruginosa*, *S. aureus* and *S. epidermidis* 1457 biofilms displayed low average similarities of  $0.12 \pm 0.13$ ,  $0.11 \pm 0.06$ ,  $0.17 \pm 0.10$  and  $0.22 \pm 0.07$ , respectively. The average angles between the neighbouring vectors were nearly orthogonal to each other at  $83.2 \pm 8.0^\circ$ ,  $83.8 \pm 3.8^\circ$ ,  $80.0 \pm 5.8^\circ$  and  $77.3 \pm 4.0^\circ$  respectively. In contrast to the other species, *S. oneidensis* (Fig. 5f) formed a thin and homogenous biofilm without any microcolonies and as such, the distance between, and regularity of, the micropillars were largely maintained. On average, the deflection vectors showed a high similarity of  $0.79 \pm 0.05$  and angle of  $37.8 \pm 5.1^\circ$  with neighbouring vectors. Thus, the similarity between neighbouring vectors can be a measure of microcolony formation, with a high similarity indicating the presence of very few or no microcolonies.

A red contour, as approximated from a Z-stack of bright-field images, was drawn along the high cell density areas and microcolonies on the deflection vector map (Fig. 5). For *E. coli* and mucoid *P. aeruginosa* biofilms, micropillars were deflected within the microcolonies as well as for the flat, undifferentiated areas of the biofilm. For *S. aureus* and *S. epidermidis* biofilms, the lengths of the deflection vectors varied widely along the contour of the biofilm. This could be due to the

different cluster expansion dynamics of the bacteria. For example, *Staphylococcus* spp. are non-motile and naturally arrange in clusters reflective of their inability to separate after division.<sup>36</sup> Biofilm growth is then primarily based on clonal expansion that would exert outward pressure on the micropillars at the growing edges, but less at the focal points of growth. In contrast, *E. coli* and *P. aeruginosa* also utilize surface motility to facilitate the spreading of a developing biofilm and differentiation of flat layer of cells.<sup>37-39</sup>

To further examine whether distinct deflection patterns generated by different species during biofilm growth could be detected, the alignment of deflection vectors at the edges of the biofilm were compared to the slope of the contour (blue line) using the cosine similarity (Fig 5). In this case, deflection vectors that point in opposite directions ( $180^\circ$  with respect to each other) but lie flat along the slope are considered parallel to the slope. Thus, cosine similarity values are absolute, with a  $\theta$  value of  $180^\circ$  also giving a similarity of 1. In order of descending similarity and wider angles, the similarities and average angle between the deflection vector and contour were  $0.69 \pm 0.04$  and  $46.6 \pm 3.5^\circ$  for *P. aeruginosa*,  $0.65 \pm 0.01$  and  $49.8 \pm 0.9^\circ$  for *E. coli*,  $0.59 \pm 0.03$  and  $53.9 \pm 2.1^\circ$  for *S. aureus*, and  $0.57 \pm 0.05$  and  $55.1 \pm 3.7^\circ$  for *S. epidermidis* respectively. Thus, deflection vectors for the *Staphylococcus* spp. biofilms were less aligned to the biofilm contour as compared to that of *E. coli* and *P. aeruginosa* biofilms, and this pattern may reflect different mechanisms of cluster expansion, i.e. clonal growth vs. surface motility.

## Conclusions

There are limited tools available to study biofilm micromechanics and dynamics, with microscopy being one of the primary instruments of investigation. Microscopy is often combined with specialist techniques such as particle<sup>11</sup> and cell-tracking,<sup>40</sup> and usually requires high-end microscopes<sup>41</sup> to probe biofilm dynamics. These techniques are labour intensive and time consuming. In addition, cell-tracking within mature biofilms is difficult because of the inaccuracy in resolving overlapping cells and long experimental times required due to slow cellular dynamics. Other micromechanical techniques used to investigate the physical properties of biological samples usually utilize equipment not available in biological laboratories. Examples of these include atomic force microscopy, optical light or laser traps, micropipette aspiration and magnetic tweezers. In this study, we present micropillar-embedded growth

chambers as an experimental tool that can be used with standard, readily available, microscopes.

Bacterial biofilms caused distinct deflection patterns of the micropillar arrays, which can be used to calculate differential pressure in biofilm, as shown here. Biomasses of the growing biofilms were physically constrained by, and exerted pressure onto the micropillars. The heterogeneous development and differentiation of the biofilms resulted in differential pressure that was responsible for deflecting the micropillars. *E. coli* biofilms had the highest average differential pressure, and *S. oneidensis* biofilms the lowest. Although *S. aureus* and *S. epidermidis* biofilms had similar average differential pressures, the distribution and range of differential pressures experienced within their biofilms were different.

The generation of pressure differences that deflected the micropillar was dependent on EPS characteristics. In general, despite increases in biomass observed in some mutants, removal of any of the major EPS components strongly reduced the pressure differences within the biofilm, as measured here. EPS components that reduce stiffness and increase biofilm malleability are expected to enhance differential pressure as they increase microbial dynamics, biofilm motion and lateral growth. In *P. aeruginosa*, the Psl polysaccharide has been shown to increase biofilm stiffness, whereas Pel is able to reduce biofilm stiffness and enhance overall malleability.<sup>11</sup> In *B. subtilis* biofilms, mechanical forces are focused on areas of weakest matrix stiffness, causing differential pressure and buckling of the biofilm layer.<sup>8</sup> EPS components that result in increased variation of biofilm mechanical properties are also expected to increase differential pressure. *P. aeruginosa* biofilms expressing Pel vary more in rheology spatially and temporally as compared to Psl.<sup>11</sup>

The formation of high cell density areas and microcolonies in the strains tested was the major factor in disrupting the array and the creation of deflection patterns. Thus, micropillars may be sensitive to the different mechanisms that drive microcolony and cluster expansion for different bacteria. Micropillars may then be coupled with the use of specific mutants to probe, among other things, how EPS components or surface motility that affect biofilm dynamics and microcolony formation translate into pressure differences to affect biofilm morphology. It has been shown that pressure from biofilm expansion in confined spaces is responsible for the wrinkled structure of pellicles at the air-liquid interface.<sup>40</sup> Differences in physical pressure generated by different bacterial strains may also have implications for the spatial organization of

bacteria in multi-species biofilm communities. Such studies can help in our understanding of the mechanics of biofilm populations or communities.<sup>42</sup>

Finally, surfaces with high aspect ratio structures (tall in height and short in width), such as micropillars, have been employed in biofilm research and industry for their anti-biofouling properties.<sup>43</sup> Thus, the methodology described here to measure differential pressure in biofilms can be coupled with such surfaces to produce a 'smart' material that detects when biofilms have eventually formed, thus having real application in cleaning and maintenance of equipment of various industries.

### Acknowledgements

We acknowledge financial support from the Singapore Centre for Environmental Life Sciences Engineering (SCELSE), whose research is funded by the National Research Foundation Singapore, Ministry of Education Singapore, Nanyang Technological University and National University of Singapore, under its Research Centre of Excellence program. We also acknowledge Singapore MIT Alliance for Research and Technology's research program in BioSystems and Micromechanics supported by the National Research Foundation Singapore.

### Notes and references

- <sup>a</sup> Singapore Centre for Environmental Life Sciences Engineering (SCELSE), Nanyang Technological University, Singapore  
<sup>b</sup> Interdisciplinary Graduate School, Nanyang Technological University, Singapore  
<sup>c</sup> BioSystems and Micromechanics IRG, Singapore-MIT Alliance for Research and Technology, National University of Singapore, Singapore  
<sup>d</sup> Department of Chemical Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA  
<sup>e</sup> School of Biological Sciences, Nanyang Technological University, Singapore  
<sup>f</sup> Centre for Marine Bio-Innovation and School of Biotechnology and Biomolecular Sciences, University of New South Wales, Australia

1. H. C. Flemming and J. Wingender, *Nat. Rev. Microbiol.*, 2010, **8**, 623-633.
2. J. W. Costerton, Z. Lewandowski, D. E. Caldwell, D. R. Korber and H. M. Lappin-Scott, *Annu. Rev. Microbiol.*, 1995, **49**, 711-745.
3. R. Singh, D. Paul and R. K. Jain, *Trends Microbiol.*, 2006, **14**, 389-397.
4. B. Q. Liao, D. M. Bagley, H. E. Kraemer, G. G. Leppard and S. N. Liss, *Water Environ. Res.*, 2004, **76**, 425-436.
5. P. S. Stewart and M. J. Franklin, *Nat. Rev. Microbiol.*, 2008, **6**, 199-210.

6. C. Douarche, J. M. Allain and E. Raspaud, *Biophys. J.*, 2015, **109**, 2195-2202.
7. A. Seminara, T. E. Angelini, J. N. Wilking, H. Vlamakis, S. Ebrahim, R. Kolter, D. A. Weitz and M. P. Brenner, *Proc. Natl. Acad. Sci. USA*, 2012.
8. M. Asally, M. Kittisopikul, P. Rue, Y. J. Du, Z. X. Hu, T. Cagatay, A. B. Robinson, H. B. Lu, J. Garcia-Ojalvo and G. M. Suel, *Proc. Natl. Acad. Sci. USA*, 2012, **109**, 18891-18896.
9. A. K. Epstein, B. Pokroy, A. Seminara and J. Aizenberg, *Proc. Natl. Acad. Sci. USA*, 2011, **108**, 995-1000.
10. K. Drescher, Y. Shen, B. L. Bassler and H. A. Stone, *Proc. Natl. Acad. Sci. USA*, 2013, **110**, 4345-4350.
11. S. C. Chew, B. Kundukad, T. Seviour, J. R. C. van der Maarel, L. Yang, S. A. Rice, P. Doyle and S. Kjelleberg, *mBio*, 2014, **5**, e01536-14.
12. O. du Roure, A. Saez, A. Buguin, R. H. Austin, P. Chavier, P. Silberzan and B. Ladoux, *Proc. Natl. Acad. Sci. USA*, 2005, **102**, 2390-2395.
13. M. T. Yang, J. P. Fu, Y. K. Wang, R. A. Desai and C. S. Chen, *Nat. Protoc.*, 2011, **6**, 187-213.
14. N. Biais, B. Ladoux, D. Higashi, M. So and M. Sheetz, *PLoS Biol.*, 2008, **6**, e87.
15. A. Cerf, J. C. Cau, C. Vieu and E. Dague, *Langmuir*, 2009, **25**, 5731-5736.
16. H. H. P. Fang, K. Y. Chan and L. C. Xu, *J. Microbiol. Meth.*, 2000, **40**, 89-97.
17. S. Kesel, S. Grumbein, I. Gumperlein, M. Tallawi, A. K. Marel, O. Lieleg and M. Opitz, *Appl. Environ. Microbiol.*, 2016, **82**, 2424-2432.
18. Y. Abe, P. Polyakov, S. Skali-Lami and G. Francius, *Biofouling*, 2011, **27**, 739-750.
19. S. M. Soto, A. Smithson, J. P. Horcajada, J. A. Martinez, J. P. Mensa and J. Vila, *Clin. Microbiol. Infect.*, 2006, **12**, 1034-1036.
20. J. R. Govan and V. Deretic, *Microbiol. Rev.*, 1996, **60**, 539-574.
21. C. F. Schierle, M. De la Garza, T. A. Mustoe and R. D. Galiano, *Wound Repair Regen.*, 2009, **17**, 354-359.
22. M. T. McCann, B. F. Gilmore and S. P. Gorman, *J. Pharm. Pharmacol.*, 2008, **60**, 1551-1571.
23. Y. A. Gorby, S. Yanina, J. S. McLean, K. M. Rosso, D. Moyles, A. Dohnalkova, T. J. Beveridge, I. S. Chang, B. H. Kim, K. S. Kim, D. E. Culley, S. B. Reed, M. F. Romine, D. A. Saffarini, E. A. Hill, L. Shi, D. A. Elias, D. W. Kennedy, G. Pinchuk, K. Watanabe, S. Ishii, B. Logan, K. H. Nealson and J. K. Fredrickson, *Proc. Natl. Acad. Sci. USA*, 2006, **103**, 11358-11363.
24. A. Folkesson, J. A. Haagensen, C. Zampaloni, C. Sternberg and S. Molin, *PLoS One*, 2008, **3**, e1891.
25. H. Wu, B. Lee, L. Yang, H. Z. Wang, M. Givskov, S. Molin, N. Hoiby and Z. J. Song, *FEMS Immunol. Med. Microbiol.*, 2011, **62**, 49-56.
26. L. Yang, Y. Liu, T. Markussen, N. Hoiby, T. Tolker-Nielsen and S. Molin, *FEMS Immunol. Med. Microbiol.*, 2011, **62**, 339-347.
27. Z. Qin, Y. Ou, L. Yang, Y. Zhu, T. Tolker-Nielsen, S. Molin and D. Qu, *Microbiol.*, 2007, **153**, 2083-2092.
28. J. P. Fu, Y. K. Wang, M. T. Yang, R. A. Desai, X. A. Yu, Z. J. Liu and C. S. Chen, *Nat. Methods*, 2010, **7**, 733-U795.
29. A. Heydorn, A. T. Nielsen, M. Hentzer, C. Sternberg, M. Givskov, B. K. Ersboll and S. Molin, *Microbiol.*, 2000, **146**, 2395-2407.
30. M. Vorregaard, Technical University of Denmark, 2008.
31. J. M. Gere and S. P. Timoshenko, *Mechanics of materials*, PWS Publishing Co., Boston, 4th edn., 1997.
32. O. Galy, P. Latour-Lambert, K. Zrelli, J. M. Ghigo, C. Beloin and N. Henry, *Biophys. J.*, 2012, **103**, 1400-1408.
33. E. Oberg, *Machinery's Handbook*, Industrial Press, New York, 27th edn., 2005.
34. K. C. Rice, E. E. Mann, J. L. Endres, E. C. Weiss, J. E. Cassat, M. S. Smeltzer and K. W. Bayles, *Proc. Natl. Acad. Sci. USA*, 2007, **104**, 8113-8118.
35. K. M. Colvin, Y. Irie, C. S. Tart, R. Urbano, J. C. Whitney, C. Ryder, P. L. Howell, D. J. Wozniak and M. R. Parsek, *Environ. Microbiol.*, 2012, **14**, 1913-1928.
36. H. Tzagoloff and R. Novick, *J. Bacteriol.*, 1977, **129**, 343-350.
37. M. Klausen, A. Aaes-Jorgensen, S. Molin and T. Tolker-Nielsen, *Mol. Microbiol.*, 2003, **50**, 61-68.
38. L. A. Pratt and R. Kolter, *Mol. Microbiol.*, 1998, **30**, 285-293.
39. G. A. O'Toole and R. Kolter, *Mol. Microbiol.*, 1998, **30**, 295-304.
40. M. Trejo, C. Douarche, V. Bailleux, C. Poulard, S. Mariot, C. Regeard and E. Raspaud, *Proc. Natl. Acad. Sci. USA*, 2013, **110**, 2011-2016.
41. Z. Kun, T. Boo Shan, B. Bernard, J. Fan, L. G. Maxsim, J. H. Joe, L. Erik, R. P. Matthew and C. L. W. Gerard, *Nature*, 2013, **497**, 388-391.
42. A. Persat, C. D. Nadell, M. K. Kim, F. Ingremeau, A. Siryaporn, K. Drescher, N. S. Wingreen, B. L. Bassler, Z. Gitai and H. A. Stone, *Cell*, 2015, **161**, 988-997.
43. A. K. Epstein, A. I. Hochbaum, P. Kim and J. Aizenberg, *Nanotechnology*, 2011, **22**, 494007.