

Chemical Science

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Acyclic Quaternary Centers from Asymmetric Conjugate Addition of Alkylzirconium Reagents to Linear Trisubstituted Enones

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Zhenbo Gao and Stephen P. Fletcher*^a

Synthetic methods for the selective formation of all carbon quaternary centres in non-cyclic systems are rare. Here we report highly enantioselective Cu-catalytic asymmetric conjugate addition of alkylzirconium species to twelve different acyclic trisubstituted enones. A variety of sp³-hybridized nucleophiles generated by in-situ hydrometallation of alkenes with the Schwartz reagent can be introduced, giving linear products bearing quaternary centres with up to 98% ee. The method is tolerant of several important functional groups and 27 total examples are reported. The method uses a new chiral nonracemic phosphoramidite ligand in a complex with copper triflate as the catalyst. This work allows the straightforward stereocontrolled formation of a valuable structural motif using only a catalytic amount of chiral reagent.


Introduction

The stereoselective formation of all-carbon quaternary centers is a challenging transformation in synthetic chemistry that has received tremendous attention. While a wide variety of methods now exist¹ the synthesis of quaternary centres in acyclic systems remains particularly difficult. In non-cyclic systems issues of steric repulsion and conformational control both need to be overcome by the catalyst in order to control enantioselectivity².

Over the past ten years copper-catalyzed asymmetric conjugate addition (ACA) of alkyl nucleophiles to α,β -unsaturated ketones has emerged as an important route to all-carbon quaternary stereocenters³. Seminal work by the Alexakis⁴ and Hoveyda⁵ groups has demonstrated that high chemo- and regio- and enantio-selectivities can be obtained using primary sp³-hybridized organometallics such as Grignard reagents,^{4d} dialkylzincs,^{5b} and aluminium reagents⁶. Many examples have now been reported using a variety of catalyst systems, and these reactions are now becoming adapted into the repertoire of synthetic chemists to make important molecules such as natural products and biologically active compounds⁷.

Cu-catalyzed ACAs to form quaternary centres normally rely on the use of cyclic α,β -unsaturated ketones, and although there are sporadic reported examples that use acyclic acceptors, these typically result in lower levels of enantioselectivity⁸. To the best of our knowledge there are only three highly

enantioselective methods for the addition of non-stabilized alkyl nucleophiles to form quaternary centres (Scheme 1 top). Two of these require highly activated electrophiles: i) addition of dialkyl zinc reagents to nitro-alkenes as developed by Hoveyda⁹ and expanded by Boehringer Ingelheim¹⁰, and ii) addition of dialkyl zinc reagents to Meldrum's acid derivatives¹¹. The third approach uses linear trisubstituted enones, which are normally activated substrates, in combination with aluminium reagents, and is limited to addition of very simple alkyl groups such as Me₃Al or Et₃Al^{6a, 12}.



Scheme 1 top: Catalytic asymmetric addition of non-stabilized nucleophiles to form non-cyclic quaternary centres. Bottom: Hydrometallation/asymmetric conjugate addition of alkenes to form quaternary centres

^a Department of Chemistry, University of Oxford. Chemistry Research Laboratory, 12 Mansfield Road, Oxford, OX1 3TA (UK). E-mail: stephen.fletcher@chem.ox.ac.uk

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

We have previously developed Cu-catalyzed ACAs using alkenes as the equivalents of alkyl nucleophiles¹³. Reacting an alkene with the Schwartz reagent (Cp_2ZrHCl) forms alkylzirconium species *in situ* which can undergo ACA without the need for cryogenic conditions. These methods allow introduction of reasonably complex alkyl nucleophiles that are normally incompatible with premade organometallic nucleophiles¹⁴.

We reported building tertiary¹³ and quaternary centers¹⁵ by addition to cyclic α,β -unsaturated ketones¹⁶. In acyclic substrates, it is much more difficult to obtain high enantioselectivity, presumably due to interconversion between *s-cis* and *s-trans* conformers. However, we were able to develop a method to construct acyclic tertiary centres with up to 91% e.e.¹⁷

In this paper, we report acyclic quaternary stereocenters via ACA of alkylzirconium species. 27 examples using a variety of linear electrophiles and a variety of alkenes are given. Overall this work expands the scope of linear α,β -unsaturated enones that can be used to form acyclic quaternary centres, and the scope and limitations of the electrophile in the current method are reported.

These reactions significantly extend the *n*-alkyl nucleophiles used to form non-cyclic all carbon quaternary products, which was previously limited to very simple groups. Here we show a wide variety of simple and functionalized alkenes can be used. These are much more readily available and easier to handle than comparable premade organometallics. Products bearing groups amenable to further modification are difficult to access using current methods, but necessary in moving asymmetric reaction development onward from simple benchmark substrates. Developing chemistry that is capable of using functionalized coupling partners in asymmetric C-C bond forming reactions is a key step toward being able to use the products in further processes such as natural product synthesis or drug discovery. Overall the work reported here significantly expands the range of non-products bearing quaternary centres that can be accessed. Our method involves a new chiral nonracemic phosphoramidite ligand, copper (I) triflate, TMSCl, and an ice-bath cooled reaction mixture that is stirred overnight.


Results and discussion

We examined the hydrometalation-ACA of ethylene (**1a**) and (E)-4-methyl-6-phenylhex-3-en-2-one (**2a**) under reactions conditions previously applied in the formation of cyclic quaternary centres¹⁵ and acyclic tertiary stereocenters¹⁷. Reaction using ligand **A** gave **3a** with 46% yield, 45% ee using conditions optimized for cyclic quaternary centres, and we obtained 70% yield, 77% ee using conditions developed for acyclic tertiary centres (Table 1, entries 1 and 2). Using ligand **D** in the addition to **2a** gave 80% yield and 84% ee (entry 3). TMSCl has long been recognized to play an important role in copper catalysed conjugate addition chemistry¹⁸ either by accelerating addition by activating the enone oxygen, or interacting with the copper species themselves. The role of

TMSCl in our reactions is not clear, but it is critical as the yield and ee both drop significantly when TMSCl is omitted (for example entry 4, 9% yield with 58% ee). We examined ligand **D** with a variety of solvents and copper sources and found that many copper sources worked poorly. Salts with electron withdrawing counterions performed well, with CuNTf_2 and CuOTf providing the best ee's (84% and 56% respectively, entries 3 and 8), these salts are generated *in situ* because the silver salts are commercially available and the isolated Cu-salts are difficult to prepare. The use of Cu(II) triflate allowed ACA (entry 9), but results were inferior to using Cu(I) directly. When different solvents were used a variety of results were obtained (entries 10-13), with ethers generally giving better yields and ee's. Overall, Et_2O gave the best results in this system (entry 3), with 80% yield and 84% ee.

In an attempt to further optimize the system, we examined new ligands; **E**, **F** and **B** were prepared and tested. These new ligands have different branched aliphatic groups attached to the amine portion of the phosphoramidite, and for reasons that are not yet clear (but are under current investigation) all of these new ligands performed better (entries 14, 15 and 16)

Table 1 Screening conditions for hydrometalation/asymmetric conjugate addition to acyclic trisubstituted enones



Entry	Copper source	Solvent	Ligand	Yield [%] ^[a]	ee [%] ^[b]
1	$\text{CuCl} + \text{AgNTf}_2$	t-BuOMe	A	46	45
2	$\text{CuCl} + \text{AgOTf}$	Et_2O	C	70	77
3	$\text{CuCl} + \text{AgOTf}$	Et_2O	D	80	84
4 ^[c]	$\text{CuCl} + \text{AgOTf}$	Et_2O	D	9	58
5	CuCl	Et_2O	D	-	-
6	$\text{CuCl} + \text{AgClO}_4$	Et_2O	D	4	2
7	$\text{CuCl} + \text{AgSbF}_6$	Et_2O	D	13	20
8	$\text{CuCl} + \text{AgNTf}_2$	Et_2O	D	20	56
9	$\text{Cu}(\text{OTf})_2$	Et_2O	D	40	69
10	$\text{CuCl} + \text{AgOTf}$	THF	D	41	75
11	$\text{CuCl} + \text{AgOTf}$	i-Pr ₂ O	D	62	52
12	$\text{CuCl} + \text{AgOTf}$	CH_2Cl_2	D	7	8
13	$\text{CuCl} + \text{AgOTf}$	Toluene	D	4	30
14	$\text{CuCl} + \text{AgOTf}$	Et_2O	E	80	86
15	$\text{CuCl} + \text{AgOTf}$	Et_2O	F	78	86
16	$\text{CuCl} + \text{AgOTf}$	Et_2O	B	80	91

Reaction conditions: ethylene balloon, $[\text{Cp}_2\text{ZrHCl}]$ (2 eq.), CH_2Cl_2 ; then (E)-4-methyl-6-phenylhex-3-en-2-one (1 eq.), copper (10 mol%), ligand (10 mol%), silver salt (11 mol%), TMSCl (5 eq.) [a] Yield of isolated product. [b] ee was determined by HPLC. [c] Without TMSCl.

than cyclohexyl-substituted **D**. Ligand **B**, bearing two alkyl chains, gave the best enantioselectivity and is capable of giving non-cyclic all carbon quaternary centres with 80% isolated yield and >90% ee (Entry 15).

Table 2 Scope of enones

Entry	Substrate	Product	Yield [%] ^[a]	ee [%] ^[b]
1			50	90
2			34	89
3			39	73
4			46	91
5			41	85
6 ^[c]			48	92
7			71	98
8			61	93
9			80	93
10 ^[d]			80	96
11 ^[d]			54	96
12		-	-	-
13		-	-	-
14			32	71

Reaction conditions: alkene (2.5 eq.), [Cp₂ZrHCl] (2 eq.), CH₂Cl₂; then enone (1 eq.), copper (10 mol%), ligand (10 mol%), silver (11 mol%), TMSCl (5 eq.)
 [a] Yield of isolated product. [b] ee was determined by HPLC. [c] ee was determined by GC. [d] The enantiomer of ligand of B was used in these examples because the order of elution of the enantiomers aided ee

Using the conditions developed above, we were able to obtain a tetra-alkyl-substituted product in good (80%) yield with high (91%) enantiomeric excess. We next studied the effect of the electrophile and examined ACA with a variety of E-substituted acyclic trisubstituted enones, where the groups adjacent to the olefin and carbonyl groups were varied (Table 2).

Generally, ee's are high. The reaction works very well when the enone is tri-*n*-alkyl-substituted, and gives lower yields (without recovery of starting material) when styrene derivatives are used. The reaction is sensitive to branching at some positions (see below) where lower yields are obtained with recovered starting material. Attempts to improve our yields by simply extending reaction times to 48 hours did not significantly change the results. It is likely a new system will be needed to overcome these limitations. The absolute stereochemistry of the products was assigned by comparison of **3b**, **3d**, **3e**, and **3i** with reported optical rotation values^{6a}, for all other products the absolute configuration is assigned by analogy.

In the case of enone **2b**, which has a phenyl ring in the β-position, moderate yield (50%) but high enantioselectivity (90% ee) was observed. Other aryl-substituted enones behaved similarly and poor results were obtained using electron rich **2d** (39%, 73% ee). We were pleased to find that a challenging heterocyclic substrate (bearing a thiophene, entry 5, 41%, 85% ee) was tolerated in this reaction. β-alkyl substituted enones **2a**, **2g-2j** (Table 1 and Table 2 entries 6-9) all performed well giving better yields and >90% ee. With a more sterically demanding branched example (**3g**, entry 6) lower yield but excellent ee (48%, 92% ee) was obtained and in this case there was unreacted **2g** remaining after the standard overnight reaction.

Somewhat surprisingly, when we increased the length of carbon chain adjacent to the enone, we obtained even higher levels of enantioselectivity (both 96% ee, entries 10 and 11). **3l** gave slightly lower yield than other tri-alkyl substituted enones (54%) but in this case there was also unreacted starting **2l**. We note that the enantiomer of ligand **B** (*ent-B*, not shown) was used to make **3k** and **3l** because it aided ee determination.

When we examined isopropyl (2*n*) and phenyl (**2n**) substituted enones (entries 12 and 13) no desired product was obtained. In these cases a secondary carbon is next to the carbonyl, and the catalytic reaction evidently does not tolerate additional substitution at this position. While it is not currently clear why these substrates do not undergo reaction, it is likely that the additional substitution restricts the conformational freedom of the enones, and this may inhibit an important part of the mechanistic pathway.

The stereochemistry of the starting enone is important; *cis*-(*Z*)-enone *cis*-**2b** (entry 14) gave much poorer results (32%, 71% ee) than *trans* **2b** (entry 1, 50%, 90% ee). This selectivity difference appears to arise from partial *cis*- to *trans*-isomerization during the course of the reaction; when we stopped the addition to *cis*-(*Z*)-enone *cis*-**2b** after one hour (*R*)-**3b** was obtained in 26% yield with 86% ee. Recovery of starting material showed that *trans*-enone **2b** was present in the mixture with the enone now having a 1:5 *trans* to *cis* ratio.

Table 2 Addition of alkenes

Entry	Substrate	Product	Yield [%] ^[a]	ee [%] ^[b]
1			53	93
2			70	90
3			69	85
4			56	92
5			59	90
6			64	90
7 ^[c]			51	84
8 ^[c]			46	87
9 ^[c]			52	78
10 ^[c]			55	74
11 ^[c]			42	88
12 ^[c]			54	82
13 ^[c]			71	91
14 ^[d]			24	88
15 ^[d]			35	92

Reaction conditions: alkene (2.5 eq.), [Cp₂ZrHCl] (2 eq.), CH₂Cl₂; then enone (1 eq.), copper (10 mol%), ligand (10 mol%), silver (11 mol%), TMSCl (5 eq.)
 [a] Yield of isolated product. [b] Enantiomeric excess was determined by HPLC. [c] Enone **2a** was used instead of **2k**. [d] Enone **2h** was used instead of **2k**.


We next used more complex nucleophiles in additions to enone **2k**. We first replaced ethylene with a range of non-

functionalized alkenes (Table 3, entries 1-6) and found that enantioselectivity was uniformly high (>90% ee) except when using *t*-Bu-substituted alkene **1d** (entry 3, 85% ee). Using more elaborate nucleophiles, which contain functional groups (Table 3, entries 7-13), was performed with enone **2a**. The formation of quaternary centres tolerates halogens (entries 7 and 8), unsaturated C-C bonds (entries 9 and 10), allyl silane (entry 11), and protected alcohols (entries 12 and 13), and aromatic rings (entries 9, 12, 14 and 15).

When attempting to introduce styrene derivatives in combination with **2a** we found it difficult to find a suitable analytical method to determine the ee. We used enone **2h** (instead of **2a**) in combination to 4-MeO-styrene **1o** to give **3z** (entry 14, 24% yield, 88% ee). Styrene did not work in our system. Alkene **1p** and enone **2h** were coupled to give **3aa** (35% yield, 92% ee, entry 15).

We next studied if we can improve our yields by simply extending reaction time from 15 hours (entry 15 in table 3) to 48 hours. However, 48-hour reaction provided almost same result (35% yield, 91% ee).

We also performed a preparative scale experiment to examine if the method was suitable for scale-up (Scheme 2). On a gram scale, we chose 1-octene **1e** and enone **2k** as reactants, and arbitrarily decreased the catalyst loading to 5 mol%, and the amount of Schwartz reagent to 1.5 equivalents. We obtained good results (0.79 g, 50% yield, 93% ee) which are similar those on small scale reactions with 10% catalyst and 2 equivalents of Cp₂ZrHCl (56% yield, 92% ee).



Scheme 2 Gram-scale catalytic acyclic asymmetric conjugate addition using 5% copper and ligand.

Experimental

CuCl (1.9 mg, 0.02mmol, 0.1 eq.), and the phosphoramidite ligand B (11.5 mg, 0.02 mmol, 0.1 eq.) were added to a flame dried round bottom flask containing Et₂O (1 mL) under an argon atmosphere. The resulting mixture was stirred at room temperature for 1 hour and then AgOTf (5.6 mg, 0.022 mmol, 0.11 eq.) was added and the mixture was stirred for an additional 15 min. To a second, flame dried, round bottom flask, Cp₂ZrHCl (103 mg, 0.4 mmol, 2.0 eq.) was added. A balloon filled with ethylene was used to purge the flask with ethylene for 5 min, and then CH₂Cl₂ (0.2 mL) was added. After stirring for 15 min under an ethylene atmosphere (balloon), a clear yellow solution was obtained. After stirring for 15 min, the stirred solution containing the copper and ligand was transferred, and filtered – using a syringe filter – to the solution containing the zirconocene species. The resulting black mixture was stirred for another 10 min before enone **2a** (38.0 mg, 0.2 mmol, 1.0 eq.) and then TMSCl (0.127 mL, 1.0 mmol, 5.0 eq.) were each added dropwise. Stirring was continued for 15 h at 0 °C, before the reaction was quenched by the addition of 1.5 mL NH₄Cl and then 3.0 mL Et₂O. The

mixture was partitioned between the aqueous and organic phases, and the aqueous layer was extracted with Et₂O (3 x 10 mL). The combined organic materials were dried with Na₂SO₄, filtered, concentrated, and the resulting yellow residue purified by flash column chromatography (Petrol: Et₂O; 90:10; SiO₂) to give the desired product. (+)-(R)-4-ethyl-4-methyl-6-phenylhexan-2-one (35 mg, 80 % yield, 91% ee) HPLC analysis indicated an enantiomeric excess of 91 % [two Chiralpak® ID in series; flow: 0.8 mL/min; hexane/i-PrOH: 99:1; λ = 210 nm; major enantiomer tR = 14.58 min; minor enantiomer, tR = 15.65 min].

Conclusions

We have developed a Cu-catalyzed ACA to generate all-carbon quaternary centers from acyclic substrates. This transformation often gives high (often over 90% ee) levels of enantioselectivity using novel phosphoramidite ligand B. A wide range of trisubstituted acyclic enones can be used, and the best results (both high yields and ee's) are obtained when forming tetra-*n*-alkyl substituted products. We can introduce a variety of non-functionalized and functionalized primary sp³-hybridized nucleophiles, these experiments significantly expand the range of nucleophiles that can be used to form quaternary centers and provided access to structures that can be further exploited in synthesis design. The nucleophilic zirconium species are generated by *in-situ* hydrozirconation of the corresponding alkene with Schwartz's reagent. Up to 98% ee and 27 working examples are reported using E-substituted enones bearing primary alkenes next to the enone-carbonyl group. Preliminary experiments show the reaction can be scaled up to at least 0.79 g using 5 mol % catalyst without compromising on the enantioselectivity.

Acknowledgements

ZG thanks the China Scholarship Council (CSC) for financial support.

Notes and references

- Selected reviews: (a) E. J. Corey and A. Guzman-Perez, *Angew. Chem. Int. Ed.* 1998, **37**, 388-401; (b) J. Christoffers and A. Baro, *Angew. Chem. Int. Ed.* 2003, **42**, 1688-1690; (c) C. J. Douglas and L. E. Overman, *Proc. Natl. Acad. Sci.* 2004, **101**, 5363-5367; (d) B. M. Trost and C. Jiang, *Synthesis*, 2006, 369-396; (e) C. Hawner and A. Alexakis, *Chem. Commun.* 2010, **46**, 7295-7306; (f) K. W. Quasdorf and L. E. Overman, *Nature*, 2014, **516**, 181-191; (g) M. Buschleb, S. Dorich, S. Hanessian, D. Tao, K. B. Schenthal and L. E. Overman, *Angew. Chem. Int. Ed.*, 2016, **55**, 4156-4186.
- For detailed discussion see: (a) J. P. Das and I. Marek, *Chem. Commun.* 2011, **47**, 4593-4623; (b) I. Marek, Y. Minko, M. Pasco, T. Mejuch, N. Gilboa, H. Chechik and J. P. Das, *J. Am. Chem. Soc.* 2014, **136**, 2682-2694.

- (a) A. Alexakis, N. Krause and S. Woodward, Eds. *Copper-Catalyzed Asymmetric Synthesis*, Wiley-VCH Verlag GmbH & Co. KGaA, 2014; (b) B. Maciá, in *Progress in Enantioselective Cu(I)-catalyzed Formation of Stereogenic Centers*, Ed. S. R. Harutyunyan, Springer, 2016, vol. 58, pp. 41-98.
- (a) M. d'Augustin, L. Palais and A. Alexakis, *Angew. Chem. Int. Ed.* 2005, **44**, 1376-1378; (b) D. Martin, S. Kehrl, M. d'Augustin, H. Clavier, M. Mauduit and A. Alexakis, *J. Am. Chem. Soc.*, 2006, **128**, 8416-8417; (c) C. Hawner, K. Li, V. Cirriez and A. Alexakis, *Angew. Chem. Int. Ed.* 2008, **47**, 8211-8214; (d) M. Tissot, D. Poggiali, H. Henon, D. Muller, L. Guenee, M. Mauduit and A. Alexakis, *Chem. Eur. J.*, 2012, **18**, 8731-8747.
- (a) A. W. Hird and A. H. Hoveyda, *J. Am. Chem. Soc.* 2005, **127**, 14988-14989; (b) K.-s. Lee, M. K. Brown, A. W. Hird and A. H. Hoveyda, *J. Am. Chem. Soc.* 2006, **128**, 7182-7184; (c) T. L. May, M. K. Brown and A. H. Hoveyda, *Angew. Chem. Int. Ed.* 2008, **47**, 7358-7362; (d) T. L. May, J. A. Dabrowski and A. H. Hoveyda, *J. Am. Chem. Soc.* 2011, **133**, 736-739.
- (a) J. A. Dabrowski, M. T. Villaume and A. H. Hoveyda, *Angew. Chem. Int. Ed.* 2013, **52**, 8156-8159; (b) K. P. McGrath and A. H. Hoveyda, *Angew. Chem. Int. Ed.* 2014, **53**, 1910-1914.
- Examples of Cu-catalyzed ACA to form quaternary centres in natural product synthesis: (a) M. Vuagnoux-d'Augustin and A. Alexakis, *Chem. Eur. J.* 2007, **13**, 9647-9662; (b) K. M. Peese and D. Y. Gin, *Chem. Eur. J.* 2008, **14**, 1654-1665; (c) M. K. Brown and A. H. Hoveyda, *J. Am. Chem. Soc.* 2008, **130**, 12904-12906; (d) S. Kehrl, D. Martin, D. Rix, M. Mauduit and A. Alexakis, *Chem. Eur. J.* 2010, **16**, 9890-9904; (e) A. Mendoza, Y. Ishihara and P. S. Baran, *Nat. Chem.* 2012, **4**, 21-25; (f) Y. Ishihara, A. Mendoza and P. S. Baran, *Tetrahedron*, 2013, **69**, 5685-5701; (g) S. G. Krasutsky, S. H. Jacobo, S. R. Tweedie, R. Krishnamoorthy and A. S. Filatov, *Org. Proc. Res. Dev.* 2015, **19**, 284-289; (h) P. Cottet, C. Bleschke, M. G. Capdevila, M. Tissot and A. Alexakis, *Adv. Synth. Catal.* 2016, **358**, 417-425.
- Y. Minko, M. Pasco, L. Lercher, M. Botoshansky and I. Marek, *Nature*, 2012, **490**, 522-526.
- J. Wu, D. M. Mampreian and A. H. Hoveyda, *J. Am. Chem. Soc.*, 2005, **127**, 4584-4585.
- X. Zeng, J. J. Gao, J. J. Song, S. Ma, J. N. Desrosiers, J. A. Mulder, S. Rodriguez, M. A. Herbage, N. Haddad, B. Qu, K. R. Fandrick, N. Grinberg, H. Lee, X. Wei, N. K. Yee and C. H. Senanayake, *Angew. Chem. Int. Ed.* 2014, **53**, 12153-12157.
- (a) E. Fillion and A. Wilsily, *J. Am. Chem. Soc.* 2006, **128**, 2774-2775; (b) S. J. Mahoney, A. M. Dumas and E. Fillion, *Org. Lett.* 2009, **11**, 5346-5349; (c) A. Wilsily and E. Fillion, *J. Org. Chem.* 2009, **74**, 8583-8594; (d) E. Fillion, A. Wilsily and T. Lou, *Synthesis*, 2009, 2066-2072; (e) A. M. Dumas and E. Fillion, *Acc. Chem. Res.*, 2010, 440-454.
- (a) K. Endo, D. Hamada, S. Yakeishi and T. Shibata, *Angew. Chem. Int. Ed.* 2013, **52**, 606-610; (b) K. Endo, S. Yakeishi, R. Takayama and T. Shibata, *Chem. Eur. J.* 2014, **20**, 8893-8897.
- R. M. Maksymowicz, P. M. Roth and S. P. Fletcher, *Nat. Chem.* 2012, **4**, 649-654.
- R. M. Maksymowicz, A. J. Bissette and S. P. Fletcher, *Chem. Eur. J.* 2015, **21**, 5668-5678.
- M. Sidera, P. M. Roth, R. M. Maksymowicz and S. P. Fletcher, *Angew. Chem. Int. Ed.* 2013, **52**, 7995-7999.
- P. M. Roth, M. Sidera, R. M. Maksymowicz and S. P. Fletcher, *Nat. Protoc.* 2014, **9**, 104-111.
- P. M. Roth and S. P. Fletcher, *Org. Lett.* 2015, **17**, 912-915.

18. (a) E. J. Corey and N. W. Boaz, *Tetrahedron Lett.* 1985, **26**, 6019-6022; (b) E. J. Corey and N. W. Boaz, *Tetrahedron Lett.* 1985, **26**, 6015-6018; (c) A. Alexakis, J. Berlan and Y. Besace, *Tetrahedron Lett.* 1986, **27**, 1047-1050; (d) E. Nakamura, S. Matsuzawa, Y. Horiguchi and I. Kuwajima, *Tetrahedron Lett.* 1986, **27**, 4029-4032; (e) E. J. Corey, F. J. Hannon and N. W. Bo, *Tetrahedron*, 1989, **45**, 545-555; (f) A. Alexakis, R. Sedrani and P. Mangeney, *Tetrahedron Lett.* 1990, **31**, 345-348. (g) B. H. Lipshutz, S. H. Dimock and B. James, *J. Am. Chem. Soc.* 1993, 9283-9284.