



Metal-Free Defluorinative Arylation of Trifluoromethyl Alkenes via Photoredox Catalysis

Journal:	<i>ChemComm</i>
Manuscript ID	CC-COM-06-2019-004265
Article Type:	Communication

SCHOLARONE™
Manuscripts

COMMUNICATION

Metal-Free Defluorinative Arylation of Trifluoromethyl Alkenes via Photoredox Catalysis

Rebecca J. Wiles, James P. Phelan, and Gary A. Molander*^a

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


Literature methods to access *gem*-difluoroalkenes are largely limited to harsh, organometallic-based methods, and known photoredox mediated processes are not amenable to aryl radical addition to trifluoromethyl alkenes. A metal-free, functional group-tolerant method for the preparation of benzylic *gem*-difluoroalkenes is described. Halogen atom abstraction from (hetero)aryl halides generates aryl radicals that undergo a defluorinative arylation of α -trifluoromethyl alkenes, tolerating electronically disparate aryl radicals and α -trifluoromethyl alkenes.

Introduction

The incorporation of fluorinated motifs into organic structures is an ever-growing field of organic chemistry. It is a well-established and often-utilized strategy to introduce fluorine into organic molecules to impart desirable pharmacological properties. These properties include increased metabolic stability, enhanced lipophilicity, and improved bioavailability.¹ As such, methods for the synthesis of new and unusual fluorinated groups continue to increase in relevance.² Most importantly, there is a great need for methods that afford the incorporation or modification of fluorinated groups under mild conditions to maximize potential feasibility in late-stage functionalization.

One functional group of interest for medicinal chemistry is the geminally-substituted difluoroalkene. These structures are postulated as isosteres for carbonyl groups, and are particularly attractive when metabolism at a carbonyl is a point of undesirable metabolic degradation.³ Although methods for the synthesis of *gem*-difluoroalkenes have been described,⁴ they rely on two-electron chemistry and can be largely classified as either the direct conversion of carbonyls into *gem*-difluoroalkenes or as the modification of an existing fluorinated group. In the context of the addition of aryl groups to

trifluoromethyl alkenes, these methods are limited to two-electron additions of organometallic species such as phenyllithium and phenylcopper reagents.⁵ To date, milder, more functional group-tolerant arylation conditions have not been described, representing an important methodological gap. Recent work by our group and others has included the development of photoredox-mediated approaches for the synthesis of *gem*-difluoroalkenes.⁶ Several classes of radical precursors have been leveraged through this approach, including alkyltrifluoroborates, alkylbis(catecholato)silicates, α -silylamines, and alkyl carboxylate salts (Scheme 1). These radical precursors are effective for the generation of alkyl radicals displaying diverse functional groups; however, these methods are not feasible for arylation, as such aryl radical precursors have elevated oxidation potentials outside the range of many common photocatalysts.⁷ The present work demonstrates the first photoredox-catalyzed arylation of trifluoromethyl alkenes using (hetero)aryl halides as a radical source.




Scheme 1 Arylation of trifluoromethyl alkenes through a new pathway

An established pathway for the formation of aryl radicals in photoredox catalysis follows a one-electron reduction of aryl halides (I), facilitated by oxidative quenching of an excited photocatalyst ([PC]*) (Scheme 2).⁸ As demonstrated by the groups of Jui, Lee, and others, these reductively generated radicals (II) can engage in Giese-type additions to alkenes.^{8,9} Then, in a redox-neutral radical polar crossover (RPC) manifold, the photocatalyst cycle is completed by oxidation of radical

^a 231 S. 34th St. Philadelphia, PA 19104.


Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

intermediate III to regenerate the photocatalyst and form cationic species IV.^{9,10} Intermediate IV could then react further by either a base-mediated elimination to re-form an alkene or by nucleophilic attack at the cationic carbon. However, a cationic intermediate would not be viable for the E1cb fluoride elimination needed to produce the desired *gem*-difluoroalkene product.



Scheme 2 Net-neutral radical polar crossover via oxidative quenching photocatalysis

To access the anionic intermediate necessary for fluoride elimination (Scheme 3, XI) we envisioned generating aryl radicals through a net reductive pathway. A recent report by the MacMillan group demonstrated the feasibility of using tris(trimethylsilyl)silanol (V) as an activator of aryl halides (Ep = +1.54 V versus SCE).¹¹ We proposed that use of V, which alters the sequence of single electron transfer (SET) events so that a reductive quenching photocatalytic cycle occurs, would generate aryl radicals in a reducing environment capable of accessing the anionic intermediate XI needed for fluoride elimination.



Scheme 3 Mechanism for silanol-mediated radical polar crossover

We proposed that the catalytic cycle would be initiated by base-mediated deprotonation of silanol V to give VI. Then, oxidation by the photocatalyst followed by a radical Brook rearrangement generates silyl radical VIII. The silyl radical next abstracts a halogen atom from aryl halide I to afford aryl radical II, which adds to the trifluoromethyl alkene IX. The photocatalytic cycle is closed by a reduction of species X to the α -trifluoromethyl anion XI, and finally an E1cb elimination of fluoride generates the desired product XII.¹²

With the aid of high throughput experimentation, the reaction conditions were evaluated (Table 1) (see Supporting Information for additional details). Photocatalyst screening

revealed the organic photocatalyst 2,4,5,6-tetrakis(3,6-dichloro-9H-carbazol-9-yl)isophthalonitrile (Cl-4CzIPN, Ep = +1.71 V versus SCE) gave the highest formation of desired product, even when compared to commonly used iridium photocatalysts.¹³ The evaluation of various solvents identified polar aprotic solvents as most suitable, with DMSO giving the highest conversion. A screening of inorganic bases saw Na₂CO₃ provide optimal product formation. Control studies confirmed the necessity of each reaction component. Without photocatalyst or silanol, no reaction, whether undesired or otherwise, was observed. Without base, which is important for silanol activation, only trace product was formed. Finally, when the reaction was allowed to run in ambient light, no reactivity was observed.

Table 1. Reaction optimization and control studies

Entry	Deviation from Standard	Conversion ^a
1	None	98
2	[Ir(dF(CF ₃)ppy) ₂ (dCF ₃ bpy)](PF ₆)	59
3	acetone	77
4	MeCN	32
5	DMF	32
6	NaOAc	92 ^b
7	K ₃ PO ₄	49
8	No Cl-4CzIPN	0
9	No TMS ₃ SiOH	0
10	No base	< 5
11	No LEDs	0

^aConversions determined by ¹⁹F NMR of crude reaction mixtures. ^bConversion to product was observed with full consumption of starting materials with the formation of unidentified reaction byproducts.

Having identified optimal conditions, the scope of the reaction was evaluated (Table 2). The reaction was optimized with the electron deficient aryl nitrile (2a), and as expected good reactivity was observed with other electron-poor substrates such as an aryl ketone (2g) and ester (2i). The reaction was also amenable to the addition of several electron-rich aryl radicals (2d, 2h, 2i). Further experiments demonstrated moderate to good yields with meta- and para-substituted aryl methoxy halides (2h and 2i), albeit with marginally lower isolated yields. More impressively, the conditions were also amenable toward a variety of heteroaryl halides, including several substituted pyridines (2b, 2c, 2f, 2j, 2k) and even an unprotected indole (2d). Importantly, both aryl iodides and -bromides are substrates, allowing facile, direct access to numerous aryl radicals from commercially available reagents. Example 2k demonstrates the selectivity observed in the presence of both aryl bromides and -iodides, with radical generation in the ratio of 7.6:1 favoring reactivity at the iodide.

Table 2. Aryl radical scope.^{a,b}

Product	Yield	Product	Yield
	75%, 70% ^c X = Br		73% ^g X = I
	71% ^{e,h} X = Br		60% ^e X = Br
	82% X = Br		63% ^d X = Br
	79% ^{d,f} X = I		67% ^{e,h} X = I
	79% ^{d,g} X = I		46% ^{f,i} X = I
	60% X = Br		78% ^d X = Br

^aAll values indicate the yield of the isolated product. ^bGeneral reaction conditions: aryl halide (2.0 equiv, 1.0 mmol), alkene (1.0 equiv, 0.50 mmol), Cl-4CzIPN (5 mol %, 0.025 mmol), TMS₃SiOH (1.5 equiv, 0.75 mmol), Na₂CO₃ (4 equiv, 2.0 mmol), DMSO (0.1 M), 18 h, irradiating with blue LEDs (30 W). See the Supporting Information for details. ^cIsolated yield on 2.5 mmol scale; reaction run for 5 d. ^dIsolated on 1.0 mmol scale. ^eIsolated yield on 0.3 mmol scale. ^fConducted using 10 mol % Cl-4CzIPN. ^gReaction run for 2 d. ^hConducted using 1 equiv Ar-X. ⁱIsolated with <5% impurity of alkene. ^jIsolated as a 7.6:1 mixture of Br:I arene.

We then sought to explore the scope of the method with respect to the trifluoromethyl alkene substrate. In addition to the dioxole substrate used in optimization, the standard conditions were applicable to other electron-rich trifluoromethylalkenes. These included dimethoxy- (2s), dimethylamino- (2t), thioether- (2o), and tert-butyl-substituted (2u) trifluoromethyl alkenes. Electron poor substrates were also reactive, affording the corresponding products in moderate to good yields. The dichloro-substituted alkene (2m) is one example of this, and also demonstrates that the method is chemoselective for the activation of aryl iodides and bromides in the presence of aryl chlorides.¹⁴ The reaction also tolerated an appended alkyne (2v) and carboxylic acid (2x), with no evidence of undesired side-reactivity. Additionally, this method is viable on substrates with unprotected alcohols (2p). Other notable examples of the substrate scope include the expansion of this method to heteroaryl trifluoromethyl alkenes (2q, 2r,

2w). The reaction was also successful when performed on a 2.5 mmol scale, although a longer reaction time was necessary (2a, Table 2). Notably, the reaction is not amenable to radical additions to trisubstituted alkenes.

Table 3. Trifluoromethyl alkene scope.^{a,b}


Product	Yield	Product	Yield
	82%		78% ^e
	72%		57% ^{c,e}
	57% ^c		59% ^g
	63%		84%
	77% ^e		53% ^g
	79% ^{d,e,g}		42% ^{d,e,f,g}

^aAll values indicate the yield of the isolated product. ^bGeneral reaction conditions: aryl halide (2.0 equiv, 1.0 mmol), alkene (1.0 equiv, 0.50 mmol), Cl-4CzIPN (5 mol %, 0.025 mmol), TMS₃SiOH (1.5 equiv, 0.75 mmol), Na₂CO₃ (4 equiv, 2.0 mmol), DMSO (0.1 M), 18 h, irradiating with blue LEDs (30 W). See the Supporting Information for details. ^cIsolated on 1.0 mmol scale. ^dConducted using 10 mol % Cl-4CzIPN. ^eReaction run for 2 d. ^fConducted using 3 equiv Ar-X and 2.0 equiv TMS₃SiOH. ^gIsolated with <5% impurity of alkene.

^aAll values indicate the yield of the isolated product. ^bGeneral reaction conditions: aryl halide (2.0 equiv, 1.0 mmol), alkene (1.0 equiv, 0.50 mmol), Cl-4CzIPN (5 mol %, 0.025 mmol), TMS₃SiOH (1.5 equiv, 0.75 mmol), Na₂CO₃ (4 equiv, 2.0 mmol), DMSO (0.1 M), 18 h, irradiating with blue LEDs (30 W). See the Supporting Information for details. ^cIsolated on 1.0 mmol scale. ^dConducted using 10 mol % Cl-4CzIPN. ^eReaction run for 2 d. ^fConducted using 3 equiv Ar-X and 2.0 equiv TMS₃SiOH. ^gIsolated with <5% impurity of alkene.

During the course of the scope exploration, interesting reactivity was noted in the case of the exceptionally electron deficient example 2b (Scheme 4). When the standard conditions using two equivalents of aryl bromide were used, a significant portion of double addition product was observed (3a). This product results from the addition of a second aryl radical to the gem-difluoroalkene, where reduction and a second fluoride elimination furnishes the monofluoroalkene. Fluorinated triaryllkenes are themselves relevant to medicinal chemists as

they are known anticancer agents in the family of tamoxifen derivatives.¹⁵ A second interesting observation was made while developing the reaction workup conditions. It was found that when electron poor compounds were eluted through a plug of KF/alumina, an olefin isomerization occurred to convert the *gem*-difluoroalkene to a difluoromethyl-substituted internal alkene (3b). Moderate amounts of this isomerization was observed even when exposure to the KF/alumina was limited to only 30 s. When the sample was stirred in KF/alumina for 3 h, complete conversion to the internal alkene was observed. These two products demonstrate the utility of the trifluoromethyl alkenes prepared here as starting materials for a diverse array of functionalization reactions.



Scheme 4 Unusual experimental observations and side reactivity.

Methods to access structurally diverse fluorinated molecules have gained significant prominence in recent years. Photoredox catalysis has offered unique opportunities for achieving such transformations with mild reaction conditions and high functional group compatibility. The reported method offers an efficient, metal-free approach to achieve the arylation of trifluoromethyl alkenes to make a new class of substituted *gem*-difluoroalkenes that were previously inaccessible under such mild conditions, and thus represents a step forward in our ability to synthesize and modify useful fluorinated motifs.

Conflicts of interest


There are no conflicts to declare.

Acknowledgements

We sincerely thank Dr. John Milligan of the University of Pennsylvania and Dr. Christopher Kelly of Virginia Commonwealth University for useful discussion. We thank Dr. Charles Hendrick for his assistance in the Penn Merck High Throughput Experimentation Center. We thank Dr. Charles W. Ross, III for assistance in obtaining HRMS data. We thank Dr. Jun Gu and Prof. Bill Dailey for helpful insight into ¹⁹F NMR data.

Notes and references

- (a) N. A. Meanwell, *J. Med. Chem.*, 2011, **54**, 2529; (b) S., Purser, P. R. Moore, S. Swallow, S., V. Gouverneur, *Chem. Soc. Rev.*, 2008, **37**, 320; (c) K. W. Hagmann, *J. Med. Chem.*, 2008, **51**, 4359; (d) K. Müeller, C. Faeh, F. Diederich, *Science*, 2007, **317**, 1881; (e) T. Liang, C. N. Neumann, T. Ritter, *Angew. Chem. Int. Ed.*, 2013, **52**, 8214; (f) C. F. Ni, J. B. Hu, *Chem. Soc. Rev.*, 2016, **45**, 5441.
- (a) G. Magueur, B. Crousse, M. Ourévitch, D. Bonnet-Delpon, J.-P. Bégué, *J. Fluorine Chem.*, 2006, **127**, 637; (b) C. Leriche, S. He, C.-w. Chang, H.-w. Liu, *J. Am. Chem. Soc.*, 2003, **125**, 6348.
- (a) P. Malátková, V. Wsól, *Drug Metab. Rev.*, 2014, **46**, 96; (b) T. Matsunaga, S. Shintani, A. Hara, *Drug Metab. Pharmacokin.*, 2006, **21**, 1; (c) O. A. Barski, S. M. Tipparaju, A. Bhatnagar, *Drug Metab. Rev.* 2008, **40**, 553.
- (a) G. Chelucci, *Chem. Rev.*, 2012, **112**, 1344; (b) S. Zhang, S. Cao, *Tetrahedron Lett.* 2017, **58**, 375; (c) M. Hu, C. Ni, L. Li, Y. Han, J. Hu, *J. Am. Chem. Soc.*, 2015, **137**, 14496; (d) X. Zhao, L. Chunmai, B. Wang, S. Cao, *Tetrahedron Lett.* 2019, **60**, 129; (e) X. Lu, X.-X. Wang, T.-J. Gong, J.-J. Pi, S.-J. He, Y. Fu, *Chem. Sci.*, 2019, **10**, 809; (f) J. Xinfei, L. Yisen, S. Hongyan, C. Song, *Tetrahedron*, 2018, **74**, 4155; (g) Y. Lan, F. Yang, C. Wang, *ACS Catal.*, 2018, **8**, 9245; (h) T. Fujita, M. Takazawa, K. Sugiyama, N. Suzuki, J. Ichikawa, *J. Org. Lett.*, 2017, **19**, 588; (i) Y. Liu, Y. Zhou, Y. Zhao, J. Qu, *Org. Lett.*, 2017, **19**, 946; (j) K. Fuchibe, H. Hatta, K. Oh, R. Oki, J. Ichikawa, *Angew. Chem. Int. Ed.*, 2017, **56**, 5890.
- (a) J.-P. Begue, D. Bonnet-Delpon, M. H. Rock, *Tetrahedron Lett.*, 1995, **36**, 5003; (b) T. Kitazume, T. Ohnogi, *Synthesis*, 1988, **8**, 614; (c) T. Miura, Y. Ito, M. Murakami, *Chem. Lett.* 2008, **37**, 1006.
- (a) L. Zhiyang, L. Yun, C. Wang, *ACS Catal.*, 2019, **9**, 775; (b) L.-H. Wu, J.-K. Cheng, L. Shen, Z.-L. Shen, T.-P. Loh, *Adv. Synth. Catal.*, 2018, **360**, 3894; (c) P.-J. Xia, Z.-P. Ye, Y.-Z. Hu, D. Song, H.-Y. Xiang, X.-Q. Chen, *Org. Lett.*, 2019, ASAP DOI: 10.1021/acs.orglett.9b00651.
- (a) C. K. Prier, D. A. Rankic, D. W. C. MacMillan, *Chem. Rev.*, 2013, **113**, 5322; (b) N. A. Romero, D. A. Nicewicz, *Chem. Rev.* 2016, **116**, 10075; (c) L. A. Shundrin, V. V. Bardin, H.-J. Frohn, *Z. Anorg. Allg. Chem.*, 2004, **630**, 1253.
- (a) I. Ghost, L. Marzo, A. Das, R. Shaikh, B. König, *Acc. Chem. Res.*, 2016, **49**, 1566; (b) G. Qiu, Y. Li, J. Wu, *Org. Chem. Front.*, 2016, **3**, 1011; (c) A. Studer, D. P. Curran, *Nat. Chem.*, 2014, **6**, 765;
- (a) R. A. Aycocock, H. Wang, N. T. Jui, *Chem. Sci.*, 2017, **8**, 3121; (b) C. P. Seath, D. B. Vogt, Z. Xu, A. J. Boyington, N. T. Jui, *J. Am. Chem. Soc.*, 2018, **140**, 15525; (c) A. J. Boyington, C. P. Seath, A. M. Zearfoss, Z. Xu, N. A. Jui, *J. Am. Chem. Soc.* 2019, **141**, 4147; (d) H. Kim, C. Lee, *Angew. Chem. Int. Ed.*, 2012, **51**, 12303.
- (a) N. A. Romero, D. A. Nicewicz, *Chem. Rev.*, 2016, **116**, 10075; (b) J. Xie, H. Jin, A. S. K. Hashmi, *Chem. Soc. Rev.*, 2017, **46**, 5193.
- (a) C. Le, T. Q. Chen, T. Liang, P. Zhang, D. W. C. MacMillan, *Science*, 2018, **360**, 1010; (b) P. Zhang, C. Le, D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2016, **138**, 8084; (c) M. Lucarini, E. Marchesi, G. F. Pedulli, *J. Org. Chem.*, 1998, **63**, 1687.
- This mechanism was supported by computational studies in our recent report, where the energy barrier for anionic fluoride elimination was calculated as ~6 kcal mol⁻¹, compared to ~63 kcal mol⁻¹ for a radical mediated fluoride elimination. J. P. Phelan, S. B. Lang, J. S. Compton, C. B. Kelly, G. A. Molander, *J. Am. Chem. Soc.* 2018, **140**, 8037.
- F. Le Vaillant, M. Garreau, S. Nicolai, G. Gryn'ova, C. Corminboeuf, J. Waser, *Chem. Sci.*, 2018, **9**, 5883.
- (a) R. J. Enemaerke, T. B. Christensen, J. Jensen, K. Daasbjerg, *J. Chem. Soc., Perkin Trans. 2*, 2001, 1620 (b) C. P. Andrieux, C. Blocman, J. M. Dumas-Bouchiat, J. M. Saveant, *J. Am. Chem. Soc.*, 1979, **101**, 3431.
- (a) C. Isanbor, D. O'Hagan, *J. Fluorine Chem.* 2006, **127**, 303; (b) S. P. Govek, J. Y. Nagasawa, K. L. Douglas, A. G. Lai, M. Kahraman, C. Bonnefous, A. M. Aparicio, B. D. Darimont, K. L. Grillo, J. D. Joseph, J. A. Kaufman, K.-J. Lee, N. Lu, M. Moon, R. Y. Prudente, J. Sensintaffar, P. J. Rix, J. H. Hager, N. D. Smith, *Bioorg. Med. Chem. Lett.*, 2015, **25**, 5163.



TOC Text:

Photoredox catalysis, using an organic photocatalyst and silanol reagent, enables the metal-free generation of aryl radicals to synthesize diverse *gem*-difluoroalkenes.