



**Environmental
Science**
Processes & Impacts

**No Evidence for Brown Carbon Formation in Ambient
Particles Undergoing Atmospherically Relevant Drying**

Journal:	<i>Environmental Science: Processes & Impacts</i>
Manuscript ID	EM-ART-10-2019-000457.R1
Article Type:	Paper

SCHOLARONE™
Manuscripts

1
2
3 **1 No Evidence for Brown Carbon Formation in Ambient Particles Undergoing Atmospherically**
4
5 **2 Relevant Drying**
6
7

8 3 Vikram Pratap¹, Michael A. Battaglia Jr.¹, Annmarie G. Carlton², Christopher J. Hennigan^{1*}

9
10 4 ¹*Department of Chemical, Biochemical and Environmental Engineering, University of Maryland,*
11 5 *Baltimore County*

12 6 ²*Department of Chemistry, University of California, Irvine*
13 7

14
15 8 **ABSTRACT:**
16
17

18 9 Recent laboratory studies have reported the formation of light-absorbing organic carbon compounds
19
20 10 (brown carbon, BrC) in particles undergoing drying. Atmospheric particles undergo cycles of
21
22 11 humidification and drying during vertical transport and through daily variations in temperature and
23
24 12 humidity, which implies particle drying could potentially be an important source of BrC globally. In this
25
26 13 work, we investigated BrC formation in ambient particles undergoing drying at a site in the eastern
27
28 14 United States during summer. Aerosol BrC concentrations were linked to secondary organic aerosol
29
30 15 (SOA) formation, consistent with seasonal expectations for this region. Measurements of water-soluble
31
32 16 organic aerosol concentrations and light absorption (365 nm) were alternated between an unperturbed
33
34 17 channel and a channel that dried particles to 41% or 35% relative humidity (RH), depending on the
35
36 18 system configuration. The RH maintained in the dry channels was below most ambient RH levels
37
38 19 observed throughout the study. We did not observe BrC formation in particles that were dried to either
39
40 20 RH level. The results were consistent across two summers, spanning ~5 weeks of measurements that
41
42 21 included a wide range of RH conditions and organic and inorganic aerosol loadings. This work suggests
43
44 22 that mechanisms aside from humidification-drying cycles are more important contributors to ambient
45
46 23 particle BrC loadings. The implications of this work on the atmospheric budget of BrC are discussed.
47
48
49
50
51 24

52
53 25 **ENVIRONMENTAL SIGNIFICANCE STATEMENT:**
54
55
56
57
58
59
60

1
2
3 26 Organic carbon in atmospheric particles can perturb solar insolation and hence, alter radiation budgets.
4
5 27 Recent laboratory studies have shown that light-absorbing organic carbon (BrC) may be produced from
6
7 28 chemical reactions in particles when dried. Atmospheric particles undergo natural drying during daily
8
9 29 cycles in relative humidity (RH) and through changes in RH encountered during vertical transport. This
10
11 30 may form brown carbon if the processes identified in the lab studies are widespread. We investigated this
12
13 31 phenomenon in the eastern United States and did not find evidence of BrC formation in atmospheric
14
15 32 particles that were dried from ambient RH to 35% and 41% RH. This suggests that other processes are
16
17 33 more important contributors to BrC formation in ambient particles.
18
19
20
21 34

22 23 24 35 1. INTRODUCTION

25
26
27 36 Brown carbon (BrC) is defined as the organic carbon that absorbs radiation in the near UV-Vis spectrum
28
29 37 (300 – 400 nm).¹⁻⁴ It has a positive radiative forcing effect on climate, and is increasingly found to be a
30
31 38 significant light-absorption component of ambient particles.^{1,5-8} On a global scale, radiative forcing by
32
33 39 BrC is estimated to be approximately 25% of the radiative forcing by black carbon, though it can have a
34
35 40 significantly higher contribution in regions strongly impacted by biomass burning emissions.^{1,7,9}

36
37
38 41 Brown carbon is primarily produced from incomplete and smoldering biomass combustion, which
39
40 42 includes wildfires, prescribed fires, wood and other biofuels used for cooking, heating, and other energy
41
42 43 demands.¹⁰ Brown carbon is also generated through secondary pathways involving a variety of precursors,
43
44 44 such as glyoxal^{11,12}, methylglyoxal¹²⁻¹⁵, isoprene¹⁶, monoterpenes¹⁷, and lignin.¹⁸ The formation of
45
46 45 secondary BrC compounds has been reported in both laboratory and ambient studies.¹⁹⁻²² For example, in
47
48 46 the southeastern United States, secondary BrC was dominated by organic aerosol from biogenic sources.²¹
49
50
51 47 On the other hand, secondary BrC in an urban location in western United States (Los Angeles) was
52
53 48 dominated by anthropogenic sources, and had several times higher absorption (normalized by carbon
54
55 49 mass) than the BrC in the Southeast, suggesting significant differences between BrC sources and/or
56
57
58
59
60

1
2
3 50 formation processes.^{19,21} Additionally, particulate water-soluble organic carbon (WSOC_p) concentrations
4
5 51 exhibited very different correlations with BrC absorbance in these locations, reinforcing geospatial
6
7 52 differences among BrC sources and formation processes in the atmosphere.²¹ The prevalence of BrC
8
9 53 throughout the troposphere provides further evidence of the importance of secondary processes
10
11 54 contributing to BrC formation.^{5,23,24} Humic-like substances (HULIS) observed in clouds are strongly
12
13 55 absorbing, and are also likely secondary in nature.^{25,26}
14
15
16 56 Glyoxal (Gly) and methylglyoxal (MeGly) are well-known precursors in secondary BrC formation. The
17
18 57 BrC formation reaction likely requires the presence of nitrogen-containing species like amines and
19
20 58 ammonium sulfate (AS).²⁷ The BrC formation occurs through an aldol-condensation reaction in the
21
22 59 aqueous phase, requires sulfate or acidic conditions, first forms imidazole oligomers eventually leading to
23
24 60 light absorbing compounds (BrC).^{11-13,15,27-31} This reaction is not considered oxidant driven and is
25
26 61 confirmed by several laboratory studies.^{11-13,15,27-31} Early studies of this pathway focused on bulk
27
28 62 reactions where Gly/MeGly and AS were mixed into an aqueous solution and analyzed for BrC
29
30 63 (absorbance) over a period of time. The reactions in the bulk phase were found to be extremely slow, as
31
32 64 light absorption increased over a period of days.^{11-13,30} Recent work focused on investigating BrC
33
34 65 formation in drying/evaporating droplets, which was more relevant to ambient aerosols.^{14,29,32} These
35
36 66 studies confirmed at least an order of magnitude faster rate of BrC formation compared to the bulk
37
38 67 reactions.^{14,29,32-34} For example, Lee et al. (2013)³⁴ reported BrC formation within a few seconds in Gly +
39
40 68 AS particles dried at an RH of $\leq 35\%$. The enhancement in BrC formation in evaporating droplets occurs
41
42 69 as the concentration of reactants in aqueous droplets increase with water evaporation, increasing reaction
43
44 70 rates. Therefore, the process of drying may also have an impact on BrC formation in ambient particles.
45
46 71 The liquid water content of particles in the atmosphere changes throughout the day with natural variations
47
48 72 in temperature and relative humidity.³⁵ Vertical transport also perturbs aerosol liquid water content due to
49
50 73 the associated temperature and RH gradients. Similarly, most clouds are non-precipitating³⁶, thus
51
52 74 particles that go through a cloud cycle experience large changes in their liquid water content and in the
53
54
55
56
57
58
59
60

1
2
3 75 aqueous phase concentrations of their components. Such humidification-drying cycles may have a
4
5 76 profound effect on secondary BrC formation.^{14,29,33,34} To our knowledge, BrC formation in ambient
6
7 77 particles undergoing drying has not been investigated previously. Herein, we characterize this
8
9 78 phenomenon in the eastern United States during the summer, a time and region where secondary BrC
10
11 79 formation is known to occur.^{20,21}

14 80 2. MATERIALS AND METHODS

16 81 Ambient particle sampling was carried out on the campus of the University of Maryland, Baltimore
17
18 82 County (UMBC).^{37,38} The experimental setup was based upon El-Sayed et al. (2015), but modified to
19
20 83 simultaneously measure WSOC_p and absorbance ($\lambda = 365$ nm) of the water-soluble aerosol components
21
22 84 (Figure 1). The experimental setup was installed inside a temperature-controlled building on UMBC's
23
24 85 campus. In addition to WSOC_p and absorbance, gas-phase ammonia and inorganic aerosol composition
25
26 86 (Na^+ , NH_4^+ , K^+ , Ca^{2+} , Mg^{2+} , Cl^- , NO_3^- , and SO_4^{2-}) were measured. Sampling was conducted in two phases.
27
28 87 Approximately 3-4 weeks of data were collected during the period of 17 July - 31 August, 2018 (Phase I),
29
30 88 and 1-2 weeks from 13 – 23 August, 2019 (Phase II). In the first phase, one dryer provided an average dry
31
32 89 channel RH of 41% (Figure 1), while two dryers were installed in the second phase to yield an average
33
34 90 dry channel RH of 35% (Figure 2). The average drying RH of $41 \pm 4\%$ was selected to simulate the
35
36 91 natural drying of particles in the atmosphere. This was in agreement with the average daytime RH of
37
38 92 $\sim 45\%$ recorded in a previous summer campaign in Baltimore.³⁹ During the present study, in Phase I, the
39
40 93 average minimum daytime RH was $55\% \pm 13\%$, and only $\sim 2\%$ of hourly data were below an RH of 41%
41
42 94 (Figure 2). During Phase II, the average minimum daytime RH was $51\% \pm 13\%$, with only $\sim 0.1\%$ of
43
44 95 hourly RH values below 35%. Therefore, the experimental results typically dried ambient particle below
45
46 96 observed minimum values experienced at the surface during our study. Even though the ambient RH
47
48 97 rarely falls below 40% in Baltimore, MD during the summer, the Phase II sampling was conducted at a
49
50 98 reduced RH of 35% in an attempt to probe the BrC formation process observed in laboratory studies.³⁴
51
52
53
54
55
56
57
58
59
60

1
2
3 99 The experimental setup is shown in Figure 1. Ambient particles were sampled at a flow rate of 12 lpm and
4
5 100 alternated through two different channels (“ambient” and “dry”) every 15 minutes using an automated
6
7 101 three-way valve (Brechtel manufacturing). In the ambient channel, particles passed unperturbed through
8
9 102 a copper tube (1/2” O.D.), while in the dry channel, they passed through a silica gel diffusion dryer.³⁴
10
11 103 Both sample paths were connected to an activated carbon (AC) denuder to remove VOCs and any
12
13 104 evaporated WSOC from the sample stream. Note that RH levels were not measured downstream of the
14
15 105 AC denuder. AC adsorbs water, which can affect the removal of VOCs, especially volatile
16
17 106 hydrocarbons.⁴⁰ However, we do not anticipate RH changes through this component of the experimental
18
19 107 setup since the gas residence time in the denuder (~1 sec) is much shorter than the equilibration timescale
20
21 108 for water vapor adsorption to AC (hours).⁴⁰ Particles were continuously collected into an aqueous
22
23 109 solution via a Particle-into Liquid Sampler (PILS, Brechtel manufacturing).⁴¹ The liquid PILS sample,
24
25 110 which was continuously generated at 1.4 mL min⁻¹, was split and sent to a TOC analyzer (Sievers 900
26
27 111 portable, GE Analytical Instruments) and a liquid waveguide capillary cell (LWCC, WPI, model 3050).
28
29 112 The TOC analyzer was operated in turbo mode and provided WSOC measurements with 4-s resolution.
30
31 113 The LWCC has an effective path length of 50 cm and was coupled to a monochromatic 365 nm UV light
32
33 114 source (Oceanoptics, LLS-LED) and a spectrometer (Oceanoptics, FLAME-S, 200-800 nm) via fiber
34
35 115 optic cables (QP400-2-SR, Oceanoptics). A longer path length of 50 cm in the LWCC provides enhanced
36
37 116 sensitivity to capture the differences in absorbance between the dry and ambient channels.⁴² Since BrC is
38
39 117 operationally defined, multiple approaches have been used in the past to measure BrC across a broad
40
41 118 spectral range of 280 – 700 nm. Studies have defined BrC based upon the absorbance at ~300 nm,^{12,15,34,43}
42
43 119 ~365 nm^{12,20,34,43} or an integrated range between 300-500 nm.^{15,44} A wide range of BrC precursors
44
45 120 (glyoxal, methylglyoxal, glycoaldehyde, hydroxy acetone, etc.) produce chromophores that absorb at
46
47 121 different wavelengths.¹² Speciation of BrC compounds formed from MeGly + AS mixtures resulted in the
48
49 122 majority of BrC compounds exhibiting peak absorption at ~300 nm, and >70% of total absorbance was
50
51 123 observed in the 300-500 nm range.¹⁵ Hecobian et al., reported average absorption in a narrow range of
52
53
54
55
56
57
58
59
60

1
2
3 124 360 – 370 nm as a measure of brown carbon.²⁰ They found that the absorbance at 365 nm was strongly
4
5 125 correlated with the integrated absorption over 300-400 nm, but avoided interferences from non-organic
6
7 126 compounds, as the basis for recommending 365 nm for the BrC measurement. We followed the approach
8
9 127 developed by Hecobian et al., which appears to be in agreement with many previous studies.^{12,34,43}
10
11 128 Absorption spectra were recorded over a range of 350-400 nm using Oceanview spectroscopy software
12
13 129 1.6.3 (Oceanoptics), and the absorbance is reported at 368 ± 5 nm. Absorbance was measured in a semi-
14
15 130 continuous mode where the PILS sample was injected into the LWCC via a syringe pump (Tricontinent,
16
17 131 C3000), providing an absorbance measurement every 15 minutes. The measurements through the dry and
18
19 132 ambient channels were termed $WSOC_p$ ($Abs_{p,368}$) and $WSOC_{p,dry}$ ($Abs_{dry,368}$), respectively. The LWCC
20
21 133 was cleaned with 0.6 M HCl and 18.2 M Ω deionized water every three days during the sampling.²⁰
22
23 134 Dynamic blanks were taken every 24 hours for both the TOC analyzer and LWCC. The ambient
24
25 135 measurements were blank-corrected. Limits of detection (LOD) were calculated as $3 \times$ the standard
26
27 136 deviation of the blanks and were $0.21 \mu\text{g-C m}^{-3}$ and 1.0×10^{-3} A.U., respectively. Calibration details of
28
29 137 the TOC analyzer and LWCC are provided in the supporting information (Text S1). The experimental
30
31 138 setup was tested with a control experiment where the sampling was performed with empty (no silica gel)
32
33 139 dry channel, with one and two dryers installed in the dry channel flow path. We did not find any
34
35 140 difference between the two channels during these control experiments (Figure S1), in agreement with our
36
37 141 past studies (see Fig. S3 in El-Sayed et al. (2016)),³⁹ indicating that differences between the two
38
39 142 measurement channels observed during ambient sampling were the result of physical and/or chemical
40
41 143 changes to the particles induced by aerosol water evaporation.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

145 3. RESULTS AND DISCUSSION

146 3.1 Drying induced BrC formation

147 The comparison of the absorbance measured for the ambient ($Abs_{p,368}$) and dry ($Abs_{dry,368}$) channels are
148 shown in Figure 3. For both dry channel configurations (one or two driers, Figure 3a and 3b,

1
2
3 149 respectively), comparison of $Abs_{p,368}$ and $Abs_{dry,368}$ yielded a slope of 1.00 and an R^2 value of 0.99. This
4
5 150 demonstrates no difference in BrC levels at either of the drying levels. Figure 4 shows a time series of the
6
7 151 absolute difference in absorbance between the two channels ($\Delta Abs_{368} = Abs_{dry,368} - Abs_{p,368}$). The ΔAbs_{368}
8
9 152 values showed no temporal pattern, and no correlation with meteorological conditions or aerosol chemical
10
11 153 components in either of the measurement phases. The mean ΔAbs_{368} values were -1.1×10^{-4} and $1.0 \times$
12
13 154 10^{-4} at 41% and 35% RHs, respectively, which are much lower than the measurement LOD (1.0×10^{-3}
14
15 155 A.U.). As discussed in Section 2, BrC is operationally defined, with measurements of absorbance at
16
17 156 different wavelengths.^{12,15,34,43} It is possible that an enhancement in absorbance at ~ 300 nm due to particle
18
19 157 drying would have been observed in our study. However, based on prior studies that have observed strong
20
21 158 correlations between absorbance at 365 nm and integrated absorbance across 300 – 400 nm, our
22
23 159 observations suggest that there was no generation of BrC in ambient particles undergoing drying.
24
25 160 To investigate further, and evaluate the production of BrC in particles undergoing drying in our
26
27 161 experimental setup, we conducted laboratory experiments with Gly and MeGly in aqueous ammonium
28
29 162 sulfate solutions (10mM + 20mM) similar to Lee et al., 2013.³⁴ The details of the experiments are
30
31 163 discussed in the supporting information (Text S2). Briefly, mixtures of glyoxal/methylglyoxal and
32
33 164 ammonium sulfate were atomized and alternated through the ambient and dry channels while Abs_{368} and
34
35 165 $WSOC_p$ measurements carried out with the same system used for the ambient sampling. The RH through
36
37 166 the bypass (ambient) channel was $\sim 85\%$, while two different RH levels were investigated for the dry
38
39 167 channel. Figure 5 shows the results of the experiments. In the case of Gly, no enhancement in absorbance
40
41 168 was observed when particles were dried to 50% RH, but drying to $\sim 33\%$ RH did produce a measurable
42
43 169 enhancement in absorbance. On the other hand, MeGly showed an enhancement of $\sim 40\%$ in A.U. at
44
45 170 drying to 50% RH. The enhancement further increased to $\sim 70\%$ when dried at $\sim 25\%$ RH. On average, we
46
47 171 found a 20% enhancement (per mass basis) in BrC absorbance for both Gly and MeGly systems when the
48
49 172 RH was reduced from 50% to $\sim 30\%$ in the dry channel. The study by Lee et al., (2013)³⁴ also
50
51 173 implemented moderate drying ($RH \leq 35\%$); however, a direct quantitative comparison with their work is
52
53 174 not possible. The kinetics of BrC formation in bulk solutions of Gly and MeGly with AS can be quite
54
55
56
57
58
59
60

1
2
3 175 slow, as absorption evolves over a period of days.¹¹⁻¹³ The delay between solution preparation and
4
5 176 atomization/particle collection would need to be standardized for such a comparison. The time-dependent
6
7 177 mass absorption coefficient (per C mass basis) is not available, which would add uncertainty to such
8
9 178 comparisons. Further, the RH and drying times would need to be standardized to enable quantitative
10
11 179 comparisons between experiments. However, these experiments demonstrate qualitative agreement with
12
13 180 prior studies that observed BrC formation (or increased rates of BrC formation) in particles undergoing
14
15 181 drying. The results also demonstrate the capability of our experimental setup to generate BrC in particles
16
17 182 undergoing drying, providing an important contrast for the ambient observations.
18
19
20 183 Overall, these results suggest that, even if BrC formation occurs in drying ambient particles, the effect is
21
22 184 quite minor. The most likely cause of this observation is the low concentrations of BrC precursors in
23
24 185 ambient particles. Gly and MeGly are formed in high abundances from the oxidation of biogenic and
25
26 186 aromatic emissions, and are thought to be important secondary BrC precursors.^{12-14,45} Based upon Gly and
27
28 187 MeGly concentrations in ambient particles (each typically in the range of 0 – 50 ng m⁻³),^{23,46,47} we
29
30 188 estimated that they may contribute a combined $\Delta\text{Abs}_{\text{p},368} = 1.0 \times 10^{-4}$ during one drying cycle, which is
31
32 189 smaller than the LOD of the LWCC employed in our study (see Supporting Information for details of this
33
34 190 estimate – Text S3).⁴⁸⁻⁵⁰ In these laboratory experiments, the WSOC_p concentration was reduced by 4%
35
36 191 (Gly) and 39% (MeGly) at 50% RH and 10% (Gly) and 43% (MeGly) at ~30% RH (Table S1). This is in
37
38 192 line with the relative WSOC_p reductions we observed in ambient sampling (Figure S4), and is consistent
39
40 193 with WSOC_p reductions due to drying in prior studies from our group.³⁸ The absolute aerosol
41
42 194 concentrations in the laboratory Gly and MeGly experiments were orders of magnitude (35 to 3600 µg m⁻³
43
44 195 ³) above ambient levels (0.1 to 10 µg m⁻³) when the mM aqueous Gly and MeGly solutions were atomized
45
46 196 (Table S1 and text S3). As discussed above, the laboratory experimental conditions were based upon
47
48 197 prior studies,³⁴ and the high concentrations employed likely contributed to the marked absorption
49
50 198 enhancements not observed in ambient particles.
51
52
53
54 199

200 3.2 WSOC and total absorbance

55
56
57
58
59
60

1
2
3 201 Previous observations during the summer have shown that some fraction of WSOC_p (5 – 30%) evaporates
4
5 202 as a result of particle drying.^{38,39} The average WSOC_{p,dry}/WSOC_p ratio was ~0.90 and 0.84 at RHs of 41%
6
7 203 and 35%, respectively (Figure S4). These results indicate the reversible uptake of WSOC to aerosol water
8
9 204 (reversible aqSOA), in agreement with our previous observations at this site.^{38,39} Gly and MeGly
10
11 205 reversibly partition to aerosol water,^{29,51} and may contribute to the reduced WSOC_p we observe in the dry
12
13 206 channel.³⁹ However, our results suggest that even though some WSOC_p is evaporating in the dry channel,
14
15 207 chemical formation of BrC is negligible.

16
17
18 208 Associations between WSOC_p and BrC can provide insight into primary emissions and secondary
19
20 209 formation processes in a given location.^{19–21} A time series of WSOC_p and Abs_{p,368} are plotted in Figure
21
22 210 S5(a). Overall, the absorbance levels were quite low (average = 0.0045), which reflects the summertime
23
24 211 conditions during which biomass burning influence was low and the source of BrC was predominantly
25
26 212 through secondary formation, consistent with previous studies.²⁰ At times, Abs_{p,368} and WSOC_p were
27
28 213 correlated, but not during periods when the WSOC_p concentration was highest. Even though elevated
29
30 214 WSOC_p was likely due to SOA formation, a process that can also be associated with BrC formation,^{13,43}
31
32 215 other competing processes like photobleaching^{52–54} during the daytime may lead to the loss of secondary
33
34 216 BrC.

35
36
37 217 The diurnal trends of WSOC_p and Abs_{p,368} are shown in Figure S5(b). The WSOC_p profile is relatively
38
39 218 flat, while absorbance appears to fall by ~ 60% in the early morning hours (00:00 to 8:00 am). A shoulder
40
41 219 is observed for both WSOC_p and absorbance during the morning rush-hour, suggesting an influence of
42
43 220 traffic.²¹ The mid-day peak of WSOC_p indicates photochemical SOA formation as a major contributor to
44
45 221 diurnal variability of WSOC_p.⁵⁵ Abs_{p,368} has a similar afternoon peak, but appears to lag behind WSOC_p,
46
47 222 suggesting the BrC formed is from secondary processes.²⁰ The lag may be due to slower kinetics of BrC
48
49 223 formation or photobleaching during mid-day hours when the actinic flux is greatest.¹² Organonitrogen
50
51 224 compounds, often formed from reactions of organic molecules with NH₄⁺ or NH₃, contribute greatly to
52
53 225 BrC levels.^{15,27} The diurnal profiles of BrC, NH₄⁺ and NH₃ are compared in Figure S6. The average
54
55 226 concentration of NH₄⁺ does not fluctuate much throughout the day, consistent with the regional nature and
56
57
58
59
60

1
2
3 227 the strong influence of sulfate on NH_4^+ concentrations in the eastern U.S.⁵⁶ Interestingly, the NH_3
4
5 228 concentration appears to be anti-correlated with BrC; however, it is unlikely that this relationship is
6
7 229 causative. The gas-phase NH_3 concentration is driven by many factors that have not been directly linked
8
9 230 to secondary BrC formation, including emissions, gas-particle partitioning, boundary layer dynamics,
10
11 231 aerosol liquid water content, and particle acidity.⁵⁷ The magnitude of concentration changes ($2\text{-}3 \mu\text{g m}^{-3}$
12
13 232 for NH_3 vs. $7\text{-}8 \times 10^{-3}$ A.U. for $\text{Abs}_{\text{p},368}$) further suggests that the apparent anti-correlation does not
14
15 233 indicate a causal relationship. Future studies may be required to confirm this hypothesis.
16
17
18 234

19 20 235 4. CONCLUSIONS

21
22
23 236 This represents, to our knowledge, the first study to characterize BrC formation in ambient particles
24
25 237 undergoing drying. We did not find evidence of BrC formation in ambient particles dried to $\sim 35\%$ RH,
26
27 238 lower than almost all ambient RH values observed at the surface during our study. The reason for this
28
29 239 observation is most likely the low concentrations of BrC precursors in ambient particles. The low
30
31 240 concentration of BrC precursor chemicals may not be a constraint in fogs and clouds. Liquid water
32
33 241 content in fog and clouds is several orders of magnitude higher than the ambient particles, enabling
34
35 242 significantly higher uptake (mass basis) of BrC producing chemicals.^{58,59} The drying of fog and cloud
36
37 243 droplets can lead to an extremely concentrated mixture that may form BrC.^{32,60} Multiple cycles of this
38
39 244 processing may lead to a buildup of BrC in an air mass, which may ultimately alter the overall
40
41 245 atmospheric BrC budget. Alternately, cloud processing in strong convective systems may be a significant
42
43 246 source of free-tropospheric BrC,²⁴ though the overall contribution of cloud processing to global BrC
44
45 247 levels remains highly uncertain.^{1,61,62} Future studies should also characterize the lifetime of secondary
46
47 248 BrC formed through drying processes, since photobleaching represents a major sink for BrC, reducing its
48
49 249 direct radiative effect (DRE).^{7,8,63} The photochemical lifetime of ambient BrC (dominated by biomass
50
51 250 burning) is estimated to be ~ 1 day, though laboratory experiments have shown that secondary BrC may
52
53 251 have shorter lifetimes.^{64,65} A significant amount of primary BrC from biomass burning (including
54
55
56
57
58
59
60

1
2
3 252 residential wood burning) is emitted in wintertime.²¹ Therefore, the summertime was appropriate to
4
5 253 investigate drying induced BrC formation when the contributions of biomass burning were minimal. The
6
7 254 effects of aerosol liquid water uptake and evaporation on primary BrC emitted from biomass burning
8
9 255 have been investigated in conjunction with photolysis (photobleaching).⁶² However, studies isolating the
10
11 256 effects of humidification and drying on the transformation of primary BrC emitted from biomass burning
12
13
14 257 have not been performed and need to be evaluated in future studies.

15
16 258 ASSOCIATED CONTENT

17
18 259 Supporting information

19
20 260 CONFLICT OF INTEREST

21
22 261 Authors declare no competing financial interests.

23
24 262 AUTHOR INFORMATION

25
26 263 Corresponding Author

27
28 264 *Phone: (410) 455-3515; fax: (410) 455-6500; e-mail: hennigan@umbc.edu.

29
30 265 ACKNOWLEDGMENTS

31
32 266 This work was funded by the National Science Foundation through awards AGS-1719252 and AGS-
33
34 267 1719245.

35
36 268
37
38 269 REFERENCES

- 39
40
41
42 270 1 A. Laskin, J. Laskin and S. A. Nizkorodov, Chemistry of Atmospheric Brown Carbon, *Chem.*
43
44 271 *Rev.*, 2015, **115**, 4335–4382.
- 45
46
47 272 2 T. W. Kirchstetter, T. Novakov and P. V. Hobbs, Evidence that the spectral dependence of light
48
49 273 absorption by aerosols is affected by organic carbon, *J. Geophys. Res. D Atmos.*, 2004, **109**, 1–12.
- 50
51
52 274 3 T. C. Bond and R. W. Bergstrom, Light absorption by carbonaceous particles: An investigative
53
54 275 review, *Aerosol Sci. Technol.*, 2006, **40**, 27–67.
- 55
56
57
58
59
60

- 1
2
3 276 4 J. Yan, X. Wang, P. Gong, C. Wang and Z. Cong, Review of brown carbon aerosols: Recent
4
5 277 progress and perspectives, *Sci. Total Environ.*, 2018, **634**, 1475–1485.
6
7
8 278 5 J. Liu, E. Scheuer, J. Dibb, L. D. Ziemba, K. L. Thornhill, B. E. Anderson, A. Wisthaler, T.
9
10 279 Mikoviny, J. J. Devi, M. Bergin and R. J. Weber, Brown carbon in the continental troposphere,
11
12 280 *Geophys. Res. Lett.*, 2014, **41**, 2191–2195.
13
14
15 281 6 C. E. Chung, V. Ramanathan and D. Decremet, Observationally constrained estimates of
16
17 282 carbonaceous aerosol radiative forcing, *Proc. Natl. Acad. Sci.*, 2012, **109**, 11624–11629.
18
19
20 283 7 X. Wang, C. L. Heald, J. Liu, R. J. Weber, P. Campuzano-jost, J. L. Jimenez, J. P. Schwarz and A.
21
22 284 E. Perring, Exploring the observational constraints on the simulation of brown carbon, 2018, 635–
23
24 285 653.
25
26
27 286 8 H. Brown, X. Liu, Y. Feng, Y. Jiang, M. Wu, Z. Lu and C. Wu, Radiative effect and climate
28
29 287 impacts of brown carbon with the Community Atmosphere Model (CAM5), 2018, 17745–17768.
30
31
32 288 9 Y. Feng, V. Ramanathan and V. R. Kotamarthi, Brown carbon: A significant atmospheric absorber
33
34 289 of solar radiation, *Atmos. Chem. Phys.*, 2013, **13**, 8607–8621.
35
36
37 290 10 P. Lin, P. K. Aiona, Y. Li, M. Shiraiwa, J. Laskin, S. A. Nizkorodov and A. Laskin, Molecular
38
39 291 Characterization of Brown Carbon in Biomass Burning Aerosol Particles, *Environ. Sci. Technol.*,
40
41 292 2016, **50**, 11815–11824.
42
43
44 293 11 C. J. Kampf, R. Jakob and T. Hoffmann, Identification and characterization of aging products in
45
46 294 the glyoxal/ammonium sulfate system - Implications for light-absorbing material in atmospheric
47
48 295 aerosols, *Atmos. Chem. Phys.*, 2012, **12**, 6323–6333.
49
50
51 296 12 M. H. Powelson, B. M. Espelien, L. N. Hawkins, M. M. Galloway and D. O. De Haan, Brown
52
53 297 carbon formation by aqueous-phase carbonyl compound reactions with amines and ammonium
54
55
56
57
58
59
60

- 1
2
3 298 sulfate, *Environ. Sci. Technol.*, 2014, **48**, 985–993.
4
5
6 299 13 N. Sareen, A. N. Schwier, E. L. Shapiro, D. Mitroo and V. F. McNeill, Secondary organic material
7
8 300 formed by methylglyoxal in aqueous aerosol mimics, *Atmos. Chem. Phys.*, 2010, **10**, 997–1016.
9
10
11 301 14 M. M. Galloway, M. H. Powelson, N. Sedehi, S. E. Wood, K. D. Millage, J. A. Kononenko, A. D.
12
13 302 Rynaski and D. O. De Haan, Secondary organic aerosol formation during evaporation of droplets
14
15 303 containing atmospheric aldehydes, amines, and ammonium sulfate, *Environ. Sci. Technol.*, 2014,
16
17 304 **48**, 14417–14425.
18
19
20 305 15 P. Lin, J. Laskin, S. A. Nizkorodov and A. Laskin, Revealing Brown Carbon Chromophores
21
22 306 Produced in Reactions of Methylglyoxal with Ammonium Sulfate, *Environ. Sci. Technol.*, 2015,
23
24 307 **49**, 14257–14266.
25
26
27 308 16 P. Zotter, I. El-haddad, Y. Zhang, P. L. Hayes, X. Zhang, Y. Lin, L. Wacker, J. Schnelle-kreis, G.
28
29 309 Abbaszade, R. Zimmermann, J. D. Surratt, R. Weber, J. L. Jimenez, S. Szidat, U. Baltensperger
30
31 310 and A. S. H. Prévôt, *Journal of Geophysical Research : Atmospheres*, 2014, 1–18.
32
33
34 311 17 T. B. Nguyen, A. Laskin, J. Laskin and S. A. Nizkorodov, Brown carbon formation from
35
36 312 ketoaldehydes of biogenic monoterpenes, *Faraday Discuss.*, 2013, **165**, 473–494.
37
38
39 313 18 A. Hoffer, G. Kiss, M. Blazsó and A. Gelencsér, Chemical characterization of humic-like
40
41 314 substances (HULIS) formed from a lignin-type precursor in model cloud water, *Geophys. Res.*
42
43 315 *Lett.*, , DOI:10.1029/2003gl018962.
44
45
46
47 316 19 X. Zhang, Y. H. Lin, J. D. Surratt and R. J. Weber, Sources, composition and absorption Ångström
48
49 317 exponent of light-absorbing organic components in aerosol extracts from the los angeles basin,
50
51 318 *Environ. Sci. Technol.*, 2013, **47**, 3685–3693.
52
53
54 319 20 A. Hecobian, X. Zhang, M. Zheng, N. Frank, E. S. Edgerton and R. J. Weber, Water-soluble
55
56
57
58
59
60

- 1
2
3 320 organic aerosol material and the light-absorption characteristics of aqueous extracts measured over
4
5 321 the Southeastern United States, *Atmos. Chem. Phys.*, 2010, **10**, 5965–5977.
6
7
8 322 21 X. Zhang, Y. H. Lin, J. D. Surratt, P. Zotter, A. S. H. Prévôt and R. J. Weber, Light-absorbing
9
10 323 soluble organic aerosol in Los Angeles and Atlanta: A contrast in secondary organic aerosol,
11
12 324 *Geophys. Res. Lett.*, 2011, **38**, 2–5.
13
14
15 325 22 J. Liu, M. Bergin, H. Guo, L. King, N. Kotra, E. Edgerton and R. J. Weber, Size-resolved
16
17 326 measurements of brown carbon in water and methanol extracts and estimates of their contribution
18
19 327 to ambient fine-particle light absorption, *Atmos. Chem. Phys.*, 2013, **13**, 12389–12404.
20
21
22 328 23 K. Mitsuishi, M. Iwasaki, M. Takeuchi, H. Okochi, S. Kato, S. I. Ohira and K. Toda, Diurnal
23
24 329 Variations in Partitioning of Atmospheric Glyoxal and Methylglyoxal between Gas and Particles
25
26 330 at the Ground Level and in the Free Troposphere, *ACS Earth Sp. Chem.*, 2018, **2**, 915–924.
27
28
29 331 24 Y. Zhang, H. Forrister, J. Liu, J. Dibb, B. Anderson, J. P. Schwarz, A. E. Perring, J. L. Jimenez, P.
30
31 332 Campuzano-Jost, Y. Wang, A. Nenes and R. J. Weber, Top-of-atmosphere radiative forcing
32
33 333 affected by brown carbon in the upper troposphere, *Nat. Geosci.*, 2017, **10**, 486.
34
35
36 334 25 M. O. Andreae and a. Gelencsér, Black carbon or brown carbon? The nature of light-absorbing
37
38 335 carbonaceous aerosols, *Atmos. Chem. Phys.*, 2006, **6**, 3131–3148.
39
40
41
42 336 26 E. R. Graber and Y. Rudich, Atmospheric HULIS: How humic-like are they? A comprehensive
43
44 337 and critical review, *Atmos. Chem. Phys.*, 2006, **6**, 729–753.
45
46
47 338 27 K. M. Updyke, T. B. Nguyen and S. A. Nizkorodov, Formation of brown carbon via reactions of
48
49 339 ammonia with secondary organic aerosols from biogenic and anthropogenic precursors, *Atmos.*
50
51 340 *Environ.*, 2012, **63**, 22–31.
52
53
54 341 28 A. Limbeck, M. Kulmala and H. Puxbaum, Secondary organic aerosol formation in the
55
56
57
58
59
60

- 1
2
3 342 atmosphere via heterogeneous reaction of gaseous isoprene on acidic particles, *Geophys. Res.*
4
5 343 *Lett.*, 2003, **30**, 4–7.
6
7
8 344 29 D. O. De Haan, a L. Corrigan, M. a Tolbert, J. L. Jimenez, S. E. Wood and J. J. Turley, Secondary
9
10 345 organic aerosol formation by self-reactions of methylglyoxal and glyoxal in evaporating droplets,
11
12 346 *Environ. Sci. Technol.*, 2009, **43**, 8184–8190.
13
14
15 347 30 G. Yu, A. R. Bayer, M. M. Galloway, K. J. Korshavn, C. G. Fry and F. N. Keutsch, Glyoxal in
16
17 348 aqueous ammonium sulfate solutions: Products, kinetics and hydration effects, *Environ. Sci.*
18
19 349 *Technol.*, 2011, **45**, 6336–6342.
20
21
22 350 31 M. Teich, D. Van Pinxteren, S. Kecorius, Z. Wang and H. Herrmann, First Quantification of
23
24 351 Imidazoles in Ambient Aerosol Particles: Potential Photosensitizers, Brown Carbon Constituents,
25
26 352 and Hazardous Components, *Environ. Sci. Technol.*, 2016, **50**, 1166–1173.
27
28
29 353 32 D. O. De Haan, L. N. Hawkins, H. G. Welsh, R. Pednekar, J. R. Casar, E. A. Pennington, A. De
30
31 354 Loera, N. G. Jimenez, M. A. Symons, M. Zauscher, A. Pajunoja, L. Caponi, M. Cazaunau, P.
32
33 355 Formenti, A. Gratien, E. Pangui and J. F. Doussin, Brown Carbon Production in Ammonium- or
34
35 356 Amine-Containing Aerosol Particles by Reactive Uptake of Methylglyoxal and Photolytic Cloud
36
37 357 Cycling, *Environ. Sci. Technol.*, 2017, **51**, 7458–7466.
38
39
40
41 358 33 T. B. Nguyen, P. B. Lee, K. M. Updyke, D. L. Bones, J. Laskin, A. Laskin and S. A. Nizkorodov,
42
43 359 Formation of nitrogen- and sulfur-containing light-absorbing compounds accelerated by
44
45 360 evaporation of water from secondary organic aerosols, *J. Geophys. Res. Atmos.*, 2012, **117**, 1–14.
46
47
48 361 34 A. K. Y. Lee, R. Zhao, R. Li, J. Liggio, S.-M. Li and J. P. D. Abbatt, Formation of Light
49
50 362 Absorbing Organo-Nitrogen Species from Evaporation of Droplets Containing Glyoxal and
51
52 363 Ammonium Sulfate, *Environ. Sci. Technol.*, 2013, **47**, 12819–12826.
53
54
55 364 35 M. A. Battaglia, S. Douglas and C. J. Hennigan, Effect of the Urban Heat Island on Aerosol pH,
56
57
58
59
60


- 1
2
3 365 *Environ. Sci. Technol.*, 2017, **51**, 13095–13103.
4
5
6 366 36 H. R. Pruppacher, ed. W. Jaeschke, Springer Berlin Heidelberg, Berlin, Heidelberg, 1986, pp.
7
8 367 133–190.
9
10
11 368 37 M. M. H. El-Sayed, Y. Wang and C. J. Hennigan, Direct atmospheric evidence for the irreversible
12
13 369 formation of aqueous secondary organic aerosol, *Geophys. Res. Lett.*, 2015, **42**, 5577–5586.
14
15
16 370 38 M. M. H. El-Sayed, D. L. Ortiz-montalvo and C. J. Hennigan, The effects of isoprene and NO_x on
17
18 371 secondary organic aerosols formed through reversible and irreversible uptake to aerosol water,
19
20 372 *Atmos. Chem. Phys.*, 2018, 1171–1184.
21
22
23 373 39 M. M. H. El-Sayed, D. Amenumey and C. J. Hennigan, Drying-Induced Evaporation of Secondary
24
25 374 Organic Aerosol during Summer, *Environ. Sci. Technol.*, 2016, **50**, 3626–3633.
26
27
28 375 40 J. E. Urbanic, E. S. Oswald, N. J. Wagner and H. E. Moore, Effect of Humid Purge Air on the
29
30 376 Performance of Commercial Activated Carbons Used for Evaporative Emission Control, *SAE*
31
32 377 *Trans.*, 1989, **98**, 723–730.
33
34
35 378 41 D. A. Orsini, Y. Ma, A. Sullivan, B. Sierau, K. Baumann and R. J. Weber, Refinements to the
36
37 379 particle-into-liquid sampler (PILS) for ground and airborne measurements of water soluble aerosol
38
39 380 composition, *Atmos. Environ.*, 2003, **37**, 1243–1259.
40
41
42 381 42 L. J. Gimbert and P. J. Worsfold, Environmental applications of liquid-waveguide-capillary cells
43
44 382 coupled with spectroscopic detection, *TrAC - Trends Anal. Chem.*, 2007, **26**, 914–930.
45
46
47 383 43 E. L. Shapiro, J. Szprengiel, N. Sareen, C. N. Jen, M. R. Giordano and V. F. McNeill, Light-
48
49 384 absorbing secondary organic material formed by glyoxal in aqueous aerosol mimics, *Atmos. Chem.*
50
51 385 *Phys.*, 2009, **9**, 2289–2300.
52
53
54 386 44 S. M. Phillips and G. D. Smith, Spectroscopic comparison of water- and methanol-soluble brown
55
56
57
58
59
60

- 1
2
3 387 carbon particulate matter, *Aerosol Sci. Technol.*, 2017, **51**, 1113–1121.
4
5
6 388 45 D. O. De Haan, E. Tapavicza, M. Riva, T. Cui, J. D. Surratt, A. C. Smith, M.-C. Jordan, S.
7
8 389 Nilakantan, M. Almodovar, T. N. Stewart, A. de Loera, A. C. De Haan, M. Cazaunau, A. Gratien,
9
10 390 E. Pangui and J.-F. Doussin, Nitrogen-Containing, Light-Absorbing Oligomers Produced in
11
12 391 Aerosol Particles Exposed to Methylglyoxal, Photolysis, and Cloud Cycling, *Environ. Sci.*
13
14 392 *Technol.*, 2018, **52**, 4061–4071.
15
16
17 393 46 R. Ortiz, S. Shimada, K. Sekiguchi, Q. Wang and K. Sakamoto, Measurements of changes in the
18
19 394 atmospheric partitioning of bifunctional carbonyls near a road in a suburban area, *Atmos. Environ.*,
20
21 395 2013, **81**, 554–560.
22
23
24 396 47 N. Sareen, E. M. Waxman, B. J. Turpin, R. Volkamer and A. G. Carlton, Potential of Aerosol
25
26 397 Liquid Water to Facilitate Organic Aerosol Formation: Assessing Knowledge Gaps about
27
28 398 Precursors and Partitioning, *Environ. Sci. Technol.*, 2017, **51**, 3327–3335.
29
30
31 399 48 Y. Chen and T. C. Bond, Light absorption by organic carbon from wood combustion, *Atmos.*
32
33 400 *Chem. Phys.*, 2010, **10**, 1773–1787.
34
35
36 401 49 H. Sun, L. Biedermann and T. C. Bond, Color of brown carbon: A model for ultraviolet and
37
38 402 visible light absorption by organic carbon aerosol, *Geophys. Res. Lett.*, 2007, **34**, 1–5.
39
40
41 403 50 J. P. S. Wong, M. Tsagaraki, I. Tsiotra, N. Mihalopoulos, K. Violaki, M. Kanakidou, J. Sciare, A.
42
43 404 Nenes and R. J. Weber, Atmospheric Evolution of Molecular Weight Separated Brown Carbon
44
45 405 from Biomass Burning, *Atmos. Chem. Phys.*, 2019, **19**, 7319–7334.
46
47
48
49 406 51 M. M. Galloway, P. S. Chhabra, A. W. H. Chan, J. D. Surratt, R. C. Flagan, J. H. Seinfeld and F.
50
51 407 N. Keutsch, Glyoxal uptake on ammonium sulphate seed aerosol: Reaction products and
52
53 408 reversibility of uptake under dark and irradiated conditions, *Atmos. Chem. Phys.*, 2009, **9**, 3331–
54
55 409 3345.
56
57
58
59
60

- 1
2
3 410 52 J. A. Faust, J. P. S. Wong, A. K. Y. Lee and J. P. D. Abbatt, Role of Aerosol Liquid Water in
4
5 411 Secondary Organic Aerosol Formation from Volatile Organic Compounds, *Environ. Sci. Technol.*,
6
7 412 2017, **51**, 1405–1413.
8
9
10 413 53 H. J. Lee, P. K. Aiona, A. Laskin, J. Laskin and S. A. Nizkorodov, Effect of solar radiation on the
11
12 414 optical properties and molecular composition of laboratory proxies of atmospheric brown carbon,
13
14 415 *Environ. Sci. Technol.*, 2014, **48**, 10217–10226.
15
16
17 416 54 R. Zhao, A. K. Y. Lee, L. Huang, X. Li, F. Yang and J. P. D. Abbatt, Photochemical processing of
18
19 417 aqueous atmospheric brown carbon, *Atmos. Chem. Phys.*, 2015, **15**, 6087–6100.
20
21
22 418 55 C. J. Hennigan, M. H. Bergin, A. G. Russell, A. Nenes and R. J. Weber, Gas/particle partitioning
23
24 419 of water-soluble organic aerosol in Atlanta, *Atmos. Chem. Phys.*, 2009, **9**, 3613–3628.
25
26
27 420 56 R. J. Weber, H. Guo, A. G. Russell and A. Nenes, High aerosol acidity despite declining
28
29 421 atmospheric sulfate concentrations over the past 15 years, *Nat. Geosci.*, 2016, **9**, 282–285.
30
31
32 422 57 R. Allen, L. T. Myles and M. W. Heuer, Ambient ammonia in terrestrial ecosystems: A
33
34 423 comparative study in the Tennessee Valley, USA, *Sci. Total Environ.*, 2011, **409**, 2768–2772.
35
36
37 424 58 H. J. Lim, A. G. Carlton and B. J. Turpin, Isoprene forms secondary organic aerosol through cloud
38
39 425 processing: Model simulations, *Environ. Sci. Technol.*, 2005, **39**, 4441–4446.
40
41
42 426 59 E. A. Betterton and M. R. Hoffmann, Henry's Law Constants of Some Environmentally Important
43
44 427 Aldehydes, *Environ. Sci. Technol.*, 1988, **22**, 1415–1418.
45
46
47 428 60 H. Kim, S. Collier, X. Ge, J. Xu, Y. Sun, W. Jiang, Y. Wang, P. Herckes and Q. Zhang, Chemical
48
49 429 processing of water-soluble species and formation of secondary organic aerosol in fogs, *Atmos.*
50
51 430 *Environ.*, 2019, **200**, 158–166.
52
53
54 431 61 Y. Desyaterik, Y. Sun, X. Shen, T. Lee, X. Wang, T. Wang and J. L. Collett, Speciation of

- 1
2
3 432 'brown' carbon in cloud water impacted by agricultural biomass burning in eastern China, *J.*
4
5 433 *Geophys. Res. Atmos.*, 2013, **118**, 7389–7399.
6
7
8 434 62 T. Moise, J. M. Flores and Y. Rudich, Optical Properties of Secondary Organic Aerosols and Their
9
10 435 Changes by Chemical Processes, *Chem. Rev.*, 2015, **115**, 4400–4439.
11
12
13 436 63 J. P. S. Wong, A. Nenes and R. J. Weber, Changes in Light Absorptivity of Molecular Weight
14
15 437 Separated Brown Carbon Due to Photolytic Aging, *Environ. Sci. Technol.*, 2017, **51**, 8414–8421.
16
17
18 438 64 H. Forrister, J. Liu, E. Scheuer, J. Dibb, L. Ziemba, K. L. Thornhill, B. Anderson, G. Diskin, A. E.
19
20 439 Perring, J. P. Schwarz, P. Campuzano-Jost, D. A. Day, B. B. Palm, J. L. Jimenez, A. Nenes and R.
21
22 440 J. Weber, Evolution of brown carbon in wildfire plumes, *Geophys. Res. Lett.*, 2015, **42**, 4623–
23
24 441 4630.
25
26
27 442 65 Q. Wang, J. Saturno, X. Chi, D. Walter, J. V. Lavric, D. Moran-Zuloaga, F. Ditas, C. Pöhlker, J.
28
29 443 Brito, S. Carbone, P. Artaxo and M. O. Andreae, Modeling investigation of light-absorbing
30
31 444 aerosols in the Amazon Basin during the wet season, *Atmos. Chem. Phys.*, 2016, **16**, 14775–
32
33 445 14794.
34
35
36
37 446
38
39
40 447
41
42 448
43
44 449
45
46 450
47
48 451
49
50 452
51
52 453
53
54 454 **Figures**
55
56
57
58
59
60

455



456

457 **Figure 1:** Schematic of the sampling setup. Note that one dryer was used in the dry channel during
458 phase-I (summer 2018), while two dryers were used during phase-II (summer 2019).

459






Figure 2: Hourly ambient relative humidity levels during a) phase-I (17-July-2018 – 31-Aug-2018), and b) phase-II (13-Aug-2019 – 23-Aug-2019) of the study. The dashed line shows the 41% RH maintained in the dry channel during phase-I sampling, and the gray shaded region represents the average $\pm 1\sigma$ RH range of 37-45%. Note, only ~2% hourly data fell below 41% RH. In phase – II sampling, the average drying RH was 35% (dashed line), and the shaded region is average $\pm 1\sigma$ RH range of 29-41%. The data were downloaded from NOAA’s Baltimore Washington International airport meteorology station, located 7 km south of UMBC (<https://www.ncdc.noaa.gov/cdo-web/search?datasetid=GHCND>).



473



474

475 **Figure 3:** Comparison of the ambient and dry channel absorbance measurements during (a) phase-I
476 sampling (dry channel RH = 41%) conducted between 17-July-2018 – 31-Aug-2018, and (b) phase-II
477 sampling (dry channel RH = 35%) conducted between 13-Aug-2019 – 23-Aug-2019. The 1:1 line (green
478 dashed line) is shown for visual reference, and the solid black line is the linear fit to the data using least-
479 squares regression analysis.

480

481


482

483


484

485

480




481



482


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



483

484 **Figure 4:** (a, b) Time series of ΔAbs_{368} ($\Delta\text{Abs}_{368} = \text{Abs}_{\text{dry},368} - \text{Abs}_{\text{p},368}$) during the phase – I and
485 phase – II sampling periods, respectively. Note the broken x-axis in panel a) corresponds to a ~10 day
486 period when the instrument was off-line for maintenance. No difference in BrC concentration was
487 observed at both RH levels. (c) Box plot of ΔAbs_{368} ($\Delta\text{Abs}_{368} = \text{Abs}_{\text{dry},368} - \text{Abs}_{\text{p},368}$) of ambient
488 samples collected at two different RHs – 41% and 35%.

489



490

491 **Figure 5:** The ratio of dry to bypass MAC is shown for Gly and MeGly experiments conducted at
492 different RH levels (RH of the dry channel is given in the figure as percentages). The dashed black
493 line is $y = 1$ line for visual guidance.

494

495

496

497

498

499

500


501

502

503

1 **Table of Contents Text and Graphic:**

2 This work shows, for the first time, that brown carbon formation is not observed in ambient
3 particles undergoing drying.



4

5