



ChemComm

Amethyrin-type expanded porphyrins that displays anti-aromatic character upon protonation

Journal:	<i>ChemComm</i>
Manuscript ID	CC-COM-06-2020-004400.R1
Article Type:	Communication

SCHOLARONE™
Manuscripts

COMMUNICATION

Amethyrin-type expanded porphyrins that displays anti-aromatic character upon protonation

Received 00th January 20xx,
Accepted 00th January 20xx

Harrison D. Root,[†] Daniel N. Mangel,[†] James T. Brewster II,[†] Hadiqa Zafar, Adam Samia, Graeme Henkelman,* and Jonathan L. Sessler*

DOI: 10.1039/x0xx00000x

The use of protonation to switch nonaromatic expanded porphyrins to their corresponding anti-aromatic forms has not been widely explored. Here, we show that free-base pyriamethyrin and dipyriamethyrin display nonaromatic character, as inferred from NMR spectroscopic analyses, their optical properties, and theoretical calculations. Addition of two protons extends the π -conjugation of these amethyrin analogues and yields formally anti-aromatic systems.

Aromaticity remains one of the most fundamental concepts in organic chemistry. Understanding the relationship between aromatic, non-aromatic, and anti-aromatic states is thus of widespread interest, as are mechanisms that can allow switching between these various limiting electronic forms. Playing a central role in these efforts are porphyrins and their analogues (collectively "porphyrinoids"). These macrocycles have proven indispensable in the context of biomedical studies, as well as material applications.^{1–3} Expanded porphyrins are a particularly interesting class of porphyrinoids that have played a key role in advancing our understanding of aromaticity due to their ability to adopt a wide range of conformations and oxidation states.⁴ Large expanded porphyrins have previously been used to access various aromatic and anti-aromatic forms, including twisted Hückel and Möbius aromatic species.^{5,6} Metalation and proton coupled electron transfer have also been used to control the electronic states of expanded porphyrins. Much less explored is the effect of simple protonation to achieve switching between non-, anti-, and aromatic forms. However, so-called pyriporphyrin derivatives have shown the ability to switch between nonaromatic and aromatic forms upon protonation.^{7,8} Additionally, some years ago we showed that cyclo[2]pyridine[4]pyrrole could be converted from its neutral non-aromatic form to the corresponding anti-aromatic species by means of diprotonation.⁹ However, to our knowledge these findings have yet to be generalized. Here we report that pyriamethyrin (**1**) and

dipyriamethyrin (**3**, **4**), analogues of amethyrin (**2**) (Figure 1), are readily converted to anti-aromatic species upon protonation with methane sulfonic acid. We have also found that the inherent anti-aromaticity of **2** is increased upon protonation.




Figure 1. Chemical structures of pyriamethyrin (**1**), amethyrin (**2**), and the dipyriamethyrins (**3** and **4**) considered in the present study.

Amethyrin and its analogues (Figure 1) have historically been of interest due to their coordination chemistry, as well as their ability to switch between various oxidation states upon metalation.^{10–16} Recent work has brought to light how the inherent properties of this framework can be modified by introducing pyridine moieties into the terpyrrolic subunit.¹⁷ This appreciation builds off of previous studies that implicate the pyridine unit as playing a key role in mediating the metal coordination chemistry relative to the parent hexapyrrolic species.^{18–21} An open question was thus whether protonation could serve to endow amethyrin and its pyridine-containing analogues with additional conjugation pathways extending over the entirety of the periphery, thereby "opening up" global (anti)aromaticity for the system as a whole.^{9,22} As detailed below we have now found that several pyridine-containing macrocycles, namely the asymmetric amethyrin (pyriamethyrin **1**) and two dipyriamethyrins (**3** and **4**), can be converted to the corresponding formal anti-aromatic forms by means of simple diprotonation (cf. e.g., Figure 2). Similar, but not identical effects, are seen in the case of amethyrin **2**.




Figure 2. Addition of 2 equivalents of methanesulfonic acid (MSA) yields the diprotonated anti-aromatic species **1b**.

Department of Chemistry, The University of Texas at Austin, 105 East 24th St., Stop A5300, Austin, USA

E-mail: ssessler@cm.utexas.edu; henkelman@utexas.edu

[†] These authors contributed equally.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

Initial UV-Vis spectroscopic studies of **1** (Figure 3) revealed a spectrum reminiscent of other amethyrin macrocycles with a strong Soret-like band at 452 nm ($\epsilon = 43162 \text{ M}^{-1} \text{ cm}^{-1}$) and two N-like bands at 373 nm and 285 nm ($\epsilon = 16721 \text{ M}^{-1} \text{ cm}^{-1}$ and $\epsilon = 13205 \text{ M}^{-1} \text{ cm}^{-1}$, respectively). Evidence that protonation leads to π -conjugation that extends throughout the macrocycle came from a UV-Vis spectroscopic titration of **1** with methane sulfonic acid (MSA) (Figure 3). Upon the addition of increasing quantities of MSA, a visible colour change from yellow to orange-red was observed. This colour change is accompanied by a red shift and an increase in the intensity of the Soret-like band (452 \rightarrow 490 nm). A growing in of new N-type absorption features at 300 nm and 410 nm, as well as a Q-type band at 577 nm, is also seen. These spectral changes are rationalized in terms of forming the diprotonated species **1b** (Figure 2).




Figure 3. UV-Vis spectral titration of **1** with MSA in CHCl_3 . The red shift in the Soret-like band and the appearance of Q-type features was taken as initial evidence of extended π -conjugation.

Macrocycles **2** and **3** also showed changes in their UV-Vis spectral features upon exposure to MSA. Upon titration with MSA, solutions of **2** in CHCl_3 a dramatic increase in the intensity of the Soret-like band at 488 nm is seen, along with the appearance of a new Q-type band at 591 nm (ESI Figure S1). The UV-Vis spectrum of macrocycle **4** revealed a red shift and an increase in the intensity of the Soret-like band (470 nm \rightarrow 516 nm), as well as the growing in of an N-type feature at 381 nm. Additionally, Q-type bands are seen at 592 nm and 646 nm upon treatment with MSA (ESI Figure S2). Again, these spectral changes were attributed to the formation of the diprotonated form of **4** and interpreted in terms of an effective expansion of the π -conjugation pathways present within this and the other amethyrin-type macrocycles considered in this study.

Further support for a proton-coupled expansion of the π -conjugation periphery came from ^1H NMR spectral studies. The free-base form of pyriamethyrin **1** in CDCl_3 displays pyrrole NH resonances that match those seen for a number of other nonaromatic porphyrinoid cores (i.e., $\delta = 10.14$ ppm, Figure 4). In contrast, upon addition of two equivalents of $\text{MSA-}d_4$, these signals shift downfield to 15.5 ppm, indicating a more strongly deshielded environment. Intermediate shift behaviour is seen upon addition of only 1 equivalent of MSA with a splitting of the pyrrole NH resonances also being observed. This latter splitting is likely due to

the asymmetry of the pyrrole NH protons that is induced upon addition of only one equivalent of MSA to **1**. However, once two equivalents of MSA are added (forming diprotonated **1b**) the NH resonances coalesce back to a single peak. This could reflect a fast exchange process or regeneration of a symmetric structure. We consider the latter explanation more likely. The downfield shift of the pyrrole NH resonances is consistent with the presence of an anti-aromatic paratropic ring current wherein the protons within the ring exist in a less shielded environment and therefore experience a downfield shift in the ^1H NMR spectrum.




Figure 4. NMR spectral titration of **1** in CDCl_3 with $\text{MSA-}d_4$. The downfield shift in the pyrrole NH resonance seen upon diprotonation is taken as evidence for conversion to an antiaromatic form.

The free-base form of amethyrin **2** displayed pyrrole NH resonances (12.10 ppm and 12.82 ppm) that are similar to what is seen in the case of **1**. Upon protonation, these pyrrole NH signals shift far downfield to 19.9 ppm (ESI Figure S3). Again, this was taken as evidence that the diprotonated form of **2** displays anti-aromatic character. As true for **1**, the addition of 1 equivalent of MSA to **3** gives rise to splitting in this signal that coalesces back to a singlet upon the addition of 2 equivalents; however, the spectra is more convoluted, making discrimination between discrete protonation states difficult. Dipyrriamethyrin **3** displays the same trend as observed for **1** and **2** (ESI Figure S4). The pyrrole NH resonances of the neutral form appear at 6.91 ppm (a chemical shift typical for non-aromatic systems). As above, the addition of 1 equivalent of MSA causes the NH resonances to split into several broad peaks that coalesce to two sharp peaks at 11.25 ppm and 10.83 ppm, respectively, upon the addition of two equivalents of MSA. These findings are again attributed to the asymmetry induced by the addition of one proton and symmetry being restored once two protons are added to the macrocycle.

To obtain further insight into the electronic nature of these amethyrin species, density functional theory (DFT) calculations were carried out using the Gaussian 16 package and the Stampede2 cluster at the Texas Advanced Computational Center (TACC) and the Halifax server at the Oden Institute for Computational Engineering and Sciences.^{23,24} Geometric relaxations were carried out using X-ray crystal structural parameters for freebase compounds **1-4** so as to obtain energy

minimized structures and simulated spectra.^{8,9,13} Agreement between experimental and theoretical data was seen in all cases.

Aromaticity was evaluated using several well-known methods. First, gauge including magnetic induced current (GIMIC) density calculations were performed on the series of expanded porphyrins.²⁵⁻³⁰ Integrated current densities were evaluated and the main current pathways densities for selected bonds were determined (ESI Figure S5). In all cases protonation induced a significant change in the current pathways. Both free-base complexes **1** and **3** displayed a minor paratropic current throughout the macrocycle, indicating that these free-base ligands experience no significant global conjugation. Furthermore, these free-base complexes displayed a high degree of local aromaticity, as observed by strong diatropic ring currents within the pyrrole and pyridine subunits. Diprotonation of complexes **1** and **3** resulted in reduction of the local aromaticity present within the pyridine and pyrrole sub-units as observed by a decrease in the current density. Loss of local aromaticity was accompanied by a gain of a global paratropic current, indicating a newly formed anti-aromatic pathway. Interestingly, in contrast to complexes **1** and **3**, free-base amethyrin **2** is characterized by a significant global paratropic current. This observation supports prior reports wherein the free-base form of amethyrin is formally antiaromatic.³¹ In analogy to what is true for **1** and **3**, protonation of **2** results in a significant increase in the global paratropic current that involves an outer conjugative pathway through the pyrrolic subunits. This increase comes at the cost of near-complete loss in local aromaticity.

Anisotropy of the induced current density (ACID) calculations provide further support for the conclusion that expanded porphyrins **1-4** display proton induced anti-aromaticity.³² For the protonated species, the ACID plots calculated with the applied magnetic field along the z-axis, show a strong dominant paratropic ring current in all cases, indicative of an anti-aromatic pathway (Figure 5).




Figure 5. Calculated anisotropy induced current density (ACID) plot for diprotonated pyriamethyrin **1b**. Current density vectors associated with a global counter-clockwise current are seen, as would be expected for an anti-aromatic compound.

In contrast, the free base forms **1** and **3** displayed little global current directionality. Rather, strong diatropic currents were seen within the local aromatic subunits. This local aromaticity is disrupted upon protonation as reflected in a reduction in the local diatropic ring currents. However, it translates directly into a net increase in global conjugation. The calculated magnetic current density data is in agreement with the experimental ¹H NMR spectral studies (Figure 4) where the observed downfield shifts in the inner NH proton signals are ascribed to the presence of an inner paratropic ring current that serves to deshield the NH proton (ESI Figures S8-S19).

Nucleus independent chemical shifts (NICS) were calculated at the center of the macrocycle, as defined by the centroid of the six nitrogen atoms, located 1 Å above the center. In all cases, significant differences in the NICS(1)_{zz} values between the free-base and protonated forms were observed.³³ The NICS(1)_{zz} values for the free-base forms of **1** and **3** were 4.68 and 3.23 ppm, respectively. While the NICS values of these metal-free porphyrinoids are characterized by slight paratropic current, free-base amethyrin **2** displayed a large positive NICS(1)_{zz} value of 20.42 ppm. This result is consistent with free-base amethyrin being an anti-aromatic compound as noted earlier.³¹ Upon protonation of the free-base forms of **1** and **3**, a significant shift in the NICS(1)_{zz} values to 16.6 and 28.2 ppm, respectively, was observed. Likewise, diprotonation of amethyrin **2b** led to a drastic shift in the NICS(1)_{zz} value to 43.2 ppm. The iso-chemical surface shielding (ICSS) plot, for ligands **1** and **3**, are shown in Figure 6. It is noteworthy that amethyrins **2** and **2b** displayed the highest anti-aromatic character, followed by pyriamethyrin **1b** and dipyriamethyrin **3b**. On this basis, we suggest that substitution of pyrrole with pyridine in the case of the amethyrin class of expanded porphyrins results in reduced through-conjugation and global anti-aromaticity. This conclusion is consistent with the differences in the chemical shift changes in the pyrrole NH resonances due to protonation seen in the NMR titration experiments discussed above, with **3** displaying the lowest chemical shift change and **2** displaying the most dramatic shift.




Figure 6. Iso-chemical surface shielding (ICSS) of pyriamethyrin **1** and **3**, dipyriamethyrin and their protonated forms. Green regions indicate non-aromatic, blue-black indicate antiaromaticity, and red indicates aromaticity.

Upon protonation, the energy gaps between the HOMO and LUMO in **1b-3b** are reduced as compared to those of the corresponding neutral forms **1-3**. The lowering of the HOMO-LUMO

gap is consistent with the idea that protonation induces anti-aromaticity. A trend is also observed where the HOMO-LUMO gap increases upon replacing pyrrole with pyridine. The HOMO-LUMO gap for dipyrriamethyrin complex **3** is 2.8 eV, while that for **2** is 2.0. This is consistent with the conclusion that the free-base macrocycle **2** already displays anti-aromatic character. However, upon protonation the HOMO-LUMO gap for complex **2b** decreases to 1.6 eV, supporting the notion that the anti-aromaticity is further increased upon protonation.

DFT calculations were performed on both the free-base and protonated forms of **1-3**. Based on these results we suggest that in all cases an increase in anti-aromatic character results from protonation. This increase in anti-aromatic character is accompanied by a significant reduction in local aromaticity in the case of **1b** and **3b**, and a complete loss of local aromaticity in the case of **2b**.

In summary, we have shown that several members of the generalized amethyrin class of macrocycles develop antiaromatic character upon diprotonation. This study provides a set of rare examples in porphyrinoid chemistry where a conversion to an anti-aromatic form occurs as the result of simple protonation. Substitution of pyrrole moieties for pyridine acts to decrease the extent to which anti-aromaticity is engendered upon protonation (i.e., increasing the number of pyridine subunits results in less antiaromaticity upon diprotonation). Support for the proposed conversion to anti-aromatic forms came from UV-Vis and NMR spectroscopic studies, as well as extensive theoretical calculations.


This work was supported by the Office of Basic Energy Sciences, U.S. Department of Energy (DOE) (Grant DEFG02-01ER15186 to J.L.S.), and the Robert A. Welch Foundation (F-1018 to J.L.S.). The authors acknowledge the Texas Advanced Computer Center (TACC) at The University of Texas at Austin for providing high performance computing resources that have contributed to the research results reported in this paper. H.D.R. thanks UT Austin for a Provost's Graduate Excellence Fellowship and the Environmental Institute at The University of Texas at Austin for a Scientist in Residence Fellowship. D.N.M. thanks UT Austin for the Graduate Research Fellowship. We thank the Laboratory Directed Research and Development (LDRD) program for a Glenn T. Seaborg Graduate Summer Student Fellowship (J.T.B.).

Conflicts of interest

There are no conflicts to declare.

Notes and references

- L. P. Cook, G. Brewer and W. Wong-Ng, *Crystals*, 2017, **7**, 223.
- M. Imran, M. Ramzan, A. K. Qureshi, M. Azhar Khan and M. Tariq, *Biosensors*, 2018, **8**, 1–17.
- H. Huang, W. Song, J. Rieffel and J. F. Lovell, *Front. Phys.*, 2015, **3**, 1–15.
- A. Osuka and S. Saito, *Chem. Commun.*, 2011, **47**, 4330–4339.
- T. Soya, H. Mori and A. Osuka, *Angew. Chemie - Int. Ed.*, 2018, **57**, 15882–15886.
- Z. S. Yoon, A. Osuka and D. Kim, *Nat. Chem.*, 2009, **1**, 113–122.
- R. Myśliborski, L. Latos-Grazyński and L. Szterenber, *European J. Org. Chem.*, 2006, 3064–3068.
- T. D. Lash, K. Pokharel, J. M. Serling, V. R. Yant and G. M. Ferrence, *Org. Lett.*, 2007, **9**, 2863–2866.
- Z. Zhang, J. M. Lim, M. Ishida, V. V. Roznyatovskiy, V. M. Lynch, H. Y. Gong, X. Yang, D. Kim and J. L. Sessler, *J. Am. Chem. Soc.*, 2012, **134**, 4076–4079.
- R. J. P. Corriu, G. Bolin, J. J. E. Moreau and C. Vernhet, *J. Chem. Soc. Chem. Commun.*, 1991, 211–213.
- F. H. Carré, R. J. P. Corriu, G. Bolin, J. J. E. Moreau and C. Vernhet, *Organometallics*, 1993, **12**, 2478–2486.
- S. Hannah, D. Seidel, J. L. Sessler and V. Lynch, *Inorganica Chim. Acta*, 2001, **317**, 211–217.
- J. T. Brewster, H. D. Root, D. Mangel, A. Samia, H. Zafar, A. C. Sedgwick, V. M. Lynch and J. L. Sessler, *Chem. Sci.*, 2019, 5596–5602.
- E. Steiner and P. W. Fowler, *Org. Biomol. Chem.*, 2006, **4**, 2473–2476.
- J. L. Sessler, A. Gebauer, A. Guba, M. Scherer and V. Lynch, *Inorg. Chem.*, 2002, **37**, 2073–2076.
- J. T. Brewster, A. Aguilar, G. Anguera, H. Zafar, M. D. Moore and J. L. Sessler, *J. Coord. Chem.*, 2018, **71**, 1808–1813.
- J. T. Brewster, Q. He, G. Anguera, M. D. Moore, X. S. Ke, V. M. Lynch and J. L. Sessler, *Chem. Commun.*, 2017, **53**, 4981–4984.
- S. Durot, J. Taesch and V. Heitz, *Chem. Rev.*, 2014, **114**, 8542–8578.
- Z. Zhou and Z. Shen, *J. Mater. Chem. C*, 2015, **3**, 3239–3251.
- Y. Liu, Q. Zhang, Q. H. Guo and M. X. Wang, *J. Org. Chem.*, 2016, **81**, 10404–10410.
- Z. Zhang, D. S. Kim, C. Y. Lin, H. Zhang, A. D. Lammer, V. M. Lynch, I. Popov, O. S. Miljanić, E. V. Anslyn and J. L. Sessler, *J. Am. Chem. Soc.*, 2015, **137**, 7769–7774.
- R. R. Valiev, I. Benkyi, Y. V. Konyshv, H. Fliegl and D. Sundholm, *J. Phys. Chem. A*, 2018, **122**, 4756–4767.
- M. J. Frisch, *et al. Gaussian 16, Revision B.01* (Gaussian Inc., Wallingford, 2016)
- Texas Advanced Computing Center (TACC), The University of Texas at Austin.
- J. Jusélius, D. Sundholm, J. Gauss, *J. Chem. Phys.* 2004, **121**, 3952–3963
- H. Fliegl, S. Taubert, O. Lehtonen, and D. Sundholm, *Phys. Chem. Chem. Phys.* 2011, **13**, 20500–20518
- H. Fliegl, J. Jusélius, and D. Sundholm, *J. Phys. Chem. A*, 2016, **120**, 5658–5664.
- M. Rauhalahhti, S. Taubert, D. Sundholm, and V. Liegeois, *Phys. Chem. Chem. Phys.* 2017, **19**, 7124–7131
- H. Fliegl, D. Sundholm, S. Taubert, J. Jusélius, and W. Klopper, *J. Phys. Chem. A*, 2009, **113**, 30, 8668–8676.
- H. Fliegl, R. Valiev, F. Pichierri and D. Sundholm, *Chem. Modell.*, 14, 1, (2018).
- E. Steiner, and P. W. Fowler, *Org. Biomol. Chem.* 2006, **4**, 2473–2476.
- R. Herges, D. Geuenich, *J. Phys. Chem. A*, 2001, **105**, 3214–3220.
- Z. Chen, C. S. Wannere, C. Corminboeuf, R. Puchta, and P. V. R. Schleyer, *Chem. Rev.* 2005, **10**, 3842–3888



228x101mm (96 x 96 DPI)