



Environmental
Science
Nano

Particle size determine the accumulation of platinum nanoparticles in the estuarine amphipod, *Leptocheirus plumulosus*

Journal:	<i>Environmental Science: Nano</i>
Manuscript ID	EN-ART-08-2021-000713.R2
Article Type:	Paper

SCHOLARONE™
Manuscripts

Environmental Significance

Many nanoparticle uptake and toxicity studies has been performed with polydispersed suspensions, that is suspensions with broad size distributions. However, the impact of nanoparticle size dispersity on uptake and toxicity have not been investigated experimentally. By tracking nanoparticles at the single particle level, this study provides insights on how nanoparticle size and size dispersity affect the uptake and elimination of polyvinylpyrrolidone coated platinum nanoparticle (PVP-PtNPs) in *Leptocheirus plumulosus*. Uptake of PtNPs increased with the decreases in PtNP size whereas and elimination of PtNPs increased with increases PtNP size. When exposed to mixtures of 40 and 70 nm PtNPs (PtNP₄₀ and PtNP₇₀, respectively) with equal masses, surface areas, or number concentrations of PtNP₄₀ and PtNP₇₀, the *L. plumulosus* accumulated higher numbers of PtNP₄₀ than PtNP₇₀. Moreover, increases in PtNP₇₀ concentration did not affect the uptake of PtNP₄₀ from the mixture. These results reveal that in a polydisperse nanoparticle suspension the uptake of a given fraction is independent of other size fractions.

1
2
3 **Particle size determine the accumulation of platinum nanoparticles in the**
4
5
6 **estuarine amphipod, *Leptocheirus plumulosus***
7
8

9 Mithun Sikder ^{1,2}, Emily Eudy ², Bo Cai ³, G. Thomas Chandler ², Mohammed Baalousha ^{1,2,*}
10
11

12 ¹ *South Carolina SmartState Center for Environmental Nanoscience and Risk (CENR), Arnold*
13 *School of Public Health, University of South Carolina, Columbia, SC, 29208, USA*
14
15

16 ² *Department of Environmental Health Sciences, Arnold School of Public Health, University of*
17 *South Carolina, Columbia, SC, 29208, USA*
18
19

20 ³ *Department of Epidemiology and Biostatistics, Arnold School of Public Health, University of*
21 *South Carolina, Columbia, SC, 29208, USA*
22
23

24 Correspondence: Dr. Mohammed Baalousha, Department of Environmental Health Sciences,
25 University of South Carolina, 921 Assembly Street, Columbia, SC, 2908, USA, Tel: +1 803-777-
26 7177, E-mail: mbaalous@mailbox.sc.edu
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Nanoparticles (NPs) typically display a wide distribution of different sizes in aquatic environments, yet little information is available on the impact of particle size dispersity on organismal uptake and elimination. This study investigated uptake and elimination of polyvinylpyrrolidone-coated platinum nanoparticles (PVP-PtNPs) of different sizes (*e.g.*, 20.0 ± 4.8 , 40.5 ± 4.1 , and 70.8 ± 4.2 nm) by the estuarine amphipod *Leptocheirus plumulosus*. Accumulation and elimination were determined by measuring total Pt body burden in amphipods exposed to PtNPs using inductively coupled plasma-mass spectroscopy, as well as the mass and number PtNP body burden using single particle-ICP-MS (sp-ICP-MS). *L. plumulosus* accumulated PtNPs of different sizes from water exposure, and mostly (*e.g.*, $> 90\%$) as NPs rather than as dissolved Pt. Mass- and number- based uptake increased with decreases in PtNP size whereas mass- and number- based elimination increased with increasing PtNP size. The residual whole-animal body burden of PtNPs after 48 h elimination increased with decreases in PtNP size, with residual body burdens approximately two-fold higher for amphipods exposed to 20 nm PtNPs than amphipods exposed to 70 nm PtNPs. PtNP influx rate (k_{inw}) increased with decreasing NP size, with k_{inw} s of 1.07 ± 0.31 , 0.82 ± 0.22 , and 0.67 ± 0.10 $\mu\text{g}\cdot\text{g}^{-1}\cdot\text{d}^{-1}$ for 20, 40, and 70 nm PtNPs, respectively. PtNP efflux rate (k_e) increased with increasing NP size, with k_e s of 0.31 ± 0.08 , 0.66 ± 0.04 , and 0.83 ± 0.07 $\cdot\text{d}^{-1}$ for 20, 40, and 70 nm NP, respectively. When exposed to mixtures of 40 and 70 nm PtNPs with equal masses, surface areas, or number concentrations of 40 and 70 nm PtNPs, *L. plumulosus* accumulated higher numbers of the 40 nm PtNPs than 70 nm PtNPs from all mixtures. The increased exposure concentration of 70 nm PtNPs in the mixture did not affect the uptake of 40 nm PtNPs, suggesting that in a polydispersed NP suspension the uptake of a given size fraction is independent of other size fractions in the mixture.

Keywords

Nanoparticle, Uptake, Elimination, Size dispersity, Single particle ICP-MS

INTRODUCTION

Although it can be assumed that NPs in an aquatic environment display a wide distribution of different sizes, little information is available on the impact of particle size polydispersity on their environmental behaviors. Numerous studies demonstrated negative impacts of NPs to aquatic organisms, but divergent results have been reported on the importance of NP size on their uptake and toxicity. For instance, several studies revealed that NPs might have size-dependent toxic effects¹⁻⁴; yet it is common to find studies attributing toxicity to a certain size fraction⁴ while others report no size-dependent toxicity⁵. Furthermore, the relationship between NP uptake and size remains uncertain. Several studies have shown decreases in NP uptake with increases in NP size⁶⁻⁸, with decreases in NP size⁹⁻¹¹, and even a lack of NP size-dependent uptake¹²⁻¹⁴. Such contradictory inferences might be due to use of different analytical techniques for NP characterization in the different studies and/or to the heterogeneity and polydispersity of the NPs most widely studied and reported in the literature¹⁵. NP size polydispersity may result in incorrect measurements of NP average size, underestimation of NP dose, masking of size-dependent properties and effects^{15, 16}.

Many early metal-based NP uptake and toxicity studies suffered from a lack of robust NP quantification and characterization during exposures, including concentration measurements and size distributions in test media, in organisms, and the relative impacts of the nanoparticulate versus dissolved metals¹⁷. Most early metal-based NP uptake studies have inferred NP uptake by measuring the total NP-metal burden within organisms exposed to NPs^{18, 19}. This approach can

1
2
3 provide accurate body burden measurements for insoluble metal NPs, but it falls short of
4
5 differentiating between dissolved metals and intact NP uptake from soluble NPs. Additionally, this
6
7 approach does not yield any information on the size distribution of the accumulated NPs^{15, 16}. The
8
9 inability to differentiate the form of measured body burden (*e.g.*, dissolved metal *vs* NPs), in
10
11 addition to the lack of concurrent characterization techniques to determine NP aggregation state in
12
13 media and tissues, limits a robust understanding of NP toxicology. Single-particle inductively-
14
15 coupled plasma mass spectrometry (sp-ICP-MS) can bridge this gap as it concurrently measures
16
17 NP size distributions, and NP number and mass concentrations²⁰ at environmentally- and
18
19 toxicologically- relevant particle concentrations (*e.g.*, ng L⁻¹ range)²¹. Recent studies
20
21 demonstrated the uptake of AuNPs in environmental organisms such as *Daphnia magna* and zebra
22
23 fish²² and *Caenorhabditis elegans* (*C. elegans*)^{23, 24} using sp-ICP-MS. The latter studies
24
25 investigated the uptake by *C. elegans* of individual AuNP size suspensions (*e.g.*, 30 nm, 60 nm,
26
27 80 nm, and 100 nm) and did not consider the uptake from mixtures of AuNPs of different sizes^{23,}
28
29 ²⁴. Other studies demonstrated the uptake of AuNPs in the alga *Pseudokirchneriella subcapitata*
30
31 using single cell (sc)-ICP-MS^{22, 25}.

32
33
34
35
36
37
38 Many studies demonstrated the toxicity of PtNPs toward environmental organisms such as
39
40 *Pseudokirchneriella subcapitata* (EC₅₀ of 15 to 200 mg Pt L⁻¹)^{26, 27}; Duckweed (*Lemna minor*),
41
42 water fleas (*Daphnia magna* - EC₅₀ of 406 to 514 µg Pt L⁻¹), bacteria (*Vibrio fischeri* - EC₅₀ of
43
44 135 to 255 µg Pt L⁻¹)²⁸; brine shrimp (*Artemia salina* - 96 EC₅₀ of 117 µg Pt L⁻¹)²⁹, and zebra fish
45
46 ³⁰. These EC₅₀ values are much higher than Pt concentrations in surface waters (*ca.* ng L⁻¹ to µg
47
48 L⁻¹)³¹. Furthermore, to date, many studies investigated the uptake of metal and metal oxide (*e.g.*,
49
50 Au, Ag, TiO₂, ZnO, CeO₂) NPs in aquatic organisms^{32, 33}. However, despite the multiple sources
51
52 of PtNPs in the environment³⁴⁻³⁹, only a few studies demonstrated the uptake and translocation of
53
54
55
56
57
58
59
60

PtNPs in environmental organisms such as zebrafish ³⁰, the freshwater snail *Lymnaea stagnalis* ⁶, aurgula and escarole plants ⁴⁰, and *Sinapis alba* and *Lepidium stavianum* plants ^{41, 42}.

Therefore, the objectives of this study are to 1) evaluate the size-dependent bioaccumulation (*i.e.*, uptake and elimination) of PtNPs in the absence of food by the common estuarine amphipod *Leptocheirus plumulosus* using sp-ICP-MS, and (2) examine the effect of NP size dispersity on the accumulation of PtNPs from polydispersed suspensions (*e.g.*, mixtures of 40 nm and 70 nm PtNPs with equal number, surface area, and mass concentrations). Throughout this report the term ‘uptake’ is used to describe particles entering and measured within the test organism and does not necessarily imply that membrane passage or tissue translocation of NPs has occurred. The term elimination is used to describe the proportion of particles excreted/eliminated *vs.* retained within the organism over time. The term ‘bioaccumulation’ represents the net outcome of all uptake and elimination processes ^{43, 44}. The study was carried out using the USEPA-approved ⁴⁵ estuarine amphipod test species *L. plumulosus* as a model particle-ingesting benthic species exposed to PtNPs of three different controlled sizes. PtNPs were chosen as a model study particle for a number of reasons: (1) PtNPs exhibit very low toxicity which if present might interfere with NP uptake and elimination ⁴⁶, (2) the low solubility of PtNPs minimizes differential uptake of dissolved versus particulate Pt ⁶, (3) there are usually undetectable background concentrations of PtNPs in aquatic environments and the test organism ⁴⁷, (4) sp-ICP-MS provides a low size-detection limit for PtNPs (*i.e.*, ~ 17 nm) ⁴⁸, and (5) the release of PtNPs from photocatalytic convertors in vehicles is growing should not be considered irrelevant in the environment without further study ³⁴, and (6) the limited number of existing studies on the uptake of PtNPs.

METHODOLOGY

Platinum nanoparticles (PtNPs)

Polyvinylpyrrolidone-coated platinum nanoparticles (PVP-PtNPs) with nominal size of 20 nm, 40 nm, and 70 nm (PtNP₂₀, PtNP₄₀, and PtNP₇₀, respectively) were synthesized using a seed-mediated growth approach⁴⁹ and briefly described in Supporting information (SI). The synthesized PtNPs were thoroughly characterized using dynamic light scattering (DLS), atomic force microscopy (AFM), energy dispersive X-ray (EDX) coupled transmission electron microscopy (TEM), and single particle inductively coupled mass spectroscopy (sp-ICP-MS) as reported elsewhere⁴⁹. The concentrations of PtNP₂₀, PtNP₄₀, and PtNP₇₀ stock suspensions were 24.5 ± 0.6 , 23.4 ± 1.2 , and 24.8 ± 0.7 mg-Pt L⁻¹, respectively.

Test organism

Leptocheirus plumulosus is a euryhaline, infaunal amphipod that reaches densities of $> 2.5 \times 10^4$ individuals m⁻² in muddy estuaries along the east coast of North America⁵⁰. *L. plumulosus* is a facultative suspension and surface-deposit feeder⁵¹ that ingests a wide range of particle types and sizes. *L. plumulosus* also serves as an important prey organism for fish and shellfish⁵², and is a sensitive species commonly used as a USEPA approved standard test species in acute⁵³ and chronic^{45, 54} sediment toxicity tests in the USA.

Leptocheirus plumulosus were cultured in muddy sediments in the laboratory following USEPA testing standard protocols⁴⁵. All cultures and treatments used 15 parts per thousand (ppt) synthetic seawater (SW) prepared with Crystal Seas® bioassay grade sea salts (Marine Enterprises International, Baltimore, MD, USA) followed by aeration of $> 90\%$ O₂ saturation and then filtered

1
2
3 twice using 0.45 μm filter. The pH of these waters was 8.1 ± 0.1 throughout the study. For each
4
5 uptake experiment, adult amphipods were harvested by passing mixed-aged cultures through a 2
6
7 mm sieve and sorting adults retained on a 500 μm sieve. Similar size/age adults were transferred
8
9 onto a 600 μm sieve partially submerged in a 2 L crystallizing dish filled with 15 ppt SW agitated
10
11 continuously by an X-shaped stir bar placed underneath the sieve. The amphipods were allowed
12
13 to depurate/defecate through the sieve for 2 h in darkness before being collected and allocated to
14
15 50-mL exposure chambers. *L. plumulosus* used in this study were sized as closely as possible to
16
17 one another using stereomicroscopy, with a mean dry weight of 1.68 ± 0.32 mg ($n = 225$).
18
19
20
21
22

23 ***Waterborne PtNP uptake experiments***

24
25 Two sets of uptake experiments were performed in this study without food additions. In
26
27 the first uptake experiment, *Leptocheirus plumulosus* ($n = 3$ amphipods in 5 replicates of each
28
29 PtNP suspension) were exposed to 45 mL of 25 $\mu\text{g L}^{-1}$ individual PtNP₂₀, PtNP₄₀, and PtNP₇₀
30
31 suspensions in 50-mL crystallizing dishes for 72 h (**Table S1**). These exposure concentrations
32
33 were selected as the lowest concentrations that allow detection by sp-ICP-MS of the accumulated
34
35 PtNPs following extraction five fold higher than the background PtNP number concentrations
36
37 detected by sp-ICP-MS (**Table S2**). In the second uptake experiment, *L. plumulosus* similarly were
38
39 exposed to mixtures of equal mass-, surface area-, number- concentrations of PtNP₄₀ and PtNP₇₀
40
41 (**Table S1**). The equal mass, surface area, and number concentration mixtures were prepared by
42
43 mixing 25 $\mu\text{g L}^{-1}$ NP₄₀ with 25, 64.3, and 98.3 $\mu\text{g L}^{-1}$ NP₇₀, respectively. Each treatment consisted
44
45 of 5 replicates (15 amphipods per treatment), and each test included an equivalent zero-dose
46
47 control (*i.e.*, no PtNP). Zero amphipod mortality occurred in any of the treatments or controls.
48
49 Each glass chamber was covered and placed in a temperature regulated incubator at 25°C with
50
51 12:12 h light:dark photoperiod according to EPA protocols⁴⁵. Amphipods were collected at 6, 24,
52
53
54
55
56
57
58
59
60

1
2
3 48, and 72 h for total (*e.g.*, sum of dissolved and PtNP) Pt concentration and PtNP number and
4 mass concentration determinations and size distribution analyses as described below. Amphipods
5 were removed from the test chambers, rinsed in 3 successive baths of 15 ppt seawater (SW), and
6 transferred to glass chambers containing 45 mL fresh 15 ppt SW for a 2 h elimination period to
7 purge gut contents before being washed thoroughly once in Ethylene Diamine Tetra-Acetic acid
8 (EDTA) and twice in ultrapure water (UPW). Amphipods were then blotted dry and immediately
9 weighed (exactly at 60 s post-removal) and suspended in 5 mL UPW for particle extraction for sp-
10 ICP-MS analysis. Dry weights of amphipods were calculated from measured wet weights using a
11 simple regression between wet and dry weights (**Figure S1**). Water samples were collected at the
12 middle point of the water column from each exposure chamber at 0 and 72 h to monitor potential
13 changes (*e.g.*, size distribution, dispersity, aggregation, and number concentration) in PtNP
14 exposure during uptake experiments. The test water and amphipods were frozen at -80 °C and later
15 thawed to room temperature before analysis using ICP-MS and sp-ICP-MS.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 ***Elimination experiments***

35
36
37
38 A separate set of experiments was performed to determine the elimination (efflux rates) of
39 PtNPs from *L. plumulosus* following a 24 h exposure to 100 µg L⁻¹ PtNPs. An intentionally higher
40 exposure concentration was used in the elimination experiment versus the uptake experiment to
41 ensure that *L. plumulosus* accumulated a sufficient amount of PtNPs within their bodies to be
42 detectable by sp-ICP-MS over the full 48 h elimination period. As previously, *L. plumulosus* of
43 similar sizes (retained on a 500 µm sieve) were collected from lab cultures, followed by gut
44 purging in freshly prepared 15 ppt SW for 2 h to empty the contents of the digestive tract.
45 Amphipods were then transferred for 24 h to 15 new glass chambers containing 45 mL each of 15
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 ppt SW spiked with 100 $\mu\text{g L}^{-1}$ PtNP₂₀, PtNP₄₀, or PtNP₇₀ without food. Each glass chamber was
4 covered and incubated in a temperature-regulated incubator at 25°C with 12:12 h light:dark
5 photoperiod. Each of 3 treatments consisted of 5 replicates and each replicate consisted of 3 *L.*
6 *plumulosus* (n=45 total). No mortality was observed in any of the experiments. After 24 h,
7 amphipods were removed from the test chambers, rinsed in 3 successive baths of 15 ppt SW,
8 followed by 2 h gut purging. The amphipods were then transferred to glass chambers containing
9 45 mL freshly-prepared 15 ppt SW without food for a 48 h dissolved Pt and PtNP elimination
10 period. The water was renewed every 12 h during the elimination experiment to minimize the
11 reuptake of depurated Pt or PtNPs from the water column. The amphipods were collected at 0, 6,
12 12, 24, 36, and 48 h, and washed thoroughly once in EDTA and twice in UPW. The amphipods
13 were immediately weighed as before, and finally suspended in 5 mL UPW. The dry weight of
14 amphipods was calculated from regressed wet weight (**Figure S1**). Water samples were also
15 collected during the elimination experiment at the middle point of the water column from each
16 elimination chamber at 0, 6, 12, 24, 36, and 48 h to monitor potential changes in NP size
17 distribution, aggregation, and number concentration. The water and amphipods were frozen to -80
18 °C upon collection and thawed to room temperature before analysis in ICP-MS and sp-ICP-MS.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 ***PtNP extraction from L. plumulosus***

42
43
44
45 PtNPs were extracted by probe sonication of amphipods in UPW using a Q700 probe-
46 sonicator (QSonica, Newtown, CT). The PtNP extraction procedure was optimized for PtNP
47 recovery by varying the number of amphipods (*e.g.*, 1, 3, 5, and 10, **Table S3a**) in each extraction
48 replicate, sonication times (*e.g.*, 1, 5, and 10 minutes, **Table S3b**), and centrifugation forces (*e.g.*,
49 100 x g and 2000 x g using Eppendorf 5810R, **Table S3c**) needed to isolate amphipod tissues.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Total Pt concentration in the supernatant and the pellet were determined by ICP-MS. The overall
4 performance of the extraction procedure was determined by comparing the Pt concentrations in
5 the supernatant and the pellet (consisting mainly of the exposed amphipods' tissue). Equal Pt
6 concentrations in the pellet and the supernatant indicates that PtNPs were fully released from the
7 organisms. Higher concentrations of PtNPs in the pellet indicates that PtNPs remained attached to
8 the organism tissues and ended up being more concentrated in the pellet. The optimal extraction
9 conditions (*i.e.*, Pt concentrations were the same in the supernatant and pellet) were determined as
10 3 *L. plumulosus* per replicate, 5 min pulse sonication with 2 seconds on and 1 second off pulsing
11 at room temperature, and low gravity centrifugation at 100 g. Under these extraction conditions,
12 the Pt concentration in the supernatant and the tissue pellet (50.0 ± 1.4 and $51.9 \pm 2.1 \mu\text{g L}^{-1}$,
13 respectively) were not statistically different (t-test, $p > 0.8$; **Table S3c**), inferring the successful
14 extraction of PtNPs from the amphipod tissue into the suspension and the lack of PtNP
15 sedimentation during the centrifugation step.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 ***PtNP concentration measurement by single particle ICP-MS***

35
36
37
38 Prior to sp-ICP-MS analysis, all extracted tissue samples were bath sonicated (Branson,
39 Model 2800, 40 kHz, USA) for 30 min to breakdown NP aggregates, followed by 100-fold dilution
40 to minimize the introduction of organic matter to the ICP-MS. Aqueous samples were diluted 100
41 fold without sonication to prevent aggregate breakup. The number and mass concentrations of the
42 extracted PtNPs and the aqueous samples were measured using sp-ICP-MS following methods
43 described elsewhere⁵⁵⁻⁵⁷. All samples were analyzed using a PerkinElmer NexION™ 350D ICP-
44 MS (Waltham, MA, USA) equipped with a Meinhard glass concentric nebulizer, a PerkinElmer
45 glass cyclonic spray chamber, and 2 mm ID quartz injector. Instrument calibration was performed
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 using a blank and at least four Au and Pt standards ranging from 0.5 to 20 $\mu\text{g L}^{-1}$. All data were
4
5 collected in single particle mode, with signals averaged for the entire analysis period (60 s). ICP-
6
7 MS nebulization efficiency was determined daily using the particle size method⁵⁶ with 60 nm
8
9 NIST™ AuNP standard reference material (actual TEM size of 56 nm; reference material 8013
10
11 manufactured by the National Institute of Standard and Technology, MD, USA) and varied
12
13 between 4% and 6% for all experiments described herein. Data were acquired at an RF power of
14
15 1600 W, a 50 μs dwell time, a 0 μs settling time, and a 60 s acquisition time. 1.2×10^6 readings
16
17 were collected for each sample during 60 s acquisition time per sample. Particle number
18
19 concentration was determined directly by counting all PtNPs measured by sp-ICP-MS in a given
20
21 sample, while PtNP mass concentration was determined by integrating the masses of all the
22
23 particles measured by sp-ICP-MS following the procedure described by Pace et al.^{55, 56}.
24
25 Recoveries, total mass determinations, and body burdens were reported as the mean and standard
26
27 error of 5 replicates.
28
29
30
31
32
33

34 ***Total Pt concentration measurement by ICP-MS***

35
36
37
38 Total Pt concentration was measured by ICP-MS following digestion of the extracted
39
40 PtNPs in a concentrated aqua regia ($\text{HNO}_3:\text{HCL} = 1 \text{ M}:3 \text{ M}$) one day before analysis. Briefly, 500
41
42 μL aliquots of the extracted PtNPs were transferred to Eppendorf tubes followed by addition of 1
43
44 mL of a freshly prepared concentrated aqua regia. The tubes were incubated under the hood at
45
46 room temperature for 24 h. Samples were diluted at least 5 times prior to ICP-MS analysis.
47
48
49
50

51 ***Biodynamic model***

52
53
54 Biodynamic modeling (BDM) is a popular approach for deconstructing metal accumulation
55
56
57

1
2
3 as it quantifies its mechanistic components⁵⁸. BDM describes bioaccumulation as the net outcome
4 of uptake (*i.e.*, influx), physiological elimination (*i.e.*, efflux), and body growth dilution (if any).
5
6 Equation 1 describes the changes in Pt concentration in an organism ($[Pt]_{org}$) overtime after a
7
8 waterborne-only exposure:
9
10

$$\frac{[Pt]_{org}}{dt} = k_{uw} \times [Pt]_{water} - k_e \times [Pt]_{org} - k_g \times [Pt]_{org} \quad (\text{Eq. 1})$$

11
12
13
14
15
16 Where, k_{uw} ($L\ g^{-1}\ d^{-1}$) is the unidirectional Pt influx rate constant from solution/suspension,
17
18 $[Pt]_{water}$ ($\mu g\ L^{-1}$) is the dispersed PtNP concentration in the solution/suspension, and k_e (d^{-1}) and k_g
19
20 (d^{-1}) are the rate constants for physiological elimination and body growth dilution, respectively. k_g
21
22 is negligible for *Leptocheirus plumulosus* for the short experimental period (**Figure S2**).
23
24 Therefore, growth dilution was not considered in determining influx rate constants. k_{uw} was
25
26 determined by dividing uptake rate by exposure concentration, assuming the uptake rate follows
27
28 first-order kinetics for the exposure concentration used⁵⁹ and k_e is determined using a nonlinear
29
30 regression that includes compartments representing fast and slow exchanging pools (**Eq. S1**).
31
32
33
34 Further details and the parameterization of these rate constants is described in the supporting
35
36 information.
37
38
39

40 ***Statistical analysis***

41
42
43 Statistical analyses were performed with SAS[®] version 9.4 software (SAS institute, Cary,
44
45 NC, United States). Correlation coefficients between PtNP₇₀ concentrations in the exposure
46
47 medium and those accumulated in the amphipods after 48 h exposure were calculated using
48
49 Pearson's correlation method. Fisher's Z-transformation was used to compute confidence intervals
50
51 for each correlation, followed by z-tests to calculate p-values for correlation comparisons between
52
53 every two mixture combinations. The particle size distribution (PSD) of PtNPs in the exposure
54
55
56
57
58
59
60

1
2
3 medium and those accumulated in the amphipods were analyzed using Kolmogorov-Smirnov (K-
4 S) tests with Bonferroni correction to compare size distributions of NPs in exposure and concurrent
5 accumulation. In all cases, statistical significance was set at p -value < 0.05 .
6
7
8
9

10 11 **RESULTS AND DISCUSSIONS**

12 13 *PtNP characterization and behavior in SW*

14
15
16 The physiochemical properties of PVP-PtNPs were determined using DLS, TEM, and sp-
17 ICP-MS⁴⁹ (**Figure S3 and Table S4**). Briefly, the mean core diameters of PtNP₂₀, PtNP₄₀, and
18 PtNP₇₀ measured by TEM were 18.5 ± 5.0 , 44.5 ± 5.0 , and 72.5 ± 3.9 nm, respectively. The energy
19 dispersive X-ray (EDX) spectra exhibited three peaks corresponding to platinum M-shell (2.05
20 keV), L α -shell (9.44 keV), and L β -shell (11 keV, **Figure S3f-h**), confirming the chemical
21 composition of PtNPs. The core diameters of PtNP₂₀, PtNP₄₀, and PtNP₇₀ measured by sp-ICP-MS
22 were 24.7 ± 1.6 , 42.9 ± 0.8 , and 77.1 ± 0.8 nm, in close agreement with those measured by TEM.
23
24 The z-average hydrodynamic diameter of PtNP₂₀, PtNP₄₀, and PtNP₇₀ were 51.0 ± 0.7 , 74.7 ± 0.2 ,
25 and 93.4 ± 1 nm, respectively. The magnitude of the zeta potential of PtNPs increased from 22.9
26 ± 1.4 to 27.2 ± 1.7 mV with the increase in PtNP size.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

42 DLS is the most commonly used analytical technique to measure NP aggregation.
43 However, the low PtNP concentration (*i.e.*, $25 \mu\text{g L}^{-1}$) used in this study hampers the use of DLS
44 to measure PtNP aggregation because it is not possible to reliably measure NP hydrodynamic
45 diameter by DLS at such low concentrations. Therefore, the aggregation behavior of PtNPs (*i.e.*,
46 PtNP₂₀, PtNP₄₀, and PtNP₇₀) in SW was monitored by measuring their number concentration and
47 number size distribution at 0 h and 72 h following mixing with SW using sp-ICP-MS (**Figure 1**).
48
49
50
51
52
53
54
55
56
57
58
59
60

PtNPs remained dispersed as single particles immediately (0 h) after mixing with SW. After 72 h of mixing PtNPs with SW, the number concentration of primary PtNPs decreased (**Figure 1, Table S5**) due to aggregate formation with aggregate sizes of 30 to 130, 70 to 200, and 90 to 250 nm for PtNP₂₀, PtNP₄₀, and PtNP₇₀, respectively. The PVP-PtNPs used in this study have been shown (using DLS and high PtNP concentrations) to aggregate in SW due to surface charge screening by high counterions (mainly Na⁺) concentrations in SW⁶⁰.

Uptake of PtNPs

Following exposure to individual PtNP size groupings, all PtNPs accumulated in *L. plumulosus* generating body burdens that were significantly greater than controls (ANOVA, $p < 0.001$). The whole-animal body burden of PtNP₂₀, PtNP₄₀, and PtNP₇₀ in *L. plumulosus* from waterborne exposure of 25 $\mu\text{g-Pt L}^{-1}$ increased exponentially as a function of exposure time (**Figure 2a**). The PtNP₂₀ showed a steeper exponential uptake than PtNP₄₀, which in turn showed steeper exponential uptake than PtNP₇₀. The whole animal body burden reached 47.4 ± 7.4 , 15.8 ± 4.5 and $8.3 \pm 1.8 \mu\text{g Pt g}^{-1}$ dry wt organism after 72 h exposures to PtNP₂₀, PtNP₄₀, and PtNP₇₀, respectively (**Table S6**). The whole-animal Pt body burden from the three PtNP exposures after 72 h was three orders of magnitude greater than the amphipod's background Pt body burden of $11.31 \pm 0.01 \text{ ng g}^{-1}$ dry weight. *L. plumulosus* did not gain weight during the 72-h uptake period (**Figure S2**). Therefore, growth dilution was not considered in determining influx rate constants. The influx rate constants (k_{in} s, mean \pm SE) from waterborne exposures were 0.042 ± 0.006 , 0.040 ± 0.009 , and $0.026 \pm 0.003 \mu\text{g g}^{-1} \text{ d}^{-1}$, respectively for PtNP₂₀, PtNP₄₀, and PtNP₇₀ (**Table S6**), indicating decreases in PtNP uptake with increase in PtNP size. The decrease in PtNP influx rate constant with increases in NP size might be attributed to the formation of larger PtNP aggregates

(**Figure S4a-c**) and correspondingly faster sedimentation from the water column, consistent with previous studies^{6, 61}. The measured size distribution of PtNPs in the exposure seawater shifted toward larger sizes and the number concentration decreased over time indicating particle aggregation (**Figure S4a-c**). Additionally, the mass concentration of PtNP₂₀, PtNP₄₀, and PtNP₇₀ in the exposure seawater decreased by 15.4 ± 3.2 , 51.3 ± 3.5 , and 72.5 ± 6.8 %, respectively after 72 h indicating increased PtNP losses with increased PtNP size due to particle uptake and/or sedimentation (**Figure S4d-f**). However, the mass balance of the test system revealed that only a small fraction (*e.g.*, 3.8 ± 0.2 , 6.8 ± 0.2 , and 3.6 ± 0.2 %) of total PtNP₂₀, PtNP₄₀, and PtNP₇₀, respectively, accumulated in *L. plumulosus* after 72 h, implying very low depletion of PtNPs from the exposure suspensions via organismal uptake (**Table S7**). Since PtNP concentrations in exposure chambers were not depleted by amphipod uptake, the decline in PtNP mass concentration can be attributed to particle sedimentation which in turn increases with increased NP size.

Pt accumulation in *L. plumulosus* was mainly due to the nano-form of Pt rather than the dissolved Pt fraction as the PtNP body burden, measured by sp-ICP-MS after 72 h exposure, and representing > 70% of the total Pt body burden for PtNP₂₀ and > 90% of the total Pt body burden for both PtNP₄₀ and PtNP₇₀ (**Table S8-10**). Additionally, the dissolved Pt concentration represented < 5% of the Pt exposure concentration⁴⁹ which is very unlikely to produce a residual sorption of Pt ions onto the surfaces of the organism sufficient to result in their detection as single particle events. The lower PtNP body burden % relative to the total Pt body burden for PtNP₂₀ compared to PtNP₄₀ and PtNP₇₀ likely is due to the size-detection limitations of sp-ICP-MS (*e.g.*, 17 nm). Only 46% of the primary PtNPs in the PtNP₂₀ stock suspension were larger than the sp-ICP-MS size detection limit (*e.g.*, 17 nm for PtNPs)⁴⁸, thus 53% of primary PtNPs in the PtNP₂₀ stock suspension were not detected by sp-ICP-MS. However, all primary PtNPs in PtNP₄₀ and

PtNP₇₀ stock suspensions were larger than the sp-ICP-MS size detection limit (**Table S4**)⁶². The higher than expected recovery of PtNP₂₀ after NP accumulation in *L. plumulosus* was due to the formation of PtNP aggregates larger than the sp-ICP-MS size detection limit in the test seawater (**Figure S5a, b**)⁶². Overall, these results indicate that the extraction procedure, which was validated for PtNP₇₀ only, provides accurate results for all PtNPs used in this study. The lower recovery of PtNP₂₀ was most likely due to the sp-ICP-MS size detection limit rather than due to the extraction procedure.

Based on mass concentration, the accumulation of PtNP₂₀ exceeded the accumulation of PtNP₄₀ and PtNP₇₀ by 3 and 5.7 fold after 72 h of exposure (**Figure 2a**). However, based on particle number concentration, the accumulation of PtNP₂₀ exceeded the accumulation of PtNP₄₀ and PtNP₇₀ by 6 and 7 orders of magnitude after 72 h of exposure. The whole-animal body burden of PtNPs (based on number concentration) also increased exponentially with time (**Figure 2b**) and the number of PtNPs accumulated in *L. plumulosus* followed the trend of PtNP₂₀ (6.5×10^{10} PtNPs₂₀ g⁻¹ dry weight) > PtNP₄₀ (1.1×10^4 PtN₄₀ g⁻¹ dry weight) > PtNP₇₀ (1.9×10^3 PtNP₇₀-g⁻¹ dry weight). For a fixed mass concentration ($25 \mu\text{g L}^{-1}$), the number of NPs in a PtNP₂₀ suspension is much higher than that in a PtNP₇₀ suspension (**Table S5**). Therefore, NPs were more abundantly available for uptake from PtNP₂₀ suspensions than from PtNP₄₀ and PtNP₇₀ suspensions, resulting in the observed higher PtNP₂₀ influx into *L. plumulosus* relative to PtNP₄₀ and PtNP₇₀ (**Figure 2b**). This relationship produced a higher body burden for the smaller PtNPs. Collectively, mass- and number-concentration based PtNP body burdens decreased with increases in PtNP size. A similar decrease in NP uptake with increases in particle size was observed for other filter feeders and deposit feeders with silver oxide, copper oxide, and gold NPs^{63,64}. Similarly, lower body burdens were observed for smaller (30 nm and 60 nm) versus larger (80 nm and 100 nm) AuNPs in *C.*

1
2
3 *elegans*^{23, 24}. The lower body burden of PtNP₇₀ compared to smaller PtNPs (*i.e.*, PtNP₄₀ and
4 PtNP₂₀) might be due to the formation of larger aggregates in SW (**Figure 1 and S4a-c**). The faster
5
6 sedimentation of these larger aggregates from the water column (**Figure S4d-f**) and/or inherently
7
8 greater difficulties passing larger aggregates through cell membranes likely render larger PtNPs
9
10 less bioavailable to *L. plumulosus*.
11
12
13

14
15 Our previous study revealed that the majority of PtNPs remained in particulate form as
16
17 shown by the low concentration of the dissolved Pt in the water relative to the amount of PtNPs
18
19 added⁴⁹. Therefore, particle ingestion is likely a major route of uptake of all PtNP sizes. Uptake
20
21 of metal oxide NPs and quantum dots via ingestion was shown previously for *L. plumulosus*^{65, 66}.
22
23 CdSe/ZnS quantum dots accumulated to a greater extent than their respective dissolved ions in *L.*
24
25 *plumulosus*⁶⁵. Estuarine benthic invertebrates accumulate metals from their surrounding
26
27 environment through two pathways: dissolved metals are accumulated *via* passage across
28
29 permeable membranes⁴³, whereas particle-associated metals are accumulated mostly by ingestion
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
67. *L. plumulosus* is capable of both suspension feeding (by ingestion of planktonic and suspended
benthic microalgae) and surface deposit feeding (by ingestion of sediment, detritus, phytodetritus,
and benthic microalgae)⁵¹. Without sediment, *L. plumulosus* feeds on suspended particles; when
burrowed, it selects food particles from the water column and from surface sediments. *L.*
plumulosus can ingest up to 3 times its body weight in sediment per day⁶⁸. *L. plumulosus* are filter
feeders, enabling them to ingest particles from the size of 70 µm down to around 200 nm⁶⁹. In
most cases, larger particles (*e.g.*, microalgae) are consumed most efficiently. The ingestion of
PtNPs by *L. plumulosus* is not surprising as all PtNP treatments formed aggregates in 15S seawater
with equivalent spherical core diameters of up to 130 to 250 nm (**Figure 1**). Aggregate
hydrodynamic diameters may actually be much larger as NPs form porous aggregates. These

1
2
3 aggregate sizes are within the lower size range of the food particles filtered/ingested by *L.*
4
5 *plumulosus*, thus promoting their passive incidental uptake and leading to measurable tissue
6
7 accumulation. The possibility that additions of algal food might influence NP dispersivity and
8
9 create unknown and variable aggregate NP:food diameters led to an *a priori* decision to eliminate
10
11 suspended food as a factor in these experiments. Future studies should manipulate food quality
12
13 and quantity as test variables in the matrix. In the present study, this additional manipulation
14
15 increased the experimental matrix considerably, and was beyond the scope of present resources.
16
17
18
19
20

21 ***Elimination of PtNPs***

22
23 The whole-animal Pt body burden following 24 h exposure to 100 $\mu\text{g L}^{-1}$ of PtNP₂₀, PtNP₄₀,
24
25 and PtNP₇₀ in seawater reached 18.3 ± 4.5 , 22.2 ± 3.8 , and 28.1 ± 3.8 $\mu\text{g Pt g}^{-1}$ dry wt organism,
26
27 respectively. After this 24 h exposure, the whole animal Pt body burden decreased with time
28
29 (**Figure 3a**). Generally, *L. plumulosus* displayed two elimination phases for PtNPs: an initial, rapid
30
31 loss (0-6 h), and a second, slower loss (6 to 48 h) (**Figure 3a**). After 6 h of depuration, the whole
32
33 animal mean body burdens dropped to 11.4 ± 3.4 , 18.7 ± 8.2 , and 14.3 ± 6.2 $\mu\text{g Pt g}^{-1}$ dry wt
34
35 organism for PtNP₂₀, PtNP₄₀, and PtNP₇₀, respectively -- a decrease of 38 ± 2 to $49 \pm 3\%$ of the
36
37 accumulated PtNPs. Between 6 and 48 h, the whole-animal body burden of PtNPs decreased
38
39 linearly and reached 8.5 ± 3.6 , 6.1 ± 1.9 , and 4.5 ± 4.6 $\mu\text{g Pt g}^{-1}$ dry wt organism for PtNP₂₀,
40
41 PtNP₄₀, and PtNP₇₀, a decrease of 54%, 72% and 84% of the initially accumulated PtNPs over 48
42
43 h of depuration, respectively. A similar depuration behavior has been observed for Cd release from
44
45 *L. plumulosus* after ingestion/accumulation of Cd from pelagic and benthic microalgae⁷⁰. These
46
47 results indicate that *L. plumulosus* depurated larger NPs more efficiently than smaller NPs. This
48
49 behavior is in good agreement with the observed increased elimination with increasing particle
50
51 size of citrate-coated AuNPs and polystyrene NPs by *D. magna*^{9, 64}. The residual (*i.e.*, non-
52
53
54
55
56
57
58
59
60

1
2
3 depurated) animal body burdens after 48 h of depuration represent $47 \pm 2 \%$, $28 \pm 1\%$, and $16 \pm$
4
5
6 2% of the accumulated PtNP₂₀, PtNP₄₀, and PtNP₇₀ after 24 h exposures to $100 \mu\text{g L}^{-1}$ of PtNP,
7
8 respectively (**Figure 3a**). It is worth noting that after 48 h of depuration, the residual body burden
9
10 was approximately two orders of magnitude higher than the measured background body burden in
11
12 non-exposed animals. The efflux rates (k_e) were 0.31 ± 0.08 , 0.66 ± 0.04 , and $0.83 \pm 0.07 \text{ d}^{-1}$,
13
14 respectively, for PtNP₂₀, PtNP₄₀, and PtNP₇₀, indicating increased depuration with increases in NP
15
16 size (**Figure 3a**).
17

18
19 Based on mass concentration, the retention of PtNP₂₀ exceeds the retention of PtNP₄₀ and
20
21 PtNP₇₀ by 1.4 and 1.9 fold after 72 h of depuration (**Figure 3a**). However, based on particle number
22
23 concentration, the retention of PtNP₂₀ exceeds the retention of PtNP₄₀ and PtNP₇₀ by 2.3 and 6.4
24
25 fold after 48 h of depuration (**Figure 3b**). Similar depuration trends were observed based on
26
27 number concentration analysis as were observed for mass concentration analysis. Initially, *L.*
28
29 *plumulosus* accumulated $2 \times 10^{10} \pm 2.5 \times 10^9 \text{ PtNP}_{20}\text{-g}^{-1}$ dry wt, $1.2 \times 10^{10} \pm 2.4 \times 10^9 \text{ PtNP}_{40}\text{-g}^{-1}$
30
31 dry wt, and $4.5 \times 10^9 \pm 1.2 \times 10^9 \text{ PtNP}_{70}\text{-g}^{-1}$ dry wt after 24 h exposure to $100 \mu\text{g L}^{-1}$ PtNPs. *L.*
32
33 *plumulosus* retained $1 \times 10^{10} \text{ PtNP}_{20}$, $4.4 \times 10^9 \text{ PtNP}_{40}$, and $1.6 \times 10^9 \text{ PtNP}_{70}\text{-g}^{-1}$ dry wt after 48 h
34
35 depuration (**Figure 3b**). *L. plumulosus* gained weight during the 48-h depuration period (**Figure**
36
37 **S6**). However, the growth rate constants (k_g) were minimal *i.e.*, 0.002 ± 0.001 , 0.005 ± 0.002 , and
38
39 $0.007 \pm 0.002 \text{ d}^{-1}$ for PtNP₂₀, PtNP₄₀, PtNP₇₀ treatments, respectively. Therefore, no adjustment
40
41 was made in k_e due to growth dilution.
42
43
44
45

46
47 The decrease in PtNP elimination with decreases in size might be attributed to increased
48
49 NP internalization and tissue/cell translocation with decreases in NP size ⁷¹ Therefore, increased
50
51 accumulation of PtNPs with decreases in size appears to be a result of a combination of increased
52
53
54
55
56
57

1
2
3 influx and decreased efflux, which may in turn contribute to increased toxicity for smaller NPs if
4 size-dependent cell and tissue translocation is real.
5
6

7
8 Depurated PtNPs were measured in the water column at the end of each water renewal
9 period during the depuration experiment. The number particle size distribution of each PtNP
10 (**Figure S7a-c**) was similar to that in the corresponding original PtNP suspension (K-S test, $p >$
11 0.22). The number and mass concentrations at the end of each water renewal period decreased over
12 time (**Figure S7d-f**). The total mass of the PtNPs detected in the water column during the
13 elimination experiment was 0.024 ± 0.008 , 0.031 ± 0.001 , and 0.103 ± 0.02 μg for PtNP₂₀, PtNP₄₀,
14 and PtNP₇₀, respectively, and accounted for 41 ± 1.9 , 40.2 ± 0.3 , and 74.9 ± 1.0 % of the total
15 depurated PtNPs (**Table S11**). The remaining fractions might have been lost from the suspensions
16 by sedimentation or organism re-uptake. In the absence of food, the depurated PtNPs may be
17 available for re-uptake, which may contribute to slower observed depuration over time.
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 ***NP uptake from mixture of PtNPs: Effect of size dispersity***

33
34

35
36 When exposed to a mixture of PtNP₄₀ and PtNP₇₀, *L. plumulosus* accumulated both PtNP
37 size distributions, and the number of accumulated PtNPs also increased with time (**Figure 4**). *L.*
38 *plumulosus*, accumulated significantly higher numbers of PtNP₄₀ than PtNP₇₀ (K-S test, $p < 0.05$)
39 from all mixtures (equal mass, surface area, or number concentrations). This is consistent with the
40 higher accumulation of PtNP₄₀ than PtNP₇₀ observed in the exposures to individual NPs (**Figure**
41 **2**). The higher accumulation of PtNP₄₀ than PtNP₇₀ from equal mass and surface area mixtures
42 might be attributed to the higher number concentration of PtNP₄₀ than for PtNP₇₀ (**Figure 5a and**
43 **b**) and/or higher accumulation of PtNP₄₀ than PtNP₇₀ in the amphipod exposures. The higher
44 accumulation of PtNP₄₀ than PtNP₇₀ from equal number concentration mixtures is attributed to
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 higher influx (**Figure 2**) and lower depuration (**Figure 3**) of PtNP₄₀ compared to the PtNP₇₀. These
4
5 results clearly indicate that in a mixture of PtNPs, smaller PtNPs accumulate to a higher extent
6
7 than larger PtNPs, even if the number concentration of both sizes are the same in each exposure.
8
9

10 The accumulation of PtNP₇₀ from equal surface area- and number-concentration mixtures
11 increased compared to the accumulation of PtNP₇₀ from equal mass concentration mixtures. This
12 can be attributed to the increased number of PtNP₇₀ in equal surface area- and number-
13 concentration mixture exposures compared to equal mass concentration mixture exposures
14 (**Figure 4a-c**). Conversely, PtNP₄₀ accumulation from these three mixtures remained constant
15 despite the increase in PtNP₇₀ number concentrations. Therefore, the increased exposure
16 concentration and accumulation of larger PtNPs (*i.e.*, PtNP₇₀) did not affect the accumulation of
17 smaller PtNPs (*i.e.*, PtNP₄₀), and the presence of larger NPs does not impact the accumulation of
18 smaller NPs in a polydispersed mixture. It is worth noting that all PtNP mixtures in our
19 experiments formed aggregates in 15S seawater as indicated by the decrease in particle number
20 concentration (Figure **S8-10**).
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 The present study demonstrated that the common estuarine amphipod *L. plumulosus* will
36 accumulate PtNPs of different sizes (*e.g.*, 20, 40, and 70 nm) in seawater-only exposures without
37 food. It also demonstrated that mass and number uptake increases with decreases in NP size. All
38 PtNPs were depurated to varying degrees, with depuration increasing with increases in NP mean
39 size. *L. plumulosus* retained/accumulated higher amounts of small PtNPs than larger PtNPs as a
40 result of a combination of increased influx and decreased efflux of smaller PtNPs. The residual
41 whole-animal body burden of PtNPs after 48 h depuration was higher for treatments of reduced
42 NP size, with residual body burdens about two-fold larger for animals exposed to PtNP₂₀ than for
43 animals exposed to PtNP₇₀. When exposed to mixtures of small and large PtNPs of equal masses,
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 surface areas, or number concentrations, *L. plumulosus* accumulated higher numbers of the small
4
5 PtNPs than large PtNPs from all mixture combinations. Additionally, an increased exposure
6
7 concentration of larger PtNPs did not affect the uptake of smaller PtNPs from a mixture of NPs of
8
9 different sizes. The increased retention of small NPs compared to larger ones might partially
10
11 explain the higher toxicity of small NPs compared to larger ones observed in other studies using
12
13 NPs of more toxic metals. Filter and suspension-feeding amphipods are an important food source
14
15 for many marine and freshwater birds, fish and macro-invertebrates and thus may contribute to
16
17 metal-NPs accumulation by predatory organisms that otherwise may not be exposed. In addition
18
19 to the inherent and largely well-studied toxic effects of most metals used in NP manufacture today,
20
21 this study further suggests that NP size spectra in aquatic environments is an important additional
22
23 parameter to consider in NP ecological risk assessment.
24
25
26
27
28
29

30 **Acknowledgment**

31
32
33
34 This research was supported by the United States National Science Foundation
35
36 ([NSF1437307](#)).
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Reference

1. Ariano, P.; Zamburlin, P.; Gilardino, A.; Mortera, R.; Onida, B.; Tomatis, M.; Ghiazza, M.; Fubini, B.; Lovisolo, D., Interaction of spherical silica nanoparticles with neuronal cells: size-dependent toxicity and perturbation of calcium homeostasis. *Small* **2011**, *7* (6), 766-74.
2. Zhang, X.-D.; Wu, D.; Shen, X.; Liu, P.-X.; Yang, N.; Zhao, B.; Zhang, H.; Sun, Y.-M.; Zhang, L.-A.; Fan, F.-Y., Size-dependent in vivo toxicity of PEG-coated gold nanoparticles. *International journal of nanomedicine* **2011**, *6*, 2071-2081.
3. Park, J.; Lim, D. H.; Lim, H. J.; Kwon, T.; Choi, J. S.; Jeong, S.; Choi, I. H.; Cheon, J., Size dependent macrophage responses and toxicological effects of Ag nanoparticles. *Chem Commun (Camb)* **2011**, *47* (15), 4382-4.
4. Choi, O.; Hu, Z., Size Dependent and Reactive Oxygen Species Related Nanosilver Toxicity to Nitrifying Bacteria. *Environmental Science & Technology* **2008**, *42* (12), 4583-4588.
5. Lin, W.; Xu, Y.; Huang, C.-C.; Ma, Y.; Shannon, K. B.; Chen, D.-R.; Huang, Y.-W., Toxicity of nano- and micro-sized ZnO particles in human lung epithelial cells. *Journal of Nanoparticle Research* **2009**, *11* (1), 25-39.
6. Sikder, M.; Croteau, M.-N.; Poulin, B. A.; Baalousha, M., Effect of Nanoparticle Size and Natural Organic Matter Composition on the Bioavailability of Polyvinylpyrrolidone-Coated Platinum Nanoparticles to a Model Freshwater Invertebrate. *Environmental Science & Technology* **2021**, *55* (4), 2452.
7. Geffroy, B.; Ladhar, C.; Cambier, S.; Treguer-Delapierre, M.; Brèthes, D.; Bourdineaud, J.-P., Impact of dietary gold nanoparticles in zebrafish at very low contamination pressure: The role of size, concentration and exposure time. *Nanotoxicology* **2012**, *6* (2), 144-160.
8. Hughes, R. N., A study of feeding in *Scrobicularia plana*. *Journal of the Marine Biological Association of the United Kingdom* **1969**, *49* (3), 805-823.
9. Rosenkranz, P.; Chaudhry, Q.; Stone, V.; Fernandes, T. F., A comparison of nanoparticle and fine particle uptake by *Daphnia magna*. *Environmental Toxicology and Chemistry* **2009**, *28* (10), 2142-2149.
10. Gaiser, B. K.; Fernandes, T. F.; Jepson, M. A.; Lead, J. R.; Tyler, C. R.; Baalousha, M.; Biswas, A.; Britton, G. J.; Cole, P. A.; Johnston, B. D.; Ju-Nam, Y.; Rosenkranz, P.; Scown, T. M.; Stone, V., Interspecies comparisons on the uptake and toxicity of silver and cerium dioxide nanoparticles. *Environmental Toxicology and Chemistry* **2012**, *31* (1), 144-154.
11. Coleman, J. G.; Johnson, D. R.; Stanley, J. K.; Bednar, A. J.; Weiss Jr, C. A.; Boyd, R. E.; Steevens, J. A., Assessing the fate and effects of nano aluminum oxide in the terrestrial earthworm, *Eisenia fetida*. *Environmental Toxicology and Chemistry* **2010**, *29* (7), 1575-1580.
12. Scown, T. M.; Santos, E. M.; Johnston, B. D.; Gaiser, B.; Baalousha, M.; Mitov, S.; Lead, J. R.; Stone, V.; Fernandes, T. F.; Jepson, M.; van Aerle, R.; Tyler, C. R., Effects of Aqueous Exposure to Silver Nanoparticles of Different Sizes in Rainbow Trout. *Toxicological Sciences* **2010**, *115* (2), 521-534.
13. Stanley, J. K.; Coleman, J. G.; Weiss Jr, C. A.; Steevens, J. A., Sediment toxicity and bioaccumulation of nano and micron-sized aluminum oxide. *Environmental Toxicology and Chemistry* **2010**, *29* (2), 422-429.
14. Unrine, J. M.; Tsyusko, O. V.; Hunyadi, S. E.; Judy, J. D.; Bertsch, P. M., Effects of Particle Size on Chemical Speciation and Bioavailability of Copper to Earthworms (*Eisenia fetida*) Exposed to Copper Nanoparticles. *Journal of Environmental Quality* **2010**, *39* (6), 1942-1953.

15. Baalousha, M.; Lead, J. R., Nanoparticle dispersity in toxicology. *Nature Nanotechnology* **2013**, *8*, 308.
16. Gaumet, M.; Vargas, A.; Gurny, R.; Delie, F., Nanoparticles for drug delivery: The need for precision in reporting particle size parameters. *European Journal of Pharmaceutics and Biopharmaceutics* **2008**, *69* (1), 1-9.
17. Kennedy, A. J.; Hull, M. S.; Bednar, A. J.; Goss, J. D.; Gunter, J. C.; Bouldin, J. L.; Vikesland, P. J.; Steevens, J. A., Fractionating Nanosilver: Importance for Determining Toxicity to Aquatic Test Organisms. *Environmental Science & Technology* **2010**, *44* (24), 9571-9577.
18. Klaine, S. J.; Alvarez, P. J. J.; Batley, G. E.; Fernandes, T. F.; Handy, R. D.; Lyon, D. Y.; Mahendra, S.; McLaughlin, M. J.; Lead, J. R., Nanomaterials in the environment: Behavior, fate, bioavailability, and effects. *Environmental Toxicology and Chemistry* **2008**, *27* (9), 1825-1851.
19. Lead, J. R.; Batley, G. E.; Alvarez, P. J. J.; Croteau, M. N.; Handy, R. D.; McLaughlin, M. J.; Judy, J. D.; Schirmer, K., Nanomaterials in the environment: Behavior, fate, bioavailability, and effects-An updated review. *Environ Toxicol Chem* **2018**, *37* (8), 2029-2063.
20. Bednar, A. J.; Poda, A. R.; Mitrano, D. M.; Kennedy, A. J.; Gray, E. P.; Ranville, J. F.; Hayes, C. A.; Crocker, F. H.; Steevens, J. A., Comparison of on-line detectors for field flow fractionation analysis of nanomaterials. *Talanta* **2013**, *104*, 140-148.
21. Gray, E. P.; Bruton, T. A.; Higgins, C. P.; Halden, R. U.; Westerhoff, P.; Ranville, J. F., Analysis of gold nanoparticle mixtures: a comparison of hydrodynamic chromatography (HDC) and asymmetrical flow field-flow fractionation (AF4) coupled to ICP-MS. *Journal of Analytical Atomic Spectrometry* **2012**, *27* (9), 1532-1539.
22. Abdolapur Monikh, F.; Chupani, L.; Arenas-Lago, D.; Guo, Z.; Zhang, P.; Darbha, G. K.; Valsami-Jones, E.; Lynch, I.; Vijver, M. G.; van Bodegom, P. M.; Peijnenburg, W. J. G. M., Particle number-based trophic transfer of gold nanomaterials in an aquatic food chain. *Nature Communications* **2021**, *12* (1), 899.
23. Johnson, M. E.; Hanna, S. K.; Montoro Bustos, A. R.; Sims, C. M.; Elliott, L. C. C.; Lingayat, A.; Johnston, A. C.; Nikoobakht, B.; Elliott, J. T.; Holbrook, R. D.; Scott, K. C. K.; Murphy, K. E.; Petersen, E. J.; Yu, L. L.; Nelson, B. C., Separation, Sizing, and Quantitation of Engineered Nanoparticles in an Organism Model Using Inductively Coupled Plasma Mass Spectrometry and Image Analysis. *ACS Nano* **2017**, *11* (1), 526-540.
24. Johnson, M. E.; Bennett, J.; Montoro Bustos, A. R.; Hanna, S. K.; Kolmakov, A.; Sharp, N.; Petersen, E. J.; Lapasset, P. E.; Sims, C. M.; Murphy, K. E.; Nelson, B. C., Combining secondary ion mass spectrometry image depth profiling and single particle inductively coupled plasma mass spectrometry to investigate the uptake and biodistribution of gold nanoparticles in *Caenorhabditis elegans*. *Analytica Chimica Acta* **2021**, *1175*, 338671.
25. Merrifield, R.; Stephan, C.; Lead, J., Quantification of Au nanoparticle biouptake and distribution to freshwater algae using single cell-ICP-MS. *Environmental science & technology* **2018**, *52* (4), 2271-2277.
26. Sørensen, S. N.; Engelbrekt, C.; Lützhøft, H.-C. H.; Jiménez-Lamana, J.; Noori, J. S.; Alatraktchi, F. A.; Delgado, C. G.; Slaveykova, V. I.; Baun, A., A Multimethod Approach for Investigating Algal Toxicity of Platinum Nanoparticles. *Environmental Science & Technology* **2016**, *50* (19), 10635-10643.
27. Książyk, M.; Asztemborska, M.; Stęborowski, R.; Bystrzejewska-Piotrowska, G., Toxic effect of silver and platinum nanoparticles toward the freshwater microalga *Pseudokirchneriella subcapitata*. *Bulletin of environmental contamination and toxicology* **2015**, *94* (5), 554-558.

- 1
2
3 28. Hlavkova, D.; Beklova, M.; Kopel, P.; Havelkova, B., Evaluation of platinum
4 nanoparticles ecotoxicity using representatives of distinct trophic levels of aquatic biocenosis.
5 *Neuroendocr. Lett* **2018**, *39*, 465-472.
- 6 29. DAĞLIOĞLU, Y.; ÇELEBİ, M. S., The evaluation of the acute toxic effects of
7 Polyvinylferrocenium supported platinum nanoparticles on *Artemia salina* Brine shrimp. *Biyolojik*
8 *Çeşitlilik ve Koruma* **2015**, *8* (3), 304-312.
- 9 30. Asharani, P. V.; Iianwu, Y.; Gong, Z.; Valiyaveetil, S., Comparison of the toxicity of
10 silver, gold and platinum nanoparticles in developing zebrafish embryos. *Nanotoxicol* **2010**, *5* (1),
11 43-54.
- 12 31. Mashio, A. S.; Obata, H.; Tazoe, H.; Tsutsumi, M.; Ferrer i Santos, A.; Gamo, T.,
13 Dissolved platinum in rainwater, river water and seawater around Tokyo Bay and Otsuchi Bay in
14 Japan. *Estuarine, Coastal and Shelf Science* **2016**, *180*, 160-167.
- 15 32. Lead, J. R.; Batley, G. E.; Alvarez, P. J.; Croteau, M. N.; Handy, R. D.; McLaughlin,
16 M. J.; Judy, J. D.; Schirmer, K., Nanomaterials in the environment: behavior, fate, bioavailability,
17 and effects—an updated review. *Environmental toxicology and chemistry* **2018**, *37* (8), 2029-2063.
- 18 33. Tripathi, D. K.; Tripathi, A.; Shweta; Singh, S.; Singh, Y.; Vishwakarma, K.; Yadav,
19 G.; Sharma, S.; Singh, V. K.; Mishra, R. K.; Upadhyay, R. G.; Dubey, N. K.; Lee, Y.; Chauhan,
20 D. K., Uptake, Accumulation and Toxicity of Silver Nanoparticle in Autotrophic Plants, and
21 Heterotrophic Microbes: A Concentric Review. *Frontiers in Microbiology* **2017**, *8*, 7.
- 22 34. Folens, K.; Van Acker, T.; Bolea-Fernandez, E.; Cornelis, G.; Vanhaecke, F.; Du Laing,
23 G.; Rauch, S., Identification of platinum nanoparticles in road dust leachate by single particle
24 inductively coupled plasma-mass spectrometry. *Science of the Total Environment* **2018**, *615*, 849-
25 856.
- 26 35. Bhattacharya, R.; Mukherjee, P., Biological properties of “naked” metal nanoparticles.
27 *Advanced Drug Delivery Reviews* **2008**, *60* (11), 1289-1306.
- 28 36. Pedone, D.; Moglianetti, M.; De Luca, E.; Bardi, G.; Pompa, P. P., Platinum nanoparticles
29 in nanobiomedicine. *Chemical Society Reviews* **2017**, *46* (16), 4951-4975.
- 30 37. Muthuswamy, N.; de la Fuente, J. L. G.; Ochal, P.; Giri, R.; Raaen, S.; Sunde, S.;
31 Rønning, M.; Chen, D., Towards a highly-efficient fuel-cell catalyst: optimization of Pt particle
32 size, supports and surface-oxygen group concentration. *Physical Chemistry Chemical Physics*
33 **2013**, *15* (11), 3803-3813.
- 34 38. Tang, J.; Tang, D., Non-enzymatic electrochemical immunoassay using noble metal
35 nanoparticles: a review. *Microchimica Acta* **2015**, *182* (13), 2077-2089.
- 36 39. Luo, X.; Morrin, A.; Killard, A. J.; Smyth, M. R., Application of nanoparticles in
37 electrochemical sensors and biosensors. *Electroanalysis: An International Journal Devoted to*
38 *Fundamental and Practical Aspects of Electroanalysis* **2006**, *18* (4), 319-326.
- 39 40. Kranjc, E.; Mazej, D.; Regvar, M.; Drobne, D.; Remškar, M., Foliar surface free energy
40 affects platinum nanoparticle adhesion, uptake, and translocation from leaves to roots in arugula
41 and escarole. *Environmental Science: Nano* **2018**, *5* (2), 520-532.
- 42 41. Asztemborska, M.; Steborowski, R.; Kowalska, J.; Bystrzejewska-Piotrowska, G.,
43 Accumulation of Platinum Nanoparticles by *Sinapis alba* and *Lepidium sativum* Plants. *Water,*
44 *Air, & Soil Pollution* **2015**, *226* (4), 126.
- 45 42. Jiménez-Lamana, J.; Wojcieszek, J.; Jakubiak, M.; Asztemborska, M.; Szpunar, J., Single
46 particle ICP-MS characterization of platinum nanoparticles uptake and bioaccumulation by
47 *Lepidium sativum* and *Sinapis alba* plants. *Journal of Analytical Atomic Spectrometry* **2016**, *31*
48 (11), 2321-2329.
- 49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 43. Rainbow, P. S., Ecophysiology of Trace Metal Uptake in Crustaceans. *Estuarine, Coastal and Shelf Science* **1997**, *44* (2), 169-176.
- 4
5 44. Petersen, E. J.; Mortimer, M.; Burgess, R. M.; Handy, R.; Hanna, S.; Ho, K. T.; Johnson, M.; Loureiro, S.; Selck, H.; Scott-Fordsmand, J. J.; Spurgeon, D.; Unrine, J.; van den Brink, N. W.; Wang, Y.; White, J.; Holden, P., Strategies for robust and accurate experimental approaches to quantify nanomaterial bioaccumulation across a broad range of organisms. *Environmental Science: Nano* **2019**, *6* (6), 1619-1656.
- 6
7 45. USEPA *Methods for assessing the chronic toxicity of marine and estuarine sediment-associated contaminants with the amphipod Leptocheirus plumulosus*; Environmental protection agency: Washington D.C., 1994.
- 8
9 46. Asharani, P. V.; Yi, L. W.; Gong, Z. Y.; Valiyaveetil, S., Comparison of the toxicity of silver, gold and platinum nanoparticles in developing zebrafish embryos. *Nanotoxicology* **2011**, *5* (1), 43-54.
- 10
11 47. Artelt, S.; Creutzenberg, O.; Kock, H.; Levsen, K.; Nachtigall, D.; Heinrich, U.; Rühle, T.; Schlögl, R., Bioavailability of fine dispersed platinum as emitted from automotive catalytic converters: a model study. *Science of The Total Environment* **1999**, *228* (2), 219-242.
- 12
13 48. Lee, S.; Bi, X.; Reed, R. B.; Ranville, J. F.; Herckes, P.; Westerhoff, P., Nanoparticle Size Detection Limits by Single Particle ICP-MS for 40 Elements. *Environmental Science & Technology* **2014**, *48* (17), 10291-10300.
- 14
15 49. Sikder, M.; Wang, J.; Chandler, G. T.; Berti, D.; Baalousha, M., Synthesis, characterization, and environmental behaviors of monodispersed platinum nanoparticles. *Journal of Colloid and Interface Science* **2019**, *540*, 330-341.
- 16
17 50. Holland, A. F.; Shaughnessy, A. T.; Scott, L. C.; Dickens, V. A.; Ranasinghe, J. A. *Progress report: long-term benthic monitoring and assessment program for the Maryland portion of Chesapeake Bay (July 1986-October 1987). Volume 2. Appendices A-Z*; United States, 1988-05-01, 1988.
- 18
19 51. DeWitt, T. H.; Redmond, M. S.; Sewall, J. E.; Swartz, R. C. *Development of a chronic sediment toxicity test for marine benthic amphipods*; United States, 1992; p 247.
- 20
21 52. Hines, A. H.; Haddon, A. M.; Wiechert, L. A., Guild structure and foraging impact of blue crabs and epibenthic fish in a subestuary of Chesapeake Bay. *Marine Ecology Progress Series* **1990**, *67* (2), 105-126.
- 22
23 53. McGee, B. L.; Schlekot, C. E.; Reinharz, E., Assessing sublethal levels of sediment contamination using the estuarine amphipod *Leptocheirus plumulosus*. *Environmental Toxicology and Chemistry* **1993**, *12* (3), 577-587.
- 24
25 54. Schlekot, C. E.; McGee, B. L.; Reinharz, E., Testing sediment toxicity in chesapeake bay with the amphipod *Leptocheirus plumulosus*: An evaluation. *Environmental Toxicology and Chemistry* **1992**, *11* (2), 225-236.
- 26
27 55. Pace, H. E.; Rogers, N. J.; Jarolimek, C.; Coleman, V. A.; Higgins, C. P.; Ranville, J. F., Determining Transport Efficiency for the Purpose of Counting and Sizing Nanoparticles via Single Particle Inductively Coupled Plasma Mass Spectrometry. *Analytical Chemistry* **2011**, *83* (24), 9361-9369.
- 28
29 56. Pace, H. E.; Rogers, N. J.; Jarolimek, C.; Coleman, V. A.; Gray, E. P.; Higgins, C. P.; Ranville, J. F., Single Particle Inductively Coupled Plasma-Mass Spectrometry: A Performance Evaluation and Method Comparison in the Determination of Nanoparticle Size. *Environmental Science & Technology* **2012**, *46* (22), 12272-12280.
- 30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 57. Gray, E. P.; Coleman, J. G.; Bednar, A. J.; Kennedy, A. J.; Ranville, J. F.; Higgins, C.
4 P., Extraction and Analysis of Silver and Gold Nanoparticles from Biological Tissues Using Single
5 Particle Inductively Coupled Plasma Mass Spectrometry. *Environmental Science & Technology*
6 **2013**, *47* (24), 14315-14323.
- 7
8 58. Luoma, S. N.; Rainbow, P. S., Why Is Metal Bioaccumulation So Variable? Biodynamics
9 as a Unifying Concept. *Environmental Science & Technology* **2005**, *39* (7), 1921-1931.
- 10 59. Luoma, S. N.; Stoiber, T.; Croteau, M.-N.; Römer, I.; Merrifield, R.; Lead, J. R., Effect
11 of cysteine and humic acids on bioavailability of Ag from Ag nanoparticles to a freshwater snail.
12 *NanoImpact* **2016**, *2*, 61-69.
- 13 60. Sikder, M.; Lead, J. R.; Chandler, G. T.; Baalousha, M., A rapid approach for measuring
14 silver nanoparticle concentration and dissolution in seawater by UV-Vis. *Science of The Total*
15 *Environment* **2018**, *618*, 597-607.
- 16 61. Skjolding, L. M.; Kern, K.; Hjorth, R.; Hartmann, N.; Overgaard, S.; Ma, G.; Veinot,
17 J. G. C.; Baun, A., Uptake and depuration of gold nanoparticles in *Daphnia magna*. *Ecotoxicology*
18 **2014**, 1-12.
- 19 62. Sikder, M.; Wang, J.; Poulin, B. A.; Tfaily, M. M.; Baalousha, M., Nanoparticle size and
20 natural organic matter composition determine aggregation behavior of polyvinylpyrrolidone
21 coated platinum nanoparticles. *Environmental Science: Nano* **2020**, *7* (11), 3318-3332.
- 22 63. Dai, L.; Syberg, K.; Banta, G. T.; Selck, H.; Forbes, V. E., Effects, Uptake, and
23 Depuration Kinetics of Silver Oxide and Copper Oxide Nanoparticles in a Marine Deposit Feeder,
24 *Macoma balthica*. *ACS Sustainable Chemistry & Engineering* **2013**, *1* (7), 760-767.
- 25 64. Skjolding, L. M.; Kern, K.; Hjorth, R.; Hartmann, N.; Overgaard, S.; Ma, G.; Veinot,
26 J. G. C.; Baun, A., Uptake and depuration of gold nanoparticles in *Daphnia magna*. *Ecotoxicology*
27 **2014**, *23* (7), 1172-1183.
- 28 65. Jackson, B. P.; Bugge, D.; Ranville, J. F.; Chen, C. Y., Bioavailability, toxicity, and
29 bioaccumulation of quantum dot nanoparticles to the amphipod *Leptocheirus plumulosus*.
30 *Environmental science & technology* **2012**, *46* (10), 5550-5556.
- 31 66. Hanna, S. K.; Miller, R. J.; Zhou, D.; Keller, A. A.; Lenihan, H. S., Accumulation and
32 toxicity of metal oxide nanoparticles in a soft-sediment estuarine amphipod. *Aquatic Toxicology*
33 **2013**, *142-143*, 441-446.
- 34 67. Louma, S., Prediction of metal toxicity in nature from bioassays: In *Metal speciation and*
35 *bioavailability in aquatic systems*, Chichester: Wiley, 1995; pp 609- 659.
- 36 68. Schlegel, C. E.; Decho, A. W.; Chandler, G. T., Bioavailability of particle-associated
37 silver, cadmium, and zinc to the estuarine amphipod *Leptocheirus plumulosus* through dietary
38 ingestion. *Limnology and Oceanography* **2000**, *45* (1), 11-21.
- 39 69. Geller, W.; Müller, H., The filtration apparatus of Cladocera: Filter mesh-sizes and their
40 implications on food selectivity. *Oecologia* **1981**, *49* (3), 316-321.
- 41 70. Yu, R.-Q.; Fleeger, J. W., Effects of nutrient enrichment, depuration substrate, and body
42 size on the trophic transfer of cadmium associated with microalgae to the benthic amphipod
43 *Leptocheirus plumulosus*. *Environmental Toxicology and Chemistry* **2006**, *25* (11), 3065-3072.
- 44 71. Foroozandeh, P.; Aziz, A. A., Insight into Cellular Uptake and Intracellular Trafficking of
45 Nanoparticles. *Nanoscale research letters* **2018**, *13* (1), 339-339.
- 46 72. Van Der Veen, I. T., Costly carotenoids: a trade-off between predation and infection risk?
47 *Journal of Evolutionary Biology* **2005**, *18* (4), 992-999.
- 48 73. Farese, R. V., Jr.; Walther, T. C., Lipid droplets finally get a little R-E-S-P-E-C-T. *Cell*
49 **2009**, *139* (5), 855-860.
- 50
51
52
53
54
55
56
57
58
59
60

1
2
3 74. Kashiwada, S., Distribution of Nanoparticles in the See-through Medaka (*Oryzias latipes*).
4 *Environmental Health Perspectives* **2006**, *114* (11), 1697-1702.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Number particle size distribution of PVP-PtNPs at 0 and 72 h post mixing of $25 \mu\text{g L}^{-1}$ PtNPs with 15 ppt synthetic seawater without amphipods for (a) PtNP₂₀, (b) PtNP₄₀, and (c) PtNP₇₀. All measurement were performed by SP-ICP-MS.




Figure 2. Platinum influx into *Leptocierious plumulosus* soft tissues after waterborne exposure to $25 \mu\text{g L}^{-1}$ PtNP₂₀, PtNP₄₀, and PtNP₇₀ for 72 h. PtNPs are presented as (a) platinum mass influx ($\mu\text{g g}^{-1}$) and (b) PtNPs number influx (NP g^{-1}). All measurement were performed by SP-ICP-MS. Data points represent mean and standard errors.




Figure 3. Retention of accumulated Pt in *L. plumulosus* after 24 h waterborne exposure to $100 \mu\text{g L}^{-1}$ PtNP₂₀, PtNP₄₀, and PtNP₇₀ presented as (a) mass based retention of Pt, and (b) NP number based retention of Pt (NP g^{-1}). All measurement were performed by SP-ICP-MS. Data points represent mean and standard errors.




Figure 4. (a-c) Number particle size distribution of and (d-f) the corresponding PtNP accumulation from mixtures of PtNP₄₀ and PtNP₇₀ containing: (a) the same mass of PtNP₄₀ and PtNP₇₀ ($25 \mu\text{g L}^{-1}$ PtNP₄₀ + $25 \mu\text{g L}^{-1}$ PtNP₇₀), (b) the same total surface area of PtNP₄₀ and PtNP₇₀ ($25 \mu\text{g L}^{-1}$ PtNP₄₀ + $64.25 \mu\text{g L}^{-1}$ PtNP₇₀), and (c) the same total number of PtNP₄₀ and PtNP₇₀ ($25 \mu\text{g L}^{-1}$ PtNP₄₀ + $98.25 \mu\text{g L}^{-1}$ PtNP₇₀). All measurement were performed by SP-ICP-MS.