



Analyst

In-situ Lipid Profiling of Insect Pheromone Glands by Direct Analysis in Real Time Mass Spectrometry

Journal:	<i>Analyst</i>
Manuscript ID	AN-ART-05-2022-000840.R1
Article Type:	Paper
Date Submitted by the Author:	05-Jun-2022
Complete List of Authors:	Cetraro, Nicolas; University of Hawai'i at Manoa, Pacific Biosciences Research Center Yew, Joanne; University of Hawai'i at Manoa, Pacific Biosciences Research Center

SCHOLARONE™
Manuscripts

1
2
3 **In-situ lipid profiling of insect pheromone glands by Direct Analysis in Real Time Mass**
4 **Spectrometry**
5

6
7
8 Nicolas Cetraro¹ and Joanne Y. Yew^{1*}
9

10
11 ¹Pacific Biosciences Research Center
12 School of Environment and Ocean Science Technology
13 University of Hawai'i at Mānoa
14 1993 East West Road
15 Honolulu, HI 96822
16
17
18

19
20 *corresponding author:
21

22 Joanne Y. Yew
23 1993 East West Rd
24 Honolulu, HI 96822
25 808-956-5870
26 jyew@hawaii.edu
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Lipid pheromones play a significant role in the behavior and ecology of many insects. The characterization of pheromone structures is a significant challenge due to their low abundance and ephemeral nature. Here we present a method for the analysis of lipid molecules from single pheromone glands of *Drosophila melanogaster* (fruit fly) using Direct Analysis in Real Time mass spectrometry (DART MS). Our results reveal that DART MS analysis of single tissues generates reproducible, species-specific lipid profiles comprised of fatty acids, wax esters, diacylglycerides and triacylglycerides. In addition, the ion source temperature and application of a solvent wash can cause significant qualitative and quantitative changes in the mass spectral profile. Lastly, we show that untargeted chemical fingerprinting of the gland can be used to accurately categorize species according to phylogenetic subgroup or genotype. Collectively, our findings indicate that DART MS is a rapid and powerful method for characterizing a broad range of lipids in tissues with minimal preparation. The application of direct tissue DART MS will expand the “secretome” of molecules produced by pheromone glands. In addition to its direct relevance to chemical ecology, the method could potentially be used in pharmaceutical studies for the screening and detection of tissue-specific drug metabolites.

Introduction

Lipid pheromones play a significant role in the behavioral and reproductive decisions of many insects¹. Considering both the beneficial and destructive impact of insects to agriculture² and their importance as vectors of disease, the identification of pheromones and their function has been an area of extensive research. The characterization of pheromones can be technically challenging due to the limited amounts that are produced (usually at or below pmol levels of individual compounds) and ephemeral nature of the molecules. Analysis by gas chromatography MS (GCMS) is the most widely used method of analysis for lipid pheromone classes including saturated and unsaturated hydrocarbons, acetate esters, epoxides, terpenoids, fatty acids, and alcohols³. However, under standard conditions, GCMS does not identify many of the heavier and more polar components without derivatization. In addition, sample preparation often requires dissection, extraction, and pooling samples from multiple individuals. For example, the analysis of drosophilid (fruit fly) pheromones typically requires 5 – 7 individuals yielding ca. 0.1 – 1 nmols of total lipids⁴⁻⁶. Purification and structural elucidation of lipid pheromones has required several thousand flies^{7,8}. Recently, laser desorption ionization (LDI) and matrix assisted laser desorption ionization (MALDI) MS have been used to characterize and image insect pheromone and lipid profiles directly from tissue^{7,9-11}. The

1
2
3 operation of LDI MS under intermediate pressure (~2 mbar) allows in-situ detection of a broad
4 range of lipid classes without derivatization or the application of a chemical matrix^{7, 9, 12, 13}. Finer
5 spatial resolution can be achieved with MALDI MSI approaches but necessitates a multi-step
6 preparation process that entails embedding, sectioning, and matrix application¹⁴.
7
8

9
10
11 Here, we describe the application of direct analysis in real time MS (DART MS), a form of
12 plasma-based ambient ionization, for pheromone gland analysis. DART MS ionization uses a
13 heated gas stream for analyte desorption through Penning ionization of atmospheric water and
14 subsequent gas-phase ionization of analytes^{15, 16}. The open air ionization geometry allows
15 individual tissue samples to be analyzed with minimal sample preparation, thus minimizing
16 artifacts, sample loss, and degradation¹⁷. The ion source produces a relatively large analysis
17 area, ranging from 5 – 40 mm², depending on the distance between the MS inlet and sample
18 and the exposure time¹⁸. Although the spatial resolution is not suitable for fine scale spatial
19 imaging of single cells, the configuration allows for rapid chemical profiling of entire insect
20 glands with minimal preparation. We show here the use of DART MS to characterize the
21 ejaculatory bulb (EB), a pheromone gland found in male *Drosophila*. Reproducible, species-
22 specific profiles of fatty acids (FAs), wax esters, diacylglycerides (DAGs), and triacylglycerides
23 (TAGs) can be generated from single pieces of tissue without derivatization or purification. In
24 addition, experimental parameters such as the ion source temperature and the application of a
25 solvent wash quantitatively and qualitatively alter the profile. Lastly, we show different
26 drosophilid subgroups or flies of different genotypes can be differentiated and classified on the
27 basis of untargeted chemical profiling of EBs.
28
29
30
31
32
33
34
35
36
37
38

39 **Results and Discussion**

40 *Single gland profiling of D. melanogaster*

41 To determine the suitability of DART MS for direct tissue analysis, we optimized sample
42 preparation and analytical parameters using the ejaculatory bulb (EB) of *D. melanogaster*. The
43 EB, located in the terminal portion of the male reproductive system, is a male pheromone-
44 producing gland that is conserved throughout the *Drosophila* family^{13, 19-21} (**Fig. 1A**). The EB
45 lipid contents have been well-characterized by GCMS and LDI MS and thus, the tissue serves
46 as a useful analytical standard. Individual EBs were dissected and placed on stainless steel
47 mesh (**Fig. 1A**). Five to seven biological replicates were analyzed using an automated linear rail
48 system that allows for uniform sample positioning and analysis time per replicate. Each gland
49 was exposed to the helium plasma for approx. 8 sec. Considering the diameter of the helium
50
51
52
53
54
55
56
57
58
59
60

1
2
3 beam (5 – 40 mm²) relative to the size of the gland (ca. 200 - 300 µm in diameter), the entirety
4 of the tissue is exposed during analysis. Direct analysis of freshly dissected EBs revealed 2
5 major peaks corresponding to two previously identified sex pheromones: 11Z-Octadecen-1-ol
6 acetate (*cis*-vacceanyl acetate; *m/z* 311.30)^{21, 22} and (3*R*,11*Z*,19*Z*)-3-acetoxy-11,19-octacosdien-
7 1-ol (CH503; *m/z* 465.43)^{7, 23} (**Fig. 1B**; **Supp. Fig. 1 & 2**). Two fragments with *m/z* reflecting the
8 loss of an acetate group from each pheromone (*m/z* 251.27 and *m/z* 405.41, respectively) are
9 also detected and are likely the result of thermal decomposition. DART MS analysis of synthetic
10 cVA and CH503 also shows the same loss of acetate in addition to CH503 dehydration (**Supp.**
11 **Fig. 2**). Other components of the gland that were identified on the basis of exact mass
12 measurements include fatty acids (FAs), DAGs, and TAGs (**Supp. Table 1**). The DAGs are
13 detected mainly as dehydrated ions ([M + H - H₂O]⁺) The loss of water observed is consistent
14 with the ionization behavior of a DAG standard analyzed under identical conditions (**Supp. Fig.**
15 **3**). Overall, the total ion chromatograms and mass spectral profiles were uniform between
16 biological replicates in terms of total intensity and composition (**Fig. 1C**; **Supp. Fig. 1**). The
17 abundance of various chemical classes changed qualitatively during the exposure period –
18 molecules below *m/z* 500 were more abundant in the initial 3 sec whilst DAGs and TAGs were
19 ionized mainly after 3-6 sec of exposure to the DART source (**Fig. 1D**). Some of the lipids,
20 particularly DAGs and TAGs, may be derived from the fat body, a storage organ that contains a
21 large amount of lipid reserves and surrounds the internal organs of insects²⁴. We have
22 previously been able to detect pheromones secreted from the gland without dissection by
23 analyzing whole flies with DART MS²⁵. However, compared to dissected tissue, reproducible
24 MS profiles are difficult to generate from whole flies due to variation in ion source placement. In
25 addition, other compounds from the external cuticle (e.g., hydrocarbons) are likely to also be
26 ionized making it difficult to identify gland-specific compounds.

27
28
29
30
31
32
33
34
35
36
37
38
39
40
41 Thermal decomposition is a common occurrence in the DART source and has been
42 observed for serum metabolites, sterols, and plant and animal oils²⁶⁻²⁸; as such, some of the FA
43 and DAG signals are also likely to be products of TAG decomposition. To differentiate
44 endogenously produced molecules from ionization artifacts, we compared the EB profile
45 generated with DART MS to *D. melanogaster* EB profiles previously obtained with laser
46 desorption ionization MS²⁰. LDI MS is a soft ionization method with minimal fragmentation²². As
47 with DART MS analysis, the wax ester pheromones cVA and CH503 in addition to multiple
48 species of FAs, DAGs, and TAGs were observed using LDI MS (**Supp. Table 1**). The detection
49 of intact DAGs by LDI MS strongly supports the endogenous presence of these molecules in the
50
51
52
53
54
55
56
57
58
59
60

1
2
3 EB but does not preclude the possibility that some of the DAG and FA signals observed in
4 DART MS spectra may also be the products of thermal decomposition (see below).
5
6

7 8 *Effect of temperature on EB tissue profile*

9 Increasing the temperature of the DART gas stream can significantly improve the ionization of
10 heavier, more polar molecules but may also lead to the rapid degradation of smaller, more
11 volatile compounds and increase the occurrence of ester hydrolysis²⁶. To characterize the
12 impact of temperature on EB lipids, we analyzed dissected tissue over a range of DART source
13 temperatures from 100 – 350 °C (**Fig. 2A**). Most compounds were reliably detected between
14 250 – 350 °C. Almost no signals were detected above baseline at temperatures below 150 °C.
15 Increasing temperature resulted in a greater abundance of deacetylated cVA and CH503 (**Fig.**
16 **2B**). The slight increase in DAG levels at higher temperatures may be the result of increasing
17 TAG decomposition. Our analysis of DAG and TAG standards confirmed that both lipid types
18 lose acyl chains with increasing temperature (**Supp. Fig. 3**). We conclude that direct tissue
19 analysis between 250 – 300 °C is optimal for detecting a breadth of lipid molecules. However,
20 products arising from the decomposition of ester lipids such as tri- and diacylglycerides and
21 acetate esters are inevitable and confound the differentiation of endogenous compounds from
22 fragment artifacts without a secondary method. Nonetheless, with mass isolation capability,
23 fragment ions can provide insight into the structural features and functional groups of lipids
24 produced in biological tissues.
25
26
27
28
29
30
31
32
33
34
35

36 *Quantitative profiling of biological tissue*

37 To investigate whether tissue profiling by DART MS can be used for quantitative analysis, we
38 added an external alkene standard, tricosene (C₂₃ H₄₆; [M+H]⁺ 323.26), directly to dissected
39 tissues at different concentrations (**Fig. 2C, D**). At an ion source temperature of 250 °C, the
40 normalized signal intensity increased linearly with sample concentration between 0.01 mM to 1
41 mM (R² = 0.99). Quantitative analysis is thus possible with the use of an appropriate internal
42 standard with similar vapor pressure to the analytes of interest. However, considering the
43 diverse array of endogenous lipid classes found within biological tissues, it will be necessary to
44 use multiple standards over a range of vapor pressures in order to match the heterogeneity of
45 compounds and their respective ionization efficiencies.
46
47
48
49
50
51
52

53 *Effect of solvent treatment on EB tissue profile*

54
55
56
57
58
59
60

1
2
3 The use of a tissue wash in MALDI imaging protocols is an effective method for removing ion
4 suppression by removing contaminants and excess salts²⁹⁻³². To test whether washing the
5 tissue prior to DART MS analysis could improve ionization and overall signal intensity by
6 removing salts and increasing membrane permeability, we applied solvents of various polarity to
7 dissected glands and assessed the responses of four categories of molecules found in the
8 gland. Our findings show that the area of the total ion chromatogram did not change significantly
9 between solvent conditions, indicating that overall ion counts were similar between each
10 condition (**Fig. 3A**). In contrast to no-solvent conditions, hexane and chloroform/ MeOH washes
11 lowered the coefficient of variation between biological replicates. It may be the case that
12 because biological tissue is heterogenous in its lipid composition and topology, the use of an
13 apolar wash helps to homogenize lipid distribution, resulting in lower variation between tissue
14 samples. Washing with hexane produced qualitative changes in the lipid profile - notably, DAG
15 signal intensity increased significantly, though some of the signals may be the products of TAG
16 fragmentation (**Fig. 3B; Supp. Fig. 4**). Overall, however, the use of solvents did not significantly
17 improve sensitivity and in fact, worsened the outcome by increasing sample-to-sample variation
18 or adding FA contamination. Analysis of solvent-only preparations showed FA signals, in some
19 cases at levels indistinguishable from measurements with tissue samples (**Supp. Fig. 5**).
20 Although the FA abundances of hexane- and MeOH-treated EBs were significantly higher than
21 background levels, nonetheless, the contamination hinders interpretation and quantitation. In
22 addition to FA contamination, EBs treated with EtOH exhibited a major signal at m/z 376.24 that
23 was found in only at low levels in other conditions. Based on predicted elemental composition
24 (C₂₆ H₃₄ O N) and high degree of conjugation (10 double bond equivalents), this molecule is
25 likely to be a contaminant rather than an endogenous lipid. These results indicate that pre-
26 treatment of the tissue with solvents provide little advantage for preliminary tissue profiling.
27 Moreover, it will be critical to use ultrapure solvents and sample preparation tools when applying
28 external standard solutions.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 *Classification by EB chemical profiles*

47 Pheromone and lipid profiles can evolve rapidly between species³³. For this reason,
48 chemotaxonomy using fatty acid profiles and other lipid features has been successfully
49 employed for the classification of bacterial and animal species³⁴⁻³⁶. Cuticular lipids have also
50 been proposed as a useful trait for taxonomic classification, especially between closely related
51 species with no obvious morphological differences^{37, 38}. We previously showed that the
52 cleavage products of unsaturated fatty acids induced by ozonolysis combined with DART MS
53
54
55
56
57
58
59
60

1
2
3 can be used for effective species classification³⁵. To test whether the EB chemical profiles
4 generated by DART MS can be applied to taxonomic classification, we used principal
5 components analysis (PCA) to classify EB profiles from 18 drosophilid species either belonging
6 to the Hawaiian *Drosophila* picturewing clade, repleta group, or melanogaster group. The MS
7 analyses from each species EB revealed a broad range of compounds ranging from m/z 150 –
8 620 (**Fig. 4A**). The members of the picturewing clade could be separated from other
9 phylogenetic groups with 100% accuracy (verified with Leave One Out Cross-Validation,
10 LOOCV) using PCA with spectral profiles. These findings indicate that endogenous lipid profiles
11 serve as a useful phenotypic feature for classifying insects and inferring phylogenetic
12 relationships. Many of the observed ions are novel molecules that have been missed by GCMS
13 and are reported for the first time here. Chemical classification will require further structural
14 elucidation by MS/MS. Moreover, in the absence of chromatographic separation and mass
15 isolation, some single peaks are likely to represent isobaric or isomeric lipids. Although the
16 former can be distinguished by pairing DART MS with detectors enabling higher mass
17 resolution, separation of the latter will require derivatization and/ or chromatographic separation.
18 Nonetheless, our results show that DART MS-generated chemical profiles can be used as
19 taxonomic markers to distinguish invertebrate species even in the absence of complete
20 structural elucidation.
21
22
23
24
25
26
27
28
29
30

31 We used the same clustering method to identify *D. melanogaster* transgenic lines with
32 aberrant EB profiles. The ability to generate mutants and screen rapidly for phenotypic
33 differences is a significant advantage of the *Drosophila* model system. We profiled the EBs of
34 transgenic lines in which the expression of individual FA biosynthesis genes was genetically
35 knocked down. Individuals with experimentally-induced changes in the EB lipid profile could be
36 differentially separated from genetic controls using linear discriminant analysis with 92.5%
37 accuracy (LOOCV; **Fig 4B**). One sample could not be classified. The results demonstrate that
38 DART MS profiling of tissues is an effective method for the rapid screening of anomalous
39 chemical profiles. This feature will allow mutants with defects in lipid metabolism to be swiftly
40 identified.
41
42
43
44
45
46
47
48

49 **Conclusions**

50 The direct analysis of tissue by DART MS generates rapid metabolite and lipid profiling from
51 single pieces of tissue. This application will be particularly useful for samples that are rare or
52 available in limited quantities. However, structural elucidation and distinguishing endogenous
53 products from thermal decomposition artifacts will require a secondary form of MS confirmation.
54
55
56
57
58
59
60

1
2
3 Both apolar and heavier, more polar molecules can be detected from single tissues with high
4 reproducibility. In principal, quantitation with direct DART MS analysis of tissue is possible with
5 the addition of an external standard although prior knowledge of the analyte concentration as
6 well as its chemical properties will be required to select an appropriate standard with similar
7 ionization properties. In addition, environmental contaminants, such as fatty acids, and signals
8 resulting from in-source decomposition are commonplace and can be difficult to distinguish from
9 endogenous products.
10
11
12
13

14 The ability to generate reproducible small molecule profiles based on relative abundances
15 allows divergent phylogenetic lineages to be separated and phenotypic changes arising from
16 genetic manipulation to be readily identified. The application of DART MS to tissue profiling will
17 help to expand the “secretome” of molecules produced by pheromone glands as well as provide
18 preliminary identification of natural products of interest produced within specialized
19 compartments. Finally, the pairing of genetic screening with DART MS profiling can be extended
20 towards metabolite and lipid profiling from single organs or biological compartments of other
21 genetically tractable model organisms and applied to pharmaceutical studies for tissue-specific
22 screening and detection of drug metabolites.
23
24
25
26
27
28
29

30 **Experimental**

31 *Materials*

32 All chemicals, unless otherwise stated, were purchased from Sigma-Aldrich (St. Louis, MO,
33 USA).
34
35
36
37

38 *Animals*

39 Hawaiian subgroup species *D. grimshawi*, *D. hemipeza* and, *D. silvestris* were obtained from
40 Hawaiian Drosophila Research Stock Center and Insectary for Scientific Training and Advances
41 in Research facility at the Univ. of Hawai'i (<https://icemhh.pbrc.hawaii.edu/instar/>). Other picture-
42 wing species (*D. pullipes*, *D. crucigera*, *D. adiantola*, *D. fasciculisetae*, *D. sproati*, and *D.*
43 *ochracea* were collected from the Koke'e State Park on Kaua'i, the Nature Conservancy
44 Waikamoi Preserve on Maui, South Kona Forest Reserve on Hawai'i Island, and the Honouliuli
45 Forest Reserve and O'ahu. All other species were obtained from the Bloomington Drosophila
46 Stock Center (Indiana, USA), National Drosophila Species Stock Center (Ithaca, NY, USA), and
47 the Evolutionary Genetics Laboratory in Tokyo Metropolitan University (Japan). When possible,
48 virgin males were used. Male reproductive status could not be confirmed with wild caught flies.
49 Hawaiian flies were maintained on Clayton-Wheeler diet ³⁹ at 19 °C and other species were
50
51
52
53
54
55
56
57
58
59
60

1
2
3 raised at 25 °C on a 12 hours light/dark cycle on a standard diet of cornmeal, agar, dextrose,
4 glucose, and 1.5% inactive yeast. Transgenic lines were generated using standard *Drosophila*
5 genetic methods by crossing *bond-GAL4* females²⁰ to one the following *UAS-RNAi* lines from
6 the Vienna Drosophila Resource Center: VDRC ID 46513, 23134, 1166, 7492, or 48662.
7
8
9

10 11 *Sample preparation for mass spectrometry*

12 Single glands from males were dissected in phosphate buffered saline (PBS; 137 mM NaCl, 2.7
13 mM KCl, 10 mM Na₂HPO₄, 1.8 mM KH₂PO₄, pH 7.4) and placed directly on a stainless steel
14 grid (DART QuickStrips; IonSense (IonSense LLC, Saugus, MA). For solvent studies, 1 µL of
15 solvent was added to the tissue surface or an empty grid (for negative controls) immediately
16 after dissection and allowed to evaporate. The following HPLC-grade solvents were tested:
17 hexane, chloroform/ methanol (2:1 v/v), methanol, and ethanol. Five to seven replicates were
18 prepared for each sample type. Immediately after dissection and placement on the sample grid,
19 the strip was placed on a DART SVP linear rail system and moved through the ion source at 1
20 mm/sec at a distance of 0.5 cm from the inlet. Synthetic tributyrin, diolein (Cayman Chemical;
21 Ann Arbor, MI, USA), *cis*-vaccenyl acetate (Cayman Chemical) and CH503 (previously
22 synthesized⁴⁰) were prepared at 1 µM in hexane and 1 µL added to the sample grid.
23
24
25
26
27
28
29
30

31 *Direct tissue quantitation of tricosene*

32 Five concentrations of tricosene diluted in hexane were tested: 0.0001, 0.001, 0.01, 0.1, and 1
33 mM. One µL of the standard was added to the EB immediately following placement on the grid.
34 Five replicate EBs from 3 day old male flies were prepared for each concentration.
35
36
37
38

39 *Sample preparation for light microscopy*

40 The EB from 3-4 day old males were dissected in PBS, fixed in 4 % paraformaldehyde for 10
41 min, then mounted on a glass slide in 50% glycerol. Digital Images were obtained with an
42 Olympus BX51 microscope equipped with a Leica DFC 7000T color digital camera.
43
44
45
46

47 *Mass spectrometry*

48 Mass spectra were acquired with an atmospheric pressure ionization time-of-flight mass
49 spectrometer (AccuTOF-DART 4G, JEOL USA, Inc., Peabody, MA) equipped with a DART SVP
50 ion source (IonSense LLC, Saugus, MA) interface, placed 1 cm away from the sampling orifice.
51 The instrument has a resolving power of 10,000 (FWHM definition) at *m/z* 500. Voltage settings
52 and acquisition parameters for positive ion mode are as previously described³⁵. Briefly, the RF
53
54
55
56
57
58
59
60

1
2
3 ion guide voltage was set at 500 V and the detector voltage set at 2200 V. The atmospheric
4 pressure ionization interface potentials were as follows: orifice 1 = -15 or -21 V, orifice 2 = -5 V,
5 ring lens = -5 V. Mass spectra were stored at a rate of one spectrum per second with an
6 acquired m/z range of 100 – 1000. Calibration for exact mass measurements was accomplished
7 by acquiring a mass spectrum of polyethylene glycol (average molecular weight 600) as an
8 external reference standard in every data file.
9
10
11
12
13

14 *Fly-assisted laser desorption ionization mass spectrometry*

15 Individual flies were attached to a custom-milled sample plate using double-sided tissue tape
16 (7111, Louis Adhesive Tapes, Thailand). No matrix was used. Mass spectra were generated
17 using a QSTAR Elite (AB Sciex, Toronto, CA) mass spectrometer equipped with a modified
18 oMALDI2 ion source (AB Sciex) and a N₂ laser ($\lambda=337$ nm) operated at a repetition rate of
19 40 Hz^{9, 20}. Two mbar of N₂ gas was used create the buffer gas environment for generation of
20 ions. [M+K]⁺ potassium-bearing compounds constitute the major ion species. Spectra were
21 internally calibrated to chitin signals at [M+K]⁺ 242.04, 648.20 and 851.28. Mass accuracy was
22 ~20 p.p.m. All spectra were acquired in positive ion mode and processed using MS Analyst
23 software (Analyst QS 2.0, AB Sciex).
24
25
26
27
28
29
30

31 *Data analysis*

32 The JEOL MassCenter software (v. 1.3.0.1) was used to calculate the area under the total or
33 extracted ion chromatograms and to generate centroided profiles. Centroided profiles of each
34 technical replicate were generated by summing signals within a 1 sec window of measurement.
35 Preliminary determination of chemical categories is based on elemental composition prediction,
36 predicted double bond equivalents, and isotope matching as performed with Mass Mountaineer
37 (v. 5.1.13.0, RBC Software). For temperature studies, raw intensity counts for all centroided
38 signals corresponding to a single chemical category were summed and normalized to the total
39 area of all signals. For fragmentation studies, fragment percentage was calculated by dividing
40 the signal abundance of the fragment by the sum of all parent ion adducts (for cVA and CH503)
41 or by the sum of fragments and parent ion adducts (for TAG and DAG standards). For tricosene
42 quantitation, the area beneath the extracted ion chromatogram for tricosene ([M+H]⁺ 323.36)
43 was used to calculate abundance. For solvent studies, the area of the extracted ion
44 chromatograms corresponding to each chemical category was normalized to the area beneath
45 the total ion chromatogram. The relative abundances from 5-7 replicates were summed and
46 averaged. Background levels of FAs were established by averaging the absolute abundances
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 from 5 replicate measurements. Statistical analysis was performed with Prism v.9.2 (GraphPad
4 Software, San Diego, CA).
5
6
7

8 *Classification analysis*

9 Centroided mass spectra were exported as text files using MassCenter for classification and
10 analysis by Mass Mountaineer. For linear discriminant analysis (LDA) and kernel principal
11 component analysis (KPCA), a signal threshold of 10% was used to extract features from
12 centroided mass spectral data. Five m/z values (for LDA) and 11 m/z values (for KPCA) with a
13 mass tolerance of 10 mmu were used in the training set. The normalized relative abundance of
14 each of the features was calculated relative to the highest intensity peak for the spectrum. Five
15 to seven biological replicates for each species or genotype were used for classification.
16 Predictive models were tested with leave on out cross validation (LOOCV).
17
18
19
20
21
22
23

24 **Figure legends**

25 **Figure 1.** DART MS analysis of the ejaculatory bulb (EB). **(A)** Image of a single EB visualized
26 under differential interference contrast illumination and placed on the stainless steel grid sample
27 holder; aed: anterior ejaculatory duct; scale bar: 100 μm . **(B)** Chemical structures and predicted
28 m/z of the two major pheromones produced in the EB: *cis*-vaccenyl acetate (cVA) and
29 (3*R*,11*Z*,19*Z*)-3-acetoxy-11,19-octacosdien-1-ol (CH503). Fragments resulting from the loss of
30 acetate groups are shown. **(C)** The total ion chromatogram from 7 EBs placed on a single
31 QuickStrip shows relatively uniform intensity between each replicate; a.u.: arbitrary units. **(D)**
32 Representative DART MS spectrum from a single gland in the first 3 sec of introduction to the
33 ion source and the subsequent 3-6 sec of analysis. Mass signals corresponding to cVA, CH503,
34 deacetylated fragments, fatty acids (FA), diacylglycerides (DAGs) and triacylglycerides (TAGs)
35 are labeled. Profiles were acquired at 300 °C. Some of the FA and DAGs may be the products
36 of TAG decomposition.
37
38
39
40
41
42
43
44
45

46 **Figure 2.** Effect of DART gas temperature on direct tissue analysis signal abundance and
47 fragmentation. **(A)** Signals corresponding to wax ester pheromones, diacylglycerides (DAG) and
48 fatty acids (FA) were detected reliably at 250 °C and above. **(B)** Fragmentation due to the loss
49 of the acetate group from cVA or CH503 was significantly higher at 350 °C compared with 150
50 °C (cVA) or 250 °C (CH503); ***: $p=0.0002$, ****: $p<0.0001$, Holm-Sidak's multiple comparisons
51 test. **(C)** Representative DART MS spectrum measured at 300 °C from an EB spiked with 100
52 nmols tricosene. **(D)** The signal intensity of tricosene spiked onto the tissue increases linearly
53
54
55
56
57
58
59
60

1
2
3 with concentration between 0.01 – 1 mM ($R^2 = 0.99$). For all graphs, bars indicate mean \pm
4 S.E.M; n=5-7 EBs per point.
5
6
7

8 **Figure 3.** Addition of a solvent wash produces qualitative changes in DART MS profile. **(A)** The
9 area beneath the total ion chromatogram did not change significantly between each of the
10 solvent conditions (ANOVA, $p=0.3773$). The coefficient of variance (CV) is shown for each
11 solvent condition; NS: no solvent, EtOH: ethanol, MeOH: methanol, Hex: hexane, CM:
12 chloroform/ methanol. **(B)** The relative abundance of various components of the EB profile
13 change with solvent conditions. The DAG abundance increased significantly with hexane
14 application whilst levels of an unknown compound (C26 H34 ON), likely a contaminant, was
15 most evident with EtOH wash. Hexane produced the least variability between replicates. The
16 mean with maximum and minimum values is shown in each plot; n=3-5 per condition. All
17 spectra were acquired at 300 °C.
18
19
20
21
22
23
24

25 **Figure 4.** Classification by DART MS EB profiles. **(A)** Principal components analysis of the EB
26 from 18 different drosophilid species differentiated between members of the Hawaiian
27 *Drosophila* picture-wing subgroup (PWD) and nonPWD flies (LOOCV: 100%). Representative
28 DART MS spectra reveal differences in chemical profile complexity from *D. pullipes* (PWD) and
29 *D. suzukii* (nonPWD). **(B)** Linear discriminant analysis separated EB profiles of genetic controls
30 from that of transgenic lines in which the expression of fatty acid-related genes in the EB have
31 been genetically repressed (LOOCV: 96%). Representative DART MS spectra from a control
32 genetic line (*bond-GAL4/ +*) and transgenic line (*bond-GAL4/ uas-bond-RNAi*) show the
33 absence of CH503 at $[M+H]^+ 465.43$ following repression of the gene *bond*, a very long chain
34 fatty acid elongase²⁰. All EBs were measured at 300 °C.
35
36
37
38
39
40
41
42

43 **Author Contributions**

44 NC: methodology and investigation.; JYY: Conceptualization, methodology, investigation,
45 validation, formal analysis, resources, and writing – original draft, review, and editing.
46 There are no conflicts to declare.
47
48
49

50 **Acknowledgements**

51 The DART MS instrument was acquired through a grant from the Department of Defense, U.S.
52 Army Research Office (W911NF1610216 to J.Y.Y). The work is supported by the National
53 Institutes of Health (Grant No. P20GM125508), Hawai'i Community Foundation (19CON-
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

95452), and the National Science Foundation (Grant No. 20256669). This article reports data obtained at the University of Hawai'i at Mānoa Microbial Genomics and Analytical Laboratory Core which is supported by the National Institutes of Health (P20GM125508). We thank Kenneth Kaneshiro, Kelvin Kanegawa, Matthew J. Medeiros, Donald Price, Kelli Konicek and the Hawaiian Drosophila Research Stock Center for providing fly samples.

References

1. Wyatt, T. D., *Pheromones and animal behavior: Chemical Signals and Signatures*. 2nd edition. ed.; Cambridge University Press: Cambridge, 2014.
2. Raven, P. H.; Wagner, D. L., Agricultural intensification and climate change are rapidly decreasing insect biodiversity. *Proc Natl Acad Sci U S A* **2021**, *118* (2), e2002548117.
3. Yew, J. Y.; Chung, H., Insect pheromones: An overview of function, form, and discovery. *Prog Lipid Res* **2015**, *59*, 88-105.
4. Foley, B. R.; Telonis-Scott, M., Quantitative genetic analysis suggests causal association between cuticular hydrocarbon composition and desiccation survival in *Drosophila melanogaster*. *Heredity* **2011**, *106* (1), 68-77.
5. Pardy, J. A.; Rundle, H. D.; Bernards, M. A.; Moehring, A. J., The genetic basis of female pheromone differences between *Drosophila melanogaster* and *D. simulans*. *Heredity* **2019**, *122* (1), 93-109.
6. Dembeck, L. M.; Böröczky, K.; Huang, W.; Schal, C.; Anholt, R. R.; Mackay, T. F., Genetic architecture of natural variation in cuticular hydrocarbon composition in *Drosophila melanogaster*. *eLife* **2015**, *4*.
7. Yew, J. Y.; Dreisewerd, K.; Luftmann, H.; Müthing, J.; Pohlentz, G.; Kravitz, E. A., A new male sex pheromone and novel cuticular cues for chemical communication in *Drosophila*. *Curr Biol* **2009**, *19* (15), 1245-54.
8. Brieger, G.; Butterworth, F. M., *Drosophila melanogaster*: identity of male lipid in reproductive system. *Science* **1970**, *167* (3922), 1262.
9. Yew, J. Y.; Soltwisch, J.; Pirkl, A.; Dreisewerd, K., Direct laser desorption ionization of endogenous and exogenous compounds from insect cuticles: practical and methodologic aspects. *J Am Soc Mass Spectrom* **2011**, *22* (7), 1273-84.
10. Kaftan, F.; Vrkoslav, V.; Kynast, P.; Kulkarni, P.; Böcker, S.; Cvačka, J.; Knaden, M.; Svatoš, A., Mass spectrometry imaging of surface lipids on intact *Drosophila melanogaster* flies. *J Mass Spectrom* **2014**, *49* (3), 223-32.
11. Vrkoslav, V.; Muck, A.; Cvačka, J.; Svatoš, A., MALDI imaging of neutral cuticular lipids in insects and plants. *J Am Soc Mass Spectrom* **2010**, *21* (2), 220-31.
12. Chin, J. S.; Ellis, S. R.; Pham, H. T.; Blanksby, S. J.; Mori, K.; Koh, Q. L.; Etges, W. J.; Yew, J. Y., Sex-specific triacylglycerides are widely conserved in *Drosophila* and mediate mating behavior. *eLife* **2014**, *3*, e01751.
13. Ng, S. H.; Shankar, S.; Shikichi, Y.; Akasaka, K.; Mori, K.; Yew, J. Y., Pheromone evolution and sexual behavior in *Drosophila* are shaped by male sensory exploitation of other males. *Proc Natl Acad Sci U S A* **2014**, *111* (8), 3056-61.
14. Niehoff, A. C.; Kettling, H.; Pirkl, A.; Chiang, Y. N.; Dreisewerd, K.; Yew, J. Y., Analysis of *Drosophila* lipids by matrix-assisted laser desorption/ionization mass spectrometric imaging. *Anal Chem* **2014**, *86* (22), 11086-92.
15. Cody, R. B., Observation of molecular ions and analysis of nonpolar compounds with the direct analysis in real time ion source. *Anal Chem* **2009**, *81* (3), 1101-7.
16. Cody, R. B.; Laramée, J. A.; Durst, H. D., Versatile new ion source for the analysis of materials in open air under ambient conditions. *Anal Chem* **2005**, *77* (8), 2297-302.
17. Chernetsova, E. S.; Morlock, G. E., Determination of drugs and drug-like compounds in different samples with direct analysis in real time mass spectrometry. *Mass Spectrom Rev* **2011**, *30* (5), 875-83.
18. Newsome, G. A.; Kayama, I.; Brogdon-Grantham, S. A., Direct analysis in real time mass spectrometry (DART-MS) of discrete sample areas without heat damage. *Anal Methods* **2018**, *10* (9), 1038-1045.
19. Bairati, A.; Depetris, S., [Research on the Nature and Ultrastructure of Keloids. li. Diffractographic and Electron Microscopic Studies]. *Boll Soc Ital Biol Sper* **1963**, *39*, 1222-4.

- 1
- 2
- 3
- 4 20. Ng, W. C.; Chin, J. S.; Tan, K. J.; Yew, J. Y., The fatty acid elongase Bond is essential for
- 5 Drosophila sex pheromone synthesis and male fertility. *Nat Commun* **2015**, *6*, 8263.
- 6 21. Guiraudie-Capraz, G.; Pho, D. B.; Jallon, J. M., Role of the ejaculatory bulb in biosynthesis
- 7 of the male pheromone cis-vaccenyl acetate in *Drosophila melanogaster*. *Integr Zool* **2007**,
- 8 *2* (2), 89-99.
- 9 22. Butterworth, F. M., Lipids of *Drosophila*: a newly detected lipid in the male. *Science* **1969**,
- 10 *163* (3873), 1356-7.
- 11 23. Shikichi, Y.; Shankar, S.; Yew, J. Y.; Mori, K., Synthesis and bioassay of the eight
- 12 analogues of the CH503 male pheromone (3-acetoxy-11,19-octacosadien-1-ol) of the
- 13 *Drosophila melanogaster* fruit fly. *Biosci Biotechnol Biochem* **2013**, *77* (9), 1931-8.
- 14 24. Arrese, E. L.; Soulages, J. L., Insect fat body: energy, metabolism, and regulation. *Annual*
- 15 *review of entomology* **2010**, *55*, 207-225.
- 16 25. Yew, J. Y.; Cody, R. B.; Kravitz, E. A., Cuticular hydrocarbon analysis of an awake
- 17 behaving fly using direct analysis in real-time time-of-flight mass spectrometry. *Proc Natl*
- 18 *Acad Sci U S A* **2008**, *105* (20), 7135-40.
- 19 26. Zhou, M.; McDonald, J. F.; Fernández, F. M., Optimization of a direct analysis in real
- 20 time/time-of-flight mass spectrometry method for rapid serum metabolomic fingerprinting. *J*
- 21 *Am Soc Mass Spectrom* **2010**, *21* (1), 68-75.
- 22 27. Yu, S.; Crawford, E.; Tice, J.; Musselman, B.; Wu, J. T., Bioanalysis without sample
- 23 cleanup or chromatography: the evaluation and initial implementation of direct analysis in
- 24 real time ionization mass spectrometry for the quantification of drugs in biological matrixes.
- 25 *Anal Chem* **2009**, *81* (1), 193-202.
- 26 28. Alberici, R. M.; Fernandes, G. D.; Porcari, A. M.; Eberlin, M. N.; Barrera-Arellano, D.;
- 27 Fernández, F. M., Rapid fingerprinting of sterols and related compounds in vegetable and
- 28 animal oils and phytosterol enriched- margarines by transmission mode direct analysis in
- 29 real time mass spectrometry. *Food Chem* **2016**, *211*, 661-8.
- 30 29. Wang, H.-Y. J.; Liu, C. B.; Wu, H.-W., A simple desalting method for direct MALDI mass
- 31 spectrometry profiling of tissue lipids. *J Lipid Res* **2011**, *52* (4), 840-849.
- 32 30. Buchberger, A. R.; Vu, N. Q.; Johnson, J.; DeLaney, K.; Li, L., A Simple and Effective
- 33 Sample Preparation Strategy for MALDI-MS Imaging of Neuropeptide Changes in the
- 34 Crustacean Brain Due to Hypoxia and Hypercapnia Stress. *J Am Soc Mass Spectrom* **2020**,
- 35 *31* (5), 1058-1065.
- 36 31. Seeley, E. H.; Oppenheimer, S. R.; Mi, D.; Chaurand, P.; Caprioli, R. M., Enhancement of
- 37 protein sensitivity for MALDI imaging mass spectrometry after chemical treatment of tissue
- 38 sections. *J Am Soc Mass Spectrom* **2008**, *19* (8), 1069-1077.
- 39 32. Thomas, A.; Patterson, N. H.; Laveaux Charbonneau, J.; Chaurand, P., Orthogonal organic
- 40 and aqueous-based washes of tissue sections to enhance protein sensitivity by MALDI
- 41 imaging mass spectrometry. *J Mass Spectrom* **2013**, *48* (1), 42-8.
- 42 33. Symonds, M. R. E.; Elgar, M. A., The evolution of pheromone diversity. *Trends Ecol Evol*
- 43 **2008**, *23* (4), 220-228.
- 44 34. Cody, R. B.; McAlpin, C. R.; Cox, C. R.; Jensen, K. R.; Voorhees, K. J., Identification of
- 45 bacteria by fatty acid profiling with direct analysis in real time mass spectrometry. *Rapid*
- 46 *Commun Mass Spectrom* **2015**, *29* (21), 2007-12.
- 47 35. Cetraro, N.; Cody, R. B.; Yew, J. Y., Carbon-carbon double bond position elucidation in
- 48 fatty acids using ozone-coupled direct analysis in real time mass spectrometry. *Analyst*
- 49 **2019**, *144* (19), 5848-5855.
- 50 36. Shaw, N., Lipid composition as a guide to the classification of bacteria. *Adv Appl Microbiol*
- 51 **1974**, *17* (0), 63-108.
- 52 37. Kather, R.; Martin, S. J., Cuticular hydrocarbon profiles as a taxonomic tool: advantages,
- 53 limitations and technical aspects. *Physiol Entomol* **2012**, *37* (1), 25-32.
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3 38. Niogret, J.; Felix, A. E.; Nicot, A.; Lumaret, J. P., Chemosystematics Using Cuticular
4 Compounds: A Powerful Tool to Separate Species in Mediterranean Dung Beetles
5 (Coleoptera: Geotrupidae). *J Insect Sci* **2019**, *19* (2).
6
7 39. Wheeler, M. R.; Clayton, F. E., A new *Drosophila* culture technique. *Dros Info Serv* **1965**,
8 *40*, 98.
9 40. Mori, K.; Shikichi, Y.; Shankar, S.; Yew, J. Y., Pheromone synthesis. Part 244: Synthesis of
10 the racemate and enantiomers of (11Z,19Z)-CH503 (3-acetoxy-11,19-octacosadien-1-ol), a
11 new sex pheromone of male *Drosophila melanogaster* to show its (S)-isomer and racemate
12 as bioactive. *Tetrahedron* **2010**, *66* (35), 7161-7168.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60