



Cite this: *Green Chem.*, 2023, **25**, 3398

Natural multi-osmolyte cocktails form deep eutectic systems of unprecedented complexity: discovery, affordances and perspectives

Marina Cvjetko Bubalo, ^{*a} Thanos Andreou, ^{*b} Manuela Panić, ^a Mia Radović, ^a Kristina Radošević^a and Ivana Radojčić Redovniković^a

While exaptation is most impressive when links between very dissimilar contexts are established, it can be even more pervasive when the previously unestablished connection seems surprisingly obvious in retrospect. We herein established such a connection between two major research fields previously advancing in parallel: osmolytes and deep eutectic solvents. Osmolytes are small molecules produced in cells as a response to external stimuli. Based on their individual interaction with macromolecules, single osmolytes are currently categorized as “kosmotropes” (stabilizing proteins) and “chaotropes” (destabilizing them). However, with two or more osmolytes, synergistic effects were also observed on top of cumulative ones. All current attempts to explain these synergistic effects have been studying osmolyte–osmolyte interactions in aqueous solutions, but none has been generally applicable so far, indicating that a new model “beyond kosmotropes and chaotropes” is needed to understand the function of osmolytes. We have gathered enough evidence to formulate a hypothesis of such a new model. First, inspired by patterns frequently observed in nature, five major stabilizing osmolytes (kosmotropes) prominent across kingdoms (trimethylamine *N*-oxide, sarcosine, glycerophosphorylcholine, dimethylsulfoniopropionate and ectoine) were for the first time employed to form novel two- and three-component DESs with all known natural perturbants/chaotropes (urea, guanidine hydrochloride and arginine). Going beyond the current three-component barrier, we mimicked the exact composition of multi-osmolyte cocktails widely observed in nature and we here report the rapid and consistent formation of deep eutectic systems of unprecedented complexity and the tunable potential of these new systems to stabilize a template protein. Based on these observations, we postulate that, *in vivo*, osmolytes form deep eutectic systems featuring new, emergent and synergistic properties which govern their interaction with macromolecules. We believe that such bio-inspired, osmolyte-based DESs can be a remarkable new tool to study complex natural systems in higher granularity and to engineer their microenvironment towards efficient and sustainable processes at scale.

Received 16th December 2022,

Accepted 6th April 2023

DOI: 10.1039/d2gc04796a

rsc.li/greenchem

Introduction

Observing nature has been one of the most successful ways to find inspiration. Humans have long used this inspiration, among other ends, to find new ways to solve complex problems through engineering, chemistry and biotechnology. Looking at the way living creatures cope with changes in the environment and understanding the functions observed in organisms and processes in nature can help us shape and create new products, processes, and systems.¹ We can learn from biological systems frequently exposed to harsh environments, from extremophilic bacteria, marine organisms, sporulating microorgan-

isms and plants, to living assemblages functioning in complex microenvironments, such as the mammal kidney. All these living systems share a similar mechanism of coping with the stressors of a harsh environment: the accumulation of small molecules commonly called osmolytes (Fig. 1). These versatile organic compounds have been attributed with several biological functions, among which the most pronounced one is the increase in the thermodynamic stability of macromolecules without compromising their native functionality. For this reason, osmolyte-induced stability of biomacromolecules has attracted considerable attention in various industrial fields. Interestingly, the exact mechanisms and interactions involved in this stabilizing effect have been a subject of debate and still remain largely unresolved.²

Meanwhile, neoteric systems that effectively mimic the natural environment for various biomolecules, the so-called Deep Eutectic Solvents (DESs), have been intensively studied

^aUniversity of Zagreb, Faculty of Food Technology and Biotechnology, Croatia.

E-mail: mcvjetko@pbf.hr

^bVIO Chemicals AG, Thessaloniki, Greece. E-mail: thanos.andreou@viochem.com






Fig. 1 Deep eutectic solvents and osmolytes: the two research fields that have been advancing in parallel. When DESs are considered as Complex Adaptive Systems, it becomes apparent that “exhibiting properties beyond the sum of their parts” is the concept that ontologically defines them. In parallel, the paradoxes observed when specific mixtures of osmolytes interact with proteins reveal that the effects of such mixtures are also beyond cumulative, leading us to explore the hypothesis of a common “lowest level of coherence” between these two research fields.

as nontoxic and highly tunable solvents in food, agrochemicals, cosmetics, and pharmaceuticals production (Fig. 1).³ Unlike conventional molecular solvents, a DES is a mixture of two or more, usually solid compounds combined in a suitable molar ratio to form a liquid at ambient temperature.⁴ A particular subgroup of these systems, Natural DESs (NADESs), only consist of compounds that occur in nature.⁵ In most cases, these compounds are strikingly similar, if not identical, to naturally occurring osmolytes, *i.e.* sugars and their derivatives, polyols, amino acids, and quaternary ammonium compounds.

Based on previous hypotheses that DES may be formed *in vivo* and may be responsible for the solubilisation and storage of biomolecules,⁶ in this paper we experimentally prove that, across kingdoms and environments, context-based combinations of osmolytes form multicomponent eutectic systems which help maintain the native conformation and functionality of proteins and other biomolecules under adverse conditions. These new findings provide an excellent opportunity to engineer new, osmolyte-based solvents and systems which are directly inspired by the natural microenvironment of biomacromolecules and can therefore mimic it effectively.

Naturally occurring osmolytes

To ensure their survival, persistence and growth, various organisms, such as animals, plants, and microorganisms, must respond in a timely manner to a myriad of stressful conditions and nutrient limitations they face in their natural environments. In particular, the loss of internal water due to drought, extreme temperatures or diseases that cause osmotic imbalance, is a common threat because it results in high concentrations of salts and organic solutes.² To maintain osmotic

balance with their environment and prevent perturbations that can cause structural changes in cellular proteins, most organisms use osmolytes, small, electrically neutral and nontoxic organic molecules.^{2,7}

While molecules acting as osmolytes vary across kingdoms, all known osmolytes can be grouped into a few main chemical categories: (i) polyols and sugar polyols (*e.g.* glycerol, sorbitol, xylitol) found in all kingdoms (ii) sugars and their derivatives (*e.g.* glucose, sucrose, trehalose) accumulated mainly in plants, insects, and polar fish; (iii) amino acids and their derivatives (*e.g.* glycine, proline, ectoine, taurine) found mainly in prokaryotic cells and plants; (iv) methylamines (*e.g.* trimethylamine *N*-oxide – TMAO, sarcosine, betaine) found mainly in marine fishes and plants; (v) methylsulfonium compounds (*e.g.* dimethylsulfoniopropionate-DMSP) found in marine organisms; (vi) γ -conjugated compounds (*e.g.* ureas and guanidines) used by mammals and marine life.^{7–9} The diversity of osmolytes, their multiple biological functions and their nontoxicity over a wide range of concentrations have led to their wide application in biotechnology, agriculture, and medicine, primarily as protein stabilizers and cell protectants (Table 1).¹⁰

Apart from their chemical structure, osmolytes have been categorized in two major groups, according to their impact on macromolecules, particularly proteins.¹¹ Urea, guanidine hydrochloride (guanidine HCl) and, although somewhat controversial,¹² arginine (Arg) are denaturing or perturbing osmolytes, commonly referred to as chaotropes, because they have been observed to disrupt the structure and function of macromolecules. On the other hand, methylamines and methylsulfonium compounds, carbohydrates, polyols, amino acids and their derivatives are referred to as kosmotropes and compatible solutes because they push the equilibrium of protein folding towards the native form in various stressful situations, often counteracting the destabilizing effect of chaotropes.¹²



Table 1 Naturally occurring osmolytes (classification, origin, function and applications)^{2,7-10}

| Osmolytes | Origin | Function | Examples of osmolytes applications |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>Polyols and sugar polyols</p> <p>Glycerol, mannitol, sorbitol, trehalose, inositol, <i>D</i>-methyl-inositol, glucosylglycerol, mannosylglycerol, arabitol, erythritol, sulfotrehalose, threitol, pinitol, <i>D</i>-ononitol, <i>L</i>-querbrachitol</p> | <p>Generally present in all kingdoms</p> | <ul style="list-style-type: none"> Freezing tolerance Cell water retention while remaining compatible with macromolecular function Stabilization of proteins Osmotic adjustment Signaling molecules Stabilization of membranes (sugars and their derivatives) | <ul style="list-style-type: none"> <i>Mannitol</i>, <i>glycerol</i> – cryoprotection <i>Glycerol</i>, <i>sorbitol</i> – treatment of neurodegenerative diseases <i>Mannitol</i> – excipient for pharmaceutical formulations, stabilization and activation of therapeutic proteins <i>Sorbitol</i> – vaccines production and stabilization <i>Erythritol</i> – treatment of dry eye syndrome <i>Sucrose</i> – excipient for pharmaceutical formulations <i>Glucose</i>, <i>sucrose</i>, <i>trehalose</i> – cryoprotection <i>Sucrose</i>, <i>trehalose</i> – vaccine production, stabilization and formulation <i>Trehalose</i> – hypothermic storage of human organs, neuroprotection |
| <p>Sugars and their derivatives</p> <p>Glucose, sucrose, fructose, trehalose, maltose, mannose, raffinose, rhamnose, xylose, α-glucosylglycerate, α-mannosylglycerate</p> | | | |
| <p>Amino acids and their derivatives</p> <p>Glycine, alanine, proline, valine, serine, isoleucine, histidine, taurine, glutamine, hypotaurine, aspartic acid, thiotaurine, ectoine, hydroxyectoine, octopine, citrulline, poly-γ-glutamic acid, <i>N</i>,γ-acetyldiaminobutyrate, <i>N</i>ϵ-acetyl-<i>L</i>-lysine</p> | <p>Found principally in prokaryotic cells and vascular plants and some mammalian organs</p> | <ul style="list-style-type: none"> Precursors for most of the osmolytes Alleviation of cytoplasmic acidosis Preventing membrane damage and ion toxicity strongly perturbing effects on enzyme catalytic velocity and protein structural stability | <ul style="list-style-type: none"> <i>Histidine</i>, <i>glycine</i>, <i>arginine</i>, <i>proline</i>, <i>L-glutamate</i> – excipients for pharmaceutical formulations <i>Ectoine</i> – skin protection, anti-inflammatory treatment, inhibitory effects in neurodegenerative diseases <i>Taurine</i> – congestive heart failure, anemia, neuroprotection <i>Proline</i> – vaccine production, flocculation and stabilization, skin healing |
| <p>Methyl-ammonium compounds</p> <p>Betaine, choline, trimethylamine oxide (TMAO), <i>N</i>-methyltaurine, sarcosine, glycerophosphorylcholine (GPC), choline sulfate, homoserine betaine</p> | <p>Found in every kingdom of life</p> | <ul style="list-style-type: none"> Protection against damage of membrane Stabilisation and activation of proteins and enzymes Regulation of ROS detoxification | <ul style="list-style-type: none"> <i>Betaine</i> – treatment of homocystinuria, neuroprotection, skin care ingredient <i>TMAO</i> – vaccine stabilisation |
| <p>Methyl-sulfonium compounds</p> <p>Dimethylsulfonium propionate (DMSP)</p> | <p>Found in phytoplankton and some halophytic vascular plants in large quantities</p> | <ul style="list-style-type: none"> osmoprotection, thermoprotection, and antioxidative activity detoxification of excess sulphur precursor of oceanic DMS in Lovelock's "Gaia hypothesis" | <ul style="list-style-type: none"> <i>DMSP</i> – one of Earth's most abundant organosulfur molecules and marine osmolyte, but no uses reported so far |
| <p>Y-conjugated compounds</p> <p>Urea, guanidine, arginine</p> | <p>Used only by relatively few types of animals (<i>e.g.</i> cartilaginous fish) and organs (mammalian kidney) (<i>urea</i>)</p> | <ul style="list-style-type: none"> Deleterious effect on protein structure and function | <ul style="list-style-type: none"> <i>Urea</i> – skin care ingredient <i>Guanidine</i> – treatment of muscle weakness caused by Eaton-Lambert syndrome <i>Arginine</i> – health supplement against chest pain, high blood pressure, erectile dysfunction and peripheral arterial disease, monoclonal antibody stabilization and formulation |



Although generally non-toxic and compatible with cytoplasmic proteins over wide concentration ranges, many kosmotropes may result harmful at high concentrations in the absence of a chaotrope. For example, at high concentrations TMAO inhibits some enzymes and enhances formation of non-functional protein aggregates *in vitro*.^{13,14}

Yancey *et al.*¹⁵ have shown that adding urea (a chaotrope) or glycine betaine (a kosmotrope) alone to the medium greatly reduced mammalian renal cells growth, while adding both types of osmolytes partly or fully restored normal growth. Also, when high protein diets dictate changes in the concentration of urea, the mammalian renal medulla appears to regulate one of its methylamine osmolytes, glycerophosphorylcholine (GPC), to maintain a constant GPC : urea concentration ratio.¹² In a totally different environment, the TMAO kosmotrope content is reported to be high in deep sea organisms only when there is an obvious chaotrope present, mostly notably urea. Furthermore, cartilaginous fish and coelacanths accumulate high concentrations of methylamine compounds (mainly TMAO, with lesser amounts of betaine and sarcosine) and certain free amino acids (mainly α -alanine and taurine) in concentrations averaging 0.2 M, which is about half the concentration of urea found in the same organism.¹⁶

The presence of osmolytes in certain combinations and proportions in nature is primarily related to the fact that, at specific molar ratios, kosmotropes, such as methylamines and amino acids, are able to counterbalance urea's deleterious effects on proteins and other macromolecules.¹² Yancey *et al.*¹⁶ were the first to conduct a series of *in vitro* experiments to demonstrate that kosmotropes (TMAO, betaine, sarcosine, β -alanine and taurine) are effective stabilizers of protein structure (*i.e.* bovine ribonuclease, rabbit and shark lactate dehydrogenases, and bovine glutamate dehydrogenase) and that these compounds largely or completely offset the denaturing effect of urea at a molar concentration ratio of 1 : 2. Ahnad and coworkers¹⁷ performed a series of studies to measure the thermal denaturation equilibrium of α -lactalbumin in the presence of urea and methylamines (TMAO and sarcosine) and found that at a molar ratio of 1 : 2 (methylamine : urea), the denaturing effect of urea on the protein was perfectly compensated by TMAO and sarcosine. In addition, Khan *et al.*¹⁸ observed that myo-inositol provides a perfect counteraction for three proteins (RNase-A, lysozyme, and α -lactalbumin) at a ratio of 1 : 2 myo-inositol to urea, whereas taurine regulates perfect counteraction in a protein-specific manner: 1.5 : 2 (RNase-A), 1.2 : 2 (lysozyme), and 1 : 2 (α -lactalbumin) taurine-urea ratio. This study has also shown that the counteraction of kosmotropes on urea is not limited to the 1 : 2 molar ratio but depends on the protein structure and its origin.

Osmolyte-protein interaction models and the over-stabilizing paradox

Over the past decade, many studies using both experimental and theoretical approaches have attempted to elucidate the

molecular mechanism by which kosmotropes, especially TMAO, stabilize proteins in the presence of the chaotrope urea. The "indirect mechanism" states that osmolytes affect the folding behavior of proteins by changing the structure of the medium through interaction with the surrounding water molecules and subsequently modulating (weakening or strengthening) the water H-bond network and its thermodynamic properties. In contrast, the "direct mechanism" approach proposes that the osmolyte interacts directly with the peptides or amino acid side chains of the protein backbone to stabilize the native folding of the protein.¹⁹ So far, studies that consider the effects of osmolyte combinations as cumulative cannot adequately account for their complex interaction with macromolecules. The culmination of this shortcoming is the over-stabilizing paradox: it has been shown that a kosmotrope (TMAO) stabilizes a protein more in the presence of a chaotrope (urea) than alone.^{20,21} A similar trend to synergy has been found with mixtures of other methylamines, *i.e.* betaine and sarcosine, with urea.^{22,23} On these grounds, it has recently been suggested that it is important to consider osmolyte-protein interactions beyond the simple notions of individual kosmotropes and chaotropes affecting the folding mechanism.

Rather, the effects of the combination of kosmotropes and chaotropes on the folding equilibrium of proteins need to be considered in a way that acknowledges not only cumulative but also synergistic effects between the system components.^{19,22} The above observations are consistent with the concept of a complex adaptive system,²⁴ which exhibits properties that extend beyond those of the sum of its parts.²⁵ These properties are often referred to as "emergent properties" and have been pivotal to a major field of research during the last two decades: the field of deep eutectic solvents.

Deep eutectic solvents: green solvents for a myriad of applications

Eutectic is a term used to define a homogeneous mixture of substances that melts at a single temperature which is lower than the melting point of any of its constituents. These mixtures have been known for more than a century.²⁶ However, when Abbott *et al.*²⁷ studied mixtures of quaternary ammonium salts (*e.g.* choline chloride, m.p. 302 °C) and urea (m.p. 133 °C) at a 1 : 2 molar ratio in 2004, they observed a substantial decrease in the solid-liquid phase transition temperature compared to the melting point of the individual components. The observed decrease was so substantially "deeper" than the one expected for an ideal mixture that these compositions remained liquid even at room temperature. These remarkable findings prompted Abbott's group to claim a novel class of such mixtures and to coin a suitable new term to describe them: Deep Eutectic Solvents (DESs). As initially suggested, such significant depression of the melting point is attributed to the interaction between the mixture components through intermolecular hydrogen bonds in a certain range of component ratios. In practice, a DES is commonly prepared



with 100% atom economy by mixing two or more, usually solid compounds, a hydrogen bond donor (HBD, *e.g.* amides, polyols, sugars, organic acids) and an acceptor (HBA, *e.g.* quaternary ammonium compounds choline chloride and betaine), in a certain molar ratio, which upon heating form a liquid at the operating temperature. While several studies on the nature of the DES component interactions have appeared,^{28–30} the most recent ones focus on how these interactions define the macroscopic properties of the resulting DES as a single, coherent entity,^{31,32} thanks to, and not despite of, the “microscopic heterogeneity” observed.³³ Along the same notion, recently Martins *et al.*³⁴ proposed a more pragmatic definition of a DES as “a mixture of two or more pure compounds for which the eutectic point temperature is below that of an ideal liquid mixture, presenting significant negative deviations from ideality”, significant enough so that “the mixture is liquid at the operating temperature for a certain composition range”. Based on these attributes, these neoteric solvents offer flexible physical properties such as low volatility, non-flammability, low toxicity, simple and solvent-free preparation from widely available natural raw materials and are therefore considered an excellent green alternative to conventional organic solvents and a promising tool for shaping numerous processes into being more efficient and sustainable.⁴ Additionally, the wide range of possible structural combinations encompassed by DES (estimated to be approximately 10⁶), their sustainability, their unique physiochemical characteristics, as well as the possibility of fine tuning their solvent properties (*e.g.* pH value, polarity, hydrophilicity/hydrophobicity, viscosity) for certain purposes, make them ideal green candidates for (bio)chemical, electrochemical, and material applications, as well as for the extraction of various compounds, both inorganic (*i.e.* metals and CO₂) and organic (*i.e.* plant metabolites, DNA, proteins).³⁵ These advantages have been reflected on the interest of the academic community in these neoteric solvents, with more than 7500 scientific papers published in the last 20 years.

Natural Deep Eutectic Solvents (NADESs), a subgroup of DESs, were introduced and defined by Choi *et al.*³⁶ as mixtures exclusively composed of two or three compounds naturally occurring in plants, often primary metabolites, such as non-toxic quaternary ammonium salts, amines, sugars, alcohols, polyols and organic acids. Precisely because of their natural origin, these solvents are expected to provide a natural, cytosol-like environment for various biomolecules, allowing them to show properties and functionality profiles that resemble the ones observed in their natural environment. Namely, in anhydrous form or with water as an additional component, these systems not only allow for excellent solubility of various biomolecules *per se*, but can also stabilise a wide range of commercially important molecules of natural origin (DNA, biologically active compounds, drugs, proteins, and enzymes) by providing a network of hydrogen bonds that favours stable structural conformations.³⁷

Given the above mentioned extractive and solubilising properties, together with their capability to stabilise diverse (bio) molecules, DESs stand out as solvents with almost endless

possibilities of applications. However, some key concerns on DESs application at the industrial level are their relatively high viscosity, possible corrosive activity, and in some cases, their high price. Thus, the choice of a DES for a particular use should be made carefully, having the production process and the final product in mind. Besides that, as DESs have a very low vapor pressure, isolation of the target compounds is considered to be one of the major challenges for their industrial application. Cautiously chosen DES can be safe for direct use in food, pharmaceutical, cosmetic and agrochemical products, and may thus widen the options during end product formulation.³⁸

Osmolytes and deep eutectic systems: what happens when the two worlds collide?

The molecules with strong H-bonding potential which have been so far considered as raw materials for the preparation of DES are mainly quaternary ammonium compounds, polyols, sugars, organic acids, and amino acids. The majority of these are in fact similar or identical to known natural osmolytes. For example, choline chloride is the most commonly used methylamine compound for the preparation of DESs. Structurally similar and naturally occurring methylamines, such as TMAO, betaine, GPC and sarcosine, are all particularly important osmolytes that stabilize macromolecules by counteracting the deleterious effects of perturbant osmolytes like urea.⁷ Out of these four natural methylamines, so far only betaine has been considered as a HBA for DES preparation, along with a few other osmolytes from other chemical classes, such as polyols (*e.g.* glycerol) and sugar polyols (*e.g.* glucose, sucrose, xylose, fructose, sorbitol, trehalose) and amino acids (*e.g.* proline, lysine, arginine and histidine) (Table 2).

As a group working in the field of DESs preparation and characterization, we were not only intrigued by the structural similarity between osmolytes and common DES components, but also by the fact that osmolytes are usually present in cells and tissues in certain combinations and often in rather strict proportions, and that these combinations and proportions are often strikingly similar or identical to those used for the preparation of synthetic DESs. Based on these observations, we decided to explore the pool of naturally occurring osmolytes and the patterns of their natural distribution in order to form novel, osmolyte-based DESs.

Eutectic systems based solely on naturally occurring osmolytes have been reported, mostly consisting of simple sugars, polyols, amino acids, and the methylamine betaine (Table 2). In general, most of these systems were prepared aiming at specific industrial applications, by combining known DES components. Here, we explored various new combinations of osmolytes, including TMAO, sarcosine, GPC, ectoine, proline, DMSP, guanidine HCl, arginine, taurine, sorbitol, and trehalose, and discovered a variety of distribution patterns that



Table 2 Reported and novel deep eutectic solvents consisting of naturally occurring osmolytes

| DES | Components | | | | Components molar ratio | Water content (wt%) | Ref. |
|----------------------------|--------------------------------|----------------------|----------------------|--------------|------------------------|---------------------|----------|
| | 1 | 2 | 3 | 4 | | | |
| B : U | Betaine | Urea | Water | — | 1 : 1 : 2 | 17 | 56 |
| B : Gly | Betaine | Glycerol | Water | — | 1 : 2 : 2 | 10 | 57 |
| B : Sol | Betaine | Sorbitol | Water | — | 1 : 1.2 : 6 | 25 | 58 |
| B : Treh | Betaine | Trehalose | Water | — | 4 : 1 | Unk. | 5 |
| B : Xyl | Betaine | Xylose | Water | — | 2 : 1 : 3 | 12 | 59 |
| B : Glc | Betaine | Glucose | — | — | 5 : 2 | — | 5 |
| B : Man | Betaine | Mannose | — | — | 5 : 2 | Unk. | 5 |
| B : Lys | Betaine | Lysine | Water | — | 1 : 1 : 27–180 | 65–93 | 39 |
| B : His | Betaine | Histidine | Water | — | 1 : 1 : 28–190 | 65–93 | 39 |
| B : Arg | Betaine | Arginine | Water | — | 1 : 1 : 30–200 | 65–93 | 39 |
| B : Arg^b | Betaine | Arginine | Water | — | 5 : 1 : 28 | 40 | <i>a</i> |
| B : G | Betaine | Guanidine HCl | Water | — | 1 : 2 : 3 | 17 | <i>a</i> |
| TMAO : U | Trimethylamine <i>N</i> -oxide | Urea | Water | — | 1 : 1 : 3.5 | 33 | <i>a</i> |
| TMAO : Gly | Trimethylamine <i>N</i> -oxide | Glycerol | Water | — | 1 : 1 : 2 | 17 | <i>a</i> |
| TMAO : Sol | Trimethylamine <i>N</i> -oxide | Sorbitol | Water | — | 1 : 1 : 2 | 12 | <i>a</i> |
| TMAO : Glc | Trimethylamine <i>N</i> -oxide | Glucose | Water | — | 5 : 2 : 12 | 23 | <i>a</i> |
| TMAO : Treh | Trimethylamine <i>N</i> -oxide | Trehalose | Water | — | 4 : 1 : 8 | 18 | <i>a</i> |
| TMAO : Xylol | Trimethylamine <i>N</i> -oxide | Xylitol | Water | — | 1 : 1 : 4 | 24 | <i>a</i> |
| TMAO : G | Trimethylamine <i>N</i> -oxide | Guanidine HCl | Water | — | 1 : 1 : 2 | 18 | <i>a</i> |
| TMAO : Arg ^b | Trimethylamine <i>N</i> -oxide | Arginine | Water | — | 15 : 4 : 56 | 37 | <i>a</i> |
| Sar : U | Sarcosine | Urea | Water | — | 2 : 5 : 9 | 25 | <i>a</i> |
| Sar : G | Sarcosine | Guanidine HCl | Water | — | 2 : 5 : 11 | 23 | <i>a</i> |
| Sar : Arg ^b | Sarcosine | Arginine | Water | — | 3 : 2 : 19 | 36 | <i>a</i> |
| Sar : Gly | Sarcosine | Glycerol | Water | — | 1 : 2 : 4 | 20 | <i>a</i> |
| GPC : U | Glycerophosphorylcholine | Urea | Water | — | 1 : 2 : 1.5 | 7 | <i>a</i> |
| GPC : G | Glycerophosphorylcholine | Guanidine HCl | Water | — | 1 : 2 : 1 | 4 | <i>a</i> |
| GPC : Arg ^b | Glycerophosphorylcholine | Arginine | Water | — | 11 : 4 : 80 | 30 | <i>a</i> |
| DMSP : U | Dimethylsulfonopropionate HCl | Urea | Water | — | 1 : 2 : 2 | 11 | <i>a</i> |
| DMSP : G | Dimethylsulfonopropionate HCl | Guanidine HCl | Water | — | 1 : 1 : 3 | 17 | <i>a</i> |
| DMSP : Arg | Dimethylsulfonopropionate HCl | Arginine | Water | — | 5 : 4 : 45 | 37 | <i>a</i> |
| DMSP : Gly | Dimethylsulfonopropionate HCl | Glycerol | Water | — | 1 : 2 : 5 | 20 | <i>a</i> |
| Glc : Fru | Glucose | Fructose | Water | — | 1 : 1 : 8 | 30 | 60 |
| Glc : Gly | Glucose | Glycerol | — | — | 1 : 4 | — | 59 |
| Glc : Treh | Glucose | Trehalose | Water | — | 1 : 2 : 13 | 21 | 59 |
| Glc : Gly:U | Glucose | Glycerol | Urea | — | 2 : 1 : 1 | — | 61 |
| Fru : Treh | Fructose | Trehalose | Water | — | 1 : 2 : 13 | 20 | 59 |
| Fru : Gly : U | Fructose | Glycerol | Urea | — | 2 : 1 : 1 | — | 61 |
| Gly : Sol | Glycerol | Sorbitol | — | — | 2 : 1 | — | <i>a</i> |
| Gly : Treh | Glycerol | Trehalose | — | — | 30 : 1 | — | 59 |
| Gly : U | Glycerol | Urea | — | — | 5 : 2 | Unk. | 62 |
| Gly : G | Glycerol | Guanidine HCl | — | — | 5 : 2 | — | 63 |
| Gly : Arg | Glycerol | Arginine | — | — | 4 : 1 | — | 64 |
| Pro : Glc | Proline | Glucose | — | — | 5 : 3 | Unk. | 5 |
| Pro : Gly | Proline | Glycerol | — | — | 1 : 2 | — | 65 |
| Pro : Sol | Proline | Sorbitol | — | — | 1 : 1 | Unk. | 5 |
| Pro : Suc | Proline | Sucrose | — | — | 2 : 1, 3 : 1 | Unk. | 5 |
| Pro : U | Proline | Urea | — | — | 2 : 1 | Unk. | 66 |
| Pro : G | Proline | Guanidine HCl | Water | — | 2 : 1 : 8 | 30 | <i>a</i> |
| Pro : Arg | Proline | Arginine | Water | — | 2 : 1 : 11 | 33 | <i>a</i> |
| Pro : GA | Proline | Glutamic acid | — | — | 2 : 1 | Unk. | 67 |
| Ser : Glc | Serine | Glucose | — | — | 5 : 4 | Unk. | 5 |
| Car : U | Carnitine | Urea | — | — | 2 : 3 | — | 68 |
| Ect : U | Ectoine | Urea | Water | — | 1 : 2 : 2 | 12 | <i>a</i> |
| Ect : G | Ectoine | Guanidine HCl | Water | — | 1 : 2 : 3 | 14 | <i>a</i> |
| Ect : Arg | Ectoine | Arginine | Water | — | 2 : 1 : 10 | 30 | <i>a</i> |
| Ect : Gly | Ectoine | Glycerol | Water | — | 1 : 2 : 4.5 | 20 | <i>a</i> |
| Sol : U | Sorbitol | Urea | Water | — | 2 : 3 : 1 | 3 | <i>a</i> |
| Sol : G | Sorbitol | Guanidine HCl | Water | — | 2 : 3 : 1 | 3 | <i>a</i> |
| Sol : Gly : U | Sorbitol | Glycerol | Urea | — | 1 : 1 : 2 | — | 61 |
| Sol : Gly : G | Sorbitol | Glycerol | Guanidine HCl | — | 1 : 1 : 2 | — | <i>a</i> |
| Suc : U | Sucrose | Urea | Water | — | 4 : 1 : 8 | 9 | <i>a</i> |
| Suc : G | Sucrose | Guanidine HCl | Water | — | 2 : 1 : 12 | 23 | <i>a</i> |
| Suc : Arg | Sucrose | Arginine | Water | — | 5 : 2 : 16 | 12 | <i>a</i> |
| Treh : Glc : Gly | Trehalose | Glucose | Glycerol | — | 1 : 1 : 5 | — | <i>a</i> |
| B : Sol : U | Betaine | Sorbitol | Urea | Water | 1 : 1.2 : 1 : 7 | 25 | 58 |
| B : Sol : G | Betaine | Sorbitol | Guanidine HCl | Water | 1 : 1.2 : 1 : 7 | 23 | <i>a</i> |
| B : Treh : Raf | Betaine | Trehalose | Raffinose | Water | 9 : 1 : 1 : 35 | 25 | 5 |
| B : Suc : Pro | Betaine | Sucrose | Proline | Water | 5 : 2 : 2 : 21 | 20 | 59 |



Table 2 (Contd.)

| DES | Components | | | | Components molar ratio | Water content (wt%) | Ref. |
|-------------------------|-------------------------------|----------------------|----------------------|--------------|------------------------|---------------------|--------------|
| | 1 | 2 | 3 | 4 | | | |
| B : Gly : Suc | Betaine | Glycerol | Sucrose | Water | 2 : 3 : 1 : 5 | 10 | 59 |
| B : Gly : Treh | Betaine | Glycerol | Trehalose | Water | 2 : 3 : 1 : 5 | 13 | 59 |
| B : Suc : Pro | Betaine | Sucrose | Proline | Water | 5 : 2 : 2 : 21 | 20 | 59 |
| B : Tau : Gly | Betaine | Taurine | Glycerol | — | 1 : 1 : 3 | — | ^a |
| B : Ect : Gly | Betaine | Ectoine | Glycerol | — | 1 : 2 : 3 | — | ^a |
| B : Ect : Sor | Betaine | Ectoine | Sorbitol | Water | 1 : 2 : 3 : 5 | 9 | ^a |
| TMAO : U : Gly | Trimethylamine N-oxide | Urea | Glycerol | Water | 1 : 2 : 2 : 2 | 8 | ^a |
| TMAO : G : Gly | Trimethylamine N-oxide | Guanidine HCl | Glycerol | Water | 1 : 1 : 2 : 2 | 9 | ^a |
| TMAO : U : Sol | Trimethylamine N-oxide | Urea | Sorbitol | Water | 1 : 2 : 2 : 2 | 6 | ^a |
| TMAO : G : Sol | Trimethylamine N-oxide | Guanidine HCl | Sorbitol | Water | 1 : 1 : 2 : 2 | 6 | ^a |
| TMAO : B : U | Trimethylamine N-oxide | Betaine | Urea | Water | 1 : 1 : 2 : 2 | 10 | ^a |
| TMAO : B : G | Trimethylamine N-oxide | Betaine | Guanidine HCl | Water | 1 : 1 : 2 : 6 | 21 | ^a |
| Suc : Glc : Fru | Sucrose | Glucose | Fructose | Water | 1 : 1 : 1 : 11 | 22 | 59 |
| Fru : Glc : Treh | Fructose | Glucose | Trehalose | Water | 1 : 1 : 1 : 11 | 21 | 59 |
| Fru : Glc : G | Fructose | Glucose | Guanidine HCl | Water | 1 : 1 : 2 : 1 | 4 | ^a |
| Fru : Gly : G | Fructose | Glycerol | Guanidine HCl | Water | 2 : 1 : 1 : 1 | 4 | ^a |
| Glc : Gly : G | Glucose | Glycerol | Guanidine HCl | Water | 2 : 1 : 1 : 6 | 16 | ^a |
| Glc : Treh : U | Glucose | Trehalose | Urea | Water | 1 : 1 : 2 : 3 | 7 | ^a |
| Glc : Treh : G | Glucose | Trehalose | Guanidine HCl | Water | 1 : 1 : 1 : 3 | 8 | ^a |
| Gly : Suc : Sor | Glycerol | Sucrose | Sorbitol | Water | 2 : 1 : 2 : 10 | 16 | 59 |
| Gly : Treh : Sor | Glycerol | Trehalose | Sorbitol | Water | 2 : 1 : 2 : 10 | 16 | 59 |
| Gly : Glc : Sor | Glycerol | Glucose | Sorbitol | Water | 1 : 1 : 1 : 3 | 12 | 59 |
| Treh : Glc : Sor | Trehalose | Glucose | Sorbitol | Water | 1 : 2 : 1 : 13 | 17 | 59 |
| Treh : Glc : Sor | Trehalose | Glucose | Sorbitol | Water | 1 : 1 : 4 : 13 | 20 | ^a |
| Treh : Pro : Gly | Trehalose | Proline | Glycerol | Water | 1 : 1 : 4 : 8 | 15 | ^a |
| Treh : Ect : Gly | Trehalose | Ectoine | Glycerol | Water | 1 : 1 : 4 : 6 | 11 | ^a |
| Pro : Fru : Gly | Proline | Fructose | Glycerol | Water | 1 : 1 : 1 : 5 | 20 | 69 |
| Pro : Glc : Gly | Proline | Glucose | Glycerol | Water | 5 : 3 : 3 : 20 | 21 | 59 |
| Pro : Gly : Sor | Proline | Glycerol | Sorbitol | Water | 1 : 1 : 1 : 3 | 12 | ^a |

^a DES were prepared as follows: the appropriate amount of the components was placed in a 25 mL round bottom flask and the mixture was heated at 60 °C with stirring for 2 hours until a clear homogeneous liquid was formed. Upon cooling to room temperature, the mixture was left on a bench for a week in order to observe possible solidification or precipitation. Before use, DES forming compounds were dried under vacuum.

^b Liquids stable at 40 °C.

form liquids stable at room temperature (Table 2). Several molar ratios of known and unknown osmolyte combinations were explored experimentally, building on the distribution patterns observed in natural systems.

All known natural methylamine osmolytes (TMAO, betaine, sarcosine and GPC), as well as DMSP, the only reported natural methylsulfonium kosmotrope, formed liquids with all known perturbant osmolytes (urea, guanidine HCl and arginine), indicating a horizontal kosmotrope: chaotrope pattern of two-component Deep Eutectic Solvent formation (Fig. 2). Considering the melting points of the individual components (sarcosine 210 °C, GPC 143 °C, TMAO 100 °C, DMSP 123 °C, urea 133 °C, guanidine HCl 182 °C, arginine 244 °C), obtaining liquids at room temperature or even at 40 °C arguably satisfies the strict criterion for the “deep” designation. This trend, also observed for glycerol with chaotropes urea and guanidine, was extended to another chaotrope, arginine, and further corroborated for sugar polyols sucrose (m.p. 180 °C) and sorbitol (m.p. 95 °C). The potential of the amino acid osmolyte proline (m.p. 252 °C) to form DES was also confirmed for all known chaotropes (urea, guanidine and arginine), together with another widely applied osmolyte and amino acid cyclic derivative, ectoine (m.p. 280 °C), which also provided stable liquid mixtures with all studied chaotropes.

As a representative of the methylamine osmolytes class, TMAO also formed stable liquids when combined with glycerol and major sugar polyols, following the example of betaine. The results previously published for betaine³⁹ prompted us to further explore high water content in selected two-component DES. In particular, betaine reportedly formed liquid mixtures with three amino acids at a molar ratio of 1 : 1, but the water content of these mixtures was mentioned to be between 65 and 93 wt%, which is substantially higher than the upper limit usually considered for water-containing DES and will be further discussed in the next section. Given the abundance and application potential of betaine, we decided to probe additional molar ratios for its combination with arginine (Arg), which is also a somewhat controversial^{40,41} but widely employed osmolyte. Indeed, in our hands betaine did form a stable liquid with arginine at a 4 : 1 molar ratio, while it also required a higher water content (40 wt%) compared to betaine : urea and betaine : guanidine DESs, which both required 17 wt%. The same trend of very particular kosmotrope: arginine ratios and relatively higher water content than usual chaotropes was observed for most of the novel Arg-based DESs. Interestingly, the arginine combinations with the other natural methylamine kosmotropes TMAO, sarcosine and GPC only remained liquid at temperatures of at least 40 °C.






Fig. 2 (a) Formation of the TMAO : urea 1 : 1 deep eutectic solvent at 33 wt% water content, (b) FT-IR and (c) ^1H NMR spectra of the TMAO : urea 1 : 1 DES, compared to the respective spectra of the individual components. Spectra were recorded using TMAO : U as a pure viscous mixture. NMR spectra of TMAO and urea were recorded using their respective solutions in D_2O , with external C_6D_6 and TMS sealed in capillary as standard at 25 °C in 5 mm NMR tubes. IR spectra were recorded on an FTIR spectrometer equipped with an attenuated total reflection module with diamond ATR crystal.

A hint to additional possibilities worth probing for the behaviour of two-component mixtures was revealed when we investigated the potential of the commercially promising TMAO–urea combinations. While these two-component systems were unstable at room temperature at the typical methylamine-to-urea molar ratio of 1 : 2 found in nature, the addition of a third osmolyte component, glycerol or sorbitol, to a final TMAO : urea : polyol molar ratio of 1 : 2 : 2, smoothly provided a stable eutectic mixture stable at the same conditions. These two polyols were also critical as a third DES component when added to the binary mixtures betaine–ectoine (molar ratio of 1 : 2) and trehalose–glucose (molar ratio of 1 : 1), once more providing three-component liquid mixtures stable at room temperature. A similar need for a third component to obtain stable liquids was observed for the binary mixture trehalose–glucose, this time with urea, affording a liquid stable at room temperature for the three-component combination Treh : Glu : U (Table 2). Following this trend, several new three-component eutectic mixtures were obtained, ranging from mixed kosmotropes : chaotrope systems, reminiscent of natural osmolyte distribution patterns, to industrially promising all-sugar and ectoine-based systems.

Some of the prepared DESs were anhydrous (*i.e.* Gly : Sol, Gly : G, B : Tau : Gly, B : Ect : Gly, Treh : Glc : U and Sor : Gly : G), while most of them required approximately 5–35 wt% of water to remain in a stable liquid form at room temperature for an extended period of time (Table 2). This was to be expected since water has been shown to play a key role in the formation of DESs by modifying the physicochemical properties of the corres-

ponding DESs supramolecular network. For instance, it has been observed that the small amount of water present in hydrated DESs can strengthen the hydrogen bond network with water monomers confined into the DES voids.^{42–45} Hammond *et al.*⁴³ demonstrated that water acts as a second small HBD when aqueous mixtures of $\text{ChCl}:\text{malic acid}$ in a 1 : 1 molar ratio contain 1–2 mol of water per mole of DES. This was also confirmed by López-Salas *et al.*⁴⁵ who showed that in the “water-in-DES” system of a ternary DES composed of resorcinol, urea and choline chloride (RUCHCl_nW , where n represents mol of water per mole of ternary DES), the tetrahedral structure of water was distorted as a consequence of its incorporation, as an additional HBD or HBA, into the hydrogen bond complexes formed among the original DES components. The same group showed that “water-in-DES” regime occurs in DES dilutions with nonaqueous hydrogen-bond-forming organic solvents (*e.g.* benzyl alcohol).⁴⁴

The point of transition from water-in-DES regime to DES-in-water regime (described as a simple aqueous solution of the individual components) depends on the DES composition, but it is generally considered that up to about 40–50 wt% of water (in some cases even up to 57%,⁴⁶ the strong interactions between the components of the DESs slowly weaken with the supramolecular structure preserved,⁴⁷ while at higher dilutions, the network can be disrupted, leading to a mixture that exhibits behavior closer to that of the individual components in an aqueous solution.⁴⁸

As mentioned, none of the newly prepared mixtures required more than 40 wt% of water, which is consistent with the strict definition of these mixtures as DESs.



Overall, we here present several major advances: TMAO, sarcosine, GPC, ectoine, DMSP, arginine and guanidine HCl, the most widely encountered osmolytes and representatives of both kosmotrope and chaotrope families, are for the first time experimentally combined to form new DESs. The potential of these seven osmolytes to form eutectic systems is further established by the preparation of 54 novel two- or three-component DESs with other osmolytes. And for the first time, these new eutectic systems are directly inspired by the distribution patterns of kosmotropes and chaotropes observed in nature. In terms of DES sustainability, choline chloride, the most widely used methylamine for DESs preparation so far, is industrially manufactured from fossil-based ethylene oxide.⁴⁹ The systems herein reported are based only on natural methylamines, and thus avoid the use of choline chloride, while exhibiting novel properties and providing new affordances.

Mimicking nature: new multicomponent DESs based on osmolyte distribution patterns in biological contexts

Single osmolytes rarely accumulate alone in a given biological context, but rather within patterns comprising one chaotrope and one or more additional kosmotropes. For example, the serum of winter-acclimatized fish contains urea, TMAO and glycerol,⁵⁰ while terrestrially hibernating amphibians accumulate urea, glucose and glycerol,⁵¹ an ubiquitous grass species use proline, betaine and sucrose to improve salt tolerance,⁵² and sharks combine urea, TMAO and betaine in their body fluids.⁵³ We have herein shown that, among others, these exact three-component combinations (TMAO–urea–glycerol, glucose–urea–glycerol, proline–betaine–sucrose and TMAO–betaine–urea) can form DESs (Table 2). We have further shown that in the case of the TMAO–urea–glycerol DES, the addition of the third osmolyte to the initial binary mixture was crucial to obtain a liquid at room temperature. Therefore, it would be reasonable to expect the formation of such ternary or even more complex osmolyte-based DESs inspired by additional and even more complex natural distribution patterns.

Encouraged by these findings and by the fact that no DES of such complexity has been reported, we decided to break the three-component barrier and explore the patterns found in the kidney of cartilaginous fish (shark) and mammals (rabbit – inner renal medulla), as well as in the muscles of sharks and skates (Fig. 3). These organs/organisms accumulate urea to high concentrations as part of their osmoregulatory strategy and a cocktail of kosmotropes to counteract the perturbing effects of the urea chaotrope on protein structure.^{7,54}

In particular, during antidiuresis (under water-deficient conditions), mammals accumulate high concentrations of compatible osmolytes in renal medullary cells, among them sorbitol, GPC, inositol, and betaine, whereas in cartilaginous fish, TMAO dominates a cocktail of urea, betaine, inositol and

α -aminoacids.⁵⁴ As for marine elasmobranchs, to be approximately isosmotic with seawater, these organisms accumulate high concentrations of urea in their body fluids and tissues, together with TMAO, betaine and taurine to “neutralise” its harmful effects.⁵⁵ Based on these observations, we calculated the molar ratios of each osmolyte cocktail directly from their concentrations in the respective tissues and subjected the mixtures to the standard DES-forming conditions (Fig. 3).

All 10 osmolyte combinations provided liquids that were stable at room temperature with water contents less than 40 wt% (Fig. 3) and most of them were rapidly formed within minutes, compared to almost 1 hour required for usual DES formation.

This is the first time that a specific molar combination of multiple osmolytes, as encountered in a specific native environment, is experimentally proven to form a deep eutectic system of unprecedented complexity (osmoDES). It is worth noting that the tested osmolyte combinations are distinct, not only in terms of which organism they belong to, but also in terms of the topology of the studied tissue (different sections of the rabbit kidney medulla, Fig. 3) as well as the specific relative localization of a given organism (skates in different sea depths, Fig. 3). Therefore, the differences observed in the osmolyte distribution patterns may reflect the response to the respective conditions, functions and stresses related to each specific microenvironment. Similar multi-osmolyte cocktails are also found in other plant, animal and human organs and tissues, such as the tissues of other marine animals (*e.g.* fish, molluscs, crustaceans, shrimps, octopods, snails and worms),^{7,70,71} plants exposed to salt stress,⁷² hibernating organisms,⁵¹ and in the human brain.⁷³ Based on this novel viewpoint, tracking osmolyte patterns in specific biological contexts could help successfully create novel, bioinspired two-, three-, and multicomponent DESs. Perhaps more significantly, such systems could be very effective in mimicking the natural microenvironment of proteins and other biomacromolecules operating within these biological contexts, by deciphering, understanding and hopefully replicating and tuning the molecular interactions between these new media and the targeted macromolecular structures.

Methylamines stabilize proteins in the presence of urea: could this phenomenon be explained by the formation of a eutectic system?

As discussed above, the molecular mechanism by which methylamines, as kosmotropes, stabilize proteins in the presence of the chaotrope urea remains controversial. At a molar ratio of 1 : 2, the thermodynamic effects of combined methylamines and urea on protein stability and function were believed to be algebraically additive.¹⁷ However, in numerous cases methylamines in the presence of urea are more potent in





Fig. 3 Bioinspired multicomponent osmoDES prepared by following osmolytes distribution patterns in the cartilaginous fish (shark *Squalus acanthias*)⁷ and mammal (rabbit *Oryctolagus cuniculus*)⁷ kidneys, and muscles of cartilaginous fish (sharks *Chiloscyllium punctatum* and *Dasyatis sabina*, and skates *Raja hollandi*).^{73,157} The molar ratios of DES forming components were calculated from their concentrations in the respective tissues and the mixtures were subjected to the standard DES-forming conditions.

stabilizing a protein than alone,²⁰ resulting in the over-stabilizing paradox mentioned earlier.

When Abbott reported the choline chloride:urea DES (ChCl:U, molar ratio 1:2),²⁷ and when later the same DES not only did not denature⁷⁴ but instead showed a strong stabilizing effect on enzymes,⁷⁵ the mechanism by which methylamines stabilize proteins in the presence of the denaturant urea was not properly discussed in the context of synergies and interferences between these two in the form of a eutectic mixture. The only study that addressed this issue showed that, as the con-

centration of betaine-urea mixtures increased up to 75 wt% (corresponding to deep eutectic conditions), the ability of the lysozyme to refold upon cooling increased accordingly, while an increase in urea concentration alone could cause denaturation of the lysozyme.⁵⁶ Zeng *et al.*⁵⁶ were the first to show that betaine as a kosmotrope and urea as a chaotrope form a DES at molar ratios in the range of 1:1 to 1:3.⁵⁶

In addition to this observation, we have here shown that the four most prominent methylamines distributed across all kingdoms of life (betaine, TMAO, sarcosine and GPC) form



eutectic systems with urea, guanidine HCl and arginine (Table 2). Furthermore, we have also shown that multicomponent DESs are consistently formed when combining prominent natural kosmotropes (betaine, TMAO, GPC, DMSP, taurine, ectoine, sarcosine, inositol and α -amino acids) with natural chaotropes urea and guanidine in molar ratios found in mammal and fish kidney and muscles (Fig. 3).

Based on these findings, we wondered whether it was possible for osmolytes to coexist in the vicinity of proteins or other macromolecules in the form of a eutectic mixture, rather than as components dissolved in water, and whether this might provide additional information on the molecular mechanism of the stabilizing effect of natural methylamine kosmotropes on proteins in the presence of a natural chaotrope.

The mechanism underlying the counteraction of synthetic choline chloride to urea under eutectic conditions is well studied. Two independent experimental and molecular dynamics studies have shown that small helix-rich protein⁷⁶ and lipase⁷⁵ exhibit excellent conformational stability in ChCl:U and that high concentrations of both choline chloride and urea (at a 1:2 molar ratio) are critical for this stabilizing effect. At these conditions, which correspond to a deep eutectic state, the authors suggest that choline chloride acts as an efficient protein stabilizer by attracting urea and excluding it from the protein surface.⁷⁶ This assumption was further corroborated by Monhemi *et al.*⁷⁵ who showed that, at deep eutectic conditions, urea molecules have a low diffusion coefficient due to hydrogen bonding with choline and chloride ions and cannot reach protein domains. Thus, the above research on synthetic ChCl with urea may serve as a basis for the hypothesis that, within cells, strong hydrogen bonds between natural kosmotropes and chaotropes under deep eutectic state are responsible for the stabilization of proteins and other biomolecules *in vivo*. Similar to synthetic ChCl, the newly discovered fact that all major natural methylamines form deep eutectic mixtures with urea and guanidine HCl can give us guidelines on how to decipher the molecular mechanism behind the protein-stabilizing effects of all-natural methylamine-based kosmotropes, specifically. At the same time, the expansive interpretation of these observations raises the question whether all osmolyte-induced changes in the conformational equilibrium of macromolecules *in vivo* are related to the formation of osmolyte-based DESs, whose emergent properties within the wider system of the cytoplasm extend beyond those of separate osmolytes in aqueous solution.

Do osmolytes form DESs in cells exposed to stress?

The cytoplasm has been traditionally viewed as a well-mixed and spatially homogeneous mixture of monomers and small complexes. However, modern molecular biology techniques are giving us unprecedented access to cell structure and dynamics at micro- and meso-length scales.⁷⁷ Recent studies

reveal that, at these scales, a number of interesting liquid phase transitions occur in two or three dimensions resulting in separate fluid phases within cells.⁷⁸ Intermolecular hydrogen bonding between osmolytes is highly likely to occur in cells at the nanoscale, as part of dynamic (to overcome various forms of stress)⁷⁹ as well as apparently static systems (*e.g.* in seeds for long-term survival).³³ High concentrations of solutes, such as sugars, alcohols, organic acids and amino acids, has provoked scientists to wonder if these solutes contribute to the nano/mesoscale organization in certain biological systems by forming DESs.^{36,80} If so, DESs formation *in vivo* should not be seen as a simple, randomly dispersed liquid phase, but as a layer or a liquid cluster in different cellular regions or subcellular assemblies such as plastids and vesicles.⁸⁰

The presence of DESs *in vivo* was first hypothesized for plant cells by Choi *et al.*,³⁶ based on the observation that plant secretions, such as sap and nectar, accumulate primary metabolites which are known DES components (choline, betaine, proline, organic acids such as malic, succinic and citric) in molar ratios typical of DES formation. The authors went on to propose that plant cells contain a third type of medium, which plays a vital role in solubilizing, storing, and transporting poorly water-soluble metabolites, adjusting the water content of plants, and protecting cells when in harsh conditions. Durand *et al.*⁸⁰ recently raised the hypothesis that DES may also be involved in cell function regulation through interactions with membranes, by affecting their properties and therefore regulating the transport and diffusion of molecules or the circulation of fluids.

Going one step further, here we experimentally show that several natural distributions of osmolytes in several species across kingdoms and environments form new multicomponent DESs of previously unimagined complexity. We thus believe that the question is now ripe: are osmolyte-based eutectic systems formed in the cytoplasm of all species to support the stability and function of biomolecules under external stress? Furthermore, could the concept of DESs formation in the cell at the mesoscale be a key to understanding the *in vivo* stabilization of biomacromolecules in general?

To support this hypothesis, we identified measurable indicators of the proposed existence of DESs *in vivo*. It has been shown that enzymes such as laccases,⁵ dehydrogenases,⁸¹ lipases and other hydrolytic enzymes^{60,82} show little or no activity in pure synthetic DESs or in DESs at low water content. However, by adding water to DESs, the activity of the enzymes increases and reaches its maximum in highly diluted mixtures (80–90 wt%). On the other hand, low water content in DES has been shown to positively affect the stability of the same enzymes.³⁶ We may postulate that mixtures of osmolytes form DESs at nano/microscale *in vivo* and in this way preserve enzymes in catalytically inactive form during cryoprotection, drought, resistance and germination. Once the stressful conditions are overcome and water enters the cell, the eutectic systems are diluted and the enzymes are activated.



Is the microenvironment a low-energy, context-aware regulator of biomacromolecule function?

Yancey *et al.*¹³ postulate that transitions between torpid and active states may not require metabolic regulation at the level of enzyme degradation or synthesis, but may rather depend on readily reversible transitions between inactive and active enzyme states in response to alterations in its microenvironment. Furthermore, this hypothesis could be relevant not only to organisms undergoing stress, but also as a general regulatory mechanism for the storage and on demand activation of enzymes. In addition, versatile enzymes (*e.g.* laccases,⁸³ lipases,⁸⁴ proteases⁸⁵) show excellent thermal stability in synthetic DESs composed of osmolytes such as betaine, polyols and sugar alcohols. Delorme *et al.*⁸³ showed that laccase is much less thermostabilized in the presence of betaine and xylitol alone than in a DES formed by those two osmolytes. This leads us to assume that formation of eutectic systems *in vivo* could contribute to the stabilization of proteins in organisms that must cope with high temperatures. Going beyond proteins, the nucleic acid-templated synthesis (NATS) of a dipeptide was studied as a simple model of non-enzymatic translation in DESs.⁸⁶ The authors showed that, compared with aqueous buffer media, various glycholine-based DESs have a positive effect on the stability of DNA-conjugated activated esters which are necessary for non-enzymatic peptide formation. In this DES, peptide synthesis was prevented due to a significant decrease in the reactivity of amines. When the DNA-conjugated activated esters were transferred back to aqueous buffers, peptide synthesis was once again observed. This suggests that the hypothesis of DES-based regulation could be relevant to the storage and “on-demand” activation of additional biomacromolecules beyond proteins.

When thinking of osmolytes forming DESs in the cytoplasm, it is worthwhile to consider the way cells react when osmolytes reach potentially toxic concentrations of up to 400 mM.⁸⁷ Jagannathan *et al.*⁸⁷ proposed that mixtures of different osmolytes in specific molar ratios may help to reduce toxicity of accumulated osmolytes. Depletion in osmolytes toxicity could be a consequence of them forming DESs. Namely, it has been shown on several occasions that phosphonium,⁸⁸ ammonium,⁸⁹ choline^{90,91} and betaine-based⁹¹ DESs show lower toxicity toward immortalized animal cell lines than the aqueous solutions of their individual components.

The accumulation of osmolytes in cells is a dynamic and reversible process. Thus, to plausibly consider that osmolytes in a cytosol form eutectic systems and thus stabilize biomacromolecules, the formation of these systems must also be reversible. As previously noted, DESs have the potential to bind up to 40–50% water and their components still interact with each other,⁴⁸ whereas further dilution behaves closer to a solution of its individual components. By performing hygroscopicity measurements of the glucose:choline chloride:water DES (molar ratio 2:5:5) Dai *et al.*⁶ proved that this process is indeed

reversible, indicating that, within cells, DES may easily arise and disappear at micro-scale level. More importantly, this inherent ability may enable fine-tuning of the cytoplasm water level as a means to control physicochemical properties (*e.g.* pH, osmolality, polarity and viscosity) which are crucial for maintaining cell homeostasis under various environmental conditions.

The way cells regulate the stabilization of macromolecules at low water content conditions (freezing and drought) could also be explained by the reversibility of DESs formation and dilution. In psychrophilic bacteria,⁹² diapausing insects^{93,94} (*e.g.* grasshoppers and spiders), amphibians⁵¹ (*e.g.* frogs) and reptiles^{95,96} (*e.g.* snakes and turtles) the accumulation of compatible solutes such as betaine, glucose, sucrose, trehalose, sorbitol, mannitol, glycerol, glycine, taurine, and urea results in the lowering of the cytoplasmic freezing point, thereby providing protection against frost. Analogous to the mechanism proposed for plants,⁶ freezing may result in the formation of a separate liquid phase (DES) that keeps both water and (macro) molecules in a stable liquid form, as discussed earlier. Indeed, the interactions between the DES components and water are strong enough to maintain water liquid in the form of a glassy DES solution even at extremely low temperatures.⁹⁷ Dilution with water, once an organism is back to ambient temperature, restores the cellular structures resulting in a living cell or organism. An analogous course of action could be applicable for organisms in dormant state. Namely, in certain anhydrobiotic organisms, ranging from bacteria and yeast (in cyst or spore stages) to nematodes and to certain arthropods that have entered a dormant state, water content may drop well below 10% and such anhydrobiotic states exhibit no detectable metabolism.⁹⁸ These organisms can remain viable in this form for years, even millennia in the case of some prokaryotes.⁹⁹ The initially obscure underlying mechanisms have been only partially clarified in the recent decades. While the effect of trehalose in anhydrobiotic states was initially touted as conclusive, recent studies indicate a more complex phenomenon,⁹⁹ which might benefit by acknowledging synergistic effects between trehalose and other osmolytes accumulated in the cytoplasm, such as glycerol, proline, ectoine, betaine and TMAO,¹⁰⁰ possibly in the form of eutectic systems. The loss of water from the cytosol at anhydrobiotic conditions could result in the formation of a DES that keeps biomolecules in an inactive form until needed. Knowing that DES easily absorb and desorb water from the environment, the existence of DESs as a third and possibly transient medium inside a cell appears as a simple and non-demanding way to tolerate water-related changes in the environment.⁶

Case study: (thermo)stabilisation of a model protein in novel osmolyte-based DESs

To showcase some of the novel osmolyte-based DESs as a liquid medium for storage of biomolecules, particularly pro-






Fig. 4 Residual lysozyme activity after incubation in osmolyte-based DES (containing 20 wt% of water) and sodium phosphate buffer solution at 25 °C (A) and 45 °C (B). Lysozyme activity was determined according to the method of Shugar *et al.*¹⁵⁸ Lysozyme solutions at a concentration of 0.1 mg ml⁻¹ were prepared in different DESs and in 10 mM sodium phosphate buffer solution (pH 7) and incubated for 7 days at 25 and 45 °C. To measure residual enzyme activity, to 525 μ l of 10 mM sodium phosphate buffer solution (pH 7) in a plastic disposable cuvette 30 μ l of *Micrococcus lysodeikticus* bacteria suspension in sterile PBS buffer (7 mg ml⁻¹) and 30 μ l of the lysozyme solution were added. Immediately after mixing, the cuvette was placed in a UV/VIS spectrophotometer and the absorbance was measured at a wavelength of 450 nm over a period of linear turbidity decline. The relative activity (%) was calculated from the initial reaction rate obtained by the enzyme after incubation, compared to the one obtained without previous exposure.

teins, we monitored the stability of a model protein, lysozyme, in these solvents (Fig. 5). The enzyme was incubated at 25 and 45 °C for 7 days in several DESs comprising natural methyl-

amines (betaine, TMAO and sarcosine), methyl sulfone DMSP and cyclic amino acid ectoine as HBA, and glycerol as HBD, as well as in two bioinspired osmoDESs (all containing 20 wt% of




Fig. 5 Mimicking nature: design, preparation and potential applications of bioinspired osmolyte-based DESs.



water). It has been previously shown that compared to buffered solutions, lysozyme activity was greatly reduced in ChCl:Gly having low water content.^{101,102} However, the enzyme activity was completely recovered upon dilution. Herein, we have also observed the same trend: in all tested osmolyte-based DESs the loss in enzyme activities compared to the reference buffer was observed (>90%, data not shown), while upon dilution in sodium phosphate buffer the enzyme activity was restored completely. Thus, for the purpose of measuring lysozyme stability in osmolyte-based DESs, the enzyme was incubated in DESs and residual activity upon its dilution in the buffer was measured. As can be seen from Fig. 4, DESs comprising sarcosine or DMSP with glycerol, at both incubation temperatures tested, enabled higher residual lysozyme activities after 7 days of incubation (in the range from 95 to 103%) in comparison to both controls, ChCl:Gly and the reference buffer¹⁰¹ with residual activities <82%. In contrast, TMAO as HBA, when paired with glycerol, did not show stabilizing effect on the tested enzyme, while the presence of ectoine was beneficial for lysozyme stabilization at 25 °C.

Two-component DESs of betaine and TMAO with urea as HBD showed similar destabilizing effects. However, the bio-inspired multicomponent osmo-DESs containing the same

osmolytes, betaine and urea, exhibited excellent protein stabilisation ability, especially at 25 °C (residual activity of 96.2%), indicating that the presence of additional osmolytes in the cocktail (sorbitol, GPC, and taurine) in the ratios replicated from a natural context drastically enhanced DES ability to stabilise lysozyme. The same was observed for TMAO:U and its multicomponent osmoDES cocktail counterpart, which besides TMAO and urea also contained betaine and taurine.

These results suggest that osmolyte-based DESs in all their variability have a potential to stabilize a template protein stored at ambient, as well as at elevated temperatures, in a significantly better and longer-lasting manner than the currently known DES stabilizing medium (ChCl:Gly) and the standard buffer for lysozyme storage. Replicating osmolyte patterns inspired by specific biological contexts can, therefore, help design novel multicomponent DESs with exquisite stabilising abilities, providing new options for the preservation of proteins, ideally without the requirement of additional excipients.

Conclusions and future perspectives

It has been argued that adaptation accounts for most incremental evolutionary changes in nature, while exaptation has




Fig. 6 OsmoDES Cynefin domains and dynamics: the exaptive connection between osmolytes and DES (point γ) triggers additional actions along the red trajectory, to discover new potential parameters which have an impact. It can also reveal new questions along the blue trajectory, which can lead to new hypotheses that are tested (point α) and may become constructors that experts can reliably and consistently apply (point β), until new practices are established at scale (point δ). Graphic adapted from cynefin.io/wiki.



been linked to the most disruptive and arguably the most fascinating ones, such as the wing or the lung.^{103,104} While exaptation is most impressive when links between very dissimilar contexts are established as a result, it can be even more pervasive when the previously unestablished connection seems surprisingly obvious in retrospect. Most importantly, when the newly formed link can act as an enabling constraint,¹⁰⁵ then a new way of thinking is revealed, based on which additional interdependencies can be envisaged and new opportunities of action can emerge.

Here we have established such a connection between the previously discrete fields of osmolytes and DESs. We first showed that osmolytes prominent across all kingdoms, TMAO, sarcosine, ectoine, GPC, DMSP, arginine and guanidine HCl, may be used to prepare new DESs. Going one step further, we followed the inspiration from several natural distributions of osmolytes in several species and environments to reveal new multicomponent DESs of unprecedented complexity and briefly showcased their tunable ability to stabilize a template protein.

These new biorelevant DESs are context-aware and context-specific snapshots of a micronarrative which reflects the nuances of the cellular microenvironment through space, time and function. This feature can provide additional and potentially orthogonal angles when looking at biological systems, whether one is trying to understand and describe them,^{9,106–114} monitor or regulate them *in vitro* and *in vivo*,^{115–120} replicate and mimic them *ex vivo* or “on-a-chip”,^{121–125} or model them *in silico*.^{126–129}

On top of that, these novel DESs are also bioinspired engineered media which may serve as new tools and open up new opportunities for action in diagnosis, monitoring, treatment and manufacture across several research fields and market sectors,^{9,130,131–138,139–143} starting from applications and current use cases where single osmolytes or aqueous mixtures have shown promise¹⁴⁴ (Fig. 5). Most importantly, this new concept can serve as a constructor¹⁴⁵ and adequate scaffolding¹⁴⁶ can be used to devise industry-relevant substrates, probe their attributes and engineer scalable processes used on legitimate input/output states. To explore this newly revealed space, we are aided and inspired by the Cynefin® Framework.^{147,148} Cynefin® is a “multi-ontology sense-making framework” which can help make sense of the systems we find ourselves in so that we can act within them.¹⁴⁹ Perhaps the most crucial confusion elucidated by this framework is the distinction between the complex and the complicated domains. Complicated is a system where we know what information we are missing (known unknowns) and we must employ the adequate expertise to obtain it. In the complex domain of “unknown unknowns”, we do not know which questions we need to answer, and we must conduct safe-to-fail probes to discover important information. Based on these experiments, the next plausible question and the next step forward is defined and a path towards a potential solution emerges.

Being aware of the properties and interdependencies of each type of system can help us respond in meaningful ways.

We have used such responses to trigger deliberate movement within or across the Cynefin domains encompassed by the osmoDES concept, creating dynamics, as we recently described elsewhere¹⁵⁰ in detail (Fig. 6). As an example, our “shallow dive into Chaos” can trigger the emergence of a set of patterns (“osmolyte cocktails form multiDESs”) as an enabling constraint. Using this constraint as a springboard can help one explore “unimagined unknowns” about, for example, empirically observed natural substances of largely unbeknownst complexity of functions, components and distributions, such as saps. Actively exploring this space may help move beyond the postulation phase about eutectic mixtures and reveal actionable experiments to understand which questions are worth asking (“unknown unknowns”- *e.g.* is there an overarching preferred ratio between total methylamine kosmotropes and total perturbants that forms a DES?). Experimenting with such exploratory questions can eventually lead to exploitable practices by resolving specific problems through expertise (“known unknowns”- *e.g.* which osmolyte distribution and ratio best stabilise a particular recombinant protein contained in that sap in order to avoid product loss due to aggregation?). To the best of our knowledge, no prior art precedent has been reported for such a complexity-based approach, neither in the field of osmolytes nor in that of Deep Eutectic Systems. To further exploit the osmoDES concept as a constructor, we first need to acknowledge both known unknowns and unknown unknowns within the current state of the system we are in. For the first, Wardley mapping¹⁵¹ can be a useful tool to assess what kind of knowledge and expertise would be most relevant for each targeted use case and which current capacities require additional resources (Fig. 7).

Very often, when we dedicate resources to linked components of a Wardley map we become more aware of the space between disciplines and the amount of potential it holds. In this liminal space, we can look for alternative answers to existing questions (known unknowns), but also find additional plausible questions worth asking (unknown unknowns). A suitable approach to explore this space is the Entangled Trios method, which involves a triad of people “*from radically different backgrounds with no prior interaction but with a common purpose. This entangling leads to a more extensive and diverse exploration of the issue at hand which may lead to new ideas that can be explored. The three are kept together by a shared purpose and by well-crafted sets of actions which ritualize the exchange of knowledge and lead to new possibilities*”.¹⁵² Potential questions within such triads which could instigate a shared purpose would be: do single osmolytes always function as DESs with additional components we do not know yet? Do other osmolyte roles exist on top of kosmo- and chaotropic? Do other DES component roles exist on top of H-bond donor and acceptor? Are current synthetic DESs simplified versions of more nuanced natural eutectic systems? What is the role of water as a DES component? Could the “three states” postulated for hydrogels¹⁵³ be relevant in DES? And how does water regulate the interaction of the deep eutectic system with biomolecules?






Fig. 7 An orientative osmoDES Wardley map. A Wardley map is the depiction of a value chain anchored to specific “user needs”. Each component in this map, including the targeted “use cases”, is classified both by its maturity (horizontal axis) and by the value it has to the user (vertical axis), with lines indicating connections and interdependencies between components. Graphic based on the template by Ben Mosior (@hiredthought) and Sue Borchardt (@contemplatethis), available by LearnWardleyMapping.com.

For additional unknown unknowns, we must first act on an attribute we can affect and have the analytical tools to sense how the system responds to the perturbation,¹⁴⁷ to become aware of the affordances, the possibilities for further action, provided within the area of each experiment. Some of these actions could be, for example:

- to replicate more natural osmolyte cocktails and monitor their ability to form eutectic systems;
- to impose specific eutectic microenvironments on known enzymes and monitor for potential dormancy/wake-up triggers;
- to force cell protein production in specific eutectic microenvironments and monitor their effect on protein aggregation;
- to induce cell proliferation in DES-induced “*meta-stress*” conditions and monitor fitness landscapes and gene evolution for protein engineering;
- to subject different varieties of nucleic acid substrates to relevant DES and monitor their ability to interact and the topologies of their interaction;
- to generate new computational modules and tools based on relevant eutectic microenvironments and monitor their applicability and functionality limits.

Questions and actions will continue to arise the more we question and act, and the more diverse minds question and act the more probable it is for these initial thoughts and experiments to

become a path. We believe that the inherent beauty of this path is the potential to change the way we understand information flows in nature¹⁵⁴ by introducing the following assemblage:

Through their constant interaction with biomacromolecules, osmolyte-based microenvironments regulate a myriad of cellular properties, enable intra- and extracellular information flows¹⁵⁵ and facilitate the “enactive” functions of the components of biological systems.¹⁵⁶ Could osmolyte-based deep eutectic media be employed to better monitor these flows in complex natural systems and allow for the emergence of new properties and affordances we can explore and exploit? And could this new knowledge be leveraged to enable information flows within engineered systems?

Author contributions

The concept for the article was jointly conceived and developed by M.C.B. and T.A. who also designed the experiments, wrote the first draft, compiled figures and tables, and provided direction. I.R.R. and K.R. helped refine the concept. T.A. was responsible for the conceptual and contextual implementation of the Cynefin® framework and the complexity-related methods. M.C.B., M.P. and M.R. performed the experiments and prepared the DESs. The paper was reviewed and edited by all authors.



Conflicts of interest

There are no conflicts to declare.

Acknowledgements

A sincere thank you to Marko Rogošić for his diligent proofreading of the paper. Vilko Smrečki is gratefully acknowledged for recording the NMR spectra. Thanks to Anna Panagiotou of Cynefin Co for the sense-making guidance and to Eleni Margoudi (@elenimargou) and Athanasia Lantouri (@_lant) for the artwork.

References

- G. Zhang, *Organogenesis*, 2012, **8**, 101–102.
- L. Czech, L. Hermann, N. Stöveken, A. A. Richter, A. Höppner, S. H. J. Smits, J. Heider and E. Bremer, *Genes*, 2018, **9**, 1–58.
- H. Vanda, R. Verpoorte, P. G. L. Klinkhamer and Y. H. Choi, in *Deep Eutectic Solvents: Synthesis, Properties, and Applications*, 2019, pp. 61–81.
- B. B. Hansen, S. Spittle, B. Chen, D. Poe, Y. Zhang, J. M. Klein, A. Horton, L. Adhikari, T. Zelovich, B. W. Doherty, B. Gurkan, E. J. Maginn, A. Ragauskas, M. Dadmun, T. A. Zawodzinski, G. A. Baker, M. E. Tuckerman, R. F. Savinell and J. R. Sangoro, *Chem. Rev.*, 2021, **121**, 1232–1285.
- Y. Dai, J. Van Spronsen and G. Witkamp, *Anal. Chim. Acta*, 2013, **766**, 61–68.
- Y. Dai, E. M. Varypataki, E. A. Golovina, W. Jiskoot, G. J. Witkamp, Y. H. Choi and R. Verpoorte, *Natural deep eutectic solvents in plants and plant cells: In vitro evidence for their possible functions*, Elsevier Ltd, 1st edn, 2021, vol. 97.
- P. H. Yancey, *J. Exp. Biol.*, 2005, **208**, 2819–2830.
- T. Hasan, K. Kumari, S. C. Devi, J. Handa, T. Rehman, N. A. Ansari and L. R. Singh, *Hum. Vaccines Immunother.*, 2019, **15**, 514–525.
- N. Kushwah, V. Jain and D. Yadav, *Biomolecules*, 2020, **10**, 132.
- M. B. Burg and J. D. Ferraris, *J. Biol. Chem.*, 2008, **283**, 7309–7313.
- T. O. Street, D. W. Bolen and G. D. Rose, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 13997–14002.
- P. H. Yancey, *Sci. Prog.*, 2004, **87**, 1–24.
- P. H. Yancey, M. E. Clark, S. C. Hand, R. D. Bowlus and G. N. Somero, *Science*, 1982, **217**(4566), 1214–1222.
- G. L. Devlin, H. Parfrey, D. J. Tew, D. A. Lomas and S. P. Bottomley, *Am. J. Respir. Cell Mol. Biol.*, 2001, **24**, 727–732.
- P. H. Yancey and M. B. Burg, *Am. J. Physiol.: Regul., Integr. Comp. Physiol.*, 1990, **258**, DOI: [10.1152/ajpregu.1990.258.1.r198](https://doi.org/10.1152/ajpregu.1990.258.1.r198).
- P. H. Yancey and G. N. Somero, *Biochem. J.*, 1979, **183**, 317–323.
- L. R. Singh, T. Ali Dar, I. Haque, F. Anjum, A. A. Moosavi-Movahedi and F. Ahmad, *Biochim. Biophys. Acta, Proteins Proteomics*, 2007, **1774**, 1555–1562.
- S. Khan, Z. Bano, L. R. Singh, M. I. Hassan, A. Islam and F. Ahmad, *PLoS One*, 2013, **8**, e72533.
- M. Mukherjee and J. Mondal, *J. Phys. Chem. B*, 2020, **124**, 11316–11323.
- J. Rösigen and R. Jackson-Atogi, *J. Am. Chem. Soc.*, 2012, **134**, 3590–3597.
- J. Rösigen, *J. Phys. Chem. B*, 2015, **119**, 150–157.
- L. M. F. Holthauzen, M. Auton, M. Sinev and J. Rösigen, *Protein stability in the presence of cosolutes*, 2011, vol. 492.
- N. Kumar and N. Kishore, *J. Chem. Phys.*, 2013, **139**, 115104.
- J. H. Holland, *J. Syst. Sci. Complexity*, 2006, **19**, 1–8.
- S. Cachel, *The Growth of Biological Thought Revisited*, 1986, vol. 88.
- F. Guthrie, *London, Edinburgh, Dublin Philos. Mag. J. Sci.*, 2010, **17**, 462–482.
- A. P. Abbott, G. Capper, D. L. Davies, R. K. Rasheed and V. Tambyrajah, *Chem. Commun.*, 2003, 70–71.
- C. Florindo, A. J. S. McIntosh, T. Welton, L. C. Branco and I. M. Marrucho, *Phys. Chem. Chem. Phys.*, 2017, **20**, 206–213.
- H. Wang, S. Liu, Y. Zhao, J. Wang and Z. Yu, *ACS Sustainable Chem. Eng.*, 2019, **7**, 7760–7767.
- M. T. Zafarani-Moattar, H. Shekaari and F. Ghaffari, *J. Mol. Liq.*, 2020, **311**, 113347.
- K. Töpfer, A. Pasti, A. Das, S. M. Salehi, L. I. Vazquez-Salazar, D. Rohrbach, T. Feurer, P. Hamm and M. Meuwly, *J. Am. Chem. Soc.*, 2022, **144**, 14170–14180.
- L. Percevault, A. Jani, T. Sohler, L. Noirez, L. Paquin, F. Gauffre and D. Morineau, *J. Phys. Chem. B*, 2020, **124**, 9126–9135.
- S. Spittle, D. Poe, B. Doherty, C. Kolodziej, L. Heroux, M. A. Haque, H. Squire, T. Cosby, Y. Zhang, C. Fraenza, S. Bhattacharyya, M. Tyagi, J. Peng, R. A. Elgammal, T. Zawodzinski, M. Tuckerman, S. Greenbaum, B. Gurkan, C. Burda, M. Dadmun, E. J. Maginn and J. Sangoro, *Nat. Commun.*, 2022, **13**, 219.
- M. A. R. Martins, S. P. Pinho and J. A. P. Coutinho, *J. Solution Chem.*, 2019, **48**, 962–982.
- Y. P. Mbous, M. Hayyan, A. Hayyan, W. F. Wong, M. A. Hashim and C. Y. Looi, *Biotechnol. Adv.*, 2017, **35**, 105–134.
- Y. H. Choi, J. van Spronsen, Y. Dai, M. Verberne, F. Hollmann, I. W. C. E. Arends, G.-J. Witkamp and R. Verpoorte, *Plant Physiol.*, 2011, **156**, 1701–1705.
- F. Legrand, C. Nguyen, L. Augis, S. Fourmentin, G. Barratt, F.-X. Legrand, F. Legrand, C. Nguyen, L. Augis, S. Fourmentin and G. Barratt, *Deep Eutectic Solvents for Innovative Pharmaceutical Formulations*, 2021.
- D. Rente, M. C. Bubalo, M. Panić, A. Paiva, B. Caprin, I. R. Redovniković and A. R. C. Duarte, *J. Cleaner Prod.*, 2022, **380**, 135147.
- Y. Liang, W. Duan, X. An, Y. Qiao, Y. Tian and H. Zhou, *Bioresour. Technol.*, 2020, **310**, 123389.



- 40 M. Ishibashi, K. Tsumoto, M. Tokunaga, D. Ejima, Y. Kita and T. Arakawa, *Protein Expression Purif.*, 2005, **42**, 1–6.
- 41 S. Sharma, S. Sarkar, S. S. Paul, S. Roy and K. Chattopadhyay, *Sci. Rep.*, 2013, **3**, 1–9.
- 42 S. Rozas, C. Benito, R. Alcalde, M. Atilhan and S. Aparicio, *J. Mol. Liq.*, 2021, **344**, 117717.
- 43 O. S. Hammond, D. T. Bowron, A. J. Jackson, T. Arnold, A. Sanchez-Fernandez, N. Tsapatsaris, V. G. Sakai and K. J. Edler, *J. Phys. Chem. B*, 2017, **121**, 7473–7483.
- 44 N. López-Salas, J. M. Vicent-Luna, E. Posada, S. Imberti, R. M. Madero-Castro, S. Calero, C. O. Ania, R. J. Jiménez-Riobóo, M. C. Gutiérrez, M. L. Ferrer and F. del Monte, *ACS Sustainable Chem. Eng.*, 2020, **8**, 12120–12131.
- 45 N. López-Salas, J. M. Vicent-Luna, S. Imberti, E. Posada, M. J. Roldán, J. A. Anta, S. R. G. Balestra, R. M. M. Castro, S. Calero, R. J. Jiménez-Riobóo, M. C. Gutiérrez, M. L. Ferrer and F. del Monte, *ACS Sustainable Chem. Eng.*, 2019, **7**, 17565–17573.
- 46 M. C. Gutiérrez, M. L. Ferrer, C. R. Mateo and F. Del Monte, *Langmuir*, 2009, **25**, 5509–5515.
- 47 A. Gutiérrez, M. Atilhan and S. Aparicio, *J. Mol. Liq.*, 2021, **339**, 116758.
- 48 L. Cicco, G. Dilauro, F. M. Perna, P. Vitale and V. Capriati, *Org. Biomol. Chem.*, 2021, **19**, 2558–2577.
- 49 Choline chloride process, <https://matthey.com/products-and-markets/chemicals/process-licensing/choline-chloride-process>, (accessed 1 March 2023).
- 50 J. A. Raymond, *Fish Physiol. Biochem.*, 1994, **13**, 13–22.
- 51 J. P. Costanzo and R. E. Lee, *J. Exp. Biol.*, 2005, **208**, 4079–4089.
- 52 G. Pittaro, L. Cáceres, C. Bruno, A. Tomás, D. Bustos, M. Monteoliva, L. Ortega and E. Taleisnik, *Grass Forage Sci.*, 2016, **71**, 683–698.
- 53 P. C. Withers, G. Morrison and M. Guppy, *Physiol. Zool.*, 1994, **67**, 693–705.
- 54 P. H. Yancey and M. B. Burg, *Am. J. Physiol.: Renal, Fluid Electrolyte Physiol.*, 1989, **257**, 602–607.
- 55 P. C. Withers, G. Morrison and M. Guppy, *Physiol. Zool.*, 1994, **67**, 693–705.
- 56 C. X. Zeng, S. J. Qi, R. P. Xin, B. Yang and Y. H. Wang, *J. Mol. Liq.*, 2016, **219**, 74–78.
- 57 M. Benlebna, M. Ruesgas-Ramón, B. Bonafos, G. Fouret, F. Casas, C. Coudray, E. Durand, M. C. Figueroa-Espinoza and C. Feillet-Coudray, *J. Agric. Food Chem.*, 2018, **66**, 6205–6212.
- 58 A. M. Syakfanaya, F. C. Saputri and A. Mun'im, *Pharmacogn. J.*, 2019, **11**, 267–271.
- 59 A. R. Jesus, L. Meneses, A. R. C. Duarte and A. Paiva, *Cryobiology*, 2021, **101**, 95–104.
- 60 M. Panić, M. Andlar, M. Tišma, T. Rezić, D. Šibalić, M. C. Bubalo and I. R. Redovniković, *Waste Manage.*, 2021, **120**, 340–350.
- 61 M. Zdanowicz, *Int. J. Biol. Macromol.*, 2021, **176**, 387–393.
- 62 S. P. Simeonov and C. A. M. Afonso, *RSC Adv.*, 2016, **6**, 5485–5490.
- 63 L. Zhong, C. Wang, G. Yang, J. Chen, F. Xu, C. G. Yoo and G. Lyu, *Bioresour. Technol.*, 2022, **343**, 126022.
- 64 S. N. Pedro, M. S. M. Mendes, B. M. Neves, I. F. Almeida, P. Costa, I. Correia-Sá, C. Vilela, M. G. Freire, A. J. D. Silvestre and C. S. R. Freire, *Pharmaceutics*, 2022, **14**, 827.
- 65 M. A. Karadendrou, I. Kostopoulou, V. Kakokefalou, A. Tzani and A. Detsi, *Catalysts*, 2022, **12**, 1–17.
- 66 S. Sut, M. Faggian, V. Baldan, G. Poloniato, I. Castagliuolo, I. Grabnar, B. Perissutti, P. Brun, F. Maggi, D. Voinovich, G. Peron and S. Dall'Acqua, *Molecules*, 2017, **22**, 1–11.
- 67 M. Faggian, S. Sut, B. Perissutti, V. Baldan, I. Grabnar and S. Dall'Acqua, *Molecules*, 2016, **21**, 1–11.
- 68 F. Ilgen and B. König, *Green Chem.*, 2009, **11**, 848–885.
- 69 T. Wang, J. Jiao, Q.-Y. Gai, P. Wang, N. Guo, L.-L. Niu and Y.-J. Fu, *J. Pharm. Biomed. Anal.*, 2017, **145**, 339–345.
- 70 R. H. Kelly and P. H. Yancey, *Biol. Bull.*, 1999, **196**, 18–25.
- 71 W. E. S. Carr, J. C. Netherton, R. A. Gleeson and C. D. Derby, *The Biological bulletin*, 1996, **190**, 149–160, DOI: [10.2307/1542535](https://doi.org/10.2307/1542535).
- 72 A. K. Parida and A. B. Das, *Ecotoxicol. Environ. Saf.*, 2005, **60**, 324–349.
- 73 P. H. Yancey, W. R. Blake and J. Conley, *Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol.*, 2002, **133**, 667–676.
- 74 J. T. Gorke, F. Srienc and R. J. Kazlauskas, *Chem. Commun.*, 2008, 1235–1237.
- 75 H. Monhemi, M. R. Housaindokht, A. A. Moosavi-Movahedi and M. R. Bozorgmehr, *Phys. Chem. Chem. Phys.*, 2014, **16**, 14882–14893.
- 76 S. Sarkar, S. Ghosh and R. Chakrabarti, *RSC Adv.*, 2017, **7**, 52888–52906.
- 77 R. P. Sear, I. Pagonabarraga and A. Flaus, *BMC Biophys.*, 2015, **8**, 4–9.
- 78 R. P. Sear, *Faraday Discuss.*, 2008, **139**, 21–34.
- 79 N. H. O'Donnell, B. L. Møller, A. D. Neale, J. D. Hamill, C. K. Blomstedt and R. M. Gleadow, *Plant Physiol. Biochem.*, 2013, **73**, 83–92.
- 80 E. Durand, P. Villeneuve, C. Bourlieu-lacanal and F. Carrière, *Adv. Bot. Res.*, 2021, **97**, 133–158.
- 81 J. P. Bittner, N. Zhang, L. Huang, P. Domínguez De María, S. Jakobtorweihen and S. Kara, *Green Chem.*, 2022, **24**, 1120–1131.
- 82 M. Panić, M. Radović, I. Maros, A. J. Tušek, M. C. Bubalo and I. R. Redovniković, *Process Biochem.*, 2021, **102**, 1–9.
- 83 A. E. Delorme, J. M. Andanson and V. Verney, *Int. J. Biol. Macromol.*, 2020, **163**, 919–926.
- 84 S. L. Cao, X. Deng, P. Xu, Z. X. Huang, J. Zhou, X. H. Li, M. H. Zong and W. Y. Lou, *J. Agric. Food Chem.*, 2017, **65**, 2084–2088.
- 85 K. P. Lin, G. J. Feng, F. L. Pu, X. D. Hou and S. L. Cao, *Front. Bioeng. Biotechnol.*, 2020, **8**, 1–8.
- 86 S. Núñez-Pertíñez and T. R. Wilks, *Front. Chem.*, 2020, **8**, 1–11.
- 87 R. Jagannathan, P. M. Abraham and P. Poddar, *J. Phys. Chem. B*, 2012, **116**(50), 14533–14540, DOI: [10.1021/jp3050516](https://doi.org/10.1021/jp3050516).



- 88 M. Hayyan, M. A. Hashim, A. Hayyan, M. A. Al-Saadi, I. M. AlNashef, M. E. S. Mirghani and O. K. Saheed, *Chemosphere*, 2013, **90**, 2193–2195.
- 89 M. Hayyan, C. Y. Looi, A. Hayyan, W. F. Wong and M. A. Hashim, *PLoS One*, 2015, **10**, 1–18.
- 90 K. Radošević, M. Cvjetko Bubalo, V. Gaurina Srček, D. Grgas, T. Landeka Dragičević and I. Radojčić Redovniković, *Ecotoxicol. Environ. Saf.*, 2015, **112**, 46–53.
- 91 S. J. Bryant, M. N. Awad, A. Elbourne, A. J. Christofferson, A. V. Martin, N. Meftahi, C. J. Drummond, T. L. Greaves and G. Bryant, *J. Mater. Chem. B*, 2022, **10**, 4546–4560.
- 92 P. De Maayer, D. Anderson, C. Cary and D. A. Cowan, *EMBO Rep.*, 2014, **15**, 508–517.
- 93 Y. Song, W. Huang, Y. Zhou, Z. Li, R. Ji and X. Ye, *Arch. Insect Biochem. Physiol.*, 2021, **108**, e21846.
- 94 K. Tanaka, *Comp. Biochem. Physiol., Part B: Biochem. Mol. Biol.*, 1995, **110**, 539–545.
- 95 K. B. Storey, J. M. Storey, S. P. J. Brooks, T. A. Churchill and R. J. Brooks, *Proc. Natl. Acad. Sci. U. S. A.*, 1988, **85**, 8350–8354.
- 96 T. A. Churchill and K. B. Storey, *Can. J. Zool.*, 1992, **70**, 99–105.
- 97 A. Jani, T. Sohler and D. Morineau, *J. Mol. Liq.*, 2020, **304**, 112701.
- 98 L. Sherwood, H. Klandorf and P. H. Yancey, *Animal Physiology: From Genes to Organisms*, 2013.
- 99 J. H. Crowe, *Subcell. Biochem.*, 2015, **71**, 263–280.
- 100 M. Manzanera, *Environ. Microbiol.*, 2021, **23**, 3351–3359.
- 101 A. Sanchez-Fernandez, S. Prevost and M. Wahlgren, *Green Chem.*, 2022, **24**(11), 4437–4442.
- 102 R. Esquembre, J. M. Sanz, J. G. Wall, F. Del Monte, C. R. Mateo and M. L. Ferrer, *Phys. Chem. Chem. Phys.*, 2013, **15**, 11248–11256.
- 103 T. Pievani and E. Serrelli, *J. Anthropol. Sci.*, 2011, **89**, 9–23.
- 104 M. Frenkel-Pinter, A. S. Petrov, K. Matange, M. Travisano, J. B. Glass and L. D. Williams, *J. Mol. Evol.*, 2022, **90**, 166–175.
- 105 E. Manning and B. Massumi, *Thought in the Act: Passages in the Ecology of Experience*, University of Minnesota Press, 2014.
- 106 W. Y. Go, X. Liu, M. A. Roti, F. Liu and S. N. Ho, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 10673–10678.
- 107 A. Thiemicke and G. Neuert, *Sci. Adv.*, 2021, **7**, 1–13.
- 108 T. Kumar, M. Yadav and L. R. Singh, *Curr. Pharm. Des.*, 2016, **22**, 3050–3057.
- 109 Y. Chen, S. Yang, J. Tavormina, D. Tampe, M. Zeisberg, H. Wang, K. K. Mahadevan, C.-J. Wu, H. Sugimoto, C.-C. Chang, R. R. Jenq, K. M. McAndrews and R. Kalluri, *Cancer Cell*, 2022, **40**, 818–834.
- 110 M. Warepam, K. Ahmad, S. Rahman, H. Rahaman, K. Kumari and L. R. Singh, *Biomolecules*, 2020, **10**, 286.
- 111 B. De Paepe, C. Merckx, J. Jarošová, M. Cannizzaro and J. L. De Blecker, *Biomolecules*, 2020, **10**, 521.
- 112 M. A. Bhat, K. Ahmad, M. S. A. Khan, M. A. Bhat, A. Almatroudi, S. Rahman and A. T. Jan, *Biomolecules*, 2020, **10**, 863.
- 113 M. Gao and R. Winter, *J. Diabetes Res.*, 2015, DOI: [10.1155/2015/849017](https://doi.org/10.1155/2015/849017).
- 114 D. C. Mikles, V. Bhat, B. J. Schuchardt, C. B. McDonald and A. Farooq, *Biopolymers*, 2015, **103**, 74.
- 115 M. Mcdermott, C. Kemper, W. Barone, G. Jost and J. Endrikat, *Br. J. Radiol.*, 2020, 93.
- 116 G. N. Gomes and Z. A. Levine, *J. Phys. Chem. B*, 2021, **125**, 1974–1996.
- 117 H. Marshall, M. Venkat, N. S. Hti Lar Seng, J. Cahn and D. H. Juers, *Acta Crystallogr., Sect. D: Biol. Crystallogr.*, 2012, **68**, 69–81.
- 118 D. Le Bihan, *Phys. Med. Biol.*, 2007, 57–90.
- 119 H. M. Taïeb, L. Bertinetti, T. Robinson and A. Cipitria, *PLoS One*, 2022, **17**, e0268297.
- 120 S. Aravindan, S. Chen, H. Choudhry, C. Molfetta, K. Y. Chen and A. Y. C. Liu, *Sci. Rep.*, 2020, **10**, 1–12.
- 121 R. Morizane, A. Q. Lam, B. S. Freedman, S. Kishi, M. T. Valerius and J. V. Bonventre, *Nat. Biotechnol.*, 2015, **33**, 1193–1200.
- 122 M. Takasato, P. X. Er, H. S. Chiu, B. Maier, G. J. Baillie, C. Ferguson, R. G. Parton, E. J. Wolvetang, M. S. Roost, S. M. C. De Sousa Lopes and M. H. Little, *Nature*, 2015, **526**, 564–568.
- 123 A. Ganguli, A. Mostafa, C. Saavedra, Y. Kim, P. Le, V. Faramarzi, R. W. Feathers, J. Berger, K. P. Ramos-Cruz, O. Adeniba, G. J. P. Diaz, J. Drnevich, C. L. Wright, A. G. Hernandez, W. Lin, A. M. Smith, F. Kosari, G. Vasmatazis, P. Z. Anastasiadis and R. Bashir, *Sci. Adv.*, 2021, **7**, DOI: [10.1126/SCIADV.ABC1323](https://doi.org/10.1126/SCIADV.ABC1323).
- 124 P. Mai, J. Hampl, M. Baca, D. Brauer, S. Singh, F. Weise, J. Borowiec, A. Schmidt, J. M. Küstner, M. Klett, M. Gebinoga, I. S. Schroeder, U. R. Markert, F. Glahn, B. Schumann, D. Eckstein and A. Schober, *Bioengineering*, 2022, **9**, 220.
- 125 B. Dorgau, M. Georgiou, A. Chaudhary, M. Moya-Molina, J. Collin, R. Queen, G. Hilgen, T. Davey, P. Hewitt, M. Schmitt, S. Kustermann, F. Pognan, D. H. Steel, E. Sernagor, L. Armstrong and M. Lako, *Stem Cells Transl. Med.*, 2022, **11**, 159–177.
- 126 J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Židek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A. Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P. Kohli and D. Hassabis, *Nature*, 2021, **596**, 583–589.
- 127 E. T. Powers and L. M. Gierasch, *J. Mol. Biol.*, 2021, **433**(20), DOI: [10.1016/J.JMB.2021.167197](https://doi.org/10.1016/J.JMB.2021.167197).
- 128 G. D. Rose, *Biochemistry*, 2021, **60**, 3753–3761.
- 129 J. P. Bittner, L. Huang, N. Zhang, S. Kara and S. Jakobtorweihen, *J. Chem. Theory Comput.*, 2021, **17**, 5322–5341.
- 130 N. B. Pham and W. S. Meng, *Int. J. Pharm.*, 2020, **585**, 119523.
- 131 J. R. Christopher, *Curr. Opin. Biotechnol.*, 2014, **30**, 211–217.



- 132 A. Singh, V. Upadhyay, A. K. Upadhyay, S. M. Singh and A. K. Panda, *Microb. Cell Fact.*, 2015, **14**, 1–10.
- 133 D. Andrijevic, Z. Vrselja, T. Lysyy, S. Zhang, M. Skarica, A. Spajic, D. Dellal, S. L. Thorn, R. B. Duckrow, S. Ma, P. Q. Duy, A. U. Isiktas, D. Liang, M. Li, S. K. Kim, S. G. Daniele, K. Banu, S. Perincheri, M. C. Menon, A. Huttner, K. N. Sheth, K. T. Gobeske, G. T. Tietjen, H. P. Zaveri, S. R. Latham, A. J. Sinusas and N. Sestan, *Nature*, 2022, **608**, 405–412.
- 134 K. Athirathinam, S. Nandakumar and R. Kandasamy, *Macromol. Res.*, 2022, **30**, 599–608.
- 135 C. H. Pi, G. Yu, A. Petersen and A. Hubel, *Sci. Rep.*, 2018, **8**, 1–13.
- 136 A. M. R. Kabir, T. Munmun, T. Hayashi, S. Yasuda, A. P. Kimura, M. Kinoshita, T. Murata, K. Sada and A. Kakugo, *ACS Omega*, 2022, **7**, 3796–3803.
- 137 A. Argiolas, G. L. Puleo, E. Sinibaldi and B. Mazzolai, *Sci. Rep.*, 2016, **6**, 1–8.
- 138 M. Zheng, Z. Wang, H. Chang, L. Wang, S. W. T. Chew, D. C. S. Lio, M. Cui, L. Liu, B. C. K. Tee and C. Xu, *Adv. Healthcare Mater.*, 2020, **9**, 1901683.
- 139 X. Wang and I. Rivière, *Mol. Ther. – Oncolytics*, 2016, **3**, 16015.
- 140 D. B. Kell, G. J. Laubscher and E. Pretorius, *Biochem. J.*, 2022, **479**, 537–559.
- 141 M. Vendruscolo and M. Fuxreiter, *Nat. Commun.*, 2022, **13**, 1–11.
- 142 M. C. Gutierrez, M. L. Ferrer, L. Yuste, F. Rojo and F. del Monte, *Angew. Chem. Int. Ed.*, 2010, **49**, 2158–2162.
- 143 S. Nardecchia, M. C. Gutiérrez, M. L. Ferrer, M. Alonso, I. M. López, J. C. Rodríguez-Cabello and F. del Monte, *Biomacromolecules*, 2012, **13**, 2029–2036.
- 144 S. R. Włodarczyk, D. Custódio, A. Pessoa and G. Monteiro, *Eur. J. Pharm. Biopharm.*, 2018, **131**, 92–98.
- 145 D. Deutsch, *Synthese*, 2013, **190**, 4331–4359.
- 146 J. Turner, D. Snowden and N. Thurlow, *Systems*, 2022, **10**(7), DOI: [10.3390/systems10010007](https://doi.org/10.3390/systems10010007).
- 147 D. Snowden, *J. Knowl. Manage.*, 2002, **6**, 100–111.
- 148 D. Snowden, Z. Goh and S. Borchardt, *Cynefin – Weaving Sense-Making into the Fabric of Our World*, Wilmington, DE, USA, 2021.
- 149 E. Commission, J. R. Centre, A. Rancati and D. Snowden, *Managing complexity (and chaos) in times of crisis : a field guide for decision makers inspired by the Cynefin framework*, Publications Office, 2021.
- 150 T. Andreou, in *Natural Resources, Green Technology And Sustainable Development*, 2022, pp. 111–116.
- 151 S. Wardley, “A Wardley Map”, <https://blog.gardeviance.org/2014/02/a-wardley-map.html>, 2017.
- 152 Entangled trios – Cynefin.io, https://cynefin.io/wiki/Entangled_trios, (accessed 9 September 2022).
- 153 B. L. Dargaville and D. W. Huttmacher, *Nat. Commun.*, 2022, **13**, 1–10.
- 154 W. S. Imari and P. C. W. Davies, *J. R. Soc., Interface*, 2013, **10**, 1–9.
- 155 H. M. Taïeb, D. S. Garske, J. Contzen, M. Gossen, L. Bertinetti, T. Robinson and A. Cipitria, *Sci. Rep.*, 2021, **11**, 13455.
- 156 T. Froese and E. A. Di Paolo, *Pragmat. Cognit.*, 2011, **19**, 1–36.
- 157 J. R. Treberg, B. Speers-Roesch, P. M. Piermarini, Y. K. Ip, J. S. Ballantyne and W. R. Driedzic, *J. Exp. Biol.*, 2006, **209**, 860–870.
- 158 D. Shugar, *Biochim. Biophys. Acta*, 1952, **8**, 302–309.

