

Dalton Transactions

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Boosting Catalyst Activity in *Cis*-Selective Semi-Reduction of Internal Alkynes by Tailoring the Assembly of All-Metal Aromatic Tri-Palladium Complexes

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Anna Monfredini,^{a†} Veronica Santacrose,^{a†} Pierre-Alexandre Deyris,^b Raimondo Maggi,^a Franca Bigi,^a Giovanni Maestri^{*a} and Max Malacria^{*b,c}


Highly symmetric $[\text{Pd}_3]^+$ clusters that present delocalized metal-metal bonds can catalyse the selective semi-reduction of internal alkynes to *cis*-alkenes. Studies on factors governing the formation of all-metal aromatics enabled the design of an optimised catalytic system that delivers *cis*-alkenes with almost complete selectivity on a gram scale with very low catalyst loadings (0.03 mol%).

All-metal aromatic molecules feature delocalised metal-metal bonds that parallel bonding of their well-known main-group counterparts.¹ While they are often synthetically challenging at best,² we recently developed a simple route³ to a family of bench-stable $[\text{Pd}_3]^+$ complexes whose core represent a prototypical well-defined sub-nanometric metal surface.⁴

These complexes could be decorated with a variety of organic fragments and represent therefore a valuable asset to assess the consequences of their bonding mode on coordination chemistry and catalysis. Internal alkynes were semi-reduced to alkenes with good *cis*-selectivity under transfer-hydrogenation conditions,⁵ which avoids the use of hazardous H_2 gas.⁶ The best first generation catalyst (I, 1 mol%) delivered full conversion of phenylpropyne **1** upon 10 hours in refluxing THF. No alkane was detected and the ratio between semi-reduced alkenes was 93:3:4 (*cis*-, *trans*- and isomerised-products respectively, Scheme 1, top).

We report herein preliminary results on factors triggering the formation of triangular all-metal aromatic $[\text{Pd}_3]^+$ complexes. Through this approach we were able to decorate their metal core with two different phosphinic ligands, a neutral and an anionic one. This in turn enabled to develop a catalytic system for the selective semi-reduction of **1** to *cis*-alkene **2** with unparalleled activity among Pd-based catalysts under transfer

hydrogenation conditions. The novel complex delivers desired *cis*-alkene almost quantitatively, in shorter times and a with thirty-three-times lower catalytic charge (Scheme 1, bottom).



Scheme 1. First vs second generation metal-aromatic transfer hydrogenation catalysts

Wondering to gain insights on factors governing the formation of delocalized metal-metal bonds, we tried to monitor these reactions by NMR. In a 50 mL Schlenk we thus mixed 0.2 mmol of $\text{Pd}(\text{dba})_2$ (dba = dibenzylideneacetone), 0.2 mmol of PPh_3 and 0.1 mmol of diphenyldisulphide in 20 mL of CDCl_3 , taking 0.5 mL samples at regular intervals to collect ^1H and ^{31}P NMR spectra (Figure 1, spectra in ESI).

Surprisingly, the broad and diagnostic ^{31}P resonance of the $[\text{Pd}_3]^+$ cation at 15.5 ppm was already visible within few minutes (Figure 1, red line) and free phosphine at -5 ppm was no longer present. From the proton spectrum, we noticed the consumption of $\text{Pd}(\text{dba})_2$ and evaluated 18% of the desired $[\text{Pd}_3]^+$ cation using dba resonances as internal standard. Two other broad ^{31}P resonances were observed at 24 and 27 ppm respectively. We were so far unable to unambiguously assign these signals, although their intensity slowly fades through

^a Dipartimento di Chimica, Università degli Studi di Parma; 17/A Parco Area delle Scienze, 43124 Parma (Italy). giovanni.maestri@unipr.it

^b ICSN CNRS (UPR2301); 1 Av. de la Terrasse, Bat. 27, 91198 Gif s/Yvette (France); max.malacria@cnsr.fr

^c UPMC Sorbonne Université, IPCM (UMR CNRS 8232); 4 place Jussieu, C. 229, 75005 Paris (France).

† These authors contributed equally.

Electronic Supplementary Information (ESI) available: reaction procedures, characterization of complexes, copies of NMR spectra. See DOI: 10.1039/x0xx00000x

time with a concomitant increase of the desired resonance at 15.5 ppm. Upon 60 minutes, the amount of the $[\text{Pd}_3]^+$ complex is around 40% (green line). Addition of 0.33 equiv. of AgSbF_6 resulted in a further increase of the desired resonance at the expense of the two unassigned species (cyan line). Upon celite filtration under nitrogen and concentration *in vacuo*, the $[\text{Pd}_3]\text{SbF}_6$ complex **II** became the almost exclusive species present in the crude mixture (Figure 1, blue line, 86% isolated yield). Taken together, these observation showed that addition of a silver salt is not necessary to trigger the formation of metal-aromatic $[\text{Pd}_3]^+$ clusters. This in turn suggested that the chlorinated solvent played a key role in the formation of the $[\text{Pd}_3]^+$ core.




Figure 1. ^{31}P spectra of crude samples from the synthesis of $[\text{Pd}_3]\text{SbF}_6$ complex highlighting the diagnostic resonance of its three equivalent PPh₃ units.

The overall process involves the formal oxidation of Pd(0) to Pd($4/3$). Considering the stoichiometry of the reaction, the disulphide can oxidise the noble metal on average to Pd(I). Radical palladium species⁷ might be sufficiently strong reductant to trigger SET (single electron transfer) on a chloroform molecule (-1.09 V vs SCE).⁸ This would form the corresponding radical anion, which eventually collapses into a radical and a chloride ion. The formal addition of a cationic Pd(II) fragment on a neutral Pd(I) dimer⁹ would eventually deliver the $[\text{Pd}_3]^+$ cation, isolable as a chloride salt, in analogy to the step-wise assembly of $[\text{Au}_3]^+$ complexes from an Au(I) cation and an Au(0) dimer reported by Bertrand.¹⁰ The $[\text{Pd}_3]^+$ cation barely forms in CH_2Cl_2 while the reaction is almost instantaneous in CCl_4 , being limited by the solubilisation of $\text{Pd}(\text{dba})_2$ only. These results fit with the trend of C-Cl BDE and reduction potentials among chlorinated solvents.⁸

Likely, the soluble AgX salt just favours anion metathesis steered by the precipitation of AgCl in chloroform. This suggests that $[\text{Pd}_3]^+$ complexes might be synthesised in any given solvent if a suitable mild stoichiometric oxidant is added. This is of particular interest for catalysis as it would allow the *in-situ* preparation of $[\text{Pd}_3]^+$ complexes in any medium.

We performed our usual reaction in degassed THF without addition of any additive. Expectedly, no traces of the desired $[\text{Pd}_3]^+$ cation were observed upon 24 hours at room temperature. On the contrary, mixing reagents for 1 hours and adding then 0.33 equiv. of AgSbF_6 allowed the mixture to turn deep red, eventually delivering complex **II** in 73% yield.

We then performed the same experiment adding a weaker Cu(II) oxidant (+0.28 V vs SCE, +0.6 V for Ag(I)).

Thus, 0.2 mmol of $\text{Pd}(\text{dba})_2$, 0.2 mmol of $\text{P}(p\text{-tol})_3$ and 0.1 mmol of dimethyldisulphide were stirred for 1 hour at RT in degassed THF. Two main resonances at 23 and 26 ppm were observed in the corresponding ^{31}P spectra, which likely correspond to species observed in chloroform presented in Figure 1 (spectra in ESI). Addition of 0.066 mmol of $\text{Cu}(\text{OTf})_2$ triggered a slow turning of the solution towards the red and, furthermore, the appearance of the diagnostic ^{31}P resonance of the $[\text{Pd}_3]\text{OTf}$ complex **III** at 15 ppm. The reaction was slower than the previous one but once again a single resonance was eventually detected upon 24 hours stirring at RT. These findings paralleled observations made using chlorinated solvents and provide preliminary insights on the mechanism of formation of all-metal aromatic $[\text{Pd}_3]^+$ complexes.

We thus attempted the catalytic semireduction of phenylpropyne **1** preparing *in-situ* the $[\text{Pd}_3]^+$ complex. Triethylammonium formate and **1** were added as a THF solution to eventually provide the same concentration previously reported (0.1 M). Sadly, the reaction proved significantly slower, requiring 24 hours to achieve full conversion. The selectivity towards desired *cis*-alkene **2** remained similar (91%). Partial decomposition to Pd black was observed through time although no alkane has been detected. The reduced activity may be due to dba, which can likely compete with the substrate slowing down its semi-reduction.

Table 1. Synthesis of $[\text{Pd}_3]^+$ complexes A-E

| Entry | R ₃ | R' | Complex | Yield (%) |
|-------|--|------------------------------------|----------|-----------|
| 1 | (4-F-C ₆ H ₄) ₃ | 4-Me-C ₆ H ₄ | A | 71 |
| 2 | (4-F-C ₆ H ₄) ₃ | Me | B | 55 |
| 3 | (4-Me-C ₆ H ₄) ₃ | Ph | C | 59 |
| 4 | (4-Me-C ₆ H ₄) ₃ | Me | D | 76 |
| 5 | (Ph) ₂ C ₆ H ₄ SO ₃ Na | Me | E | 72 |


Reaction conditions: on 0.2 mmol scale, 0.01 M in degassed chloroform.

We then turned to another strategy to design better catalysts for this reaction. Reasoning that $[\text{Pd}_3]^+$ is the major species in chloroform even without the addition of any other metal salts, we run the reaction with 0.33 equiv. of a sodiumsulfonate phosphine. This should provide a zwitterionic complex by filtering off coproduced NaCl. We prepared accordingly a small library of complexes (Table 1).

Several organic fragments varying both electronic and steric features could be easily introduced. Products **A-E** were purified through a series of solvent washings³ (55-76% yield). They were characterized *via* HRMS analyses and multinuclear NMR experiments. In contrast to C_3 -symmetric complexes, as salt **I**, which showed $[Pd_3]^+$ cations in ESI⁺ experiments, we invariably observed protonated molecular adducts only for **A-D**. This confirms their zwitterionic nature. Regarding NMR, two pattern of resonances in a 2:1 ratio were observed for bridging thiolate fragments in ¹H spectra. ³¹P signals followed suite. For instance, a broad triplet and a broad doublet in a 1:2 ratio were observed for **A** (at 17 and 13 ppm respectively).

We then tested these new complexes on the catalytic semi-reduction of phenylpropyne **1** in the presence of a five-fold excess of triethylammonium formate as hydrogen source (Table 2). This excess ensures that the chemoselectivity towards semi-reduction of alkynes is due to the catalytic system rather than to a mere shortage of hydrogen donor.¹¹

Table 2. Catalytic semi-reduction of phenylpropyne **1** with $[Pd_3]^+$ complexes **A-E**



| Entry | R | R' | t (h) | Conv. of 1 (%) ^a | Sel. to 2 (%) ^a |
|-------|--|------------------------------------|-------|------------------------------------|-----------------------------------|
| 1 | (4-F-C ₆ H ₄) ₃ | 4-Me-C ₆ H ₄ | 8 | 97 | 85 |
| 2 | (4-F-C ₆ H ₄) ₃ | Me | 5 | quant. | 87 |
| 3 | (4-Me-C ₆ H ₄) ₃ | Ph | 6 | quant. | 78 |
| 4 | (4-Me-C ₆ H ₄) ₃ | Me | 3 | quant. | 77 |
| 5 | (Ph) ₂ C ₆ H ₄ SO ₃ Na | Me | 3 | quant. | 77 |

Conditions: 0.35 mmol of **1**, 0.1 M in THF, 1.75 mmol of triethylammonium formate, 1 mol% of complexes **A-E**; a: 0.1 mL samples were periodically taken and analysed by GC and GC-MS using *p*-xylene (0.3 mmol) as internal standard.

Complexes with fluorinated phosphines proved active with either aromatic or aliphatic bridging thiolates (entries 1 and 2). The latter proved a better substitution pattern providing complete conversion of substrate upon five hours. In both cases, *cis*-selectivity resulted lower than that previously observed (85% and 87% respectively). No traces of alkane were detected. Alkenes **3** and **4** formed in a comparable amount (6-8%). Complexes bearing two *p*-tolylphosphines were even more active (entries 3 and 4). In combination with bridging methylthiolates, complete conversion of **1** was achieved within 3 hours, albeit at the expense of selectivity towards **2** (77%). **3** and **4** formed in 10% and 13% yield respectively. The same outcome was observed using complex **E** (entry 5, 77%). Interestingly, the catalytic system using any complex of the series **A-E** proved both more active and less selective than the best one previously developed based on C_3 -symmetric $[Pd_3]^+$ complexes (Scheme 1, top).

We reasoned that this might be a *trans*-like effect¹² of the anionic ligand that would make more labile the phosphine on

the opposite edge of the metal kernel. It can correlate with (slight) upfield shifts of the latter observed by ³¹P NMR and could rationalise the activity of complexes with *p*-fluorinated phosphines (**A** and **B**, entries 1-2). Their corresponding C_3 -symmetric $[Pd_3]^+$ peers are indeed fairly active only for the semi-reduction of **1**, not even attaining full conversion upon 96 hours.⁵ DFT modeling supports this hypothesis. The Pd-P distance with the anionic phosphine shortens in **D** compared to **I**, while those with neutral ones are slightly longer (Figure 2, details in ESI).





Figure 2. Comparison of optimised structures of cation **I** and zwitterion **D**; the Pd-P distances in **D** at the M11/lacvp(d) level vary by +0.0041 and -0.0078 Å with the neutral and anionic phosphines respectively.

In all cases we noticed formation of traces of Pd black when conversion of **1** passed 70%, suggesting that zwitterionic complexes **A-E** could be less robust than their cationic analogues. Nonetheless, no trace of over-reduced alkane has been ever detected, even prolonging warming after full conversion. In these cases, partial isomerization to **3** and **4** was however noticed (below 5% upon 24 hours).

Table 3. Addition of ligands on the semi-reduction of **1** with complex **D**



| Entry | L | mol% | t (h) | Conv. of 1 (%) ^a | Sel. to 2 (%) ^a |
|-------|---|------|-------|------------------------------------|-----------------------------------|
| 1 | P(4-Me-C ₆ H ₄) ₃ | 3 | 4 | quant. | 96 |
| 2 | P(4-Me-C ₆ H ₄) ₃ | 10 | 24 | quant. | 96 |
| 3 | dba | 6 | 24 | quant. | 92 |
| 4 | P(Ph) ₂ C ₆ H ₄ SO ₃ Na | 3 | 4 | quant. | 96 |


Conditions: as Table 2; a: by GC using *p*-xylene as internal standard.

Taken together, these results confirmed the complete chemoselectivity of the catalytic system towards alkynes and let us speculate that isomerisation of **2** could be due to Pd species generated from the decomposition of the actual catalyst in the reducing reaction medium.¹³ We wondered

whether addition of a suitable ligand might slow down these undesired processes without quenching activity (Table 3). We were delighted that addition of 3 mol% of *p*-tolylphosphine proved decisive (entry 1). Full conversion of **1** was achieved upon 4 hours and the selectivity towards **2** was almost complete (96%). Isomers **3** and **4** formed in 2% yield each. Furthermore, the homogeneous orange solution of complex **D** did not show any trace of formation of black metal colloids before consumption of **1** (photos in ESI). We then added 10 mol% of phosphine (entry 2). Conversion of **1** required 24 hours, in agreement with a stiffer competition between substrates and ligands for metal coordination. Selectivity towards **2** remained 96% despite the prolonged warming. As in the previous case, no traces of visible heterogeneous aggregates appeared throughout this (longer) experiment. The use of a coordinating electron-poor olefin as dba confirmed the outcome previously observed generating a $[\text{Pd}_3]^+$ complex *in-situ* (vide supra), requiring 24 hours to achieve full conversion of **1** (entry 3). In this case, traces of Pd black were observed together with a slightly more pronounced formation of **3** and **4** (4% each). Addition of 3 mol% of the sodium salt of a sulfonated phosphine gave **2** with 96% selectivity within 4 hours (entry 4) and the reaction mixture remained once more a red homogeneous solution.

clearly visible throughout the reaction, indicating that the complex remains stable in solution until full conversion of **1**. These results suggest that addition of a suitable ligand can be a tool to stabilise the active species and to minimise undesired isomerization of *cis*-olefins. We thus tried to exploit these hypotheses by reducing the catalyst loading (Table 4).

Table 4. Reduction of catalyst loadings on semi-reduction of **1**



| Entry | X (mol%) | Y (mol%) | t (h) | Conv. of 1 (%) ^a | Sel. to 2 (%) ^a | TOF (h ⁻¹) ^b |
|------------------|----------|----------|-------|------------------------------------|-----------------------------------|-------------------------------------|
| 1 | 1 | - | 3 | quant. | 77 | 18 |
| 2 | 0.3 | - | 4 | quant. | 90 | 137 |
| 3 | 0.1 | - | 7 | quant. | 96 | 329 |
| 4 ^c | 0.05 | - | 6 | 77 | 98 | 582 |
| 5 ^c | 0.05 | 0.15 | 5 | quant. | 98 | 1026 |
| 6 ^{c,d} | 0.03 | 0.09 | 6 | quant. | 98 | 1242 |

Conditions: 5.0 mg of complex **D**, 0.0035 mmol; THF was progressively reduced among the series to keep a fixed concentration of **D** of 0.001 M; *a*: by GC; *b*: determined from the linear slope of a time/moles of **2** curve, typically between 0.5 and 2-4 hours; *c*: without THF; *d*: without *p*-xylene as internal GC standard.




Figure 3. UV-Vis absorption spectra of samples collected from the catalytic reaction of Table 3, entry 1; 100 μl samples were taken via syringe and diluted with 4.9 mL of degassed THF prior to analyses; the right box presents absorption spectra of pure **D** (blue) and **D** with added tritylphosphine (green) in THF.

We performed ESI-MS analyses on samples collected at regular intervals to check the stability of **D** during the reaction. This method proved useful to detect a fading concentration of cation **I** throughout the reaction, in agreement with the formation of visible Pd black.⁵ However, no Pd-containing ion has been detected in this case, likely owing to a difficult ionisation of zwitterion **D** (spectra in ESI). We thus resorted to UV-Vis absorption to monitor the diagnostic bands of **D** in the region between 300 and 550 nm (Figure 3, box), which is not affected by any reagent during the catalytic reaction. Figure 3 shows absorption spectra of reaction samples. Both bands in the near UV (320-380 nm) and the saddle at 450 nm remains

Keeping in mind the practical viability of the synthetic method, we decided to fix the concentration of complex **D** in the mixture, reducing progressively the amount of THF to hold for a higher amount of **1**. This is possible thanks to the increased solubility in apolar media of zwitterion **D** compared to salt **I**. Reducing the catalyst loading to 0.3 mol% proved beneficial for selectivity (90%, entry 2) and full conversion was attained upon four hours. Further reduction of **D** to 0.1 mol% provided 96% of **2** upon 7 hours (entry 3). The reaction with 0.05 mol% did not reach full conversion, and the consumption of **1** ended at 77% upon 6 hours (entry 4). In all these cases, formation of traces of Pd black were observed. For comparison, first generation complexes, as **I**, were no longer active before full conversion of alkyne at catalyst loadings lower than 1 mol%.

We then repeated our last attempt adding 0.15 mmol% of phosphine (entry 5). To our delight, the solution remained homogeneous and **1** was completely converted upon 5 hours, delivering almost exclusively **2** (98%). We further reduced the loading of **D** to 300 ppm, observing a comparable outcome (entry 6). Full conversion of 1.20 g of **1** was observed upon 6 hours, with 98% selectivity using commercial reagents without any previous purification. Pure **2** has been isolated adsorbing the crude on a 2 cm silica pad and eluting with 5 mL of pentane (1.15 g, 96% yield). This highlights the practical and environmental viability of present method.¹⁴ We are unaware of Pd-based catalytic systems active at so-low loadings under transfer-hydrogenation conditions.^{6,11,13-14}


Conclusions

We reported preliminary studies on the mechanism of formation of triangular Pd complexes with delocalized metal-metal bonds. Through this approach, the panel of readily accessible structures has been further expanded, allowing to access zwitterionic tripalladium complexes. These clusters form in one-step from commercial reagents, are definitely stable at the solid state and proved remarkable precatalysts under transfer-hydrogenation conditions, highlighting the great potential of all-metal aromatics in catalysis.

Notes and references

‡ We thank MIUR, UPCM and CNRS for funding. Research on metal aromaticity is funded through SIR Project "AROMA-TriP" (Grant No.: RBSI14NKFL, GM).

- 1 J. M. Mercero, A. I. Boldyrev, G. Merino and J. M. Ugalde, *Chem. Soc. Rev.*, 2015, **44**, 6519; I. Fernández, G. Frenking and G. Merino, *Chem. Soc. Rev.*, 2015, **44**, 6452.
- 2 R. Hoffmann, *Am. Sci.*, 2015, **103**, 18.
- 3 Y. Wang, P.-A. Deyris, F. Blanchard, Y. Li, F. Bigi, R. Maggi, S. Blanchard, G. Maestri and M. Malacria, *Chem. Eur. J.*, 2015, **21**, 12271.
- 4 S. Blanchard, L. Fensterbank, G. Gontard, E. Lacôte, G. Maestri and M. Malacria, *Angew. Chem. Int. Ed.*, 2014, **53**, 1897.
- 5 P.-A. Deyris, T. Caneque, Y. Wang, P. Retailleau, F. Bigi, R. Maggi, G. Maestri and M. Malacria, *Chem. Cat. Chem.*, 2015, **7**, 3266.
- 6 D. Wang and D. Astruc, *Chem. Rev.*, 2015, **115**, 6621; C. Oger, L. Balas, T. Durand and J.-M. Galano, *Chem. Rev.*, 2013, **113**, 1313; S. Gladiali and E. Alberico, *Chem. Soc. Rev.*, 2006, **35**, 226.
- 7 G. Maestri, M. Malacria and E. Derat, *Chem. Commun.*, 2013, **49**, 10424; T. Troadec, S.-y. Tan, C. J. Wedge, J. P. Rourke, P. R. Unwin and A. B. Chaplin, *Angew. Chem. Int. Ed.*, 2016, **55**, 3754; G. Manolikades and P. Knochel, *Angew. Chem. Int. Ed.*, 2009, **48**, 205; Q. Liu, X. Dong, J. Li, J. Xiao, Y. Dong and H. Liu, *ACS Catalysis*, 2015, **5**, 6111.
- 8 A. Studer and D. Curran, *Nat. Chem.*, 2014, **6**, 765; J. Bertrand, I. Gallardo, M. Moreno and J. M. Saveant, *J. Am. Chem. Soc.*, 1992, **114**, 9576; X.-Q. Hu, X. Qi, J.-R. Chen, Q.-Q. Zhao, Q. Wei, Y. Lan and W.-J. Xiao, *Nat. Commun.*, 2016, **7**, 11188.
- 9 R. Vilar, D. M. P. Mingos and C. J. Cardin, *J. Chem. Soc., Dalton Trans.*, 1996, 4313.
- 10 L. Jin, D. S. Weinberger, M. Melaimi, C. E. Moore, A. L. Rheingold and G. Bertrand, *Angew. Chem. Int. Ed.*, 2014, **53**, 9059.
- 11 P. Hauwert, R. Boerleider, S. Warsink, J. J. Weigand and C. J. Elsevier, *J. Am. Chem. Soc.*, 2010, **132**, 16900; R. Shen, T. Chen, Y. Zhao, R. Qiu, Y. Zhou, S. Yin, X. Wang, M. Goto and L.-B. Han, *J. Am. Chem. Soc.*, 2011, **133**, 17037; J. Broggi, V. Jurcik, O. Songis, A. Poater, L. Cavallo, A. M. Z. Slawin and C. S. J. Cazin, *J. Am. Chem. Soc.*, 2013, **135**, 4588.
- 12 B. J. Coe and S. J. Glenwright, *Coord. Chem. Rev.*, 2000, **203**, 5; B. Pinter, V. Van Speybroeck, M. Waroquier, P. Geerlings and F. De Proft, *Phys. Chem. Chem. Phys.*, 2013, **15**, 17354.
- 13 R. M. Drost, V. Rosar, S. Dalla Marta, M. Lutz, N. Demitri, B. Milani, B. de Bruin and C. J. Elsevier, *Chem. Cat. Chem.*, 2015, **7**, 2095; N. T. S. Phan, M. Van Der Sluys and C. W. Jones, *Adv. Synth. Catal.*, 2006, **348**, 609.
- 14 E. D. Slack, C. M. Gabriel and B. H. Lipshutz, *Angew. Chem. Int. Ed.*, 2014, **53**, 14051.



282x140mm (72 x 72 DPI)

For Table of Contents

Zwitterionic all-metal aromatic tri-Palladium complexes can catalyse the *cis*-selective semi-reduction of internal alkynes at ppm levels