



Cite this: *Chem. Sci.*, 2019, 10, 3413

All publication charges for this article have been paid for by the Royal Society of Chemistry

5,10-Dimesityldiindeno[1,2-*a*:2',1'-*i*]phenanthrene: a stable biradicaloid derived from Chichibabin's hydrocarbon†

Marcin A. Majewski,^a Piotr J. Chmielewski,^a Alan Chien,^b Yongseok Hong,^c Tadeusz Lis,^a Maciej Witwicki,^a Dongho Kim,^a Paul M. Zimmerman^{*b} and Marcin Stępień^{ID *a}

A diindenophenanthrene biradicaloid, formally derived from Chichibabin's hydrocarbon, is obtained in a short, scalable synthesis. The present system is electron-rich and devoid of conjugated substituents, and still exhibits very good stability under ambient conditions. The introduction of the diindeno[1,2-*a*:2',1'-*i*] phenanthrene ring framework results in a singlet biradicaloid system with an easily accessible triplet state ($\Delta E_{S-T} = -1.30$ kcal mol⁻¹) and a small electronic bandgap (1.39 V). The stability limits of the title hydrocarbon were explored systematically in the solid state, to reveal an unusual thermally initiated hydrogen-scrambling oligomerization process.

Received 11th January 2019
Accepted 5th February 2019


DOI: 10.1039/c9sc00170k

rsc.li/chemical-science

Introduction

The use of proaromatic motifs is a commonly applied strategy to create open-shell π -conjugated organic molecules.¹ In particular, quinoidal structures, which regain benzenoid aromaticity upon formal unpairing of two electrons, have been explored in a variety of molecular settings.² A classical example of a quinoidal biradicaloid is found in Chichibabin's hydrocarbon (**1**, Scheme 1),^{3,4} which recovers two Clar sextets in its open-shell resonance form. Simple biradicaloid structures, such as **1** itself,⁴ tend to be unstable under ambient conditions, and extensive effort has been given to developing chemically robust open-shell systems. The diindeno[1,2-*b*:1',2'-*i*]anthracene (DIAn) derivative **3**, reported recently by Haley *et al.*^{5,6} exemplifies key strategies used to stabilize π -biradicaloids, namely extended ring fusion, steric blocking, and the use of conjugated substituents, such as ethynyl groups. **3** represents the family of stabilized diindenoarene biradicaloids,⁷⁻⁹ which comprises indenofluorenes¹⁰⁻¹⁴ and their larger congeners, *e.g.* ones containing naphthalene,¹⁵⁻¹⁸ pyrene,¹⁹ perylene,²⁰ and bischrysenes²¹ cores, or extended by benzenoid fusion.^{9,22-24}

Here we show that by a judicious yet simple modification of the classical structure of Chichibabin's hydrocarbon **1**, it is possible to create a much more stable biradicaloid system. This effect is achieved by (a) fusion of an extra benzene ring in the



Scheme 1 5,10-Dimesityldiindeno[1,2-*a*:2',1'-*i*]phenanthrene (**2**) and its structural relationship to Chichibabin's hydrocarbon (**1**) and Haley's diindenoanthracene (**3**). Mes = mesityl, TIPS = tri(isopropyl)silyl.

^aWydział Chemii, Uniwersytet Wrocławski, ul. F. Joliot-Curie 14, 50-383 Wrocław, Poland. E-mail: marcin.stepien@chem.uni.wroc.pl; Web: <http://www.mstepien.edu.pl>

^bDepartment of Chemistry, University of Michigan, 930 N. University Ave, Ann Arbor, MI 48109, USA. E-mail: paulzim@umich.edu

^cDepartment of Chemistry, Yonsei University, 50 Yonsei-ro, Seoul 120-749, Korea. E-mail: dongho@yonsei.ac.kr

† Electronic supplementary information (ESI) available: Experimental procedures, spectra of new compounds, and details of computational studies. CCDC 1861241. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc00170k



biphenyl section of **1**, (b) pentannulation of two phenyl substituents, and (c) steric blocking of the remaining two phenyl groups in the form of bulky mesityl substituents. The resulting molecule, 5,10-dimesityldiindeno[1,2-*a*:2',1'-*f*]phenanthrene **2** features a fused ring system (DIPh) isomeric to the DIAn framework of **3**. Advantageously, **2** achieves ambient stability in the absence of bulky TIPS-acetylene substituents. This structural simplification results in a pure hydrocarbon system with a significantly lower molecular weight. The type of pentannulation introduced in **2** is also more effective than earlier modifications of **1** in which adjacent phenyl groups were fused to form terminal fluorene subunits. Those systems would either spontaneously polymerize^{25–27} or required multiple benzo fusion to achieve ambient stability.²⁸

Results and discussion

Compound **2** was obtained in three steps from the previously reported 1,8-dibromophenanthrene-2,7-dicarbaldehyde **4** (ref. 29) (Scheme 2), and isolated as an air-stable dark-blue crystalline powder. The overall route from commercially available materials (six steps) is efficient and high-yielding: **2** is easily obtained in quantities exceeding 100 mg. In comparison with **3**, the absence of TMS-ethynyl substituents in **2** makes the present design structurally simpler, furnishing the DIPh ring system as a heteroatom-free hydrocarbon derivative. The use of the phenanthrene ring system for elaboration of open-shell aromatics has so far been limited, with a single example provided by the recently reported [4]chrysaorene, a fully conjugated coronoid ring system with a tetraradicaloid character.^{30,31} Diffraction-quality crystals of **2** were grown without difficulty, and the structure of the new hydrocarbon was unambiguously confirmed using X-ray crystallographic analysis.

In the solid state, the diindenophenanthrene core of **2** is found to deviate slightly from planarity, apparently as a result of steric repulsions in the two cove-like regions of the fused ring system (Fig. 1). DFT-optimized geometries of **2** are however planar, indicating that the experimentally observed distortion might be promoted by packing forces in the crystal. No π -





Fig. 1 Molecular structure of **2** determined using X-ray crystallographic analysis. An *n*-hexane solvent molecule and hydrogen atoms are omitted for clarity.

stacking of DIPh cores was observed in the solid-state structure, in line with the bulky Mes substitution of the fused ring system. The “quinomethane” bonds in **2** (1.395(4) and 1.408(4) Å) are longer than expected of formally double C=C bonds, a feature that correlates with the increasing biradicaloid character of diindenoarenes.⁷ The above bond lengths are similar to the corresponding distances determined for **3** (ref. 5) and **1** (ref. 4) (1.406 and 1.415(3) Å, respectively), in line with the presumed biradicaloid character of **2**.

In the electronic spectrum of **2**, the most intense band (λ_{\max} = 600 nm, ϵ = 4.1×10^4 M⁻¹ cm⁻¹, Fig. 2) is responsible for the



Scheme 2 Synthesis of 5,10-dimesityldiindeno[1,2-*a*:2',1'-*f*]phenanthrene (**2**). Reagents and conditions: (a) PhB(OH)₂, Pd(PPh₃)₄, Na₂CO₃ (aq), dioxane, reflux, overnight. (b) (1) MesMgBr, THF, rt, overnight. (2) BF₃·Et₂O, DCM, rt, 10 min. (c) 2.5 equiv. DDQ, toluene, rt, overnight.




Fig. 2 (Top) Electronic absorption spectrum of **2** (dichloromethane, ca. 2×10^{-5} M, blue trace) and **7** (dichloromethane, green traces). The photo shows the appearance of **2** in a dichloromethane solution. (Bottom) Formation of the radical cation **2**^{•+} (purple) upon oxidation with magic blue. Intermediate titration stages are shown in gray.



deep blue color of the compound, and appears at a longer wavelength than that reported for **1** ($\lambda_{\max} = 574 \text{ nm}$).⁴ While the main absorption in **2** is hypsochromically shifted relative to the corresponding band of **3** (690 nm), the absorption edge is observed at longer wavelengths (1050 nm *vs. ca.* 900 nm for **3**), indicating a reduced optical bandgap of **2**. According to a broken-symmetry TD-DFT calculation (ESI[†]), the first transition, dominated by the HOMO–LUMO excitation, is dipole-forbidden, in accord with the weak intensity of the corresponding experimental absorption.

In cyclic voltammetry measurements (Fig. 3), **2** undergoes three one-electron oxidations (0.26, 0.66, and 1.11 V *vs.* Fc^+/Fc), of which only the first one is fully reversible. Good reversibility was however observed for two reductions (−1.13 and −1.42 V). All of the above redox events are shifted by up to 0.15 V to higher potentials relative to those determined for **3**. The electrochemical bandgap of 1.39 V determined for **2** is smaller than the already uniquely narrow gap of **3** (1.45 V). Quinoidal molecules containing isomeric ring fusion-patterns often yield large variations of optical signatures and biradicaloid characters.^{12,14,32} In particular, the number of Clar sextets in the open-shell valence structure of DIPh is increased relative to DIAn (four in the open-shell formula **2'** *vs.* three in **3'**, *cf.* Scheme 1), and such a change has been demonstrated to enhance open-shell characters of hydrocarbon biradicaloids.³³ In the present case, the bandgap change is likely induced not only by the different topologies of the ring systems in **2** and **3**, but also by the difference in substitution patterns. TIPS-ethynyl groups were previously shown to reduce the optical bandgap of pentacene by 0.27 eV, an effect attributed to disproportionate lowering of the LUMO energy.³⁴ The influence of substituents on the bandgap of **3** has not been investigated experimentally; however, if the response of **3** to ethynyl substitution is qualitatively similar to that observed in pentacene, the bandgap-reducing effect of replacing anthracene with phenanthrene in **2** would be even greater than indicated by electrochemistry data.

At 220 K, **2** produces a sharp ¹H NMR spectrum in the CDCl_3 solution, enabling unambiguous assignment of all signals




Fig. 3 Cyclic voltammetry (top) and differential pulse voltammetry scans (bottom) obtained for **2** (dichloromethane, tetrabutylammonium hexafluorophosphate as a supporting electrolyte).

(Fig. 4). When the sample was heated to 360 K, the spectrum showed progressive yet reversible line broadening, accompanied by shortening of T_1 relaxation times, observed already at 300 K (Fig. S10 and S11[†]). These spectral changes were indicative of chemical exchange with a paramagnetic species, which was presumed to be the thermally populated triplet state of **2**, in analogy to the reported behavior of **3** and related systems.

Solid-state samples of **2** yielded a well-defined ESR line ($g = 2.00248$), however, without resolved triplet components. The lack of fine structure is apparently consistent with the relatively large distance between the two quinomethane carbons in **2** (9.84 Å *vs.* 10.32 Å in **3** (ref. 5) and 10.05 Å in **1** (ref. 4)). In the point-dipole approximation,³⁵ this distance gives a value of the zero-field splitting (ZFS) parameter $|D| = 29 \text{ G}$ (0.003 cm^{-1}). This estimate ignores electron delocalization^{36,37} and can be taken as the lower bound of the actual ZFS interaction. DFT calculations performed for **2** at various levels of theory produced somewhat variable, but noticeably higher estimates of D (50 to 100 G, 0.005 to 0.009 cm^{-1} , see ESI[†] for computational details). These higher values reflect the effect of spin delocalization but nevertheless suggest that the ZFS in **2** may indeed be at the limit of measurability.³⁸ Changes of the doubly integrated ESR signal intensity recorded in the 143–333 K range could be fitted with the Bleaney–Bowers equation to provide a singlet–triplet gap of $\Delta E_{S-T} = -1.30 \text{ kcal mol}^{-1}$. The measurement of bulk magnetic susceptibility using the Evans method (for a toluene- d_8 solution), showed a similar increase of the triplet contribution, yielding a ΔE_{S-T} value of $-1.19 \text{ kcal mol}^{-1}$, respectively. It was however observed that prolonged heating of the samples above 340 K resulted in further non-reversible increase of the magnetism, which was linked to the thermally induced reactivity of **2**, discussed below.

2 can be chemically oxidized to a stable radical cation $\mathbf{2}^+$ using $[\text{N}(p\text{-C}_6\text{H}_4\text{Br})_3][\text{SbCl}_6]$ (magic blue, Scheme 3, Fig. 2 and





Fig. 4 Temperature-dependent ¹H NMR spectra of **2** (CDCl_3 , 600 MHz). The signals were assigned at 220 K using 2D spectroscopic methods (see the ESI[†]).



S20†). Further oxidation was observable when a stronger oxidant, (NO)[SbF₆], was employed, but the resulting product was not stable. When treated with an excess of sodium metal in THF-*d*₈, **2** cleanly produced the brownish dianion 2²⁻,^{6,14} which was extremely unstable to traces of moisture and oxygen. The dianion yielded clearly defined, diamagnetic ¹H and ¹³C NMR spectra, which were analyzed in detail with the help of 2D correlation methods (Fig. 5, S13 and S14†). The ¹H resonance pattern of 2²⁻ is characterized by significant downfield relocations of the a, b, and c resonances relative to their positions in the spectrum of **2**, suggesting that the aromaticity of the phenanthrene core is restored in the dianion. ¹³C NMR signals of quinomethane carbons in 2²⁻ were unambiguously identified at 99.0 ppm on the basis of ¹H-¹³C HMBIC correlations. This upfield shift, reproduced in a GIAO calculation (95.4 ppm), is characteristic of π-conjugated carbanions,³⁹ and provides strong evidence for the charged nature of 2²⁻. Interestingly, the bond length pattern obtained for 2²⁻ in a DFT calculation is very similar to that observed experimentally in the fluorenyl anion,⁴⁰ indicating a possible similarity of the electronic structures of these two systems. When traces of water were present during the Na-induced reduction of **2**, the protonated monoanion (2-H)⁻ was obtained instead of 2²⁻. This new species, identified *in situ* on the basis of its 1D and 2D ¹H NMR spectra (Fig. S12†), could be partly converted back to the neutral **2** by exposure to air.

Femtosecond transient absorption (fs-TA) measurements performed for **2** (Fig. 6A) revealed ultrafast relaxation dynamics, similar to those observed in other quinodimethane biradicaloids.^{7,8} Evolution-associated spectra, obtained by global analysis (Fig. 6B and S18†), showed an initial decay with a time constant τ₁ = 200 ± 20 fs corresponding to an internal conversion to the S₁ state, followed by rapid deactivation to the ground state (τ₂ = 7.0 ± 0.7 ps), possibly admixed with vibrational cooling. The short excited-state lifetime of **2** is comparable with those of other diindenoarenes,⁴¹ and may be similarly linked to non-radiative decay *via* an S₁/S₀ conical intersection.



Scheme 3 Redox reactivity of **2**. Reagents and conditions: (a) [N(*p*-C₆H₄Br)₃][SbCl₆] (magic blue). (b) Na metal, THF-*d*₈. (c) Na metal, THF-*d*₈, traces of moisture. (d) Air (partial recovery).




Fig. 5 ¹H NMR spectrum (THF-*d*₈, 600 MHz, 300 K, bottom) and DFT-optimized geometry of 2²⁻ (PCM(THF)/B3LYP/6-31G(d,p), top).

The biradicaloid nature of **2** was explored computationally using the spin-flip approach.⁴² A RAS(2,2)-SF/cc-pVTZ calculation^{43,44} was performed on the substituent-free DIPh core of **2** optimized at the CAM-B3LYP/6-311++G(d,p) level of theory. A value of the Δ*E*_{S-T} gap of -1.43 kcal mol⁻¹ was obtained, in remarkably good agreement with the results of ESR and Evans measurements. The strong biradicaloid character of the singlet state is indicated by the RAS-SF natural orbital occupancies, which gave a diradical index γ₀ of 0.69. In both the S and T states, the odd-electron density, obtained from the RAS-SF calculation, is localized principally at the five-membered rings (Fig. 7B), in line with the open-shell configuration 2'. The validity of 2' was further verified using 1D and 2D NICS scans performed on the DIPh core (Fig. 7A and S17†), which showed that the aromaticity of the central phenanthrene unit increases in the triplet state. In the singlet state, the terminal six-membered rings of DIPh remain aromatic, whereas the five-membered rings show a weak antiaromatic ring current. This





Fig. 6 fs-TA spectra (A) and evolution-associated spectra (B) of **2** in THF with the excitation of 600 nm.





Scheme 5 Potential oligomerization pathways for **2**. Gibbs free energies (red labels, kcal mol⁻¹, UCAM-B3LYP/6-31G(d,p), gas phase, 298 K) are given relative to S-2 + S-2, unless indicated otherwise. S, D, and T indicate respectively the broken-symmetry-singlet, doublet and triplet wavefunctions. *r* and *m* indicate the *racemic* pair and the *meso* isomer, respectively.

Looking for a more feasible pathway, we considered addition of the neutral **2** to the radical cation **2**^{•+}, which is expected to form in small amounts in bulk samples of **2**. Such an addition would produce the radical-cation intermediate **8**^{•+} (Scheme 5), which yields broken-symmetry geometries characteristic of a Robin–Day class-I or -II mixed-valence (MV) system.⁵³ The calculated barrier for this process ($\Delta G^\ddagger = 31.82$ kcal mol⁻¹, doublet hypersurface) is significantly lower than those obtained for dimerization of the neutral **2**. While the value is still relatively high, it does not bear a quantitative significance because the accuracy of the calculation is potentially limited by the MV character of the system,⁵³ the neglect of solvation and dispersion phenomena, and the choice of the standard state.

The calculated ΔG value for the electron transfer process **8**^{•+} + **2** → **2**^{•+} + **8** are low (0 to 2 kcal mol⁻¹) indicating that in principle, the electron can be transferred between chains of various lengths. The oligomerization can thus propagate *via* a sequence of hydrogen, proton, and electron transfer steps. Importantly, while the initial adduct **8** is high in energy, the formation of its rearranged isomers, such as **9a–b**, which contain hydrogenated 5-membered rings, is predicted to be significantly exergonic. A similar stabilization is predicted for **9a**^{•+} (ΔG of ca. -25 kcal mol⁻¹ vs. S-2 + 2^{•+}). The overall reactivity described above would be distinct from reported cases of biradicaloid oligo- and cyclooligomerizations, in which the

addition typically involves sites with highest spin densities, sometimes relatively hindered,^{26,27,52,54–56} or conjugated acetylene substituents.⁵⁷ DFT shows that, while the initial barrier to dimerization may be relatively high, the proposed oligomerization is a strongly exergonic process, and should be feasible for neat **2** at high temperatures. Further growth of the oligomer chain can produce a variety of species containing random sequences of closed-shell and open-shell sections. It is worth noting that in such sequences, terminal fluorenyl radicals may form (as seen in the idealized oligomer **11**), which can be expected to produce a doublet ESR spectrum, qualitatively similar to that observed for **2** at the beginning of its decomposition (*vide supra*).

Conclusions

We have reported here a simple analogue of Chichibabin's hydrocarbon, characterized by very good ambient stability. The present design is simpler than those explored previously,^{28,58–61} and achieves the stabilizing effect without increasing the electronic bandgap or introduction of heteroatoms. The hydrogen-scrambling oligomerization, described herein, is relevant as a prospective method of synthesizing open-shell polymeric materials, but it also provides a guiding principle for designing stable electron-rich biradicaloids.



Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Financial support from the National Science Center of Poland (DEC-2012/07/E/ST5/00781 to M. S. and DEC-2015/19/N/ST5/00760 to M. A. M.) and the Foundation for Polish Science (START fellowship to M. A. M.) is gratefully acknowledged. Quantum chemical calculations were performed in the Wrocław Center for Networking and Supercomputing. P. M. Z. thanks the Alfred P. Sloan Foundation for support of this project. The work at Yonsei University was supported by Strategic Research (NRF-2016R1E1A1A01943379) through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT (Information and Communication Technologies) and Future Planning. We thank Prof. Frank Würthner (Würzburg), Dr Yi-Lin Wu (Northwestern University), and anonymous referees for helpful comments.

Notes and references

- Z. Zeng, X. Shi, C. Chi, J. T. L. Navarrete, J. Casado and J. Wu, *Chem. Soc. Rev.*, 2015, **44**, 6578–6596.
- J. Casado, *Top. Curr. Chem.*, 2017, **375**, 73.
- A. E. Tschitschibabin, *Ber. Dtsch. Chem. Ges.*, 1907, **40**, 1810–1819.
- L. K. Montgomery, J. C. Huffman, E. A. Jurczak and M. P. Grendze, *J. Am. Chem. Soc.*, 1986, **108**, 6004–6011.
- G. E. Rudebusch, J. L. Zafra, K. Jorner, K. Fukuda, J. L. Marshall, I. Arrechea-Marcos, G. L. Espejo, R. Ponce Ortiz, C. J. Gómez-García, L. N. Zakharov, M. Nakano, H. Ottosson, J. Casado and M. M. Haley, *Nat. Chem.*, 2016, **8**, 753–759.
- G. E. Rudebusch, G. L. Espejo, J. L. Zafra, M. Peña-Alvarez, S. N. Spisak, K. Fukuda, Z. Wei, M. Nakano, M. A. Petrukhina, J. Casado and M. M. Haley, *J. Am. Chem. Soc.*, 2016, **138**, 12648–12654.
- C. K. Frederickson, B. D. Rose and M. M. Haley, *Acc. Chem. Res.*, 2017, **50**, 977–987.
- Y. Tobe, *Top. Curr. Chem.*, 2018, **376**, 12.
- T. Kubo, *Chem. Lett.*, 2015, **44**, 111–122.
- D. T. Chase, B. D. Rose, S. P. McClintock, L. N. Zakharov and M. M. Haley, *Angew. Chem., Int. Ed.*, 2011, **50**, 1127–1130.
- A. Shimizu and Y. Tobe, *Angew. Chem., Int. Ed.*, 2011, **50**, 6906–6910.
- A. Shimizu, R. Kishi, M. Nakano, D. Shiomi, K. Sato, T. Takui, I. Hisaki, M. Miyata and Y. Tobe, *Angew. Chem., Int. Ed.*, 2013, **52**, 6076–6079.
- A. G. Fix, P. E. Deal, C. L. Vonnegut, B. D. Rose, L. N. Zakharov and M. M. Haley, *Org. Lett.*, 2013, **15**, 1362–1365.
- J. J. Dressler, Z. Zhou, J. L. Marshall, R. Kishi, S. Takamuku, Z. Wei, S. N. Spisak, M. Nakano, M. A. Petrukhina and M. M. Haley, *Angew. Chem., Int. Ed.*, 2017, **56**, 15363–15367.
- B. D. Rose, C. L. Vonnegut, L. N. Zakharov and M. M. Haley, *Org. Lett.*, 2012, **14**, 2426–2429.
- H. Miyoshi, S. Nobusue, A. Shimizu, I. Hisaki, M. Miyata and Y. Tobe, *Chem. Sci.*, 2013, **5**, 163–168.
- J. E. Barker, C. K. Frederickson, M. H. Jones, L. N. Zakharov and M. M. Haley, *Org. Lett.*, 2017, **19**, 5312–5315.
- H. Miyoshi, M. Miki, S. Hirano, A. Shimizu, R. Kishi, K. Fukuda, D. Shiomi, K. Sato, T. Takui, I. Hisaki, M. Nakano and Y. Tobe, *J. Org. Chem.*, 2017, **82**, 1380–1388.
- D. Hibi, K. Kitabayashi, K. Fujita, T. Takeda and Y. Tobe, *J. Org. Chem.*, 2016, **81**, 3735–3743.
- K. Sbagoud, M. Mamada, J. Marrot, S. Tokito, A. Yassar and M. Frigoli, *Chem. Sci.*, 2015, **6**, 3402–3409.
- J. Ma, J. Liu, M. Baumgarten, Y. Fu, Y.-Z. Tan, K. S. Schellhammer, F. Ortmann, G. Cuniberti, H. Komber, R. Berger, K. Müllen and X. Feng, *Angew. Chem., Int. Ed.*, 2017, **56**, 3280–3284.
- C. K. Frederickson, L. N. Zakharov and M. M. Haley, *J. Am. Chem. Soc.*, 2016, **138**, 16827–16838.
- J. Liu, J. Ma, K. Zhang, P. Ravat, P. Machata, S. Avdoshenko, F. Hennersdorf, H. Komber, W. Pisula, J. J. Weigand, A. A. Popov, R. Berger, K. Müllen and X. Feng, *J. Am. Chem. Soc.*, 2017, **139**, 7513–7521.
- J. Melidonie, J. Liu, Y. Fu, J. J. Weigand, R. Berger and X. Feng, *J. Org. Chem.*, 2018, **83**, 6633–6639.
- W. Theilacker, H. Schulz, U. Baumgarte, H.-G. Drössler, W. Rohde, F. Thater and H. Uffmann, *Angew. Chem.*, 1957, **69**, 322–333.
- J. Ipaktschi, R. Hosseinzadeh and P. Schlaf, *Angew. Chem., Int. Ed.*, 1999, **38**, 1658–1660.
- D. Beaudoin, O. Levasseur-Grenon, T. Maris and J. D. Wuest, *Angew. Chem., Int. Ed.*, 2016, **55**, 894–898.
- Z. Zeng, Y. M. Sung, N. Bao, D. Tan, R. Lee, J. L. Zafra, B. S. Lee, M. Ishida, J. Ding, J. T. López Navarrete, Y. Li, W. Zeng, D. Kim, K.-W. Huang, R. D. Webster, J. Casado and J. Wu, *J. Am. Chem. Soc.*, 2012, **134**, 14513–14525.
- M. Shimizu, Y. Tomioka, I. Nagao, T. Kadowaki and T. Hiyama, *Chem.-Asian J.*, 2012, **7**, 1644–1651.
- X. Lu, T. Y. Gopalakrishna, H. Phan, T. S. Heng, Q. Jiang, C. Liu, G. Li, J. Ding and J. Wu, *Angew. Chem., Int. Ed.*, 2018, **57**, 13052–13056.
- H. Gregolińska, M. Majewski, P. J. Chmielewski, J. Gregoliński, A. Chien, J. Zhou, Y.-L. Wu, Y. J. Bae, M. R. Wasielewski, P. M. Zimmerman and M. Stępień, *J. Am. Chem. Soc.*, 2018, **140**, 14474–14480.
- A. Konishi, Y. Okada, M. Nakano, K. Sugisaki, K. Sato, T. Takui and M. Yasuda, *J. Am. Chem. Soc.*, 2017, **139**, 15284–15287.
- Z. Sun, S. Lee, K. H. Park, X. Zhu, W. Zhang, B. Zheng, P. Hu, Z. Zeng, S. Das, Y. Li, C. Chi, R.-W. Li, K.-W. Huang, J. Ding, D. Kim and J. Wu, *J. Am. Chem. Soc.*, 2013, **135**, 18229–18236.
- I. Kaur, W. Jia, R. P. Koproški, S. Selvarasah, M. R. Dokmeci, C. Pramanik, N. E. McGruer and G. P. Miller, *J. Am. Chem. Soc.*, 2008, **130**, 16274–16286.
- M. Abe, *Chem. Rev.*, 2013, **113**, 7011–7088.
- P. Bertrand, C. More, B. Guigliarelli, A. Fournel, B. Bennett and B. Howes, *J. Am. Chem. Soc.*, 1994, **116**, 3078–3086.



- 37 C. Riplinger, J. P. Y. Kao, G. M. Rosen, V. Kathirvelu, G. R. Eaton, S. S. Eaton, A. Kutateladze and F. Neese, *J. Am. Chem. Soc.*, 2009, **131**, 10092–10106.
- 38 C. Wenstrup, M. J. Regimbald-Krnel, D. Müller and P. Comba, *Angew. Chem., Int. Ed.*, 2016, **55**, 14600–14605.
- 39 K. Müllen, *Chem. Rev.*, 1984, **84**, 603–646.
- 40 S. Filippini, J. N. Jones, J. A. Johnson, A. H. Cowley, F. Grepioni and D. Braga, *Chem. Commun.*, 2003, 2716–2717.
- 41 B. D. Rose, L. E. Shoer, M. R. Wasielewski and M. M. Haley, *Chem. Phys. Lett.*, 2014, **616–617**, 137–141.
- 42 A. I. Krylov, *Chem. Phys. Lett.*, 2001, **338**, 375–384.
- 43 P. M. Zimmerman, F. Bell, M. Goldey, A. T. Bell and M. Head-Gordon, *J. Chem. Phys.*, 2012, **137**, 164110.
- 44 F. Bell, P. M. Zimmerman, D. Casanova, M. Goldey and M. Head-Gordon, *Phys. Chem. Chem. Phys.*, 2013, **15**, 358–366.
- 45 H. Sakurai, H. Tobita, M. Kira and Y. Nakadaira, *Angew. Chem., Int. Ed. Engl.*, 1980, **19**, 620.
- 46 A. Kostenko, B. Tumanskii, M. Karni, S. Inoue, M. Ichinohe, A. Sekiguchi and Y. Apeloig, *Angew. Chem., Int. Ed.*, 2015, **54**, 12144–12148.
- 47 J. J. Dressler, M. Teraoka, G. L. Espejo, R. Kishi, S. Takamuku, C. J. Gómez-García, L. N. Zakharov, M. Nakano, J. Casado and M. M. Haley, *Nat. Chem.*, 2018, **10**, 1134–1140.
- 48 J. L. Zafra, L. Qiu, N. Yanai, T. Mori, M. Nakano, M. P. Alvarez, J. T. L. Navarrete, C. J. Gómez-García, M. Kertesz, K. Takimiya and J. Casado, *Angew. Chem., Int. Ed.*, 2016, **55**, 14563–14568.
- 49 K. Uchida, S. Ito, M. Nakano, M. Abe and T. Kubo, *J. Am. Chem. Soc.*, 2016, **138**, 2399–2410.
- 50 P. Ravat, T. Šolomek, M. Rickhaus, D. Häussinger, M. Neuburger, M. Baumgarten and M. Juríček, *Angew. Chem., Int. Ed.*, 2016, **55**, 1183–1186.
- 51 T. Šolomek, P. Ravat, Z. Mou, M. Kertesz and M. Juríček, *J. Org. Chem.*, 2018, **83**, 4769–4774.
- 52 A. Kikuchi, F. Iwahori and J. Abe, *J. Am. Chem. Soc.*, 2004, **126**, 6526–6527.
- 53 C. Sutton, T. Körzdörfer, V. Coropceanu and J.-L. Brédas, *J. Phys. Chem. C*, 2014, **118**, 3925–3934.
- 54 T. Itoh, *Prog. Polym. Sci.*, 2001, **26**, 1019–1059.
- 55 W. S. Trahanovsky and S. P. Lorimor, *J. Org. Chem.*, 2006, **71**, 1784–1794.
- 56 J. Inoue, K. Fukui, T. Kubo, S. Nakazawa, K. Sato, D. Shiomi, Y. Morita, K. Yamamoto, T. Takui and K. Nakasuji, *J. Am. Chem. Soc.*, 2001, **123**, 12702–12703.
- 57 X. Fu and D. Zhao, *Org. Lett.*, 2015, **17**, 5694–5697.
- 58 H. Kataoka and M. Nakagawa, *Bull. Chem. Soc. Jpn.*, 1963, **36**, 799–806.
- 59 T. Kawase, N. Ueno and M. Oda, *Tetrahedron Lett.*, 1992, **33**, 5405–5408.
- 60 A. Lichtblau, H. D. Hausen, W. Schwarz and W. Kaim, *Inorg. Chem.*, 1993, **32**, 73–78.
- 61 C. Jiang, Y. Bang, X. Wang, X. Lu, Z. Lim, H. Wei, S. El-Hankari, J. Wu and Z. Zeng, *Chem. Commun.*, 2018, **54**, 2389–2392.

