



Cite this: *Org. Biomol. Chem.*, 2019, **17**, 1206

Silent catalytic promiscuity in the high-fidelity terpene cyclase δ -cadinene synthase†

Marianna Loizzi, David J. Miller  and Rudolf K. Allemann  *

δ -Cadinene synthase (DCS) is a high-fidelity sesquiterpene synthase that generates δ -cadinene as the sole detectable organic product from its natural substrate (*E,E*)-FDP. Previous work with this enzyme using substrate analogues revealed the ability of DCS to catalyse both 1,10- and 1,6-cyclisations of substrate analogues. To test whether this apparent promiscuity was an artefact of alternate substrate use or an inherent property of the enzyme, aza analogues of the proposed α -bisaboyl cation intermediate were prepared since this cation would be formed after an initial 1,6-cyclisation of FDP. In the presence of 250 μ M inorganic diphosphate both (*R*)- and (*S*)-aza-bisaboyl cations were potent competitive inhibitors of DCS ($K_i = 2.5 \pm 0.5$ mM and 3.44 ± 1.43 μ M, respectively). These compounds were also shown to be potent inhibitors of the 1,6-cyclase amorpho-4,11-diene synthase but not of the 1,10-cyclase aristolochene synthase from *Penicillium roquefortii*, demonstrating that the 1,6-cyclase activity of DCS is most likely an inherent property of the enzyme even when the natural substrate is used and not an artefact of the use of substrate analogues.

Received 13th November 2018,
Accepted 10th December 2018

DOI: 10.1039/c8ob02821d

rs.c.li/obc

Introduction

Terpene synthases catalyse some of the most complex reactions in the natural world. From a small pool of isoprenyl diphosphates they generate a myriad of hydrocarbons and alcohols that are often processed into thousands of terpenoids with diverse biological activities with many potential applications for instance as agrochemicals or therapeutic agents.¹

The details of terpene synthase chemistry² have been investigated by site directed mutagenesis and with non-natural amino acids,³ analogues of substrates,⁴ and putative reaction intermediates,⁵ X-ray crystallography,^{2b,c,6} and computational modelling.^{2d,7} Together these investigations revealed a fascinating, yet still incomplete picture. A series of X-ray crystal structure of aristolochene synthase from *Aspergillus terreus* in both closed and open conformations along with complexes containing the complete substrate (or analogue), diphosphate anion and/or Mg^{2+} co-factors⁸ revealed the physical steps of the catalytic cycle. Binding of a Mg^{2+} -ion is followed by coordination of the prenyl diphosphate substrate and a second Mg^{2+} ion; coordination of a third Mg^{2+} ion triggers active site closure to form the Michaelis complex.⁸ Diphosphate cleavage is then triggered to form an initial carbocation and the hydro-

phobic active site shelters this high energy intermediate from bulk solvent.^{2b,c} The active site, lined with hydrophobic and aromatic amino acid residues then steers the initial carbocation through a series of ring closures and rearrangements prior to quench of the final carbocation either by proton loss or nucleophilic attack by water.^{2b,c,6e} Usually this is tightly controlled by the enzyme, with a single enantiomer dominating the product pool whereby several rings and stereocentres are often generated in a single chemical step from an achiral precursor. Control of this process is thought to arise from a product-like active site contour in combination with direction of carbocation location in the intermediates through the negative charge on the diphosphate anion and aromatic amino acid side chains that can stabilise carbocations at certain locations through cation- π interaction.^{2b,c,4a} A small subset of terpene synthases, on the other hand, exhibit significant promiscuity, presumably through having a less structured and/or flexible active site that allows the intermediates to sample a large number of reactive conformations prior to final carbocation quench. For example, δ -selinine synthase and γ -humulene synthases from *Abies grandis* generate 34 and 52 products from farnesyl diphosphate (**1**), respectively.⁹ Terpene synthases have been postulated to evolve through such promiscuous intermediates prior to further evolution into high-fidelity synthases.^{3e} The modern δ -cadinene synthase (DCS) from *Gossypium arboreum* is a high-fidelity sesquiterpene synthase that catalyses the formation of the bicyclic hydrocarbon (+)- δ -cadinene (**7**),¹⁰ the first committed step in the biosynthesis of the phytoalexin gossypol.¹¹ The catalytic domain

School of Chemistry, Cardiff University Main Building, Park Place, Cardiff, CF10 3AT, UK. E-mail: allemannrk@cardiff.ac.uk

† Electronic supplementary information (ESI) available: General experimental procedures, enzyme preparation and purification, kinetics data, gas chromatograms, mass spectra and NMR spectra. See DOI: 10.1039/c8ob02821d



is situated in the C-terminal domain and adopts the α -helical fold domain, typical of class 1 terpene synthases.^{2b,c,12} It contains the conserved aspartate rich motif D³⁰⁷DTYD³¹¹ on helix D, but instead of the usual characteristic NSE/DTE Mg²⁺ binding motif, DCS has a second aspartate rich motif D⁴⁵¹DVAE⁴⁵⁵ on helix H.^{6e} Despite only generating a single detectable hydrocarbon product, extensive mechanistic analysis of the DCS-catalysed reaction pathway has not unambiguously defined the chemical steps of its catalytic cycle. Moreover, conversion of fluorinated and stereochemically altered FDP analogues with DCS revealed an underlying mechanistic promiscuity with products arising from an initial 1,10-, 1,6- or 1,11-ring closure depending upon the substrate analogue used (*vide infra*).¹² Two chemical mechanisms remain plausible for the formation of δ -cadinene from FDP (Scheme 1). Both pathways involve initial formation of (3*R*)-nerolidyl diphosphate ((3*R*)-NDP, (2)) as an enzyme-bound intermediate. In pathway (a), a 1,10-macrocyclisation occurs to generate *cis*-germacradienyl cation (4). A subsequent [1,3]-hydride shift is followed by a 1,6-electrophilic ring closure to cadinenyl cation (6), from which δ -cadinene (7) is formed after proton loss from C6. In pathway (b), a 1,6 ring-closure of 2 is followed by a [1,3]-hydride shift from C1 to C7; subsequently a second ring closure and a [1,5]-hydride shift lead to cadinenyl

cation, an intermediate common to both pathways. In previous work, using substrate analogues we were unable to definitively rule out pathway (b) and indeed when 6-fluorofarnesyl diphosphate (6F-FDP) was used as a substrate analogue it proved to be a potent inhibitor ($K_i = 2.4 \mu\text{M}$), giving no detectable pentane-extractable products when incubated with DCS. This result is consistent with an initial 1,6-cyclisation pathway since it would be expected to undergo 1,10-ring closure and give an abortive product rather than inhibit the enzyme in the latter scenario. On the other hand, 2-fluorofarnesyl diphosphates (2F-FDP) and 10-fluorofarnesyl diphosphate (10F-FDP) gave products arising from 1,10- and 1,11 ring-closures, respectively, consistent with an initial 1,10-ring closure mechanism.¹²

Hence examination of the catalytic mechanism of DCS using FDP analogues has led to inconclusive, yet intriguing results, showing that this enzyme has the potential to use alternative reaction pathways. Yet the question arises, is this simply an artefact of the substrate used or is this an inherent property of the enzyme? The work described here provides alternative mechanistic data for the DCS-catalysed transformation of FDP to δ -cadinene using aza-analogues of putative carbocation intermediates. Although the highly unstable carbocationic intermediates formed during terpene synthase catalysis, cannot be isolated, it is possible to replace the sp^2 hybridised carbocationic carbon of a given intermediate with an sp^3 hybridised nitrogen in an amine analogue or with a sp^2 hybridised nitrogen in an iminium ion. Although the tetrahedral tertiary ammonium ions inherently are imperfect geometric analogues of the planar carbocations, these aza-terpenoids are thought to mimic the topological and electrostatic properties of carbocations generated by these enzymes.⁵ However, since they cannot be processed by the enzyme, they often act as tightly bound competitive inhibitors of terpene synthases.^{5c,13}

Hence, the use of strategically designed aza-analogues may enable the disentanglement of the possible reaction mechanisms catalysed by DCS. Here we report the stereoselective synthesis of the two enantiomers of aza-bisabolyl cation and their kinetic evaluation as inhibitors of DCS. Comparison of their effect upon catalysis by AS and amporpha-4,11-diene synthase (ADS), enzymes that follow 1,10- and 1,6-ring-closure mechanisms, validate the result that DCS has inherent 1,6- as well as 1,10 ring closure activity.

Results and discussion

If α -bisabolyl cation 8 is a reaction intermediate on the pathway to δ -cadinene (7), one or both of enantiomeric aza-analogues of 11 (Fig. 1) should act as competitive inhibitors of DCS.

Both enantiomers of 11 have previously been prepared.^{14a} Here we report an alternative synthesis that is more concise and avoids the use of harsh reaction conditions. Key to the synthesis of both enantiomers is an enantioselective synthesis of the two enantiomers of carboxylic acid 18 (Scheme 2). This was achieved through asymmetric Diels–Alder reaction of an



Scheme 1 Possible chemical steps for DCS catalysed production of δ -cadinene from FDP (1).






Fig. 1 Chemical structures of the α -bisabolyl cation (**8**) and corresponding aza analogues (*R*)-**11** and (*S*)-**11**.



Scheme 2 Synthesis of aza-analogues (*R*)- and (*S*)-**11**. Reagents and conditions: (i) acryloyl chloride, BuLi, THF, 35%. (ii) Et_2AlCl , CH_2Cl_2 , -100°C , 54%. (iii) Acryloyl chloride, NEt_3 , CH_2Cl_2 , 60%. (iv) TiCl_4 , CH_2Cl_2 , -10°C , 84%. (v) LiOH, THF, H_2O , MeOH, 50°C , 99%. (vi) DPPA, NEt_3 , (vii) *p*-Methoxybenzyl alcohol, toluene, 60% over two steps. (viii) LiAlH_4 , Et_2O , 50%, (ix) 4-methylpent-3-enoic acid, EtNPr_2 , HBTU, DMF, 70%. (x) LiAlH_4 , Et_2O . (xi) HCl in Et_2O , 55% two steps.

acrylate derivatised with a chiral auxiliary with a butadiene.¹⁵ Oxazolidinone **12** was alkylated with acryloyl chloride after deprotonation with *n*-butyl lithium with 35% yield. The resulting ester **13** was then subjected to an asymmetric Diels-Alder reaction with 2-methylbutadiene.¹⁵ The enantioselectivity and yield were optimal at -100°C in CH_2Cl_2 (52%, ee >95%, de >95% (see ESI† for details). This produced the key compound to generate the *S* enantiomer of the aza-analogue **11**. The equivalent *R* configured ester was generated using *D*-pantolactone (**15**) as a chiral auxiliary.¹⁵ After alkylation with acryloyl chloride, using NEt_3 as the base in CH_2Cl_2 , diester **16** was isolated in 60% yield. Again, an asymmetric Diels-Alder reaction with 2-methylbutadiene was carried out, this time at -10°C in CH_2Cl_2 using TiCl_4 as a Lewis acid catalyst yielding the *R* ester in 84% yield (ee = 92% and de = 97%).¹⁶ The latter procedure

was in fact optimal for both enantiomers but due to the high cost of *L*-pantolactone not used for bulk preparation for the *R*-enantiomer of **18**. Optical purity of all subsequent compounds was checked using chiral GC, HPLC and/or polarimetry and in all cases no loss of optical purity was detected in later synthetic steps.

Both syntheses now proceeded in identical manner and Scheme 2 only illustrates the synthesis of the *S*-enantiomer of **11**. Hydrolysis of **14** using LiOH in an equivolume mixture of THF, water and methanol for 1 h at 50°C gave carboxylic acid **18** in near quantitative yield. **18** was converted to *p*-methoxybenzyl urethane derivative **19** by treatment with diphenylphosphoryl azide (DPPA) followed by a Curtius rearrangement in the presence of *p*-methoxybenzyl alcohol, which proceeded with strict retention of stereochemistry.^{15b} The overall yield of the urethane product **19** was 60% over the two steps. Final conversion to (*S*)-**11** was achieved first through reduction with LiAlH_4 in anhydrous Et_2O (50%) then HBTU mediated coupling to 4-methylpent-3-enoic acid (70%) followed by a second reduction with LiAlH_4 in Et_2O . To prevent air oxidation upon storage the product was converted to its hydrochloride salt with HCl in ether, yielding (*S*)-**11**·HCl in 55% yield over the final two steps. The optical purity of (*S*)-**11** was estimated to be $\geq 98\%$ by comparison with previously reported data.^{14a} Similar results were obtained for the synthesis of (*R*)-**11**.

To validate any results obtained for these compounds as inhibitors of DCS, they were tested as inhibitors of aristolochene synthase from *Penicillium roqueforti* (AS) and amorphadiene synthase (ADS). These two enzymes are known to proceed via 1,10- and 1,6-cyclisations of the initial carbocation during their catalytic cycle (Scheme 3).^{1g} Hence aza-bisabolyl cations **11** should act as poor inhibitors of AS and potent inhibitors of ADS, as they closely resemble a reaction intermediate in the latter case only.

Recombinant AS and ADS were prepared and purified according to previously published procedures^{17,18} and both



Scheme 3 Initial catalytic chemical steps leading to (c) (+)-aristolochene (**22**) and (d) amorphadiene (**23**) follow 1,10- and 1,6-cyclisation of FDP, respectively.



(*R*)-**11** and (*S*)-**11** were tested as inhibitors using a standard radiolabelled assay involving conversion of tritium labelled FDP by each enzyme and scintillation counting of the pentane extractable products.^{5b} Terpene synthases are known to efficiently bind cation-PP_i pairs and inhibition was assessed both in the presence and absence of 250 μM diphosphate (Table 1). Synergistic inhibition of aza-analogues **11** with diphosphate has been observed previously for a variety of other terpene synthases.^{5d,13c,14}

Kinetic data were fitted by non-linear regression to the Michaelis-Menten equation ($v_0 = k_{\text{cat}}[E][S]/(K_M + [S])$). The mode of inhibition was determined by inspection of double reciprocal plots and observed to be competitive in all cases where inhibition was significant at low concentrations of **11**. K_I was determined from a plot of inhibitor concentration versus $K'_M/(k_{\text{cat}}[E])$ where $K'_M = K_M(1 + [I]/K_I)$.

The inhibition data for AS and ADS validate both of these compounds as valuable mechanistic probes for the present investigation since they are poor inhibitors of AS and potent inhibitors of ADS. PP_i had little effect on the ability to inhibit AS ($K_I > 200$ μM in both the presence and absence of PP_i for AS). Both enantiomers of **11** acted as competitive inhibitor of ADS, showing that they are able to compete effectively with the natural substrate FDP at the active site. As these aza-compounds cannot be turned over by ADS, these results support the intermediacy of an α-bisabolyl cation in the biosynthesis of amorpha-4,11-diene, in agreement with the findings of Picaud *et al.*^{18b} who used deuterated farnesyl diphosphate and deuterium exchange experiments to suggest that the *R*-enantiomer of the α-bisabolyl cation is the sole intermediate formed in the biosynthesis of amorpha-4,11-diene. Therefore, only the *R* enantiomer of **11** would be expected to inhibit ADS; however, if the *S*-enantiomer was a slightly more potent inhibitor ($K_I = 50$ μM for (*R*)-**11** versus 25 μM for (*S*)-**11**) Table 1. This is consistent with a flexible model for sesquiterpene active sites, according to which an active site can accommodate a variety of intermediates of different shape and charge distribution without being rigidly complementary to a single intermediate or transition state species. For example, work by Cane *et al.* showed that both enantiomers of the aza-analogue **11** were equally effective inhibitors of trichodiene synthase.^{14a} It is also notable that the presence of PP_i enhanced inhibition of ADS

by both enantiomers, improving the K_I approximately 20-fold ($K_I = 1.5$ and 3.7 μM for the *S* and *R* enantiomers respectively) demonstrating that the active site of ADS prefers a cation-anion pair in its active site.^{5d,13}

Recombinant DCS was generated with a C-terminal hexahistidine tag (DCS-His₆) as previously described.¹⁹ Inhibition assays were carried out using the same protocol used for AS and ADS. Both aza analogues were found to be competitive inhibitors of DCS-His₆ in the presence of PP_i but only poor inhibitors in its absence (Table 1). DCS clearly requires a cation-anion pair in its active site for effective inhibition by aza-analogues. Our results provide strong evidence for 1,6-cyclase activity for DCS.

Conclusions

The aza-bisabolyl cations **11** were potent competitive inhibitors of ADS, a 1,6-cyclase yet were much poorer inhibitors of PR-AS, a known 1,10-cyclase. When DCS was challenged with these aza-analogues in the presence of diphosphate anion they were potent inhibitors of the conversion of FDP to (+)-δ-cadinene (**7**), which would only be expected if DCS had a 1,6-cyclase activity. The use of a variety of substrate analogues possessing different stereochemistry and heteroatoms did not lead to clear results regarding whether DCS follow a 1,6 or 1,10 pathway.¹² If the proposed initial isomerism of the substrate to nerolidyl diphosphate (**2**) was suppressed using a fluorine atom at C2 then a 1,10 cyclisation was observed (Fig. 2). 2-Fluorogemacrene A (**25**) was the DCS catalysed product from the transoid (2*Z*,6*E*)-2-fluorofarnesyl diphosphate (**24**) while the cisoid substrate analogue **26** gave the cisoid product 2F-helminthogermacrene A (**27**).¹² However, in nearly every other case involving the use of substrate analogues with DCS, 1,6-cyclisation was observed at least in-part (Fig. 2).¹² These results may simply reflect the use of different substrates rather than an inherent ability of DCS to catalyse the conversion of FDP to **7** along two distinct reaction paths.²⁰ The observation that **11** acts as a competitive inhibitor of the DCS catalysed conversion of FDP provides strong evidence that DCS can efficiently use a 1,6-cyclisation pathway.

The fact that inorganic diphosphate led to a more tightly bound active site carbocation/diphosphate ion pair is consistent with previous work where often the active site recognises a cation-PP_i pair more effectively than the cation alone.^{5b,c-7a,14} The fact that (*R*)-**11** acts as a weak inhibitor in the absence of PP_i is more difficult to explain. It was previously suggested that, in the DCS active site pocket, the alkenyl chain of (3*R*)-nerolidyl diphosphate (**2**) is ideally positioned to ensure the formation of the α-bisabolyl cation with an *R* configuration at C6 (Scheme 1).¹² The C1–C7 hydride shift from **8** to **9** then occurs to the same *Si* face of C7 in cation **8**, therefore a (7*R*)-**9** formation is expected (Scheme 1). Hence the (*R*)-**11** should mimic better the α-bisabolyl cation generated by this enzyme, and therefore act as competitive inhibitor with higher binding affinity when compared with the *S*-enantiomer. The evidence

Table 1 Kinetic data for inhibition of ADS, AS and DCS by (*R*)-(**11**) and (*S*)-**11**. Uninhibited kinetic data for each enzyme: ADS $K_M = 2 \pm 0.15$ μM $k_{\text{cat}} = 1.19 \times 10^{-2} \pm 52 \times 10^{-5}$ s⁻¹. AS $K_M = 2.42 \pm 0.11$ μM, $k_{\text{cat}} = 1 \times 10^{-2} \pm 2 \times 10^{-5}$ s⁻¹. DCS-His₆ $K_M = 0.58 \pm 11$ μM $k_{\text{cat}} = 1.26 \times 10^{-3} \pm 5 \times 10^{-6}$ s⁻¹

Enzyme	Aza-analogue	K_I (μM) (+250 μM PP _i)	K_I (μM)
ADS	(<i>S</i>)- 11	1.5 ± 0.5	25 ± 5
	(<i>R</i>)- 11	3.7 ± 1.9	50 ± 17
AS	(<i>S</i>)- 11	255 ± 23	295 ± 23
	(<i>R</i>)- 11	489 ± 62	472 ± 48
DCS	(<i>S</i>)- 11	3.44 ± 1.43	273 ± 77
	(<i>R</i>)- 11	2.5 ± 0.5	1700 ± 300






Fig. 2 Summary of 1,10 and 1,6 cyclisation products generated through DCS catalysis.

that both enantiomers of **11** are equally as effective in the presence of PP_i is consistent with a permissive model of the active site structure, according to which an active site should accommodate a variety of rearranged intermediates of different shape and charge distribution without being rigidly complementary to a single intermediate. On the other hand, their lack of inhibitory effects on the 1,10-cyclase PR-AS shows that a major difference in the connectivity of the aza-analogue compared to the carbocation intermediate (*i.e.* bisaboyl cation rather than the 10-membered ring containing germacrene cation) renders them ineffective as inhibitors; hence the 1,6-cyclase activity of DCS postulated previously¹² is intrinsic to the enzyme.

Terpene synthases can generate great structural and stereochemical complexity in one synthetic step and have therefore potential as powerful synthetic biocatalysts for the generation of many bioactive compounds.^{4f,18a,21,22} A clear understanding of the catalytic strategies employed by these enzymes can aid

their redesign to produce nature-like compounds that are not found in the biosphere.^{23,24}

Experimental

General experimental procedures, enzyme preparation and purification are described in ESI† along with kinetics data, gas chromatograms, mass spectra and NMR spectra.

(R)-4,4-Dimethyl-2-oxotetrahydrofuran-3-yl acrylate (**16**)

Freshly distilled propenoyl chloride (0.41 mL, 5 mmol) was added over 1 h to a stirred solution of (*R*)-pantolactone (500 mg, 3.84 mmol) and Et_3N (583 mg, 5.76 mmol) in anhydrous CH_2Cl_2 (10 mL) at -24°C . The resulting mixture was stirred for 5 h at -24°C , and subsequently washed with aqueous 1 M HCl (10 mL). The aqueous phase was then extracted with CH_2Cl_2 (3×20 mL). The combined organic phases were washed with saturated NaHCO_3 solution (3×20 mL), water (3×20 mL) and brine (3×20 mL). The organic phase was dried over MgSO_4 , concentrated under reduced pressure and the residue was purified by flash chromatography on silica ($\text{EtOAc}:\text{hexane } 4:6$) to yield the pure compound as a yellow oil (375 mg, 53%). δ_{H} (300 MHz, CDCl_3) 6.48 (1 H, d, $J = 17.5$, $\text{CHH}=\text{C}$), 6.18 (1H, dd, $J = 17.5, 10.5$ Hz, $\text{CH}=\text{CH}_2$), 5.93 (1H, d, $J = 10.5$ Hz, $\text{CHH}=\text{C}$), 5.40 (1H, s, $\text{C}=\text{OCH}-\text{O}$), 4.03 (2 H, s, $\text{CH}_2-\text{OC}=\text{O}$), 1.17 (3 H, s, CH_3), 1.08 (s, 3 H, CH_3); δ_{C} (63 MHz, CDCl_3) 174.7 ($\text{OC}=\text{OCHO}$), 172.5 ($\text{OC}=\text{OC}=\text{H}_2$), 134.0 ($\text{H}_2\text{C}=\text{CHC}=\text{O}$), 118.7 ($\text{H}_2\text{C}=\text{CHC}=\text{O}$), 76.1 ($\text{OC}=\text{OCHO}$), 74.6 (OCH_2CH), 40.2 ($\text{C}-(\text{CH}_3)_2$), 23.4 (CH_3), 23.0 (CH_3). $\alpha_{\text{D}} +10^\circ$ (CH_2Cl_2 , $c = 17$). Data are in agreement with previous work.^{15a}

(R)-4,4-Dimethyl-2-oxotetrahydrofuran-3-yl-(S)-4-methylcyclohex-3-ene-1-carboxylate (**17**)

To a solution of **16** (302 mg, 1.67 mmol) in anhydrous CH_2Cl_2 (5 mL) at -10°C , TiCl_4 (0.82 mL, 0.82 mmol, 1.0 M solution in CH_2Cl_2) was added, and the resulting solution was stirred under argon at -10°C for 1 h. 2-Methylbutadiene (0.23 mL, 2.3 mmol) was then added over 5 min and the mixture was left stirring for 3 h at -10°C . The reaction was quenched by addition of 10% Na_2CO_3 in water (5 mL). The aqueous phase was then extracted with CH_2Cl_2 (3×20 mL). The organic layers were combined, washed with H_2O (3×10 mL), brine (3×10 mL), dried over anhydrous MgSO_4 , filtered and concentrated under reduced pressure. The residue was purified by flash chromatography on silica ($\text{EtOAc}:\text{hexane } 2:8$) to yield pure **17** as a colourless oil (353 mg, 85% yield). δ_{H} (300 MHz, CDCl_3) 5.32 (1 H, s, $\text{OCHC}=\text{O}$), and (1 H, br $\text{CH}=\text{C}$), 3.98 (2 H, s, CHCH_2O), 2.69–2.51 (1 H, m, $\text{CH}-\text{C}=\text{O}$), 2.19 (2 H, m broad, $\text{CHO}-\text{CH}_2\text{C}=\text{C}$), 1.98 (3 H, m, $\text{CHH}-\text{CH}_2$), 1.73 (1H, m, $-\text{CHH}-\text{CH}_2$), 1.58 (s, 3H, $\text{CH}_3\text{C}=\text{C}$), 1.13 (3H, s, CH_3), 1.04 (3 H, s, CH_3). δ_{C} (75 MHz, CDCl_3) 174.7 ($\text{OC}=\text{OCHO}$), 172.5 ($\text{OC}=\text{OC}=\text{H}_2$), 134.0 ($\text{HC}=\text{CCH}_3$), 118.7 ($\text{HC}=\text{CCH}_3$), 76.1 ($\text{OC}=\text{OCHO}$), 74.6 (OCH_2C), 40.2 ($\text{C}-(\text{CH}_3)_2$), 38.9 ($\text{CH}_2\text{CH}-\text{C}=\text{O}$), 28.9 ($\text{HC}=\text{CCH}_2$), 27.7 ($\text{C}=\text{CHCH}_2$), 25.2



(HC=CCH₂CH₂), 23.4 (CCH₃), 23.0 (CCH₃), 19.8 (CH₃C=CH). LRMS (EI⁺) *m/z*: 252.13 (50%), 122.07 (20), 94.08 (100), 79.05 (30), 67.05 (10). α_D -51.3 (CHCl₃, *c* = 1). Data are in agreement with previous work.¹⁶

(S)-3-Acryloyl-4-benzoyloxazolidin-2-one (13)

To a solution of (S)-4-benzoyloxazolidin-2-one (1.00 g, 5.60 mmol) in anhydrous THF (12 mL) at -78 °C, *n*-BuLi (2.1 M in THF, 3.14 mL, 6.59 mmol) was added dropwise over 30 minutes and the mixture stirred for a further 3 h at -78 °C. Freshly distilled acryloyl chloride (557 mg, 6.16 mmol) was added dropwise over 20 minutes and the reaction stirred for 2 h at -78 °C. The reaction was then allowed to warm to room temperature overnight. The reaction was quenched with sat. NH₄Cl (20 mL) and extracted with diethyl ether (3 × 30 mL). The organic layer was washed with water (3 × 40 mL), saturated aqueous NaHCO₃ (3 × 40 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure. Flash chromatography on silica gel (hexane : ethyl acetate 6 : 4) afforded **13** as a colorless solid (452 mg, 35%). δ_H (CDCl₃, 300 MHz) 7.45 (dd, 1H, *J* = 6.0, 18.0, CH=CH₂), 7.23 (m, 5H, aromatic Hs), 6.54 (dd, 1H, *J*_{H,H} = 18.0, 18.0, CHH=CH₂), 5.87 (dd, 1H, *J* = 9.0, 9.0, CHH=C), 4.68 (m, 1H, CHN), 4.14 (m, 2H, CH₂O), 3.29 (dd, 1H, *J* = 9.0, 9.0, C=CHHPh), 2.74 (dd, 1H, *J* = 12.0, 12.0, CHHPh) α_D -86° (CH₂Cl₂, *c* = 0.65). Data are in agreement with previous work.^{14a}

(R)-4-Benzyl-3-((S)-4-methylcyclohex-3-enecarbonyl)oxazolidin-2-one (14)

To a stirred solution of **13** (200 mg, 0.86 mmol) at -100 °C, were added 2-methylbutadiene (1.72 mL, 17.2 mmol) in anhydrous CH₂Cl₂ (5.0 mL) and Et₂AlCl (1.2 mL, 1.5 eq.). The reaction was stirred at -100 °C for 30 min then the mixture was poured into ice cold aqueous hydrochloric acid (1 M, 20 mL). The mixture was extracted with CH₂Cl₂ (2 × 10 mL). The combined organic layers were dried over anhydrous MgSO₄, filtered, and concentrated under reduced pressure. The product was purified by flash chromatography on silica (EtOAc : hexane : Et₃N 92 : 7 : 1) to yield the Diels Alder adduct (**14**) as a white crystalline solid (139 mg, 54%). δ_H (300 MHz, CDCl₃) 7.34–7.10 (5 H, m, ArCH), 5.36 (1 H, br., CH=C), 4.63 (1 H, dt, *J* = 16.6, 6.9, CHN), 4.23–3.95 (2 H, m, CH₂O), 3.60 (1 H, t, *J* = 8.8, CHC=O), 3.20 (1 H, dd, *J* = 13.2, 3.3, CHHPh), 2.70 (1 H, dd, *J* = 13.3, 9.5, CHHPh), 2.20–1.6 (6 H, m, CH₂CH₂CHCH₂C=C), 1.64 (3H, s, CH₃) α_D +79 (CH₂Cl₂, *c* = 1.4). LR-MS (EI⁺) *m/z*: 299.15 (100% M⁺), 300.16 (15), 269.06 (18), 267.07 (50), 232.10 (20), 178.08 (100), 146.07 (30), 140.03 (55), 122.07 (20), 91.00 (65), 63.00 (30). Data are in agreement with previous work.^{14a}

(R)- and (S)-4-Methylcyclohex-3-ene-1-carboxylic acid (18)

To a solution of **14** or **17** (0.32 mmol) in THF : MeOH : H₂O (1 : 1 : 1, 1.5 mL), LiOH (67 mg, 1.6 mmol) was added, and the resulting mixture was vigorously stirred for 1 h at 50 °C. The reaction was then cooled to room temperature and concentrated under reduced pressure. The resulting slurry was dis-

solved in H₂O (10 mL) and extracted with CH₂Cl₂ (3 × 5 mL). The resulting aqueous phase was acidified to pH = 2 at 0 °C with 15% HCl, extracted with a mixture of *n*-pentane : CH₂Cl₂ (98 : 2 3 × 10 mL), dried over anhydrous Na₂SO₄, and concentrated under reduced pressure to give **18** as a white powder (35 mg, 80%). δ_H (CDCl₃, 300 MHz) δ 5.32 (1H, s, CH=), 2.54–2.39 (1 H, m, CH-COOH), 2.17 (2H, m, -CH₂CH=C), 1.93 (3 H, m, CHH-CH₂), 1.69 (1H, m, CHH-CH₂), 1.59 (3H, s, CH₃). δ_C (100 MHz, CDCl₃) δ 182.3 (HOC=O), 133.8 (HC=CCH₃), 119.0 (CH₂-HC=CCH₃), 39.0 (HC-COOH), 29.13 (CH₂C=CH₃), 27.3 (CH₂CH=CCH₃), 25.5 (CH₂CH₂C=CH), 23.5 (CH₃). (S)-**18**: α_D -80.6 (CHCl₃, *c* = 0.5); -106.4 (95% EtOH, *c* = 4). (R)-**18**: α_D +93 (CHCl₃, *c* = 0.5); +105.5 (95% EtOH, *c* = 4). M.p. 82–92 °C. LRMS (EI⁺) *m/z* 140.06 (100%, M⁺), 136.06 (15), 125.05 (40), 122.06 (100), 95.87 (100), 94.06 (100), 93.07 (80), 79.04 (100), 77.03 (100), 68.06 (90), 67.04 (100) Data are in agreement with previous work.^{15,16}

(R)- and (S)-4-Methoxybenzyl [(4-methylcyclohex-3-en-1-yl)methyl]-carbamate (19)

To a solution of **18** (1.3 g, 9.3 mmol) in anhydrous toluene (20 mL) at 0 °C, diphenylphosphoryl azide (2.2 mL, 10.2 mmol) and Et₃N (3.9 mL, 27.8 mmol) were added. The resulting mixture was left stirring for 3 h at 100 °C before 4-methoxybenzyl alcohol (1.27 mL, 10.2 mmol) was added, and the reaction was left to stir for 16 h at 100 °C. The reaction was then allowed to cool to room temperature and the solution was concentrated under reduced pressure. The residue was the purified by flash chromatography on silica gel (EtOAc : *n*-hexane 1 : 9) to yield **19** as a yellow crystalline solid (1.28 g, 80%). δ_H (300 MHz, CDCl₃) δ 7.23 (2 H, dt, *J* = 2.9 and 5.3 CH ArCH), 6.81 (2 H, dt, *J* = 2.9 and 5.3, ArCH), 5.21 (1 H, br, HC=C), 4.95 (2 H, s, -OCH₂Ph), 4.67–4.60 (1 H, m, CHN), 3.74 (3 H, s, -OCH₃), 2.29–2.20 (2 H, m, CH₂-CH₂CHN), 1.93 (2 H, m, CH₂-CH₂CHN), 2.01–1.86 (2 H, m, CH-CH₂CHN), 1.55 (3 H, s, CH₃C=CH). δ_C (75 MHz, CDCl₃) δ 159.5 (=CO-CH₃), 155.8 (NHC=O), 134.1 (C=CCH₃), 130.0 (C=C aromatic), 128.7 (CCH₂O), 118.3 (C=CCH₃), 113.9 (C=C aromatic), 66.3 (CCH₂O), 62.8 (CHN), 55.3 (OCH₃), 31.9 (CH₂CHN), 28.4 (CH₂C=C), 28.0 (CH₂C=C), 23.4 (=CCH₃). ν_{max} (thin film, cm⁻¹) 3300 (N-H stretch), 2900–2700 (C-H stretch), 1650 (C=O ester stretch), 1250 (C-N stretch), 830 (aromatic CH bending); (S)-**19**: α_D -9.3, (*c* = 0.6, CHCl₃) (R)-**19**: α_D +12 (*c* = 0.6, CHCl₃) m.p. 69–71 °C LRMS (EI⁺) *m/z*: 275.15 (100% M⁺), 276.15 (20), 259.12 (18), 258.12 (60), 231.12 (25), 228.1128(12), 214.16 (100). HRMS (EI⁺) 275.1522; C₁₆H₂₁NO₃ requires 275.1521.

(R)- & (S)-N-Methyl-1-(4-methylcyclohex-3-en-1-yl)methanamine (20)

To a stirred solution of carbamate **19** (100 mg, 0.4 mmol) in anhydrous diethyl ether (7 mL) at 0 °C, was added LiAlH₄ (50 mg, 1.28 mmol). The mixture was then heated to reflux for 5 h. The reaction was cooled to 0 °C before it was quenched by the addition of water (6 mL) and an excess of 15% NaOH solution (6 mL). The resulting mixture was left to stir at 0 °C for



1 h and the precipitate was removed by filtration through a Celite pad. The organic phase was extracted with water (2 × 10 mL) and the pooled organic layers were then washed with 10% HCl (2 × 10 mL) and the organic fraction was discarded. The combined aqueous layers were adjusted to pH 12 by dropwise addition of 10% NaOH (15 mL). The product was extracted with diethyl ether (4 × 15 mL), then dried over anhydrous MgSO₄, and filtered. The product was then concentrated carefully under reduced pressure to give **20** as a volatile colorless oil (20 mg, 40%). δ_{H} (300 MHz, CDCl₃) 5.24 (1 H, m br, CH=C), 2.55 (1 H, dt, J = 16.6 and 8.0, CHNH₂), 2.37 (3 H, s, HNCH₃), 2.25–2.10 (1 H, m, NH), 1.99–1.87 (2 H, m, CH₂CH=C), 1.87–1.66 (2 H, m, CH₂CH₂C=CH), 1.61 (1 H, broad m, CH₂CH₂C=CH), 1.60 (3 H, s, H₂C=CCH₃), 1.45–1.26 (1 H, m, CH₂CH₂C=CH). δ_{C} (75 MHz, CDCl₃) 134.0 (C=CCH₃), 119.1 (C=CCH₃), 54.8 (CHNH), 32.7 (CH₂CHNH), 32.1 (C=CHCH₂CH), 29.7 (CH₃N), 29.0 (CH₂C=), 23.4 (CH₃C=) (*S*)-**20**: α_{D} –79 (c 1.00, CHCl₃) (*R*)-**20**: α_{D} +84 (c 1.00, CHCl₃) Data in agreement with previous work.^{14a}

(*R*)- and (*S*)-*N*,4-Dimethyl-*N*-(4-methylcyclohex-3-en-1-yl)pent-3-enamide (21)

To a stirred solution of **20** (172 mg, 1.5 mmol) and DIPEA (775 mg, 6.0 mmol) in anhydrous DMF (6 mL), HBTU (1.15 g, 3.0 mmol) was added and the resulting mixture was stirred at room temperature for 20 min before **20** (358 mg, 1.5 mmol) was added. The reaction was then stirred for 24 h at room temperature. The mixture was concentrated under reduced pressure and the residue was dissolved in diethyl ether (20 mL). The solution was washed with water (2 × 25 mL), 10% NaHCO₃ (2 × 25 mL), 10% HCl (2 × 10 mL), and brine (25 mL) before it was dried over anhydrous MgSO₄ and concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel (EtOAc:hexane 4:6) yielding **21** as a colorless oil (166 mg, 50%). δ_{H} (400 MHz, CDCl₃) 5.24 (2 H, dd, J = 14.6 and 8.0, NCHCH₂CH= and =CHCH₂C=O), 4.70–4.54 (0.5 H, m, CHN), 3.75 (0.5 H, dd, J = 12.6 and 9.7, CHN), 3.01 (2 H, dd, J = 12.6, 6.8 Hz, CH₂C=O), 2.75 (1.5 H, s, CH₃N), 2.72 (1.5 H, s, CH₃N), 2.20–1.86 (5 H, m, CH₂CH₂CHCH₂), 1.71–1.88 (1 H, m, CH₂CH₂CHCH₂), 1.75 (s, 3 H, CNCH₂=CHCH₃), 1.68 (3 H, s, =CCH₃), 1.66 (3 H, s, =CCH₃). (*S*)-**21**: α_{D} –9.5, (CHCl₃, c = 0.9) (*R*)-**21**: α_{D} +10, (CHCl₃, c = 0.9) HRMS (EI⁺) 221.1779; C₁₄H₂₃NO requires 221.1780. Data in agreement with previous work.^{14a}

(*R*)- and (*S*)-*N*,4-Dimethyl-*N*-(4-methylpent-3-en-3-yl)cyclohex-3-en-1 ammonium chloride (11)

To a stirred solution of **21** (41 mg, 0.19 mmol) in anhydrous diethyl ether at 0 °C, was added LiAlH₄ (33 mg, 0.87 mmol). The mixture was heated to reflux for 6 h then allowed to cool to room temperature and stirred for a further 12 h. The reaction was quenched by the addition of water (6 mL) and 15% NaOH (6 mL) at 0 °C and stirred for 1 h at 0 °C. The white precipitate was removed by filtration on Celite and the filtrate was extracted with diethyl ether (2 × 25 mL). The combined

organic layers were dried over anhydrous Na₂SO₄, concentrated under reduced pressure, and the residue was purified by flash chromatography on silica (Et₂O:MeOH 1:9) to yield the amine as a yellow oil (32 mg, 65% yield). δ_{H} (300 MHz, CDCl₃) 5.27 (1 H, d, J = 2.4, H₃CC=CH), 5.03 (1H, t, J = 5.6, (CH₃)₂C=CH), 2.64–2.44 (1 H, m, CHN), 2.38 (2 H, ddd, J = 7.6, 5.8 and 2.6, CH₂N), 2.23 (3 H, s, CH₃N), 2.08 (2 H, dd, J = 15.5 and 7.2, CH₂CH₂N), 2.04–1.85 (4 H, m, CH₂CH₂CHN), 1.85–1.67 (2 H, m, C=CHCH₂CHN), 1.62 (3 H, s, CH₃C=CH), 1.55 and 1.57 (2 × 3 H, 2 × s, (CH₃)₂C=CH). δ_{C} (75 MHz, CDCl₃) δ 133.9 (HC=CCH₃), 132.6 (CH=CCH₃), 122.1 (CH=CCH₃), 120.0 (NCH₂CH₂CH=CCH₃), 58.9 (CHN), 53.5 (CH₂N), 37.9 (CH₂CH₂CHN), 30.8 (=CHCH₂N), 27.2 (CH₂CCH₃), 26.5 (CH₃N), 25.7(CH₂CH₂N), 25.6 (CH₃CCH₃), 23.2 (CH₃CCH₃), 17.8 (CH₃C=). HRMS (EI⁺) 207.1990; C₁₄H₂₅N requires 207.1987. (*S*)-**11**: α_{D} –61 (CHCl₃, c = 1) (*R*)-**11** α_{D} +63 (CHCl₃, c = 1). Data are in agreement with previous work.^{14a} The amine was then dissolved in Et₂O (1 mL) and HCl (1 M in anhydrous Et₂O) was added slowly. A light yellow precipitate was formed. The ether was concentrated under reduced pressure and the salt stored in 1.2 mL of deionised water. δ_{H} (300 MHz, MeOD) 5.38 (1 H, br, H₃CC=CH), 5.14 (1 H, t, J = 5.0, (CH₃)₂C=CH), 3.58–3.42 (1 H, m, CHN), 3.32 (1 H, dd, J = 4.9, 1.6, CHHCH₂N), 3.25–2.97 (1 H, m, CHHCH₂N), 2.85 (3 H, s, CH₃N), 2.59–2.39 (2 H, m, CH₂CH₂CHN), 2.39–2.26 (2 H, m, CH₂CH₂CHN), 2.26–2.05 (2 H, m, C=CHCH₂CH₂N), 1.84 (2 H, dd, J = 12.2, 10.4 Hz, CH₃C=CHCH₂), 1.76 (3 H, s, CH₃C=CH), 1.72 (6 H, 2 × s, (CH₃)₂C=CH). δ_{C} (75 MHz, MeOD) δ 136.1 (HC=CCH₃), 134.3 (CH=C(CH₃)₂), 117.6 (CH=CCH₃), 116.5 (NCH₂CH₂CH=CCH₃), 61.8, 61.5 (CHN), 53.0, 52.5 (CH₂N), 35.9, 35.2 (CH₃N), 29.1, 29.0 (CH₂CH₂CHN), 25.7 (=CHCH₂N), 24.5 (CH₂CCH₃), 24.2, 24.0 (NCH₂CH₂), 23.25, 22.64 (CH₃C=), 21.64 (CH₃CCH₃), 16.65 (CH₃CCH₃). ν_{max} (neat, cm^{–1}) 2972 (broad, N–H stretch), 1379 (C–N stretch), 1161, 1051 and 1022 (C–N stretch), 950, 879, 815; HRMS (APCI⁺) 208.2057, C₁₄H₂₆N requires 208.2065; m.p. 131–133 °C, (*S*)-**11** α_{D} –62.2 (MeOH, c = 0.09), (*R*)-**11** α_{D} +57.1 (MeOH, c = 0.09). Data are in agreement with previous work.^{14a}

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

We thank Dr Robert Jenkins and Simon Waller for assistance with GC-MS experiments and Dr Juan A. Faraldos (all Cardiff University) for valuable scientific discussions during this work. This work was supported by the UK's Biotechnology and Biological Sciences Research Council (BBSRC) through grants (BB/H01683X/1, BB/M022463/1, BB/N012526/1). We thank Cardiff University for the provision of a postgraduate studentship to M. L.



Notes and references

- 1 (a) J. D. Connolly and R. A. Hill, *Dictionary of Terpenoids*, Chapman and Hall, London, 1991; (b) J. S. Glasby, *Encyclopedia of Terpenoids*, Wiley, Chichester, 1982; (c) E. M. Davis and R. Croteau, in *Biosynthesis: Aromatic Polyketides, Isoprenoids, Alkaloids*, Springer, Berlin Heidelberg, 2000, vol. 209, p. 53; (d) E. M. Davis, in *Comprehensive Natural Products II*, ed. L. Mander and H.-W. Liu, Elsevier, 2010, vol. 1, p. 585; (e) T. Toyomasu and T. Sassa, in *Comprehensive Natural Products II*, ed. L. Mander and H.-W. Liu, Elsevier, 2010, vol. 1, p. 643; (f) T. Kushiro and Y. Ebizuka, in *Comprehensive Natural Products II*, ed. L. Mander and H.-W. Liu, Elsevier, 2010, vol. 1, p. 673; (g) D. J. Miller and R. K. Allemann, *Nat. Prod. Rep.*, 2012, **29**, 60; (h) Y. Gao, R. B. Honzatko and R. J. Peters, *Nat. Prod. Rep.*, 2012, **29**, 1153.
- 2 (a) D. Cane, *Chem. Rev.*, 1990, **90**, 1089; (b) D. W. Christianson, *Chem. Rev.*, 2006, **106**, 3412; (c) D. W. Christianson, *Chem. Rev.*, 2017, **117**, 11570; (d) D. J. Tantillo, *Angew. Chem., Int. Ed.*, 2017, **56**, 10040.
- 3 (a) B. Felicetti and D. E. Cane, *J. Am. Chem. Soc.*, 2004, **126**, 7212; (b) M. Seemann, G. Z. Zhai, J. W. de Kraker, C. M. Paschall, D. W. Christianson and D. E. Cane, *J. Am. Chem. Soc.*, 2002, **124**, 7681; (c) J. A. Aaron, X. Lin, D. E. Cane and D. W. Christianson, *Biochemistry*, 2010, **49**, 1787; (d) B. T. Greenhagen, P. E. O'Maille, J. P. Noel and J. Chappell, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 9826; (e) Y. Yoshikuni, T. E. Ferrin and J. D. Keasling, *Nature*, 2006, **440**, 1078; (f) V. González, S. Touchet, D. J. Grundy, J. A. Faraldos and R. K. Allemann, *J. Am. Chem. Soc.*, 2014, **136**, 14505; (g) J. A. Faraldos, A. K. Antonczak, V. González, R. Fullerton, E. M. Tippmann and R. K. Allemann, *J. Am. Chem. Soc.*, 2011, **133**, 13906.
- 4 (a) J. Degenhardt, T. G. Kollner and J. Gershenzon, *Phytochemistry*, 2009, **70**, 1621; (b) D. E. Cane, P. C. Prabhakaran, J. S. Oliver and D. B. McIlwaine, *J. Am. Chem. Soc.*, 1990, **112**, 3209; (c) S. H. Kim, K. Heo, Y. J. Chang, S. H. Park, S. K. Rhee and S. U. Kim, *J. Nat. Prod.*, 2006, **69**, 758; (d) C. D. Poulter, J. C. Argyle and E. A. Mash, *J. Biol. Chem.*, 1978, **253**, 7227; (e) R. Croteau, *Arch. Biochem. Biophys.*, 1986, **251**, 777; (f) D. J. Miller, F. L. Yu, D. W. Knight and R. K. Allemann, *Org. Biomol. Chem.*, 2009, **7**, 962; (g) J. A. Faraldos, Y. X. Zhao, P. E. O'Maille, J. P. Noel and R. M. Coates, *ChemBioChem*, 2007, **8**, 1826.
- 5 (a) J. Y. Jiang, X. F. He and D. E. Cane, *Nat. Chem. Biol.*, 2007, **3**, 711; (b) J. A. Faraldos, B. Kariuki and R. K. Allemann, *J. Org. Chem.*, 2010, **75**, 1119; (c) J. A. Faraldos and R. K. Allemann, *Org. Lett.*, 2011, **13**, 1202; (d) A. S. Narula, A. Rahier, P. Beneviste and F. Shuber, *J. Am. Chem. Soc.*, 1981, **103**, 2408; (e) C. D. Poulter, *Acc. Chem. Res.*, 1990, **23**, 70.
- 6 (a) J.-X. Li, X. Fang, Q. Zhao, J.-X. Ruan, C.-Q. Yang, L. J. Wang, D. J. Miller, J. A. Faraldos, R. K. Allemann, X. Y. Chen and P. Zhang, *Biochem. J.*, 2013, **451**, 417; (b) D. J. Grundy, M. Chen, V. González, S. Leoni, D. J. Miller, D. W. Christianson and R. K. Allemann, *Biochemistry*, 2016, **55**, 2112; (c) K. Y. Shishova, F. Yu, D. J. Miller, J. A. Faraldos, Y. Zhao, R. M. Coates, R. K. Allemann, D. E. Cane and D. W. Christianson, *J. Biol. Chem.*, 2008, **283**, 15431; (d) M. Chen, N. Al-lami, M. Janvier, E. L. D'Antonio, J. A. Faraldos, D. E. Cane, R. K. Allemann and D. W. Christianson, *Biochemistry*, 2013, **52**, 5441; (e) H. A. Gennadios, V. González, L. Di Costanzo, A. Li, F. L. Yu, D. J. Miller, R. K. Allemann and D. W. Christianson, *Biochemistry*, 2009, **48**, 6175.
- 7 (a) J. A. Faraldos, D. J. Grundy, O. Cascon, S. Leoni, M. W. van der Kamp and R. K. Allemann, *Chem. Commun.*, 2017, **52**, 14027; (b) M. W. van der Kamp, J. Sirirak, J. Zurek, R. K. Allemann and A. J. Mulholland, *Biochemistry*, 2013, **52**, 8094; (c) R. K. Allemann, N. J. Young, S. Ma, D. G. Truhlar and J. Gao, *J. Am. Chem. Soc.*, 2007, **129**, 13008; (d) S. R. Hare and D. J. Tantillo, *Beilstein J. Org. Chem.*, 2017, **13**, 1669; (e) D. J. Tantillo, *Nat. Prod. Rep.*, 2011, **28**, 1035.
- 8 C. L. Steele, J. Crock, J. Bohlmann and R. Croteau, *J. Biol. Chem.*, 1998, **273**, 2078.
- 9 G. D. Davis and M. Essenberg, *Phytochemistry*, 1995, **39**, 553.
- 10 (a) C. R. Benedict, J. L. Lu, D. W. Pettigrew, J. Liu, R. D. Stipanovic and H. J. Williams, *Plant Physiol.*, 2001, **125**, 1754; (b) S. Opitz, G. Kunert and J. Gershenzon, *J. Chem. Ecol.*, 2008, **34**, 508.
- 11 (a) G. M. Bianchini, R. D. Stipanovic and A. A. Bell, *J. Agric. Food Chem.*, 1999, **47**, 4403; (b) F. Chen, D. Tholl, J. Bohlmann and E. Pichersky, *Plant J.*, 2011, **66**, 212.
- 12 J. A. Faraldos, D. J. Miller, V. González, Z. Yoosuf-Aly, O. Cascón, A. Li and R. K. Allemann, *J. Am. Chem. Soc.*, 2012, **134**, 5900.
- 13 (a) J. A. Faraldos, B. M. Kariuki and R. M. Coates, *Org. Lett.*, 2011, **13**, 836; (b) S. C. Karcher and S. A. Laufer, *J. Med. Chem.*, 2009, **52**, 1778; (c) A. N. van Nhien, C. Tomassi, C. Len, J. L. Marco-Contelles, J. Balzarini, C. Pannecouque, E. de Clercq and D. Postel, *J. Med. Chem.*, 2005, **48**, 4276; (d) M. Taton, M. Ceruti, L. Cattel and A. Rahier, *Phytochemistry*, 1996, **43**, 75; (e) A. Steiger, H.-J. Pyun and R. M. Coates, *J. Org. Chem.*, 1992, **57**, 3444; (f) D. A. Evans, K. T. Chapman and J. Bisaha, *J. Am. Chem. Soc.*, 1988, **110**, 1238; (g) D. A. Evans, K. T. Chapman and J. Bisaha, *J. Am. Chem. Soc.*, 1984, **106**, 4261.
- 14 (a) D. E. Cane, G. Yang and R. M. Coates, *J. Org. Chem.*, 1992, **57**, 3454; (b) M. M. Ravn, R. J. Peters, R. M. Coates and R. Croteau, *J. Am. Chem. Soc.*, 2002, **124**, 6998; (c) K. Zhou, R. J. Peters and R. M. Coates, *J. Am. Chem. Soc.*, 2007, **124**, 12453.
- 15 (a) T. Poll, A. Sobczak, H. Hartmann and G. Helmchen, *Tetrahedron Lett.*, 1985, **26**, 3095; (b) R. M. Sandifer, M. D. Thompson, R. G. Gaughan and C. D. Poulter, *J. Am. Chem. Soc.*, 1982, **4**, 7376.



- 16 (a) L. Argenti, F. Bellina, A. Carpita and R. Rossi, *Synth. Commun.*, 1995, **25**, 1909; (b) W. J. Watkins, Preparation of thiophencarboxylic acid derivatives for use as flaviviridae viruses inhibitors, *US Patent* 20130052161, 2013.
- 17 M. Calvert, S. E. Taylor and R. K. Allemann, *Chem. Commun.*, 2002, 2384.
- 18 (a) M. Demiray, X. Tang, J. A. Faraldos, T. Wirth and R. K. Allemann, *Angew. Chem., Int. Ed.*, 2017, **56**, 4347; (b) S. Picaud, P. Mercke, X. He, O. Sterner, M. Brodelius, D. E. Cane and P. E. Brodelius, *Arch. Biochem. Biophys.*, 2006, **448**, 150.
- 19 M. Loizzi, V. González, D. J. Miller and R. K. Allemann, *ChemBioChem*, 2018, **19**, 100–105.
- 20 D. J. Tantillo, *Angew. Chem., Int. Ed.*, 2017, **56**, 10040.
- 21 X. Tang, M. Demiray, T. Wirth and R. K. Allemann, *Bioorg. Med. Chem.*, 2017, **19**, 100.
- 22 D. J. Miller, F. Yu and R. K. Allemann, *ChemBioChem*, 2007, **15**, 1819.
- 23 S. Touchet, K. Chamberlain, C. M. Woodcock, D. J. Miller, M. A. Birkett, J. A. Pickett and R. K. Allemann, *J. Am. Chem. Soc.*, 2015, **136**, 7550.
- 24 F. Huynh, D. J. Miller and R. K. Allemann, *ChemBioChem*, 2018, **19**, 1834.

