


PCCP

Physical Chemistry Chemical Physics

rsc.li/pccp



ISSN 1463-9076




PAPER

Milan Ončák, Martin K. Beyer *et al.*
Size-dependent H and H₂ formation by infrared multiple
photon dissociation spectroscopy of hydrated vanadium
cations, V⁺(H₂O)_n, *n* = 3–51



Cite this: *Phys. Chem. Chem. Phys.*,
2022, 24, 14699

Size-dependent H and H₂ formation by infrared multiple photon dissociation spectroscopy of hydrated vanadium cations, V⁺(H₂O)_n, n = 3–51^{†‡}

Jakob Heller, Ethan M. Cunningham,  Jessica C. Hartmann, 
Christian van der Linde,  Milan Ončák * and Martin K. Beyer 

Infrared spectra of the hydrated vanadium cation (V⁺(H₂O)_n; n = 3–51) were measured in the O–H stretching region employing infrared multiple photon dissociation (IRMPD) spectroscopy. Spectral fingerprints, along with size-dependent fragmentation channels, were observed and rationalized by comparing to spectra simulated using density functional theory. Photodissociation leading to water loss was found for cluster sizes n = 3–7, consistent with isomers featuring intact water ligands. Loss of molecular hydrogen was observed as a weak channel starting at n = 8, indicating the advent of inserted isomers, HVOH⁺(H₂O)_{n–1}. The majority of ions for n = 8, however, are composed of two-dimensional intact isomers, concordant with previous infrared studies on hydrated vanadium. A third channel, loss of atomic hydrogen, is observed weakly for n = 9–11, coinciding with the point at which the H and H₂O calculated binding energies become energetically competitive for intact isomers. A clear and sudden spectral pattern and fragmentation channel intensity at n = 12 suggest a structural change to inserted isomers. The H₂ channel intensity decreases sharply and is not observed for n = 20 and 25–51. IRMPD spectra for clusters sizes n = 15–51 are qualitatively similar indicating no significant structural changes, and are thought to be composed of inserted isomers, consistent with recent electronic spectroscopy experiments.

Received 18th February 2022,
Accepted 6th April 2022

DOI: 10.1039/d2cp00833e

rsc.li/pccp

Introduction

Hydrated ions are ubiquitous to many important biological and chemical processes.¹ Studying gas-phase hydrated ions provides a well-defined system to investigate fundamental ion–water behavior at the molecular level.^{2–6} Important insights include solvation, hydrogen production, and corrosion mechanisms in which gas-phase studies offer the promise of bridging the gap between small clusters and bulk aqueous solution.^{7–14}

Early experiments by Armentrout investigated the successive binding energies of water molecules to first row transition metals by collision induced dissociation (CID).¹⁵ Size-dependent fragmentation behavior of a range of hydrated metal ions has been studied by blackbody infrared radiative dissociation (BIRD),^{16–19} along with proton transfer reactions employing H₂O/D₂O exchange.^{20,21} BIRD experiments on V⁺(H₂O)_n complexes

(n = 5–30) reveal strongly size-dependent intracluster redox reactivity. Vanadium was found in oxidation state +II for cluster sizes n = 9–12, resulting in V(OH)⁺(H₂O)_n ions and loss of atomic hydrogen, and the +III state for cluster sizes n = 9–23, leading to V(OH)₂⁺(H₂O)_n ions and loss of H₂; all other cluster sizes show only water fragmentation.¹⁹

A wealth of spectroscopic studies have been performed on gas-phase solvated metal ions, including electronic spectroscopy,^{22–34} with many more employing infrared spectroscopy, a powerful technique used to reveal structural trends of metal–ligand clusters.^{1,35–58} The hydrated vanadium cation, however, has taken center stage and has been studied in detail by many groups using different experimental techniques. Laser ablated vanadium, when co-deposited with water molecules in an argon matrix, reacts spontaneously forming the inserted complex, HVOH.⁵⁹ Sasaki *et al.* investigated small V⁺(H₂O)_n complexes (n = 2–8) employing the inert-messenger technique, revealing a square-planar coordination of the vanadium center with four water molecules.⁵⁸ Duncan and coworkers have performed many infrared studies on solvated vanadium complexes, the first in 2003 on V⁺(H₂O)Ar_n and V⁺(D₂O)Ar_n complexes in the O–H stretching region.⁶⁰ The doubly-charged species, V²⁺(H₂O)Ar_n (n = 2–7), was also studied, revealing a red-shift

Institut für Ionenphysik und Angewandte Physik, Universität Innsbruck,
Technikerstraße 25, 6020 Innsbruck, Austria. E-mail: milan.oncak@uibk.ac.at,
martin.beyer@uibk.ac.at

[†] Dedicated to Professor Wolfgang E. Ernst for his seminal contributions to the physics of molecules, clusters, metallic nanoparticles and material surfaces.

[‡] Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2cp00833e>



To confirm the presence of specific structural isomers inferred from the UV/Vis spectra, this study utilizes infrared multiple photon dissociation (IRMPD) spectroscopy in the O–H stretching region investigating the solvation evolution of $V^+(H_2O)_n$ ions up to $n = 51$. The goal is to establish the presence of inserted isomers at particular clusters sizes and any possible infrared-driven H and H_2 loss. The infrared-driven elimination of molecular hydrogen was recently observed for particular $HAHO^+(H_2O)_{n-1}$ clusters, $n = 9$ –14, probed in the 1400–2250 cm^{-1} region.⁶⁴ Presence of a hydride was found to be crucial in the elimination of molecular hydrogen, and found to occur only for cluster sizes $n = 12, 13$ and 14.

The experiments are performed on a 4.7 Tesla Fourier-Transform Ion Cyclotron Resonance Mass Spectrometer (FT-ICR MS) Bruker Spectrospin CMS47X.^{65–68} The mass spectrometer is equipped with a Bruker infinity cell⁶⁹ along with a laser vaporization source,^{70,71} where a solid disk of vanadium is vaporized by a frequency doubled Litron Nano S 60-30 Nd:YAG laser (532 nm, 5 mJ per pulse, 30 Hz). The plasma containing V^+ is entrained in a pulse of helium seeded with water vapor created *via* a homebuilt piezoelectric valve following the design of Proch and Trickl.⁷² The ensuing pulse is cooled *via* supersonic expansion, whereby hydrated vanadium complexes, $V^+(H_2O)_n$, are created. The clusters are transferred into the ICR cell where they are stored and mass-selected within the 4.7 T magnetic field⁷³ under ultra-high vacuum conditions ($p \approx 5 \times 10^{-10}$ mbar).

To complement the experimental photodissociation spectra, density functional theory (DFT) calculations were performed generating structures of $V^+(H_2O)_n$ at the B3LYP/aug-cc-pVDZ level of theory. Simulated infrared bands are scaled by a factor of 0.96 to compensate for anharmonicity and method deficiency, and spectra are generated by implementing Gaussian functions to band positions, each with a full-width-half-maximum (FWHM) of 20 cm^{-1} . All calculations were performed using the Gaussian 16 package.⁸³ Relative energies of isomers are given in kJ mol^{-1} , inclusive of zero-point energy. All isomers of the intact $V^+(H_2O)_n$ structure considered in this study are present in the quintet ground state, with other multiplicities lying higher in energy. The inserted $HVOH^+(H_2O)_n$ structures were found in triplet multiplicity, and are more stable than their singlet and quintet analogues. Wave function stabilization was performed prior to each structure optimization.

In Fig. 1, we compare the measured IRMPD spectra of $V^+(H_2O)_n$, $n = 3-51$, Fig. 2 includes fragmentation intensities of the observed dissociation channels. Infrared spectra of Ar and N_2 tagged as well as untagged $V^+(H_2O)_n$ have been published previously in two separate studies by the groups of Duncan and Ohashi, and the original spectra from the Duncan group are included in Fig. 1.^{58,62} Our photodissociation spectra up to $n = 7$ are concordant with these previous studies. Since no tagging technique is used in this work, investigations of smaller systems ($n \leq 2$) are not possible due to inefficient elimination




Fig. 1 Infrared multiple photon dissociation spectra of the $V^+(H_2O)_n$ complexes ($n = 3-12, 14, 15, 20, 25, 30, 42$ and 51) presented in black. For comparison, infrared spectra from Duncan and coworkers are also presented in red for cluster sizes $n = 3-5, 7, 8, 11, 14, 20, 25$ and 30 .⁶²

of H_2O . For cluster sizes $n \geq 8$, differences start to emerge between the IRMPD spectra measured in the present work (black spectra) and those recorded by Duncan and coworkers (red spectra).⁶² Most notably, the broad and weak red-shifted feature at $2500-3000\text{ cm}^{-1}$ seen in our IRMPD spectrum for $n = 11$ is not present in the $n = 11$ spectrum recorded by Duncan. Going larger to $n = 30$, the strongest feature is a broad absorption at $2800-3650\text{ cm}^{-1}$, with weaker, “free” O–H bands at 3700 and 3720 cm^{-1} . Duncan’s spectrum shows the opposite, the “free” O–H bands are the most intense, while the absorption at $3200-3650\text{ cm}^{-1}$ is weaker and spectrally narrower. These spectral differences can be attributed to the disparities in the two experimental methods employed. In particular, a time-of-flight mass spectrometry instrument is used in the

study by Duncan, employing single-photon infrared photon dissociation (IRPD) methods. Upon increasing the number of water molecules, the heat capacity of the cluster increases, which means water evaporation from the cluster becomes more demanding energetically. Thus, under a single-photon regime, the photofragmentation rate decreases upon increasing cluster size. In contrast, clusters in the present study are trapped in the center of an ICR cell, whereby infrared multiple photon dissociation (IRMPD) spectroscopy is employed. Therefore, any increases in heat capacity can be readily overcome by introducing more photons, causing significant photodissociation.

We start our discussion with the smallest clusters, $n = 3-7$, for which we see structured spectra with clearly defined bands. The only observed dissociation channel is water evaporation.






Fig. 2 Fragmentation channel intensities for all investigated clusters. The points are connected with lines to guide the eye.

In Fig. 3, we analyze the experimental as well as theoretical spectra of $V^+(H_2O)_5$ as an example (spectra of other small clusters are analyzed in Fig. S1–S3, ESI†). The experimental spectrum is in excellent agreement with the untagged spectra measured by Duncan, as well as the spectra recorded by Ohashi.^{58,62} As shown in Fig. 3, all three low-lying isomers of the intact $V^+(H_2O)_5$ present a good agreement with the bands in the observed spectrum, possessing both the four- (**Va**, **Vb**) and three-coordinate (**Vc**) vanadium cation. The bands observed at 3510 and 3690 cm^{-1} can be explained by isomer **Va**, consistent with Duncan's results of the $V^+(H_2O)_5\text{-Ar}$ cluster.⁶² The feature at 3345 cm^{-1} can be ascribed to the single-acceptor water molecule of isomer **Vb**, and both features at 3345 and 3230 cm^{-1} are in reasonable agreement with the two single-acceptor water molecules in **Vc**. In principle, the feature at 3230 cm^{-1} can also be assigned to the bending overtone of isomer **Va**, as proposed by Ohashi and coworkers.⁵⁸ However, our anharmonic frequency analysis shows that, while there are indeed bending overtones in the 3100–3200 cm^{-1} region, their intensity is two orders of magnitude lower compared to the intensity of the most intense O–H stretch. The four bands present in the observed IRMPD spectrum can be thus ascribed to contributions from only intact isomers.

The inserted species, $HVOH^+(H_2O)_4$ (**iVa**–**iVc**), are considerably more stable than the intact clusters (Fig. 3d). However, they show intense red-shifted bands below 3000 cm^{-1} which are not observed experimentally. Thus, the barrier to insertion must be too high to attain at these temperatures for this cluster size. This is also consistent with previous BIRD investigations on $V^+(H_2O)_5$, which presented only dissociation of water molecules.¹⁹

Starting with $n = 8$, H_2 evaporation appears as a new reaction channel, although sequential loss of water molecules is still the main channel (Fig. 2), consistent with previous BIRD studies of $V^+(H_2O)_8$.¹⁹ The number, and composition, of ligands lost during the IRMPD process for each cluster size is given in




Fig. 3 (a) Infrared multiple photon dissociation spectrum of the $V^+(H_2O)_5$ complex along with the simulated spectra of low-lying (b) intact (**Va**–**Vc**) and (c) inserted (**iVa**–**iVc**) isomers calculated at the B3LYP/aug-cc-pVDZ level of theory. (d) Structural isomers of intact (**Va**–**Vc**) and inserted isomers (**iVa**–**iVc**) with relative energies given in kJ mol^{-1} inclusive of zero-point energy. In each case the observed photodissociation events in a) are due to the sequential loss of intact water molecules, mH_2O .

the ESI† (see Table S1). In each case, m denotes the sequential number of water molecules, x denotes the number of waters following H dissociation, and y the number of waters following H_2 dissociation. Fig. 4a presents the IRMPD spectrum measured for $V^+(H_2O)_8$, showing a qualitatively similar IRMPD spectrum as observed for $n = 5$ – 7 . As discussed by Duncan,⁶² upon increasing cluster size the structural evolution of $V^+(H_2O)_n$ clusters retains a quasi-planar, square structural motif up to $n = 8$, presenting a metal coordination number of four. The two-dimensional structure is broken with inserted isomers. Different to the experimental spectrum recorded by




Fig. 4 Infrared multiple photon dissociation spectra of (a) $V^+(H_2O)_8$ and (b) $V^+(H_2O)_9$ along with the simulated photodissociation spectra of low-lying intact and inserted isomers calculated at the B3LYP/aug-cc-pVDZ level of theory. In each case, photodissociation events are due to sequential loss of either (i) loss of intact water molecules mH_2O , (ii) loss of $H + xH_2O$, and (iii) loss of $H_2 + yH_2O$, represented as dashed red, blue, and green lines, respectively.

Duncan, the spectrum in Fig. 4a shows the presence of a weak feature between 2790–3110 cm^{-1} .

IR spectra provided by quantum chemical calculations offer two possible explanations for the low-intensity tail of the spectrum. The band could correspond to inserted isomers (**iVIIIa–iVIIId**, see Fig. 5a) as these spectra present red-shifted bands in this region. However, the intact isomer **VIIId** also presents a band in this region at 3078 cm^{-1} . This vibration is represented by a green arrow in Fig. 5a, showing the O–H stretch in a second solvation sphere water molecule, in a double-acceptor site bound to another third solvation sphere water molecule. The observed red-shift signifies significant weakening of the O–H bond, and could indicate the onset of H atom transfer from the second sphere water molecule, forming H_3O^+ in the third sphere. Isomer **VIIId** is also only 7 $kJ\ mol^{-1}$ less stable than the most stable intact isomer found, **VIIIa**.

The discrepancy between the spectrum recorded in the present study and that by Duncan could be rationalized by temperature effects; for clusters thermalized in an ICR cell, warmer clusters are present, where entropic effects dominate enthalpic effects. More entropic water binding motifs with

red-shifted bands, such as single-acceptor sites observed in **VIIId**, are present. This effect has been observed before in our own infrared studies on hydrated zinc cations^{55,84} and in IR studies on $M^+(H_2O)_n$ complexes elsewhere.^{85,86}

The main contribution to the red-shifted feature between 2790–3110 cm^{-1} is the $H_2 + H_2O$ fragmentation channel (dashed green spectrum), and is slightly more intense than the intact water loss channel (dashed red spectrum). Based on previously calculated H_2 loss mechanisms of $Al^+(H_2O)_{20}$, pathways involving inserted isomers are more energetically favorable.^{87,88} Briefly, a proton moves from a first to a second shell water molecule leading to H_3O^+ . Through consecutive proton transfer through a “water wire”, the proton reaches a site near the hydride statistically. When the proton is pointing towards the hydride at the metal center, the hydride and the proton can recombine leading to the release of an H_2 molecule. The energy released from this recombination step leads to evaporation of multiple H_2O molecules, see calculated reaction energies in Table 1. Thus, based on the weakly observed $H_2 + H_2O$ fragment channel, there is a small contribution from inserted isomers. However, the majority of the ions present in the experiment are intact structures, $V^+(H_2O)_8$, as can be






Fig. 5 Calculated intact and inserted isomers of (a) $V^+(H_2O)_8$ and (b) $V^+(H_2O)_9$. All isomers were calculated at the B3LYP/aug-cc-pVDZ level of theory with relative energies given in kJ mol^{-1} inclusive of zero-point energy. Arrows indicate the O–H stretch of the most red-shifted band, in each case a water molecule in a double acceptor binding motif in the second solvation sphere.

Table 1 Calculated insertion energies (E_{ins}) and dissociation energies (all in kJ mol^{-1}) for intact and inserted isomers of $V^+(H_2O)_n$ stoichiometry. Calculated at the B3LYP/aug-cc-pVDZ level of theory

n	E_{ins}	Intact ($V^+(H_2O)_n$)			Inserted ($HVOH^+(H_2O)_{n-1}$)		
		H_2O	H	H_2	H_2O	H	H_2
5	–79	50	116	–142	63	195	–63
6	–90	42	88	–156	54	178	–66
7	–103	39	63	–174	52	165	–71
8	–111	40	45	–185	48	156	–74
9	–119	35	34	–196	43	153	–77

deduced from the low-intensity red tail of the spectrum. The energetically less stable intact isomers are generated in the source chamber, while the inserted isomers, $HVOH^+(H_2O)_7$, are generated *via* insertion and consecutive water fragmentation of larger clusters. A possible $2H + H_2O$ dissociation channel can be discarded as it lies by 4.48 eV (H_2 dissociation energy) higher in energy and is unlikely to contribute.

The IRMPD spectrum of $n = 9$ is presented in Fig. 4b, showing that the dissociation of water molecules is the dominant fragmentation channel, followed by $H + H_2O$, then the

$H_2 + H_2O$ channel. Similar fragmentation channels were observed in BIRD experiments, suggesting the same reactions take place with different heating mechanisms.¹⁹ Based on the discussion above, we can attribute H_2 loss to the presence of inserted isomers, $HVOH^+(H_2O)_8$. The H loss, on the other hand, can be assigned only to intact isomers as it is energetically too demanding for the inserted species (153 kJ mol^{-1} , Table 1). In addition, the energy required to dissociate an H atom from an intact cluster (34 kJ mol^{-1}) becomes isoenergetic with H_2O evaporation for $n = 9$ (Table 1). The $H + H_2O$ dissociation channel, along with the water dissociation channel, is thus assigned to intact isomers, $V^+(H_2O)_9$. In other words, the thermochemical considerations point at the co-existence of both cluster types. The intact and inserted isomers of $V^+(H_2O)_9$, along with the atomic and molecular hydrogen dissociation energy pathways, are presented in Fig. 6. The lowest energy pathway starts with an intact isomer $^5[V(H_2O)_9]^+$, with H lost from a water molecule (*i.e.* without insertion, resulting in product $^4[VOH(H_2O)_8]^+$). The insertion step is exothermic by 119 kJ mol^{-1} , which would certainly result in evaporation of multiple water molecules. Nevertheless, atomic H loss from the inserted species $^3[HVOH(H_2O)_8]^+$ requires 257 kJ mol^{-1} , resulting in the inserted product $^2[HV(OH)_2(H_2O)_7]^+$. Atomic H loss from the inserted species $^3[HVOH(H_2O)_8]^+$, resulting in $^4[VOH(H_2O)_8]^+$, is also energetically demanding, calculated as 153 kJ mol^{-1} . Based on these energies, the photodissociation spectrum of $H + H_2O$ in Fig. 4b originates from intact isomers.

Similar to $n = 8$, the spectrum of $n = 9$ also shows a broad structureless absorption at $2410\text{--}3640 \text{ cm}^{-1}$, the “free” O–H band at 3690 cm^{-1} and a broad red-shifted feature at $2410\text{--}3000 \text{ cm}^{-1}$ which gains intensity and an increased spectral width for $n = 9$. Fig. 4b also shows the simulated spectra of calculated intact isomers, **IXa–IXe**, and inserted isomers, **iIXa–iIXd**, the structures of which are shown in Fig. 5b. The broad absorption at $2410\text{--}3640 \text{ cm}^{-1}$ is the hydrogen bonding region composed of many different O–H environments with varying degrees of red-shift, with both intact and inserted isomers presenting agreement. The “free” O–H band at 3690 cm^{-1} also shows agreement with both isomer classes.

The red-shifted feature at $2410\text{--}3000 \text{ cm}^{-1}$ with all three reaction channels present provides particular insight into the cluster structure. As discussed above, we attribute the feature to the presence of both inserted isomers **iIXb** and **iIXd** (showing bands at 2820 and 2965 cm^{-1} , respectively) and the intact isomers, namely **IXc**, **IXd**, and **IXe** with bands at 3095 , 2990 , and 2895 cm^{-1} , respectively. The O–H stretch of the latter bands are shown as arrows in Fig. 5b, in each case the O–H bond of a second-sphere water molecule in a double-acceptor motif, bound to a third-sphere water molecule. Isomer **IXe** shows the most red-shifted band, which is caused by two fourth-sphere water molecules bound to the third-sphere water molecule acting on the second-sphere water. Induced polarisation effects onto the second-sphere water molecule from the metal center, and from the third-sphere water molecule, remove electron density from the bonding orbitals in the O–H bond, which weaken the bond causing the red-shift.






Fig. 6 Calculated intact and inserted isomers of $V^+(H_2O)_9$, along with insertion energy (ins.), molecular hydrogen and atomic hydrogen dissociation energy pathways. All isomers and energy pathways were calculated at the B3LYP/aug-cc-pVDZ level of theory with relative energies given in kJ mol^{-1} inclusive of zero-point energy.

We conclude that the majority of the ions for $n = 9$ are composed of intact isomers, however $n = 9$ represents the cluster size at which the abundance of inserted isomers starts to increase. Our conclusions are consistent with our previous investigation of $V^+(H_2O)_n$ in the ultraviolet/visible range, with a decrease in photodissociation cross section starting at $n \approx 9$, decreasing further upon increasing water until $n \geq 15$ where no photoinduced fragmentation was observed, indicating an oxidation state change from V(I) to V(III).⁶³

For $n = 11$, bands assigned to single-acceptor water motifs around 3200 cm^{-1} increase in intensity, as presented in Fig. 7. In terms of fragmentation channels, the $H_2 + H_2O$ channel (green dashed spectrum) is as intense as the $H + H_2O$ channel (blue dashed spectrum). However, the onset of H_2 loss starts at lower photon energies (2890 cm^{-1}) than H loss (3120 cm^{-1}). Below 2890 cm^{-1} , only the H_2O fragment channel is detected. The dissociation patterns suggest that intact and inserted isomers still co-exist for this cluster size.

Going to $V^+(H_2O)_{12}$, differences in fragmentation channels are observed (Fig. 7). In the whole spectral range, the $H_2 + H_2O$ channel is the most intense with no $H + H_2O$ loss observed and water loss observed as a weaker channel. Additionally, the spectral band intensities change, indicating a completed structural change to the inserted clusters. The band at 2900 cm^{-1} for $n = 9, 10$ and 11 , seems to decrease in intensity but it is still observable above the noise level. Since $V^+(H_2O)_{12}$ shows the second highest BIRD rate for $H_2 + H_2O$ loss, the signal-to-noise ratio, especially in the lower wavelength region, is lower due to BIRD dissociation, even at low temperatures.¹⁹ It is possible that this band becomes lost in the noise and thus, is undetectable. Given that $H_2 + H_2O$ is the most intense channel, the majority of ions present for $n = 12$ are composed of inserted isomers. This would indicate that 12 water molecules are enough to form three dimensional structures, whereby the




Fig. 7 Infrared multiple photon dissociation spectra of $V^+(H_2O)_n$ complexes ($n = 10-12, 14, 15$). In each case photodissociation events are due to sequential loss of either (i) loss of intact water molecules mH_2O , (ii) loss of $H + xH_2O$, and (iii) loss of $H_2 + yH_2O$, represented as dashed red, blue, and green lines, respectively.





- 11 R. Brimblecombe, G. F. Swiegers, G. C. Dismukes and L. Spiccia, *Angew. Chem., Int. Ed.*, 2008, **47**, 7335.
- 12 K. Fuke, K. Hashimoto and S. Iwata, *Adv. Chem. Phys.*, 1999, **110**, 431.
- 13 G. Niedner-Schatteburg and V. E. Bondybey, *Chem. Rev.*, 2000, **100**, 4059.
- 14 W. A. Donald, R. D. Leib, J. T. O'Brien, A. I. S. Holm and E. R. Williams, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 18102.
- 15 N. F. Dalleska, K. Honma, L. S. Sunderlin and P. B. Armentrout, *J. Am. Chem. Soc.*, 1994, **116**, 3519.
- 16 O. P. Balaj, C. B. Berg, S. J. Reitmeier, V. E. Bondybey and M. K. Beyer, *Int. J. Mass Spectrom.*, 2009, **279**, 5.
- 17 M. Beyer, C. Berg, H. W. Görlitzer, T. Schindler, U. Achatz, G. Albert, G. Niedner-Schatteburg and V. E. Bondybey, *J. Am. Chem. Soc.*, 1996, **118**, 7386.
- 18 M. Beyer, U. Achatz, C. Berg, S. Joos, G. Niedner-Schatteburg and V. E. Bondybey, *J. Phys. Chem. A*, 1999, **103**, 671.
- 19 B. S. Fox, I. Balteanu, O. P. Balaj, H. C. Liu, M. K. Beyer and V. E. Bondybey, *Phys. Chem. Chem. Phys.*, 2002, **4**, 2224.
- 20 C. van der Linde and M. K. Beyer, *J. Phys. Chem. A*, 2012, **116**, 10676.
- 21 C. van der Linde and M. K. Beyer, *Phys. Chem. Chem. Phys.*, 2011, **13**, 6776.
- 22 B. Scharfschwerdt, C. van der Linde, O. P. Balaj, I. Herber, D. Schütze and M. K. Beyer, *Low Temp. Phys.*, 2012, **38**, 717.
- 23 D. E. Lessen, R. L. Asher and P. J. Brucat, *J. Chem. Phys.*, 1990, **93**, 6102.
- 24 M. Rosi and C. W. Bauschlicher, *J. Chem. Phys.*, 1989, **90**, 7264.
- 25 M. Rosi and C. W. Bauschlicher, *J. Chem. Phys.*, 1990, **92**, 1876.
- 26 J. M. Farrar, *Int. Rev. Phys. Chem.*, 2003, **22**, 593.
- 27 C. S. Yeh, K. F. Willey, D. L. Robbins, J. S. Pilgrim and M. A. Duncan, *Chem. Phys. Lett.*, 1992, **196**, 233.
- 28 C. T. Scurlock, S. H. Pullins, J. E. Reddic and M. A. Duncan, *J. Chem. Phys.*, 1996, **104**, 4591.
- 29 M. Ončák, T. Taxer, E. Barwa, C. van der Linde and M. K. Beyer, *J. Chem. Phys.*, 2018, **149**, 44309.
- 30 T. Taxer, M. Ončák, E. Barwa, C. van der Linde and M. K. Beyer, *Faraday Discuss.*, 2019, **217**, 584.
- 31 J. S. Daluz, A. Kocak and R. B. Metz, *J. Phys. Chem. A*, 2012, **116**, 1344.
- 32 A. Kocak, G. Austein-Miller, W. L. Pearson, G. Altinay and R. B. Metz, *J. Phys. Chem. A*, 2013, **117**, 1254.
- 33 Y. Abate and P. D. Kleiber, *J. Chem. Phys.*, 2005, **122**, 84305.
- 34 M. Sanekata, F. Misaizu and K. Fuke, *J. Chem. Phys.*, 1996, **104**, 9768.
- 35 T. Iino, K. Ohashi, K. Inoue, K. Judai, N. Nishi and H. Sekiya, *Eur. Phys. J. D*, 2007, **43**, 37.
- 36 B. Bandyopadhyay, K. N. Reishus and M. A. Duncan, *J. Phys. Chem. A*, 2013, **117**, 7794.
- 37 J. M. Lisy, *Int. Rev. Phys. Chem.*, 1997, **16**, 267.
- 38 J. P. Beck and J. M. Lisy, *J. Chem. Phys.*, 2011, **135**, 44302.
- 39 N. R. Walker, R. S. Walters, M.-K. Tsai, K. D. Jordan and M. A. Duncan, *J. Phys. Chem. A*, 2005, **109**, 7057.
- 40 R. S. Walters, N. R. Brinkmann, H. F. Schaefer and M. A. Duncan, *J. Phys. Chem. A*, 2003, **107**, 7396.
- 41 R. S. Walters, E. D. Pillai and M. A. Duncan, *J. Am. Chem. Soc.*, 2005, **127**, 16599.
- 42 T. D. Vaden, J. M. Lisy, P. D. Carnegie, E. D. Pillai and M. A. Duncan, *Phys. Chem. Chem. Phys.*, 2006, **8**, 3078.
- 43 P. D. Carnegie, A. B. McCoy and M. A. Duncan, *J. Phys. Chem. A*, 2009, **113**, 4849.
- 44 P. D. Carnegie, B. Bandyopadhyay and M. A. Duncan, *J. Chem. Phys.*, 2011, **134**, 14302.
- 45 P. D. Carnegie, B. Bandyopadhyay and M. A. Duncan, *J. Phys. Chem. A*, 2008, **112**, 6237.
- 46 P. D. Carnegie, B. Bandyopadhyay and M. A. Duncan, *J. Phys. Chem. A*, 2011, **115**, 7602.
- 47 B. Bandyopadhyay and M. A. Duncan, *Chem. Phys. Lett.*, 2012, **530**, 10.
- 48 T. Iino, K. Ohashi, Y. Mune, Y. Inokuchi, K. Judai, N. Nishi and H. Sekiya, *Chem. Phys. Lett.*, 2006, **427**, 24.
- 49 Y. Inokuchi, K. Ohshimo, F. Misaizu and N. Nishi, *J. Phys. Chem. A*, 2004, **108**, 5034.
- 50 Y. Inokuchi, K. Ohshimo, F. Misaizu and N. Nishi, *Chem. Phys. Lett.*, 2004, **390**, 140.
- 51 T. Iino, K. Ohashi, K. Inoue, K. Judai, N. Nishi and H. Sekiya, *J. Chem. Phys.*, 2007, **126**, 194302.
- 52 K. Furukawa, K. Ohashi, N. Koga, T. Imamura, K. Judai, N. Nishi and H. Sekiya, *Chem. Phys. Lett.*, 2011, **508**, 202.
- 53 M. F. Bush, R. J. Saykally and E. R. Williams, *J. Am. Chem. Soc.*, 2008, **130**, 9122.
- 54 J. T. O'Brien and E. R. Williams, *J. Phys. Chem. A*, 2008, **112**, 5893.
- 55 E. M. Cunningham, T. Taxer, J. Heller, M. Ončák, C. van der Linde and M. K. Beyer, *Phys. Chem. Chem. Phys.*, 2021, **23**, 3627.
- 56 K. R. Asmis, *Phys. Chem. Chem. Phys.*, 2012, **14**, 9270.
- 57 H. Schwarz and K. R. Asmis, *Chem. – Eur. J.*, 2019, **25**, 2112.
- 58 J. Sasaki, K. Ohashi, K. Inoue, T. Imamura, K. Judai, N. Nishi and H. Sekiya, *Chem. Phys. Lett.*, 2009, **474**, 36.
- 59 M. Zhou, J. Dong, L. Zhang and Q. Qin, *J. Am. Chem. Soc.*, 2001, **123**, 135.
- 60 N. R. Walker, R. S. Walters, E. D. Pillai and M. A. Duncan, *J. Chem. Phys.*, 2003, **119**, 10471.
- 61 T. B. Ward, E. Miliordos, P. D. Carnegie, S. S. Xantheas and M. A. Duncan, *J. Chem. Phys.*, 2017, **146**, 224305.
- 62 P. D. Carnegie, J. H. Marks, A. D. Brathwaite, T. B. Ward and M. A. Duncan, *J. Phys. Chem. A*, 2020, **124**, 1093.
- 63 J. Heller, T. F. Pascher, D. Muß, C. van der Linde, M. K. Beyer and M. Ončák, *Phys. Chem. Chem. Phys.*, 2021, **23**, 22251–22262.
- 64 J. Heller, W. K. Tang, E. M. Cunningham, E. G. Demissie, C. van der Linde, W. K. Lam, M. Ončák, C.-K. Siu and M. K. Beyer, *Angew. Chem., Int. Ed.*, 2021, **60**, 16858–16863.
- 65 A. Akhgarnusch, W. K. Tang, H. Zhang, C.-K. Siu and M. K. Beyer, *Phys. Chem. Chem. Phys.*, 2016, **18**, 23528.
- 66 M. Allemann, H. Kellerhals and K. P. Wanczek, *Int. J. Mass Spectrom. Ion Process.*, 1983, **46**, 139.
- 67 C. Berg, T. Schindler, G. Niedner-Schatteburg and V. E. Bondybey, *J. Chem. Phys.*, 1995, **102**, 4870.



- 68 A. Akhgarnusch, R. F. Höckendorf and M. K. Beyer, *J. Phys. Chem. A*, 2015, **119**, 9978.
- 69 P. Caravatti and M. Allemann, *Org. Mass Spectrom.*, 1991, **26**, 514.
- 70 V. E. Bondybey and J. H. English, *J. Chem. Phys.*, 1981, **74**, 6978.
- 71 T. G. Dietz, M. A. Duncan, D. E. Powers and R. E. Smalley, *J. Chem. Phys.*, 1981, **74**, 6511.
- 72 D. Proch and T. Trickl, *Rev. Sci. Instrum.*, 1989, **60**, 713.
- 73 A. G. Marshall, C. L. Hendrickson and G. S. Jackson, *Mass Spectrom. Rev.*, 1998, **17**, 1.
- 74 N. Polfer, B. G. Sartakov and J. Oomens, *Chem. Phys. Lett.*, 2004, **400**, 201.
- 75 J. Heller, E. M. Cunningham, C. van der Linde, M. Ončák and M. K. Beyer, *J. Phys. Chem. Lett.*, 2022, **13**, 3269.
- 76 R. L. Wong, K. Paech and E. R. Williams, *Int. J. Mass Spectrom.*, 2004, **232**, 59.
- 77 D. Thölmann, D. S. Tonner and T. B. McMahon, *J. Phys. Chem.*, 1994, **98**, 2002.
- 78 R. C. Dunbar, *Mass Spectrom. Rev.*, 2004, **23**, 127.
- 79 T. Schindler, C. Berg, G. Niedner-Schatteburg and V. E. Bondybey, *Chem. Phys. Lett.*, 1996, **250**, 301.
- 80 P. D. Schnier, W. D. Price, R. A. Jockusch and E. R. Williams, *J. Am. Chem. Soc.*, 1996, **118**, 7178.
- 81 M. Sena and J. M. Riveros, *Rapid Commun. Mass Spectrom.*, 1994, **8**, 1031.
- 82 B. S. Fox, M. K. Beyer and V. E. Bondybey, *J. Phys. Chem. A*, 2001, **105**, 6386.
- 83 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 16 Revision A.03*, 2016.
- 84 E. M. Cunningham, T. Taxer, J. Heller, M. Ončák, C. van der Linde and M. K. Beyer, *Int. J. Mol. Sci.*, 2021, **22**, 6026.
- 85 J. Kim, S. Lee, S. J. Cho, B. J. Mhin and K. S. Kim, *J. Chem. Phys.*, 1995, **102**, 839.
- 86 K. Ohashi, J. Sasaki, G. Yamamoto, K. Judai, N. Nishi and H. Sekiya, *J. Chem. Phys.*, 2014, **141**, 214307.
- 87 B. M. Reinhard and G. Niedner-Schatteburg, *J. Phys. Chem. A*, 2002, **106**, 7988.
- 88 C.-K. Siu, Z.-F. Liu and J. S. Tse, *J. Am. Chem. Soc.*, 2002, **124**, 10846.
- 89 C. J. Johnson, L. C. Dzugan, A. B. Wolk, C. M. Leavitt, J. A. Fournier, A. B. McCoy and M. A. Johnson, *J. Phys. Chem. A*, 2014, **118**, 7590.

