

Catalysis Science & Technology

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Exploring the Mechanism of the Hydroboration of Alkenes by Amine-Boranes Catalysed by $[\text{Rh}(\text{Xantphos})]^+$

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/


Heather C. Johnson,^a Rebecca Torry-Harris,^a Laura Ortega,^b Robin Theron,^c J. Scott McIndoe^c and Andrew S. Weller^{a*}

The $[\text{Rh}(\text{Xantphos})]^+$ fragment acts as an effective catalyst for the hydroboration of the alkene TBE (tert-butyl ethene) using the amine-borane $\text{H}_3\text{B}\cdot\text{NMe}_3$ at low (0.5 mol%) catalyst loadings to give the linear product. Investigations into the mechanism using the initial rate method and labelling studies show that reductive elimination of the linear hydroboration product is likely the rate-limiting step at the early stages of catalysis, and that alkene and borane activation (insertion into a Rh-H bond and B-H oxidative addition) are reversible. The resting state of the system has also been probed using electrospray ionization mass spectrometry (ESI-MS) using the pressurised sample infusion (PSI) technique. This system is not as effective for hydroboration other alkenes such as 1-hexene, or using phosphine borane $\text{H}_3\text{B}\cdot\text{PCy}_3$, with decomposition or P-B bond cleavage occurring respectively.

Introduction

Hydroboration, the addition of a B-H bond across an unsaturated C-C bond, is a versatile methodology that affords organoboranes, from which subsequent functionalisation leads to products of use in organic synthesis.¹⁻⁴ Non-metal catalysed hydroboration generally yields the anti-Markovnikov product, whereas transition metal catalysts enable control over the regioselectivity of hydroboration. Such selectivity (*i.e.* linear *versus* branched products) has been shown to vary with different catalysts, alkenes and even reaction conditions.⁵⁻⁸ Historically metal-catalysed hydroborations have used three-coordinate boron substrates such as catechol (HBCat) or pinacol borane.^{2, 3} By contrast four-coordinate amine-boranes (prototypically $\text{H}_3\text{B}\cdot\text{NMe}_3$) have traditionally been used in uncatalysed hydroboration where N-B cleavage is proposed to afford a reactive trivalent BH_3 molecule,⁹ although iodine-induced hydroboration is proposed to operate *via* a three coordinate intermediate that retains the B-N bond.¹⁰ Amine-boranes have, instead, received much recent attention due to their potential as hydrogen storage systems and as precursors to oligomeric or polymeric B-N materials *via* dehydrocoupling,¹¹ and we,^{12, 13} alongside others,^{9, 11} have been exploring the role of the metal catalyst in these processes. Recognising that B-H oxidative cleavage from a bound sigma complex to form a metal boryl hydride (Scheme 1a) is closely related to the same mode of activation of a B-H bond at a metal in hydroboration (Scheme 1b), we reported in 2011 that the addition of the alkene tert-butylethene (TBE) to the sigma amine-borane complex $[\text{Rh}(\text{P}^t\text{Bu}_2^t\text{Bu})_2(\eta^2\text{-H}_3\text{B}\cdot\text{NMe}_3)][\text{BAR}^F_4]$ resulted in the formation of the linear hydroboration product $[\text{Rh}(\text{P}^t\text{Bu}_2^t\text{Bu})_2(\eta^2\text{-H}_2\text{B}(\text{CH}_2\text{CH}_2^t\text{Bu})\cdot\text{NMe}_3)][\text{BAR}^F_4]$ ¹⁴ $[\text{Ar}^F = 3,5\text{-(CF}_3)_2\text{C}_6\text{H}_4]$. $[\text{Rh}(\text{P}^t\text{Bu}_2^t\text{Bu})_2][\text{BAR}^F_4]$ also slowly (94 hrs, 5 mol%)

catalysed this process to form free $\text{H}_2\text{B}(\text{CH}_2\text{CH}_2^t\text{Bu})\cdot\text{NMe}_3$ (**I**), Scheme 1c.



Scheme 1. Rh-catalysed hydroboration using amine-borane (a) and catechol borane (b); Hydroboration of TBE using $\text{H}_3\text{B}\cdot\text{NMe}_3$, (b).


Kinetic experiments allowed for a mechanism to be proposed in which the hydroborated product inhibited catalytic turnover and reductive elimination of the product was also suggested to be slow.¹⁵ Independently, in 2012, a similar methodology using *N*-heterocyclic carbene-boranes and chiral Rh-based catalysts was reported for intramolecular hydroborations of alkenes.¹⁶ Very recently we briefly communicated that by using a $[\text{Rh}(\kappa^2_{\text{P,P}}\text{-Xantphos})]^+$ based catalyst,¹⁷ TBE can be hydroborated to give **I**. In the absence of this alkene, dehydrogenative homocoupling of the borane occurs (see Scheme 4), a process suggested to occur *via* the B-H activated

intermediate that is no longer intercepted by coordination of alkene.¹⁸ We now report in detail on this hydroboration, including kinetic data that support a proposed mechanism, as well as assessing the scope of this catalyst with regard to other alkenes and phosphine–boranes.

Results and Discussion


Preliminary stoichiometric and catalytic studies

Addition of excess TBE to the Rh(III) sigma–borane complex $[\text{Rh}(\kappa^3\text{P}_2\text{O}_2\text{-Xantphos})(\text{H})_2(\eta^1\text{-H}_3\text{B}\cdot\text{NMe}_3)][\text{BAR}^{\text{F}}_4]$ **1** resulted in the rapid formation (less than 5 minutes) of the Rh(I) complex $[\text{Rh}(\kappa^2\text{P}_2\text{-Xantphos})(\eta^2\text{-H}_2\text{B}(\text{CH}_2\text{CH}_2^t\text{Bu})\cdot\text{NMe}_3)][\text{BAR}^{\text{F}}_4]$ (**2**) as the sole metal-containing product (Scheme 2), presumably by initial hydrogenation of one equivalent of alkene to form a Rh(I) species, followed by hydroboration of another equivalent. The solid-state structure and NMR spectroscopic data for **2** have previously been communicated.¹⁸ In a similar manner, addition of trimethylvinyl silane to **1** gives the equivalent complex **3**, $[\text{Rh}(\kappa^2\text{P}_2\text{-Xantphos})(\eta^2\text{-H}_2\text{B}(\text{CH}_2\text{CH}_2\text{SiMe}_3)\cdot\text{NMe}_3)][\text{BAR}^{\text{F}}_4]$, in which $\text{H}_2\text{B}(\text{CH}_2\text{CH}_2\text{SiMe}_3)\cdot\text{NMe}_3$ (**II**) is bound to the metal centre.



Scheme 2 Formation of **2** (E = C) and **3** (E = Si). $[\text{BAR}^{\text{F}}_4]^-$ anions not shown.

Complex **3** was characterised by NMR spectroscopy, ESI-MS (electrospray ionisation mass spectrometry) and microanalysis, which together show similar analytical data to **2** and closely related $[\text{Rh}(\text{P}^i\text{Bu}_2^t\text{Bu})_2(\eta^2\text{-H}_2\text{B}(\text{CH}_2\text{CH}_2^t\text{Bu})\cdot\text{NMe}_3)][\text{BAR}^{\text{F}}_4]$.¹⁴ The alkyl borane binds to the metal centre through two sigma Rh–H–B interactions, evident by single ¹¹B quadrupolar-broadened signal at δ –6.54 in the ¹H NMR spectrum of relative integral 2 H, which collapses to an overlapping doublet of doublets (virtual triplet) on decoupling to ¹¹B [$J(\text{RhH}) = 36$ Hz, $J(\text{P}_{\text{trans}}\text{H}) = 36$ Hz]. Two, relative integral 2H, multiplets were observed at δ 1.17 and δ 0.78, indicating that the anti-Markovnikov (i.e. linear) product of hydroboration is bound to the metal centre. A ²⁹Si-¹H HMBC NMR experiment showed a correlation between silicon [δ (²⁹Si) 2.1] and the alkyl protons at δ 1.17, assigning these to those α to Si. The Xantphos methyl groups are observed as two separate environments (δ 1.73 and δ 1.67). In the ¹¹B NMR spectrum a broad resonance is observed at δ 37, typical for η^2 -coordination of an amine–borane to a Rh^I centre,^{19,20} which has shifted 45.7 ppm downfield from that in **1** (δ –8.7).¹⁸ Similar changes in ¹¹B chemical shift have been noted in related systems on moving between Rh(I) and Rh(III) oxidation states.^{19,21} The ³¹P{¹H} NMR spectrum shows a single environment δ 26.7 [d, $J(\text{RhP}) = 182$ Hz]. The solid-state structure of complex **3** supports the solution data (Figure 1), in particular a close Rh⋯B distance of 2.179(7) Å, which is the same within error to that found in **2**, 2.162(5) Å,¹⁸ and the formation of the linear hydroboration product. Complexes such as **2** and **3** are valence isoelectronic analogs of sigma alkane complexes,^{22–24} while related alkyl sigma amine–borane complexes have previously been prepared.²⁵



Scheme 3 Rhodium-catalysed hydroboration of TBE by $\text{H}_3\text{B}\cdot\text{NMe}_3$.





Figure 1 Solid-state structure of the cationic portion of **3**. Displacement ellipsoids are drawn at the 50% probability level. $[\text{BAR}^{\text{F}}_4]^-$ anion and all carbon-bound H atoms are omitted for clarity. Only the major component of the disordered NMe_3 and SiMe_3 groups are shown. Selected bond lengths (Å) and angles (°): Rh1–P1, 2.2398(18); Rh1–P2, 2.2670(17); Rh1–O1, 3.2342(73); Rh1–B1, 2.179(7); B1–N1, 1.603(4); P1–Rh1–P2, 98.23(6).

With complexes **2** and **3** in hand the catalytic hydroboration of $\text{H}_3\text{B}\cdot\text{NMe}_3$ with TBE was explored using these as precatalysts (Scheme 3). As previously reported,¹⁸ complex **2** (5 mol%) catalyses the complete conversion to $\text{H}_2\text{B}(\text{CH}_2\text{CH}_2^t\text{Bu})\cdot\text{NMe}_3$ (**I**) from TBE and $\text{H}_3\text{B}\cdot\text{NMe}_3$ within 3 hours. The catalysis was conducted with a 2:1 ratio of alkene: $\text{H}_3\text{B}\cdot\text{NMe}_3$ as the $\{\text{Rh}(\text{Xantphos})\}^+$ fragment has been reported to promote the slow dehydrogenative homocoupling of $\text{H}_3\text{B}\cdot\text{NMe}_3$ to form $[\text{Rh}(\kappa^2\text{P}_2\text{-Xantphos})(\eta^2\text{-H}_4\text{B}_2\cdot 2\text{NMe}_3)][\text{BAR}^{\text{F}}_4]$ (**4**) alongside **1** (Scheme 4),¹⁸ and a two-fold excess of alkene prevents the formation of **4** in detectable quantities (*vide infra*). During catalysis, complex **2** was the only observed resting state by ¹H and ³¹P{¹H} NMR spectroscopy.²⁶ The ¹H NMR spectrum of isolated **I** confirms anti-Markovnikov regioselectivity, with two, integral 2H, multiplets at δ 1.42 and δ 0.55 assigned to the methylene groups. The ¹¹B NMR spectrum shows a triplet at δ –0.83 [$J(\text{BH}) = 96$ Hz].¹⁴



Scheme 4 Formation of complex **4** from addition of excess $\text{H}_3\text{B}\cdot\text{NMe}_3$ to **2**. $[\text{BAR}^{\text{F}}_4]^-$ anions not shown.

Kinetic Studies

Given the promising rate of hydroboration of TBE with catalyst **2** to afford **I**, the catalyst loading was reduced to 0.5 mol%, relative to $\text{H}_3\text{B}\cdot\text{NMe}_3$. Under these conditions ($[\text{H}_3\text{B}\cdot\text{NMe}_3] = 0.19$ M, $[\text{TBE}] = 0.38$ M, 1,2- $\text{F}_2\text{C}_6\text{H}_4$ solvent), consumption of $\text{H}_3\text{B}\cdot\text{NMe}_3$ to yield **I** proceeded to 85% completion after 12 hours as monitored by ¹¹B

NMR spectroscopy, with the balance being made by unreacted $\text{H}_3\text{B}\cdot\text{NMe}_3$. Longer reaction times did not result in further reaction, suggesting either product inhibition and/or catalyst decomposition. When starting from precatalyst **3** a very similar overall temporal profile was observed, suggesting the identity of the initially bound amine–borane (**I** or **II**) does not affect the overall rate of catalysis.

The potential for product inhibition,¹⁴ and the parallel homocoupling reaction with excess $\text{H}_3\text{B}\cdot\text{NMe}_3$, suggested that the method of initial rates was most appropriate to probe the reaction orders with respect to substrates and the catalyst.²⁷ After 300 s at 5 mol% loading of **3** *ca.* 50% substrate conversion had occurred, while at 0.5 mol% this was now only *ca.* 10% conversion, making the lower loading suitable for study by the initial rate method. We further chose to study catalyst **3** as this would also give additional information as to the evolution of the likely resting states. Table 1 presents the data from this study, and Figure 2 presents some of these data in graphical format.

Comparison of entries 1, 2 and 3 (Figure 2a) show that the reaction is essentially first order in [TBE]. Entries 1 and 4 demonstrate a first order relationship in catalyst **3** (Figure 2b). Entries 2, 5 and 6 show that increasing the concentration of $[\text{H}_3\text{B}\cdot\text{NMe}_3]$ moves from an approximate first order relationship to an inhibition of catalysis at higher concentrations of amine–borane (Figure 2c), we presume as homocoupling to form **4** becomes competitive. In complex **4**, the diborane(4) is relatively strongly bound to the metal centre, remaining intact even with the addition of MeCN,¹⁸ and thus is unlikely to be as active in catalysis. Indeed, use of **4** as a catalyst (0.5 mol%) resulted in reduced turnover. Addition of excess product **I** (~70 equivalents, entry 7) results in a significant slowing of the initial rate – consistent with strong product inhibition, as observed to a lesser degree with $[\text{Rh}(\text{P}^i\text{Bu}_2\text{Bu})_2][\text{BAR}^F_4]$ system.¹⁴ Catalyst **3** and catalyst **2** operated at the same initial rate, within error (entries 1 and 8), suggesting that the identity of the bound primary borane (i.e. **I** or **II**) does not influence initial rate of turnover.

Table 1 Initial rates obtained from variation of concentration of **3**, $\text{H}_3\text{B}\cdot\text{NMe}_3$ and TBE, 295 K, 1,2- $\text{F}_2\text{C}_6\text{H}_4$ solvent. ^a Calculated from the pseudo zero-order region of the temporal evolution of **I** as measured by ^{11}B NMR spectroscopy over the first 300 s of catalysis. ^b With an additional 70 equiv. **I** at the start of catalysis. ^c Using $\text{D}_3\text{B}\cdot\text{NMe}_3$ instead of $\text{H}_3\text{B}\cdot\text{NMe}_3$.

Entry	[3] (10^{-4} M)	$[\text{H}_3\text{B}\cdot\text{NMe}_3]$ (M)	[TBE] (M)	Initial Rate ^a (10^{-5} M s ⁻¹)
1	9.5	0.19	0.38	6.81 ± 0.12
2	9.5	0.19	0.19	3.64 ± 0.27
3	9.5	0.19	0.76	12.98 ± 0.38
4	19.0	0.19	0.38	13.25 ± 0.56
5	9.5	0.38	0.19	7.41 ± 0.53
6	9.5	0.76	0.19	6.33 ± 0.40
7 ^b	9.5	0.19	0.38	2.07 ± 0.11
8	[2] 9.5	0.19	0.38	7.44 ± 0.64
9	9.5	0.19 ^c	0.38	5.09 ± 0.07

Hydroboration of TBE and $\text{H}_3\text{B}\cdot\text{NMe}_3$ catalysed using **3**




Figure 2. Initial rate experiments: (a) Variation [TBE]; (b) variation of [**3**]; (c) variation of $\text{H}_3\text{B}\cdot\text{NMe}_3$. See Table 1 for more details.

enables more information to be gleaned about possible resting states. At 0.5 mol% loading, the catalyst concentration is too low to be observable by NMR spectroscopy under the conditions used. However, at 5 mol% loading the rhodium-containing species can be probed by ^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy. The diagnostic, broad, hydride signals for **2** and **3** appear at similar chemical shifts in 1,2- $\text{F}_2\text{C}_6\text{H}_4$ solvent [δ – 6.85¹⁸ and δ – 6.54 respectively]. In the early stages of catalysis (~20% conversion), the ^1H NMR spectrum shows a mixture of **2** and **3**, evident by broad overlapping hydride peaks. As catalysis progresses, this broad overlapping resonance sharpens and **2** becomes the dominant species demonstrating that **I** displaces **II** in the resting state during catalysis. Under the conditions of excess $\text{H}_3\text{B}\cdot\text{NMe}_3$ (cf. entry 6), using 5 mol% **3** to enable monitoring by ^1H NMR spectroscopy, complex **4** grows in over time, whereas under conditions of excess TBE it not observed. This is consistent with the kinetic data that suggest removal from the system of active catalyst at high $[\text{H}_3\text{B}\cdot\text{NMe}_3]$, leading to inhibition.

The change in resting state from **3** to **2** has also been probed using electrospray ionization mass spectrometry (ESI–MS) using the pressurised sample infusion (PSI) technique.^{28–30} The particular advantage of this technique is that it allows for very high data density over a wide dynamic range, and is thus ideal for analysing evolving mixtures during catalysis. Figure 3 shows the temporal profile of the catalysis using [**3**]. This experiment was run at 15 mol%, which was determined to be the best conditions for the optimal (low) concentration necessary for PSI-ESI-MS. Immediately at the start of catalysis the resting state moves from **3** to **2**, consistent with the NMR experiments. These ESI–MS experiments also reveal the presence, at early stages of the reaction of three other species. The first is identified as $[\text{Rh}(\text{Xantphos})(\text{H}_3\text{B}\cdot\text{NMe}_3)]^+$, (m/z = 754.24; calc. 754.20), although we cannot comment on the precise structure: it could be a Rh(I) sigma-bound amine–borane complex, or a Rh(III) B–H activated hydrido–boryl. Both structural forms have precedent^{19,31} and are likely to be in equilibrium with one another.³² Indeed both have been calculated to be accessible, but thermodynamically unfavoured, compared with **2**.¹⁸ $[\text{Rh}(\text{Xantphos})(\text{H}_2)]^+$ (m/z = 683.15; calc. 683.11) and [**1**]⁺ are also observed, which we suggest both come from a small amount of [**1**]⁺ formed parallel with **4** during catalysis (Scheme 4). That we do not observe any of these species by ^1H NMR spectroscopy (hydride region) suggests that ESI-MS is particularly sensitive to their observation. These species decay at a very similar rate to [**3**]⁺, which suggests that the build-up of **I** during catalysis pushes any equilibria operating to favour of **2**. This observation is also consistent with product inhibition from initial rate experiments.





Figure 3. ESI-MS under PSI conditions²⁹ of the reaction of TBE with $\text{H}_3\text{B}\cdot\text{NMe}_3$ catalysed by **3**. Conditions: $\text{H}_3\text{B}\cdot\text{NMe}_3$, 0.006 M, TBE 0.013 M; [**3**] 0.001 M, 1,2- $\text{F}_2\text{C}_6\text{H}_4$. Under these conditions of concentration and experiment catalysis proceeded to 80% conversion.

Labelling Studies

Complex **3** (and **2**) are initially produced under conditions of excess TBE (Scheme 2), suggesting that the alkene does not bind competitively with **II** (or **I**), while the dependence of the rate law upon both $[\text{TBE}]$ and $[\text{H}_3\text{B}\cdot\text{NMe}_3]$ indicates that irreversible B–H oxidative addition prior to alkene coordination is not rate-determining. The potential for reversibility of the binding of both $\text{H}_3\text{B}\cdot\text{NMe}_3$ and TBE to the metal centre was further probed using $\text{D}_3\text{B}\cdot\text{NMe}_3$ instead of $\text{H}_3\text{B}\cdot\text{NMe}_3$ during catalysis. Initial rate experiments (Table 1, entry 9) showed a KIE of 1.34 ± 0.04 , consistent with irreversible B–H activation not being rate determining. However, due to the H/D exchange observed between the amine–borane and alkene (*vide infra*) the absolute magnitude of this measurement should be treated with a degree of caution.



Scheme 5 The products observed after 1 hour of catalysis using $\text{D}_3\text{B}\cdot\text{NMe}_3$. Conditions: $[\text{D}_3\text{B}\cdot\text{NMe}_3] = 0.19$ M, $[\text{TBE}] = 0.38$ M, 1,2- $\text{F}_2\text{C}_6\text{H}_4$, 0.5 mol% **3**.


After 1 hour of catalysis under conditions of excess alkene (28% conversion, Scheme 5) ^2H NMR spectroscopy showed incorporation of deuterium into the internal position of the free, unreacted, alkene (δ 5.89), while the corresponding signal in the ^1H NMR spectrum decreased by ca. 25% relative to the other alkene signals at δ 4.99 and 4.89. This demonstrates that H/D exchange occurs only occurs at the internal alkene proton. H/D exchange in free amine–borane was evidenced by the ^{11}B NMR spectrum that at early stages of catalysis showed a broad peak corresponding to $\text{D}_3\text{B}\cdot\text{NMe}_3$ and evolved with time to show significant signs of B–H coupling.³³ The final product *d*-**I** showed no H/D exchange α -to the borane, and $\sim 40\%$ H/D exchange at the β position (i.e. 60% D).

These data suggest that coordination of $\text{H}_3\text{B}\cdot\text{NMe}_3$, B–H activation, coordination and insertion of the alkene into the Rh–H bond are all reversible, to ultimately give the linear product. Moreover the lack of H/D exchange in this final product at the α -position, and a similar lack of exchange in the terminal positions of the free alkene, suggests that insertion to form the branched product is not occurring. We also suggest that hydride migration to the alkene, rather than boryl migration to form an intermediate such as **F** (Scheme 6), is by far dominant.³⁴ Intermediates such as **F** have been

postulated in dehydrogenative borylation reactions,^{5, 8, 35, 36} the products of which are not observed here. Although we cannot fully discount that boryl migration from **F** is reversible but the barrier to reductive elimination from **F** is high, we consider that this scenario is less likely based upon literature precedent.^{6–8, 35}

Bringing these data and observations together leads us to propose the catalytic pathway shown in Scheme 6 for the hydroboration of TBE using $\text{H}_3\text{B}\cdot\text{NMe}_3$ and **3**. This pathway is similar to that reported for using the $[\text{Rh}(\text{P}^i\text{Bu}_2\text{Pr})_2]^+$ catalyst system,¹⁴ as well as late transition metal hydroboration systems that use, for example, HBCat.^{3, 5, 6, 8}

The elementary steps in this cycle are thus: complex **2** does not react with TBE but undergoes reversible B–H activation with $\text{H}_3\text{B}\cdot\text{NMe}_3$, (i) and (ii), as shown by H/D exchange into free $\text{D}_3\text{B}\cdot\text{NMe}_3$ during the early phases of catalysis. Monitoring by ESI-MS shows a species consistent with **A** or **B** ($m/z = 754.24$) before bound **2** becomes the only species observed. TBE binding and insertion into Rh–H is reversible, (iii), as demonstrated by H/D exchange into the free alkene during catalysis. No branched product is observed³⁷ and no H/D exchange at the α -position of the linear product is measured, showing that insertion from **C** to form **D** (vi) is neither kinetically competent nor reversible. Insertion from **C** to give the linear intermediate **E** is reversible (iv), as there is significant (40%) H/D exchange at the β -position in the final product, as well as into the free alkene when $\text{D}_3\text{B}\cdot\text{NMe}_3$ is used, that suggests that β -H-elimination from **E** occurs. Overall these H/D labelling experiments suggest that reductive elimination (v) is the turnover-limiting step during the early stages of catalysis. As reductive elimination would be expected to have a small (close to unity) KIE, the modest measured value might reflect a system at equilibrium before the turnover limiting step (as postulated), i.e. an equilibrium isotope effect.³⁸




Scheme 6 Proposed mechanism for the catalytic hydroboration using data from the early phase of catalysis. $[\text{Rh}] = [\text{Rh}(\text{Xantphos})]^+$.

Hydroboration of alkenes other than TBE

Under stoichiometric conditions the hydroboration of alkenes other than TBE was explored. With the hindered alkene 2,3-dimethyl-2-

butene (2 equiv.) no evidence of hydroboration using $1/\text{H}_3\text{B}\cdot\text{NMe}_3$ was observed, with **1** remaining the major organometallic species in solution over an hour. After 24 hours, a 2:1 mixture of **1** and **4** were present, suggesting that the B–B homocoupling was occurring. Cyclohexene behaved similarly, as previously reported, with no hydroboration observed, instead it promotes loss of H_2 from **1**, driving the formation of **4**.¹⁸ To probe the possibility of, non-productive, coordination of cyclohexene at the metal centre, a mixture of $\text{D}_3\text{B}\cdot\text{NMe}_3$ (0.19 M), cyclohexene (0.38 M) and **3** (5 mol%, relative to $\text{D}_3\text{B}\cdot\text{NMe}_3$) was monitored by ^2H NMR spectroscopy. This showed, as well as deuterium incorporation into signals for cyclohexane that arise from deuteration of **1** (δ 1.58 and δ 1.43), incorporation of deuterium into the alkene signal (δ 5.89) after 90 minutes (Scheme 7). These data suggest that reversible alkene coordination and deuteride insertion can occur to give an intermediate similar to **E**; while the lack of hydroborated product suggests that the reductive elimination, as proposed to be the rate determining step for catalysis with TBE, is slow compared with overall dehydrogenative homocoupling from an intermediate **B** to form **4**. This is presumably related to the relative rates of reductive C–B coupling of primary and secondary alkyl–boryls, which in turn is likely related to barriers to re-orientation of the sp^3 alkyl and boryl groups prior to reductive coupling,³⁹ which is expected to be greater for locally bulkier substituents. Thus TBE undergoes hydroboration, while cyclohexene does not.




Scheme 7. Reversible deuterio-insertion with cyclohexene.

With 1-hexene, catalysis (5 mol% **3** relative to $\text{H}_3\text{B}\cdot\text{NMe}_3$) reached 37% conversion after 30 minutes yielding a product consistent with $\text{Me}(\text{CH}_2)_5\text{H}_2\text{B}\cdot\text{NMe}_3$, as shown by a triplet [$J(\text{HB}) = 91$ Hz] in the ^{11}B NMR spectrum at $\delta -1.4$, although we were not able to isolate this material pure and thus cannot comment on the linear:branched ratio. After 1 week, a maximum conversion of 63% is reached. However, significant decomposition of the catalyst was observed, for which we cannot definitively provide a structure derived from the spectroscopic data. Thus, the hydroboration of alkenes with **3** appears to work best with TBE, with other alkenes only of limited utility.

Hydroboration with phosphine-boranes


The addition of H_2 to $[\text{Rh}(\text{Xantphos})(\text{NBD})][\text{BAR}^{\text{F}}_4]$ (NBD = norbornadiene) in the presence of the tertiary phosphine–borane $\text{H}_3\text{B}\cdot\text{PCy}_3$ afforded $[\text{Rh}(\kappa^3_{\text{P,O,P}}\text{-Xantphos})(\text{H})_2(\eta^1\text{-H}_3\text{B}\cdot\text{PCy}_3)][\text{BAR}^{\text{F}}_4]$ (**5**) in quantitative yield by NMR spectroscopy (Scheme 8). Complex **5** was characterised *in situ* using NMR spectroscopy, and presents very similar data to the analogous complex **1**.¹⁸ It is also related to the sigma phosphine–borane complex $[\text{Ru}(\text{Xantphos})(\text{H})(\text{PPh}_2\text{H})(\text{H}_3\text{B}\cdot\text{PPh}_2\text{H})][\text{BAR}^{\text{F}}_4]$.⁴⁰ The ^1H NMR spectrum shows 3 hydride environments in a 3:1:1 ratio, consistent with the phosphine–borane bound in an η^1 fashion and two mutually *cis* hydrides: $\delta -1.42$ [br, BH_3 , sharpens on ^{11}B decoupling], $\delta -14.62$ [br dtd, RhH] and $\delta -19.13$ [dtd, RhH]. Two ^{31}P environments

are observed in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum at δ 41.8 [d, $J(\text{RhP}) = 114$ Hz] and δ 20.1 (br, PCy_3) in a 2:1 ratio respectively. Attempts to crystallise **5** were unsuccessful, resulting in decomposition.



Scheme 8 Formation of complex **5**. $[\text{BAR}^{\text{F}}_4]^-$ anion not shown.

Addition of 2 equivalents of TBE to pale yellow **5** resulted in the formation of a dark green solution of a new compound formulated as $[\text{Rh}(\kappa^2_{\text{P,P}}\text{-Xantphos})(\eta^2\text{-H}_3\text{B}\cdot\text{PCy}_3)][\text{BAR}^{\text{F}}_4]$ (**6**), Scheme 9, in quantitative yield by NMR spectroscopy. Removal of volatiles allowed the isolation of **6** as a dark green solid. The NMR data for **6** are consistent with η^2 binding of the phosphine–borane; in the ^1H NMR spectrum, a quadrupolar broadened, integral 3H signal is observed at $\delta -2.38$, which sharpens on ^{11}B decoupling, while the ^{11}B NMR spectrum shows a broad signal at $\delta -3.0$. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum is also consistent with a Rh(I) phosphine–borane complex, with two signals observed at δ 28.9 [d, $J(\text{RhP}) = 190$ Hz] and δ 17.3 (br, PCy_3). These data are consistent with those reported for other $[\text{Rh}(\text{chelating phosphine})(\text{H}_3\text{B}\cdot\text{PR}_3)]^+$ complexes.⁴¹ Addition of H_2 (4 atm) to a CD_2Cl_2 solution of **6** reforms complex **5** in quantitative yields by NMR spectroscopy. Degassing a CD_2Cl_2 solution of **5** and placing under static vacuum for 4 hours resulted in an approximately 1:1 ratio of **5**:**6**, suggestive of an equilibrium between the two species. Interestingly, for the $[\text{Rh}(\text{Xantphos})]^+$ fragment we cannot isolate, or observe by NMR spectroscopy, the equivalent Rh(I) $\text{H}_3\text{B}\cdot\text{NMe}_3$ complex to **6**, as **4** forms instead from homocoupling.¹⁸



Scheme 9 Formation of **6**, **7** and **8**. $[\text{BAR}^{\text{F}}_4]^-$ anions are not shown.

The ability of **6** to mediate hydroboration was probed by addition of excess (2.5 equiv.) TBE in $1,2\text{-F}_2\text{C}_6\text{H}_4$ solvent, by addition of the alkene to *in situ* generated **6**, Scheme 9. After 45 minutes a new peak is apparent in the ^1H NMR spectrum at $\delta -5.58$ that is assigned to an Rh–HB interaction, consistent with the slow formation of $[\text{Rh}(\kappa^2_{\text{P,P}}\text{-Xantphos})(\eta^2\text{-H}_2\text{B}(\text{CH}_2\text{CH}_2\text{tBu})\text{-PCy}_3)][\text{BAR}^{\text{F}}_4]$ (**7**), similar to **2** and **3**. After 16 hours the ratio of **7** had increased relative to **6** (~5:1 **7**:**6**). However small amounts (*ca.* 5% by $^{31}\text{P}\{^1\text{H}\}$ NMR

spectroscopy) of a parallel product resulting from P–B cleavage $[\text{Rh}(\kappa^3_{\text{P,O,P}}\text{-Xantphos})(\text{PCy}_3)]^+[\text{BAR}^{\text{F}}_4]^-$ (**8**) were also observed by $^{31}\text{P}\{^1\text{H}\}$ NMR spectroscopy at δ 61.3 [dt, $J(\text{RhP}) = 192$, $J(\text{PP}) = 34$ Hz] and δ 37.8 [dd, $J(\text{RhP}) = 155$, $J(\text{PP}) = 34$ Hz]. P–B bond cleavage has been noted previously during metal-catalysed dehydrocoupling of phosphine-boranes.^{40,42,43} After a further 12 hours all of **6** was consumed, but a greater proportion of **8** (*ca.* 33%) was also present. Recrystallisation of the reaction mixture after several hours afforded a small crop of green crystals of **7** suitable for X-ray diffraction, and although the resulting data quality was poor and only the gross connectivity can thus be discussed, the solid-state structure of **7** suggests anti-Markovnikov hydroboration, as with **2** and **3** (see ESI). The bulk composition could not be reliably determined by NMR spectroscopy as the alkyl region of the ^1H NMR spectrum is dominated by the cyclohexyl peaks from the mixture of **6** and **8**.

From such mixtures, several orange crystals of **8** also grew, confirming the solid-state structure of **8** (Figure 4) as a square planar Rh(I) complex [sum of angles around Rh = 361.5°] with Xantphos coordinated in a *mer* manner, similar to the closely related $[\text{Rh}(\kappa^3_{\text{P,O,P}}\text{-Xantphos})(\text{PCyp}_3)]^+[\text{BAR}^{\text{F}}_4]^-$ (Cyp = cyclopentyl),⁴⁴ and $[\text{Rh}(\kappa^3_{\text{P,O,P}}\text{-Xantphos})(\text{CO})]^+[\text{BF}_4]^-$.⁴⁵ Complex **8** has been independently synthesised from addition of PCy_3 to **2** (see Experimental).




Figure 4 Solid state structure of the cationic portion of **8**. $[\text{BAR}^{\text{F}}_4]^-$ anion and H atoms are omitted for clarity. Displacement ellipsoids are drawn at the 50% probability level. Selected bond lengths (Å) and angles ($^\circ$): Rh1–P1, 2.2854(7); Rh1–P2, 2.2923(7); P3–Rh1, 2.2611(7); Rh1–O1, 2.2395(19); P2–Rh1–P1, 159.70(3); P3–Rh1–O1, 170.44(6).

Conclusions

We have described that the $[\text{Rh}(\text{Xantphos})]^+$ fragment acts as an effective catalyst for the hydroboration of TBE using the amine–borane $\text{H}_3\text{B}\cdot\text{NMe}_3$ at low (0.5 mol%) catalyst loadings. Investigations into the mechanism using the initial rate method and labelling studies show that reductive elimination of the linear hydroboration product is likely the rate-limiting step at the early stages of catalysis. This system is not as effective for other alkenes such as 1-hexene, or using the phosphine–borane $\text{H}_3\text{B}\cdot\text{PCy}_3$; with decomposition or P–B bond cleavage occurring respectively.

Experimental Section

All manipulations, unless otherwise stated, were performed under an argon atmosphere using standard Schlenk and glove-box techniques. Glassware was oven dried at 130°C overnight and flamed under vacuum prior to use. Pentane, hexanes, CH_2Cl_2 and MeCN were dried using a Grubbs type solvent purification system (MBraun SPS-

800) and degassed by successive freeze-pump-thaw cycles. $1,2\text{-F}_2\text{C}_6\text{H}_4$ (pre-treated with alumina) and CD_2Cl_2 were dried over CaH_2 , vacuum distilled and stored over 3 Å molecular sieves. $\text{H}_3\text{B}\cdot\text{NMe}_3$ was purchased from Aldrich and sublimed prior to use (5×10^{-2} Torr, 298 K). Cyclohexene and 3,3-dimethyl-1-butene (TBE) were purchased from Aldrich, dried over sodium, vacuum distilled and stored over 3 Å molecular sieves. $[\text{Rh}(\text{Xantphos})(\text{NBD})]^+[\text{BAR}^{\text{F}}_4]^-$ was prepared by the literature method.⁴⁶ NMR spectra were recorded on a Bruker AVIII-500 spectrometer at room temperature, unless otherwise stated. In $1,2\text{-F}_2\text{C}_6\text{H}_4$, ^1H NMR spectra were pre-locked to a sample of C_6D_6 (25%) and $1,2\text{-F}_2\text{C}_6\text{H}_4$ (75%) and referenced to the centre of the downfield solvent multiplet, $\delta = 7.07$. ^{31}P and ^{11}B NMR spectra were referenced against 85% H_3PO_4 (external) and $\text{BF}_3\cdot\text{OEt}_2$ (external) respectively. Chemical shifts (δ) are quoted in ppm and coupling constants (J) in Hz. ESI-MS were recorded on a Bruker MicroTOF instrument interfaced with a glove-box,²⁸ or using the PSI-ESI technique as described previously,^{29,30} and in detail below. Microanalyses were performed by Elemental Microanalysis Ltd.

ESI-MS reaction monitoring using pressurized sample infusion.

A Schlenk flask under nitrogen containing **3** (4.7 mg, 0.0028 mmol) and $\text{H}_3\text{B}\cdot\text{NMe}_3$ (1.4 mg, 0.019 mmol) was pressurized to 1.5 psi using 99.998% purity argon gas and connected to the mass spectrometer via a short length of PEEK tubing. A solution of TBE (4.8 μL , 0.038 mmol) in $1,2\text{-F}_2\text{C}_6\text{H}_4$ (3 mL) was injected into the pressurized Schlenk flask through a septum and collection on the mass spectrometer was initiated. Mass spectra were collected on a Micromass Q-ToF mass spectrometer in positive ion mode using pneumatically assisted electrospray ionization: capillary voltage, 2900 V; sample cone voltage, 15 V; extraction voltage, 0.5 V; Source temperature, 92°C ; desolvation temperature, 192°C ; cone gas flow, 100 L/h; desolvation gas flow, 200 L/h; collision voltage, 2 V; MCP voltage, 2400 V. No smoothing of the data was performed. Aliquots of the reaction mixture were removed via syringe during the reaction for analysis by ^{11}B NMR spectroscopy.

General procedure for catalytic hydroboration. The alkene and $1,2\text{-F}_2\text{C}_6\text{H}_4$ (0.6 mL) were mixed in a Young's NMR tube and transferred to a new NMR tube containing **2** or **3** and $\text{H}_3\text{B}\cdot\text{NMe}_3$. The samples were immediately frozen in liquid N_2 , and monitored *in situ* by ^{11}B NMR spectroscopy on warming. See Table 1 for more details of relative concentrations.

Synthesis and characterisation of new complexes. Synthesis of $[\text{Rh}(\kappa^2_{\text{P,P}}\text{-Xantphos})(\eta^2\text{-H}_2\text{B}(\text{CH}_2\text{CH}_2\text{SiMe}_3)\cdot\text{NMe}_3)]^+[\text{BAR}^{\text{F}}_4]^-$ (**3**).

$[\text{Rh}(\text{Xantphos})(\text{nbd})]^+[\text{BAR}^{\text{F}}_4]^-$ (100 mg, 0.06 mmol) and $\text{H}_3\text{B}\cdot\text{NMe}_3$ (4.4 mg, 0.06 mmol) were dissolved in $1,2\text{-F}_2\text{C}_6\text{H}_4$ in a Young's flask, the contents immediately frozen in liquid N_2 , and the argon headspace replaced with H_2 (*ca.* 4 atm), yielding **1** *in situ* upon warming to room temperature. The flask was degassed (3 freeze-pump-thaw cycles), opened to an argon atmosphere, and TMVS (40 μL , 0.272 mmol) was added. The solution turned from pale yellow to dark green. After 10 minutes, the volatiles were removed *in vacuo*, and the solid washed twice with pentane (2×5 mL) with sonication. The solid was dried *in vacuo*, affording a blue/green powder, mass 86 mg (82% yield). Crystals suitable for X-ray diffraction were grown from recrystallisation from $1,2\text{-F}_2\text{C}_6\text{H}_4$ and pentane at -30°C . ^1H NMR (500 MHz, $1,2\text{-F}_2\text{C}_6\text{H}_4$): δ 8.33 (s, 8H, $[\text{BAR}^{\text{F}}_4]^-$), 7.69 (s, 4H, $[\text{BAR}^{\text{F}}_4]^-$), 2.61 (s, 9H, NMe_3), 1.73 (s, 3H, Xantphos CH_3), 1.67 (s, 3H, Xantphos CH_3), 1.17 (m, 2H, CH_2SiMe_3), 0.78 (br m, 2H, $\text{CH}_2\text{BH}_2\text{NMe}_3$), 0.06 (s, 9H, SiMe_3), -6.54 (br, 2H, BH_2). The peak at -6.54 ppm sharpens into an overlapping doublet of doublets (virtual triplet) upon decoupling to ^{11}B [$J(\text{RhH}) = 36$ and $J(\text{PH}) = 36$ Hz]. Signals from the Xantphos

aryl ligand were not observed, presumably obscured by the solvent. ^{29}Si - ^1H HMBC NMR (500 MHz, 1,2- $\text{F}_2\text{C}_6\text{H}_4$): correlation observed between silicon at δ 2.1 and protons at δ 1.17 and 0.06. $^{31}\text{P}\{^1\text{H}\}$ NMR (202 MHz, 1,2- $\text{F}_2\text{C}_6\text{H}_4$): δ 26.7 [d, $J(\text{RhP}) = 182$ Hz]. ^{11}B NMR (160 MHz, 1,2- $\text{F}_2\text{C}_6\text{H}_4$): δ 37 (br, BH_2), -6.2 (s, $[\text{BAR}^{\text{F}}_4]$). ESI-MS (1,2- $\text{F}_2\text{C}_6\text{H}_4$, 60 °C, 4.5 kV): m/z 854.28 $[\text{M}]^+$ (calc. 854.28). Peak displays the expected isotopic pattern. Elemental microanalysis: Calc. $\text{RhP}_2\text{OC}_{79}\text{H}_{68}\text{B}_2\text{F}_{24}\text{NSi}$ (1717.94 gmol^{-1}): C, 55.23; H, 3.99; N, 0.82. Found: C, 55.16; H, 4.03; N, 0.88.

Synthesis of $[\text{Rh}(\kappa^3_{P,O,P}\text{-Xantphos})(\text{H})_2(\eta^1\text{-H}_3\text{B-PCy}_3)][\text{BAR}^{\text{F}}_4]$ (5**).** $[\text{Rh}(\text{Xantphos})(\text{nbd})][\text{BAR}^{\text{F}}_4]$ (20 mg, 0.01 mmol) and $\text{H}_3\text{B-PCy}_3$ (2.5 mg, 0.01 mmol) were dissolved in 1,2- $\text{F}_2\text{C}_6\text{H}_4$ in a high pressure NMR tube, the contents immediately frozen in liquid N_2 , and the argon headspace replaced with H_2 (ca. 4 atm), yielding **5** *in situ* upon warming to room temperature and shaking. **5** could not be isolated due to loss of dihydrogen upon removal from the H_2 atmosphere. Attempts to recrystallise under H_2 resulted in impure oil, as measured by NMR spectroscopy. However, the following NMR spectroscopic data were obtained from the hydrogenation of pre-formed **6**. ^1H NMR (500 MHz, CD_2Cl_2): δ 8.10 – 7.27 (m, 26H, Xantphos aryl signals), 7.73 (s, 8H, $[\text{BAR}^{\text{F}}_4]$), 7.56 (s, 4H, $[\text{BAR}^{\text{F}}_4]$), 2.05 – 0.81 (m, 39H, overlapping Cy and Xantphos CH_3 signals), -1.63 (br, 3H, BH_3), -14.80 (br, 1H, RhH), -19.40 (br, 1H, RhH). ^1H NMR (500 MHz, CD_2Cl_2 , 200 K, selected data): δ 1.87 (s, 3H, Xantphos CH_3 signal), 1.47 (s, 3H, Xantphos CH_3 signal), -14.34 (br m, 1H, RhH), -19.40 [br dtd, $J(\text{RhH}) = 26$, $J(\text{PH}) = 14$, $J(\text{HH}) = 7$ Hz, RhH]. $^{31}\text{P}\{^1\text{H}\}$ NMR (202 MHz, CD_2Cl_2): δ 41.8 [d, $J(\text{RhP}) = 116$ Hz], 20.2 (br, PCy_3). ^{11}B NMR (160 MHz, CD_2Cl_2): δ -6.6 (s, $[\text{BAR}^{\text{F}}_4]$), -43.5 (br, BH_3).

Synthesis of $[\text{Rh}(\kappa^2_{P,P}\text{-Xantphos})(\eta^2\text{-H}_3\text{B-PCy}_3)][\text{BAR}^{\text{F}}_4]$ (6**).** **5** (55 mg, 0.03 mmol) was formed *in situ* under H_2 (4 atm) in a Young's crystallisation flask. The flask was degassed (3 freeze-pump-thaw cycles), opened to an argon atmosphere, and TBE (8 μL , 0.06 mmol) was added. The solution turned from pale yellow to dark green. After 5 minutes, the volatiles were removed *in vacuo* to prevent onward reactivity, and washed twice with pentane (3 mL) with sonication. The resulting oily solid was redissolved in the minimum amount of 1,2- $\text{F}_2\text{C}_6\text{H}_4$ and layered with pentane, affording green needle-like crystals (not suitable for X-ray diffraction) at -30 °C. mass 40 mg (73% yield) ^1H NMR (500 MHz, CD_2Cl_2): δ 7.73 (s, 8H, $[\text{BAR}^{\text{F}}_4]$), 7.56 (s, 4H, $[\text{BAR}^{\text{F}}_4]$), 7.64 – 6.40 (m, 26H, Xantphos aryl signals), 1.86 – 1.08 (m, 39H, PCy_3 and Xantphos CH_3 signals), -2.38 (br, 3H, BH_3). Upon decoupling to ^{11}B , the signal at -2.38 sharpens. $^{31}\text{P}\{^1\text{H}\}$ NMR (202 MHz, CD_2Cl_2): δ 28.9 [d, $J(\text{RhP}) = 190$ Hz], 17.3 (br, PCy_3). ^{11}B NMR (160 MHz, CD_2Cl_2): δ -3.0 (br, BH_3), -6.6 (s, $[\text{BAR}^{\text{F}}_4]$). ESI-MS (1,2- $\text{F}_2\text{C}_6\text{H}_4$, 60 °C, 4.5 kV) was attempted but decomposition resulted under these conditions. Elemental microanalysis: Calc. $\text{RhP}_3\text{OC}_{89}\text{H}_{80}\text{B}_2\text{F}_{24}$ (1839.03 gmol^{-1}): C, 58.13; H, 4.38. Found: C, 57.43; H, 4.59.

Attempted synthesis of $[\text{Rh}(\kappa^2_{P,P}\text{-Xantphos})(\eta^2\text{-H}_2\text{B}(\text{CH}_2\text{CH}_2^t\text{Bu})\text{-PCy}_3)][\text{BAR}^{\text{F}}_4]$ (7**).**

Complex **5** (0.02 mmol) was formed *in situ* under H_2 (4 atm) in a Young's crystallisation flask as described above. The flask was degassed (3 freeze-pump-thaw cycles), opened to an argon atmosphere, and TBE (7 μL , 0.05 mmol) was added. The solution turned from pale yellow to dark green, as complex **6** is formed immediately. Upon leaving the mixture for 45 minutes, a mixture of **6** and **7** are present. Standing for longer periods (12 hours) resulted in a decreased amount of **6**, and a new

complex, **8**, was now observed (see below for an independent synthesis of **8**). Due to this mixture, our attempts to isolate pure **7** were unsuccessful. A small number of green single crystals suitable for diffraction were obtained by leaving the reaction mixture for several hours to form a mixture of **6** and **7**, layering the solution with pentane and storing at -30 °C. Analysis by NMR spectroscopy of these crystals dissolved in CD_2Cl_2 indicated a mixture of **6** and **7**. Due to the mixture, only selected NMR data for **7** are available as the alkyl and aryl regions are obscured by **6** signals. ^1H NMR (500 MHz, CD_2Cl_2): δ -5.92 (br, Rh- H_2B). The signal sharpens into an overlapping doublet of doublets (virtual triplet) upon decoupling to ^{11}B [$J(\text{RhH}) = 33$ and $J(\text{PH}) = 33$ Hz]. $^{31}\text{P}\{^1\text{H}\}$ NMR (202 MHz, CD_2Cl_2): δ 29.5 [d, $J(\text{RhP}) = 190$ Hz], 8.8 (br, PCy_3). ^{11}B NMR (160 MHz, CD_2Cl_2): δ 18.5 (br, BH_3), -6.6 (s, $[\text{BAR}^{\text{F}}_4]$). ESI-MS (1,2- $\text{F}_2\text{C}_6\text{H}_4$, 60 °C, 4.5 kV) was attempted but decomposition resulted under these conditions. Upon repeating the reaction a small number of orange crystals of **8** formed which could be mechanically separated from green **6** and **7**, whose spectroscopic data match that from independently synthesised **8** (see below).

Synthesis of $[\text{Rh}(\kappa^3_{P,O,P}\text{-Xantphos})(\text{PCy}_3)][\text{BAR}^{\text{F}}_4]$ (8**).**

1,2- $\text{F}_2\text{C}_6\text{H}_4$ (0.4 mL) was added to **2** (17 mg, 0.010 mmol) and PCy_3 (0.011 mmol) in a small Schlenk flask. The colour changed from blue to orange immediately on mixing. The solvent was removed *in vacuo*, and the remaining orange oil was washed three times with pentane (2 mL) with sonication. Stirring under vacuum afforded an orange powder, mass 13 mg (71% yield). ^1H NMR (500 MHz, CD_2Cl_2): δ 7.72 (s, 8H, $[\text{BAR}^{\text{F}}_4]$), 8.05 – 7.71 (m, 30H, Xantphos aryl signals and second $[\text{BAR}^{\text{F}}_4]$ peak), 1.66 (s, 6H, Xantphos methyl signals), 1.84 – 0.50 (m, 33H, PCy_3). $^{31}\text{P}\{^1\text{H}\}$ NMR (202 MHz, CD_2Cl_2): δ 55.4 [dt, $J(\text{RhP}) = 192$, $J(\text{PP}) = 34$, PCy_3], 31.6 [dt, $J(\text{RhP}) = 155$, $J(\text{PP}) = 34$ Hz, Xantphos]. ESI-MS (1,2- $\text{F}_2\text{C}_6\text{H}_4$, 60 °C, 4.5 kV): m/z 961.33 $[\text{M}]^+$ (calc. 961.33). Peak displays the expected isotopic pattern.

Acknowledgements

NSERC (Discovery and Discovery Accelerator Supplements) for operational funding and CFI and BCKDF for infrastructure (JSM); The University of Victoria for graduate scholarships (RT); the Royal Society for an International Exchange Award (ASW and JSM), the EPSRC for a DTG award (HCJ).

Notes and references

^a Department of Chemistry, University of Oxford, Mansfield Road, Oxford, UK, OX1 3TA.

^b Visiting Student from Departamento de Química Inorgánica, Instituto de Investigaciones Químicas (IIQ), Universidad de Sevilla, Consejo Superior de Investigaciones Científicas, Avda. Américo Vespucio 49, 41092 Sevilla, Spain

^c Department of Chemistry, University of Victoria, P.O. Box 3065, Victoria, BC V8W 3V6, Canada.

Electronic Supplementary Information (ESI) available: X-ray crystallographic data collection and refinement details. Crystallographic data have been deposited with the Cambridge Crystallographic Data

Center (CCDC), CCDC 1001382 (3) 1001383 (8), can be obtained via www.ccdc.cam.ac.uk/data_request/cif See DOI: 10.1039/b000000x/

1. D. Mannig and H. Noth, *Angew. Chem.-Int. Edit. Engl.*, 1985, **24**, 878-879.
2. J. Hartwig, *Organotransition metal chemistry: from bonding to catalysis*, University Science Books, Sausalito, 2010.
3. C. M. Crudden and D. R. Edwards, *Eur. J. Org. Chem.*, 2003, **2003**, 4695-4712.
4. T. Davan, E. W. Corcoran and L. G. Sneddon, *Organometallics*, 1983, **2**, 1693-1694.
5. K. Burgess, W. A. Van der donk, S. A. Westcott, T. B. Marder, R. T. Baker and J. C. Calabrese, *J. Am. Chem. Soc.*, 1992, **114**, 9350-9359.
6. D. A. Evans, G. C. Fu and B. A. Anderson, *J. Am. Chem. Soc.*, 1992, **114**, 6679-6685.
7. D. R. Edwards, Y. B. Hleba, C. J. Lata, L. A. Calhoun and C. M. Crudden, *Angew. Chem. Int. Ed.*, 2007, **46**, 7799-7802.
8. J. M. Brown and G. C. Lloyd Jones, *J. Am. Chem. Soc.*, 1994, **116**, 866-878.
9. A. Staubitz, A. P. M. Robertson, M. E. Sloan and I. Manners, *Chem. Rev.*, 2010, **110**, 4023-4078.
10. M. Scheideman, G. Wang and E. Vedejs, *J. Am. Chem. Soc.*, 2008, **130**, 8669-8676.
11. E. M. Leitao, T. Jurca and I. Manners, *Nat Chem*, 2013, **5**, 817-829.
12. L. J. Sewell, M. A. Huertos, M. E. Dickinson, A. S. Weller and G. C. Lloyd-Jones, *Inorg. Chem.*, 2013, **52**, 4509-4516.
13. L. J. Sewell, G. C. Lloyd-Jones and A. S. Weller, *J. Am. Chem. Soc.*, 2012, **134**, 3598-3610.
14. L. J. Sewell, A. B. Chaplin and A. S. Weller, *Dalton Trans.*, 2011, **40**, 7499-7501.
15. In this report we did not describe H/D labelling studies. Subsequent studies using D₃B-NMe₃ showed that alkene coordination and deuteride insertion to form the linear product where all reversible, as found in this current study and consistent with reductive elimination being the rate-determining step at the early stages of catalysis. L. J. Sewell, DPhil Thesis, University of Oxford, 2013.
16. M. Toure, O. Chuzel and J.-L. Parrain, *J. Am. Chem. Soc.*, 2012, **134**, 17892-17895.
17. M. Kranenburg, Y. E. M. van der Burgt, P. C. J. Kamer, P. W. N. M. van Leeuwen, K. Goubitz and J. Fraanje, *Organometallics*, 1995, **14**, 3081-3089.
18. H. C. Johnson, C. L. McMullin, S. D. Pike, S. A. Macgregor and A. S. Weller, *Angew. Chem. Int. Ed.*, 2013, **52**, 9776-9780.
19. T. M. Douglas, A. B. Chaplin, A. S. Weller, X. Yang and M. B. Hall, *J. Am. Chem. Soc.*, 2009, **131**, 15440-15456.
20. G. Alcaraz and S. Sabo-Etienne, *Coord. Chem. Rev.*, 2008, **252**, 2395-2409.
21. A. B. Chaplin and A. S. Weller, *Inorg. Chem.*, 2010, **49**, 1111-1121.
22. W. H. Bernskoetter, C. K. Schauer, K. I. Goldberg and M. Brookhart, *Science*, 2009, **326**, 553-556.
23. M. D. Walter, P. S. White, C. K. Schauer and M. Brookhart, *J Am Chem Soc*, 2013, **135**, 15933-15947.
24. S. D. Pike, A. L. Thompson, A. G. Algarra, D. C. Apperley, S. A. Macgregor and A. S. Weller, *Science*, 2012, **337**, 1648-1651.
25. Y. Kawano, K. Yamaguchi, S. Y. Miyake, T. Kakizawa and M. Shimoi, *Chem.-Eur. J.*, 2007, **13**, 6920-6931.
26. Related Rh(PR₃)₂(η²-carborane) complexes have been reported as the resting states during cage B-H hydroboration reactions with alkenes: J. D. Hewes, C. W. Kreimendahl, T. B. Marder and M. F. Hawthorne, *J. Am. Chem. Soc.*, 1984, **106**, 5757-5759; E. Molinos, S. K. Brayshaw, G. Kociok-Köhn and A. S. Weller, *Organometallics*, 2007, **26**, 2370-2382.
27. R. B. Jordan, *Reaction Mechanisms of Inorganic and Organometallic Systems*, Oxford University Press, 2007.
28. A. T. Lubben, J. S. McIndoe and A. S. Weller, *Organometallics*, 2008, **27**, 3303-3306.
29. K. L. Vikse, M. P. Woods and J. S. McIndoe, *Organometallics*, 2010, **29**, 6615-6618.
30. L. P. E. Yunker, R. L. Stoddard and J. S. McIndoe, *J. Mass Spectrom.*, 2014, **49**, 1-8.
31. A. B. Chaplin and A. S. Weller, *Angew. Chem. Int. Ed.*, 2010, **49**, 581-584.
32. A. G. Algarra, L. J. Sewell, H. C. Johnson, S. A. Macgregor and A. S. Weller, *Dalton Trans*, 2014.
33. Reliable integration to quantify the degree of H/D exchange in the borane was frustrated by the quadrupolar-broadened signals.
34. R. T. Baker, J. C. Calabrese, S. A. Westcott, P. Nguyen and T. B. Marder, *J. Am. Chem. Soc.*, 1993, **115**, 4367-4368.
35. D. E. Kadlecsek, P. J. Carroll and L. G. Sneddon, *J. Am. Chem. Soc.*, 2000, **122**, 10868-10877.
36. E. Molinos, S. K. Brayshaw, G. Kociok-Köhn and A. S. Weller, *Organometallics*, 2007, **26**, 2370-2382.
37. T. C. Morrill, C. A. D'Souza, L. Yang and A. J. Sampognaro, *J. Org. Chem.*, 2002, **67**, 2481-2484.
38. E. M. Simmons and J. F. Hartwig, *Angew. Chem. Int. Ed.*, 2012, **51**, 3066-3072.
39. R. Ghosh, T. J. Emge, K. Krogh-Jespersen and A. S. Goldman, *J Am Chem Soc*, 2008, **130**, 11317-11327.
40. A. E. W. Ledger, C. E. Ellul, M. F. Mahon, J. M. J. Williams and M. K. Whittlesey, *Chem-Eur J*, 2011, **17**, 8704-8713.
41. M. A. Huertos and A. S. Weller, *Chem Sci*, 2013, **4**, 1881.
42. M. A. Huertos and A. S. Weller, *Chem Commun*, 2012, **48**, 7185.
43. T. N. Hooper, M. A. Huertos, T. Jurca, S. D. Pike, A. S. Weller and I. Manners, *Inorg. Chem.*, 2014.
44. R. Dallanegra, A. B. Chaplin and A. S. Weller, *Organometallics*, 2012, **31**, 2720-2728.
45. A. J. Sandee, L. A. van der Veen, J. Reek, P. Kamer, M. Lutz, A. L. Spek and P. van Leeuwen, *Angew. Chem. Int. Ed.*, 1999, **38**, 3231-3235.
46. R. J. Pawley, G. L. Moxham, R. Dallanegra, A. B. Chaplin, S. K. Brayshaw, A. S. Weller and M. C. Willis, *Organometallics*, 2010, **29**, 1717-1728.



5x2mm (600 x 600 DPI)