

Cite this: *RSC Adv.*, 2014, 4, 30462Received 5th February 2014
Accepted 27th June 2014

DOI: 10.1039/c4ra01027b

www.rsc.org/advances

Click chemistry facilitates direct labelling and super-resolution imaging of nucleic acids and proteins†

Anika Raulf,^a Christoph K. Spahn,^a Patrick J. M. Zessin,^a Kieran Finan,^b
Stefan Bernhardt,^c Alexander Heckel^c and Mike Heilemann^{*a}

We demonstrate high-density labelling of cellular DNA and RNA using click chemistry and perform confocal and super-resolution microscopy. We visualize the crescent and ring-like structure of densely packed RNA in nucleoli. We further demonstrate click chemistry with unnatural amino acids for super-resolution imaging of outer-membrane proteins of *E. coli*.

Various fluorescence microscopy techniques which can bypass the spatial resolution limit have been developed in recent years. Commonly summarized as “super-resolution microscopy” techniques, they constitute a new and valuable toolbox for cell biology (for details on the different techniques, we refer to some recent reviews^{1–3}). All fluorescence microscopy-based methods require suitable and specific strategies for labelling a target structure. An additional request for super-resolution microscopy is a particularly high labelling density, which is reasoned by the sampling theorem.⁴

An elegant strategy to covalently bind small synthetic fluorophores to cellular target structures with high efficiency and specificity makes use of copper(I)-catalysed Huisgen 1,3-dipolar cycloadditions,⁵ which is one example of a chemical reaction today referred to as “click chemistry”.^{6,7} This technique was initially introduced to fluorescence microscopy for labelling nucleic acids^{8,9} and phospholipids.¹⁰ In the recent years, the field has evolved dramatically, and new developments include alternative chemical strategies for click-like reactions, copper-free click chemistry, live cell labelling as well as dual-colour labelling with two orthogonal click reactions.^{11–17} Click chemistry has also entered the field of super-resolution microscopy,

and has been applied to visualize chromosomal DNA in eukaryotes^{18,19} and *Escherichia coli* (*E. coli*)²⁰ as well as proteins in the plasma membrane.¹³ A recent approach has also reported optical mapping of DNA sequences by methyltransferase-directed click chemistry and sequence readout by super-resolution microscopy.²¹

Single-molecule localization microscopy demands bright fluorophores that allow robust and well controllable photo-switching.²² Synthetic fluorophores exhibit a high brightness, yet require chemical functionalizations that allow labelling target structures. In addition, the chemical nano-environment (and thus also the target-specific label) of the fluorophore influences the photophysical properties²³ or may induce quenching.²⁴ So far, azide-substituted and red-emitting carbocyanines (e.g. Alexa Fluor 647)¹⁸ or custom-synthesized rhodamines¹⁹ were used for super-resolution imaging of DNA. Biological applications, however, require multiple targets labelled. Here, we present click chemistry labelling of cellular nucleic acids using a variety of spectrally distinct fluorophores and demonstrate confocal and single-molecule localization-based super-resolution fluorescence microscopy,^{25–28} facilitating new dye combinations for multi-colour imaging. Furthermore, we present a simple strategy to label the outer membrane proteins of *E. coli* using click chemistry in combination with an unnatural amino acid, and visualize the boundaries of bacterial cells with super-resolution imaging.

First, we explored a series of synthetic fluorophores for click chemistry labelling and aimed to find a suitable candidate that is photoswitchable if conjugated to DNA and not red-emitting. High-density labelling of DNA was achieved by addition of an alkyne-modified nucleobase (typically 5-ethynyl-2-deoxyuridine, EdU) to the growth medium, followed by natural uptake by the cells and incorporation into the chromosomal DNA during replication.⁸ After fixation and permeabilisation, the alkyne group of EdU is conjugated to an azide-substituted fluorophore through a copper-catalysed click reaction (Fig. 1A). Here, we prepared (see Materials and methods and Fig. S1 and S2†) or purchased various azide-substituted fluorophores and labelled

^aInstitute of Physical & Theoretical Chemistry, Goethe-University Frankfurt, Max-von-Laue-Str. 7, 60438 Frankfurt/Main, Germany. E-mail: heilemann@chemie.uni-frankfurt.de

^bUCL Medical School, Gower St., London, UK

^cInstitute for Organic Chemistry and Chemical Biology, Goethe-University Frankfurt, Max-von-Laue-Str. 9, 60438 Frankfurt/Main, Germany

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c4ra01027b




Fig. 1 Confocal and super-resolution imaging of chromosomal DNA. (A) Alkyne-modified nucleic acids (R_1 = thymidine in EdU, uracil in EU) or unnatural amino acids react in a copper-catalyzed reaction with an azide-conjugated fluorophore in a [3 + 2] cycloaddition. (B–E) Confocal images of chromosomal DNA in HeLa cells which were pulsed with EdU for 15 min, fixed and permeabilized and labelled with the fluorophore–azides ATTO 488 (B), Alexa Fluor 488 (C), Rhodamine 6G (D) or Alexa Fluor 647 (E) (scale bar 5 μ m). (F) Super-resolution image of chromosomal DNA labelled with Alexa Fluor 488 in a replicating HeLa cell (scale bar 5 μ m) and magnified view (G) (scale bar 0.5 μ m).

chromosomal DNA in eukaryotic cells. We recorded fluorescence images of cells with confocal microscopy (Fig. 1B–E) and report a specific and high-density labelling of DNA in the cell nucleus for all tested rhodamine (ATTO 488, Alexa Fluor 488, Rhodamine 6G) and one carbocyanine fluorophore (Alexa Fluor 647). We explored the specificity as well as the efficiency of the target labelling by varying the reaction times with the fluorophore–azide-conjugates (Fig. S3†).

So far, super-resolution imaging of DNA using bright and photoswitchable fluorophores was only demonstrated using the red-emitting carbocyanine fluorophore Alexa Fluor 647^{18,20} (Fig. S3†). Having fluorophores emitting at different wavelengths available for super-resolution imaging is however desirable, but often complicated by photophysical interactions with the nanoenvironment.^{23,24} Here, we demonstrate redox-induced photoswitching and single-molecule super-resolution imaging of pulse-labelled DNA labelled with the photoswitchable rhodamine fluorophore Alexa Fluor 488 (Fig. 1F and G).^{29,30} The pattern visualized in Fig. 1F is formed by newly synthesized DNA and typical for cells in the middle of DNA replication (S-phase).³¹ These results are consistent with previous work that used BrdU for pulse labelling and diffraction-limited microscopy.³² We determined a localization error using nearest-neighbour analysis³³ and obtained values of 4.4 nm (Alexa Fluor 488, Fig. S4†) and of 12.3 nm (Alexa Fluor 647, Fig. S3 and S4†) (note that the higher localization error of Alexa Fluor 647 is reasoned by a lower number of photons detected in the shorter integration time of 20 ms, compared to 100 ms for Alexa Fluor 488; see Materials and methods and ref. 33). This finding suggests Alexa Fluor 488 as a suitable photoswitchable fluorophore for super-resolution imaging of nucleic acids.

Click chemistry was also reported for RNA labelling and applied to determine RNA transcription and turnover with confocal microscopy.⁹ Here, we introduce for the first time click-chemistry labelling of RNA with fluorophores compatible with single-molecule super-resolution imaging. We applied 5-ethynyluridine (EU) to pulse-label RNA of HeLa cells for 10 min and used click chemistry to label nascent RNA with Alexa Fluor 647. We visualized the nuclear RNA using confocal (Fig. 2A) and super-resolution microscopy (Fig. 2B–C). In the confocal microscopy image (Fig. 2A), high-density RNA is found within the nucleoli, where ribosomal genes are transcribed, whereas the fluorescence signal from the nucleoplasm is lower and appears rather uniform. The super-resolution image (Fig. 2B) confirms that nascent RNA transcribed by RNA polymerases II and III is distributed in distinct spots that are localized all over the nucleoplasm. The magnified image (Fig. 2C) provides additional structural information on the “hot-spots” of RNA polymerase I transcription sites visualized in the nucleoli. These structures seem to be crescent (due to optical sectioning) and rather circularly shaped around darker centres (fibrillar centres). The ribosomal RNA is surrounded by dark areas representing the condensed nucleolar chromatin. This is much expected and known from earlier work on nucleolar transcription *e.g.* visualized using electron microscopy and BrUTP immunostaining (for an in-depth discussion, we refer to ref. 34).

In comparison to other labelling techniques for nucleic acids, click chemistry with alkyne-modified nucleobases has several advantages. First, incorporation of EdU and EU leads to specific labelling only of newly synthesized DNA or RNA; nucleic acids which were already synthesized prior to EdU/EU addition will not be labelled, allowing studies of nascent replication or transcription sites similar as with BrdU or BrUTP.^{34–37} In






Fig. 2 Confocal and super-resolution imaging of nuclear RNA. Living HeLa cells were exposed to EU for 10 min, then fixed with formaldehyde and permeabilized. RNA was labelled via click chemistry with Alexa Fluor 647 azide and imaged with confocal microscopy (A) (scale bar 5 μm) and super-resolution microscopy (B and C). (B) HeLa nucleus showing nucleoplasmic and nucleolar RNA (scale bar 5 μm). (C) The magnified inset of the cell in (B) shows crescent structures of the nucleolar RNA (scale bar 1 μm).

contrast to BrdU/BrUTP, no immunostaining and additional antigen retrieval steps (e.g. 4 M HCL or DNase treatment) are required, which limit the application in crowded or densely packed structures such as nuclei and bacteria. Second, high labelling densities as they are required for sub-diffraction microscopy techniques can be achieved. Third, virtually any fluorophore can be conjugated to nucleic acids. This allows users to choose the optimal fluorophore for the particular application, e.g. photoswitchable fluorophores for single-molecule localization microscopy. DNA labelling with intercalator fluorophores compatible with super-resolution imaging has also been demonstrated.³⁸ However, the photon yields and photostabilities of intercalators are lower than those of bright organic fluorophores. In some cases, the low quantum yield of unbound intercalators can also be the source of additional background signal.

An important aspect of labelling techniques applied in living organisms or cells is the biocompatibility. Both EdU and EU

together were found to be non-toxic in eukaryotic cells even after exposure times of up to 20 hours⁹ and are widely used for confocal microscopy of eukaryotic cells. Here, we applied EdU/EU treatment in eukaryotic cells for 10 to 15 minutes. Furthermore, we monitored the growth rate of *E. coli* in the presence of EdU (Fig. S5[†]). We found no significant perturbation of the growth rate, which again suggests a rather high biocompatibility.

Click chemistry has also entered the field of protein labelling through the use of unnatural amino acids.^{15–17} This opens a novel and simple route to label the outer membrane of cells with photoswitchable fluorophores, and to visualize the boundaries of a cell with sub-diffraction resolution. We demonstrated this by visualizing the outer membrane of *E. coli*, which previously was only achieved at high contrast by phase contrast imaging.³⁹ We grew *E. coli* cells in medium that contained the unnatural amino acid homopropargylglycine (HPG). During cell growth, HPG is incorporated into newly synthesized proteins, replacing methionine, which facilitates the




Fig. 3 Super-resolution imaging of the *E. coli* outer-membrane proteins. (A) The methionine-analogue homopropargylglycine (HPG) is incorporated into newly synthesized proteins. Click reaction with an azide exclusively labels HPG residues outside the cell, resulting in an outer-membrane (OM) stain. Cytosolic or inner-membrane (IM) proteins are separated by the outer membrane and the peptidoglycan layer (PG), and are not labeled. (B) 3D dSTORM image of a HPG-treated *E. coli* cell, labeled with Alexa Fluor 647. The z-position is color-coded and indicates the cylindrical shape of the bacterial cell. (C) Cell diameters can be determined from the intensity profile of a cross-section (white rectangle in (B)) (0.8 μm for the cell in (B)) (scale bar 0.5 μm).



subsequent labelling of all newly synthesized proteins with an azide-fluorophore¹⁷ (Fig. 3A). In order to specifically label the proteins on the outer membrane, cell integrity has to be maintained. We therefore chose to apply a fixation protocol which does not disrupt the outer membrane, so that staining of cytosolic proteins is prevented. Hence, we labelled newly synthesized proteins on the outer membrane which expose an alkyne group. Tagged with a suitable fluorophore for single-molecule super-resolution microscopy, this simple approach allows visualizing of the bacterial outer membrane with high resolution, thus providing a sharp image of the contour of a bacterial cell (Fig. 3B). The resulting image allows determination of the bacterial length and width. 3D-imaging would allow the visualization of the cylindrical shape, which would be a viable reference structure.⁴⁰ In addition, this approach is in principle live-cell compatible; incorporated azide-containing amino acids could be labelled in copper-free click-reactions.^{11,12}

In summary, we have demonstrated click chemistry as versatile tool for super-resolution fluorescence microscopy of cellular structures. Click chemistry on the one hand can provide the high labelling densities typically required for super-resolution microscopy. On the other hand, it is a flexible strategy to label nucleic acids with virtually any synthetic fluorophore, which allows choosing the best-suited one for a specific super-resolution technology. Furthermore, this approach can be combined with pulse labelling and thus visualize particular states of the cell cycle. Finally, a whole new area of applications opened with the introduction of click chemistry into protein labelling through unnatural amino acids. In the future, click chemistry will certainly develop into a widespread and generally used toolbox for cell and tissue labelling.

Acknowledgements

We thank Jana Juli and Martin Held for help with the experiments. We acknowledge funding by the German Research Foundation (SFB 902 and EXC115).

References

- 1 C. G. Galbraith and J. A. Galbraith, Super-resolution microscopy at a glance, *J. Cell Sci.*, 2011, **124**(Pt 10), 1607–1611. Epub 2011/05/04.
- 2 M. Heilemann, Fluorescence microscopy beyond the diffraction limit, *J. Biotechnol.*, 2010, **149**(4), 243–251. Epub 2010/03/30.
- 3 L. Schermelleh, R. Heintzmann and H. Leonhardt, A guide to super-resolution fluorescence microscopy, *J. Cell Biol.*, 2010, **190**(2), 165–175. Epub 2010/07/21.
- 4 C. E. Shannon, Communication in the presence of noise (Reprinted from the Proceedings of the IRE, vol 37, pg 10–21, 1949), *Proc. IEEE*, 1998, **86**(2), 447–457.
- 5 C. W. Tornøe, C. Christensen and M. Meldal, Peptidotriazoles on solid phase: [1,2,3]-triazoles by regioselective copper(I)-catalyzed 1,3-dipolar cycloadditions of terminal alkynes to azides, *J. Org. Chem.*, 2002, **67**(9), 3057–3064. Epub 2002/04/27.
- 6 H. C. Kolb, M. G. Finn and K. B. Sharpless, Click Chemistry: Diverse Chemical Function from a Few Good Reactions, *Angew. Chem., Int. Ed.*, 2001, **40**(11), 2004–2021. Epub 2001/07/04.
- 7 M. Meldal and C. W. Tornøe, Cu-catalyzed azide-alkyne cycloaddition, *Chem. Rev.*, 2008, **108**(8), 2952–3015. Epub 2008/08/14.
- 8 A. Salic and T. J. Mitchison, A chemical method for fast and sensitive detection of DNA synthesis *in vivo*, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**(7), 2415–2420. Epub 2008/02/15.
- 9 C. Y. Jao and A. Salic, Exploring RNA transcription and turnover *in vivo* by using click chemistry, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**(41), 15779–15784. Epub 2008/10/09.
- 10 C. Y. Jao, M. Roth, R. Welti and A. Salic, Metabolic labeling and direct imaging of choline phospholipids *in vivo*, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**(36), 15332–15337. Epub 2009/08/27.
- 11 G. de Almeida, E. M. Sletten, H. Nakamura, K. K. Palaniappan and C. R. Bertozzi, Thiacycloalkynes for copper-free click chemistry, *Angew. Chem., Int. Ed.*, 2012, **51**(10), 2443–2447. Epub 2012/01/28.
- 12 J. C. Jewett, E. M. Sletten and C. R. Bertozzi, Rapid Cu-free click chemistry with readily synthesized biarylazacyclooctynones, *J. Am. Chem. Soc.*, 2010, **132**(11), 3688–3690. Epub 2010/03/02.
- 13 I. Nikic, T. Plass, O. Schraidl, J. Szymanski, J. A. Briggs, C. Schultz, *et al.* Minimal Tags for Rapid Dual-Color Live-Cell Labeling and Super-Resolution Microscopy, *Angew. Chem., Int. Ed.*, 2014, **53**(8), 2245–2249.
- 14 T. Plass, S. Milles, C. Koehler, C. Schultz and E. A. Lemke, Genetically encoded copper-free click chemistry, *Angew. Chem., Int. Ed.*, 2011, **50**(17), 3878–3881. Epub 2011/03/25.
- 15 K. E. Beatty, J. C. Liu, F. Xie, D. C. Dieterich, E. M. Schuman, Q. Wang, *et al.* Fluorescence visualization of newly synthesized proteins in mammalian cells, *Angew. Chem., Int. Ed.*, 2006, **45**(44), 7364–7367. Epub 2006/10/13.
- 16 K. E. Beatty, J. Szychowski, J. D. Fisk and D. A. Tirrell, A BODIPY-cyclooctyne for protein imaging in live cells, *ChemBioChem*, 2011, **12**(14), 2137–2139. Epub 2011/08/11.
- 17 K. E. Beatty, F. Xie, Q. Wang and D. A. Tirrell, Selective dye-labeling of newly synthesized proteins in bacterial cells, *J. Am. Chem. Soc.*, 2005, **127**(41), 14150–14151. Epub 2005/10/13.
- 18 P. J. Zessin, K. Finan and M. Heilemann, Super-resolution fluorescence imaging of chromosomal DNA, *J. Struct. Biol.*, 2012, **177**(2), 344–348. Epub 2012/01/10.
- 19 L. M. Wysocki, J. B. Grimm, A. N. Tkachuk, T. A. Brown, E. Betzig and L. D. Lavis, Facile and general synthesis of photoactivatable xanthene dyes, *Angew. Chem., Int. Ed.*, 2011, **50**(47), 11206–11209. Epub 2011/09/29.
- 20 C. Spahn, U. Endesfelder and M. Heilemann, Super-resolution imaging of *Escherichia coli* nucleoids reveals highly structured and asymmetric segregation during fast growth, *J. Struct. Biol.*, 2014, **185**(3), 243–249.
- 21 C. Vranken, J. Deen, L. Dirix, T. Stakenborg, W. Dehaen, V. Leen, *et al.* Super-resolution optical DNA Mapping via DNA methyltransferase-directed click chemistry, *Nucleic Acids Res.*, 2014, **42**(7), e50.



- 22 A. Furstenberg and M. Heilemann, Single-molecule localization microscopy-near-molecular spatial resolution in light microscopy with photoswitchable fluorophores, *Phys. Chem. Chem. Phys.*, 2013, **15**(36), 14919–14930. Epub 2013/08/09.
- 23 S. Nangneri, B. Flottmann, F. Herrmannsdorfer, K. Thomas and M. Heilemann, Single-molecule super-resolution imaging by tryptophan-quenching-induced photoswitching of phalloidin-fluorophore conjugates, *Microsc. Res. Tech.*, 2014, **77**(7), 510–516.
- 24 S. Doose, H. Neuweiler and M. Sauer, Fluorescence quenching by photoinduced electron transfer: a reporter for conformational dynamics of macromolecules, *ChemPhysChem*, 2009, **10**(9–10), 1389–1398. Epub 2009/05/29.
- 25 E. Betzig, G. H. Patterson, R. Sougrat, O. W. Lindwasser, S. Olenych, J. S. Bonifacino, *et al.* Imaging intracellular fluorescent proteins at nanometer resolution, *Science*, 2006, **313**(5793), 1642–1645. Epub 2006/08/12.
- 26 M. Heilemann, S. van de Linde, M. Schuttpelz, R. Kasper, B. Seefeldt, A. Mukherjee, *et al.* Subdiffraction-resolution fluorescence imaging with conventional fluorescent probes, *Angew. Chem., Int. Ed.*, 2008, **47**(33), 6172–6176. Epub 2008/07/23.
- 27 S. T. Hess, T. P. Girirajan and M. D. Mason, Ultra-high resolution imaging by fluorescence photoactivation localization microscopy, *Biophys. J.*, 2006, **91**(11), 4258–4272. Epub 2006/09/19.
- 28 M. J. Rust, M. Bates and X. Zhuang, Sub-diffraction-limit imaging by stochastic optical reconstruction microscopy (STORM), *Nat. Methods*, 2006, **3**(10), 793–795. Epub 2006/08/10.
- 29 G. T. Dempsey, J. C. Vaughan, K. H. Chen, M. Bates and X. Zhuang, Evaluation of fluorophores for optimal performance in localization-based super-resolution imaging, *Nat. Methods*, 2011, **8**(12), 1027–1036. Epub 2011/11/08.
- 30 M. Heilemann, S. van de Linde, A. Mukherjee and M. Sauer, Super-resolution imaging with small organic fluorophores, *Angew. Chem., Int. Ed.*, 2009, **48**(37), 6903–6908. Epub 2009/08/12.
- 31 H. Nakayasu and R. Berezney, Mapping replicational sites in the eukaryotic cell nucleus, *J. Cell Biol.*, 1989, **108**(1), 1–11. Epub 1989/01/01.
- 32 H. Ma, J. Samarabandu, R. S. Devdhar, R. Acharya, P. C. Cheng, C. Meng, *et al.* Spatial and temporal dynamics of DNA replication sites in mammalian cells, *J. Cell Biol.*, 1998, **143**(6), 1415–1425. Epub 1998/12/16.
- 33 U. Endesfelder, S. Malkusch, F. Fricke and M. Heilemann, A simple method to estimate the average localization precision of a single-molecule localization microscopy experiment, *Histochem. Cell Biol.*, 2014, **141**(6), 629–638.
- 34 A. Papantonis and P. R. Cook, Transcription factories: genome organization and gene regulation, *Chem. Rev.*, 2013, **113**(11), 8683–8705. Epub 2013/04/20.
- 35 B. K. Kennedy, D. A. Barbie, M. Classon, N. Dyson and E. Harlow, Nuclear organization of DNA replication in primary mammalian cells, *Genes Dev.*, 2000, **14**(22), 2855–2868. Epub 2000/11/23.
- 36 D. S. Dimitrova and R. Berezney, The spatio-temporal organization of DNA replication sites is identical in primary, immortalized and transformed mammalian cells, *J. Cell Sci.*, 2002, **115**(Pt 21), 4037–4051. Epub 2002/10/03.
- 37 A. Z. Canals-Hamann, R. P. das Neves, J. E. Reittie, C. Iniguez, S. Soneji, T. Enver, *et al.* A biophysical model for transcription factories, *BMC Biophys.*, 2013, **6**, 2. Epub 2013/02/12.
- 38 C. Flors, Photoswitching of monomeric and dimeric DNA-intercalating cyanine dyes for super-resolution microscopy applications, *Photochem. Photobiol. Sci.*, 2010, **9**(5), 643–648. Epub 2010/05/06.
- 39 S. J. Holden, T. Pengo, K. L. Meibom, C. Fernandez Fernandez, J. Collier and S. Manley, High throughput 3D super-resolution microscopy reveals *Caulobacter crescentus* *in vivo* Z-ring organization, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**(12), 4566–4571. Epub 2014/03/10.
- 40 M. D. Lew, S. F. Lee, J. L. Ptacin, M. K. Lee, R. J. Twieg, L. Shapiro, *et al.* Three-dimensional superresolution colocalization of intracellular protein superstructures and the cell surface in live *Caulobacter crescentus*, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**(46), E1102–E1110. Epub 2011/10/28.

