



Cite this: *Chem. Commun.*, 2017, 53, 1214

Long range electrostatic forces in ionic liquids

Matthew A. Gebbie,^{*a} Alexander M. Smith,^b Howard A. Dobbs,^c Alpha A. Lee,^d
Gregory G. Warr,^e Xavier Banquy,^f Markus Valtiner,^g Mark W. Rutland,^{*hi}
Jacob N. Israelachvili,^{*c} Susan Perkin^{*b} and Rob Atkin^{*j}

Ionic liquids are pure salts that are liquid under ambient conditions. As liquids composed solely of ions, the scientific consensus has been that ionic liquids have exceedingly high ionic strengths and thus very short Debye screening lengths. However, several recent experiments from laboratories around the world have reported data for the approach of two surfaces separated by ionic liquids which revealed remarkable long range forces that appear to be electrostatic in origin. Evidence has accumulated demonstrating long range surface forces for several different combinations of ionic liquids and electrically charged surfaces, as well as for concentrated mixtures of inorganic salts in solvent. The original interpretation of these forces, that ionic liquids could be envisioned as "dilute electrolytes," was controversial, and the origin of long range forces in ionic liquids remains the subject of discussion. Here we seek to collate and examine the evidence for long range surface forces in ionic liquids, identify key outstanding questions, and explore possible mechanisms underlying the origin of these long range forces. Long range surface forces in ionic liquids and other highly concentrated electrolytes hold diverse implications from designing ionic liquids for energy storage applications to rationalizing electrostatic correlations in biological self-assembly.

Received 3rd November 2016,
Accepted 9th December 2016

DOI: 10.1039/c6cc08820a

www.rsc.org/chemcomm

Ions impact all aspects of modern life. The smart phones that link much of the Earth's population are powered by batteries that depend on the directed motion of large collections of ions.¹ Collective ion motion will also play a key role in energy storage technologies that will enable societies to transition away from fossil fuels to greener energy sources, such as supercapacitors,² and the correlated behaviour of ions is responsible for many biological processes that support life.³ Consequently, a complete understanding of the fundamental

principles that govern the behaviour of large collections of ions would impact many branches of science and engineering, and allow ion collective motion to be predictably tuned.

This is a tremendous challenge, in large part due to the large potential energy of electrostatic interactions and long range of electrostatic interactions. In particular, analytical theories describing disordered soft matter systems, like liquid-based energy storage devices and biological systems, are sparse, and such systems are too complex to be reliably simulated.⁴ Increasingly, researchers are exploring the physics of ionic liquids to provide general insight into the fundamental nature of ionic correlations, as ionic liquids are comparatively simple systems that exhibit the key features of electrostatic correlations in disordered soft matter.

Ionic liquids are pure salts that are liquids at ambient conditions.⁵ Strong electrostatic interactions determine the key properties of ionic liquids, as for traditional high melting point salts. In contrast with traditional salts, ionic liquids are typically composed of large, asymmetric polyatomic ions with ionic radii that are about 5–10 times larger than those of monatomic ions like Li^+ or K^+ . The large size of ionic liquid ions increases the average separation distance between cation and anion charge centers, decreasing the strength of electrostatic interactions. Concurrently, the presence of flexible and/or asymmetric ionic structures, such as alkyl chains on cations or conformational flexibility in bis(trifluoromethylsulfonyl)imide

^a Geballe Laboratory for Advanced Materials, Stanford University, Stanford, CA 94305, USA. E-mail: magebbie@stanford.edu

^b Department of Chemistry, Physical & Theoretical Chemistry Laboratory,
University of Oxford, Oxford, OX1 3QZ, UK. E-mail: susan.perkin@chem.ox.ac.uk

^c Department of Chemical Engineering, University of California, Santa Barbara, CA 93106, UK. E-mail: jacob@engineering.ucsb.edu

^d School of Engineering and Applied Sciences, Harvard University, Cambridge, MA 02138, USA

^e School of Chemistry, F11, The University of Sydney, NSW 2006, Australia

^fFaculty of Pharmacy, Université de Montréal, Montréal, QC, Canada

^gInterface Chemistry and Surface Engineering, Max Planck Institut für Eisenforschung GmbH, Düsseldorf, 40237, Germany

^h Surface and Corrosion Science, KTH Royal Institute of Technology, SE-10044 Stockholm, Sweden. E-mail: mark@kth.se

ⁱ *SP Chemistry Materials and Surfaces, SE 114 86 Stockholm, Sweden*

*^j Priority Research Centre for Advanced Fluid Interfaces, Newcastle Institute for Energy and Resources, The University of Newcastle, Australia.
E-mail: rob.atkin@newcastle.edu.au*

anions, introduce significant conformational and configurational entropy in ionic liquids, leading to a large number of energetically similar, and thus thermally accessible, states. Both aspects drastically suppress the melting points of ionic liquids; all of the ionic liquids discussed in this article are liquids at room temperature.

Ionic liquids exhibit a collection of properties that are of technological interest. For example, ionic liquids have extremely low vapour pressures, modest intrinsic ionic conductivities, high thermal and electrochemical stability, and are known to dissolve a wide range of chemical species, including lithium salts and polyaromatic compounds.^{5,6} As a consequence of this unusual range of properties, ionic liquids have significant promise for many emerging applications such as: electrochemical energy storage devices, high temperature and/or vacuum phase materials synthesis, semiconductor and superconductor gating applications, and novel self-assembly media.^{5–10}

For many applications, the structure and properties of the electric double layer formed by ionic liquids at solid surfaces is of key importance, and has thus been the focus of a major research effort during the past decade.⁴ As pure salts, the consensus has traditionally been that ionic liquids have exceedingly high ionic strengths and thus very short Debye lengths.^{4,11} However, in 2013 Gebbie *et al.*¹² presented force-distance data for two surfaces separated by an ionic liquid which revealed attractive forces between gold and mica surfaces extending to significantly greater distance than anticipated (Fig. 1A). Subsequently, a repulsive force of similar range was measured between two mica surfaces (Fig. 1B).¹³ The authors concluded that these forces are electrostatic in origin, with the implication that the ionic liquids behave as dilute electrolytes, challenging the traditional notion that ionic liquids exhibit Debye lengths that are on the order of ionic radii.

Repulsive long range surface forces have since been measured between surfaces with the same charge in several laboratories around the globe.^{13–16}

Similar exponentially decaying forces have also been measured for ionic liquids mixed with a polar solvent,¹⁶ and in simple metal halide electrolytes in water at high concentration.¹⁷ Critically, this shows that the force is not a unique feature of pure ionic liquids. Thus, it is clear that long range surface forces operate between electrically charged surfaces across many different ionic liquids and concentrated electrolytes, although their origin remains the subject of debate (Fig. 2).^{13,16,18–20}

In this article, we review experimental evidence for the existence of long range surface forces in pure ionic liquids and identify areas of consensus and division. Cases where prior experimental results stand in contrast to the recent reports of long range forces in ionic liquids are discussed, and the mechanisms that may give rise to these long range forces are described phenomenologically. Critical open questions are highlighted and potential experiments aimed at a deeper understanding of long range surface forces in ionic liquids are proposed.


Ionic liquid interfacial nanostructure

Solid surfaces perturb the nanostructure and dynamics of all liquids. In the bulk, aprotic ionic liquids with alkyl chain lengths longer than 3 carbon units, and most protic ionic liquids, have sponge-like nanostructures; only ionic liquids with very short alkyl chains exhibit charge ordering nanostructures that are reminiscent of high-temperature molten salts.^{21,22} Very long chain surfactant-like ionic liquids form liquid crystalline phases,²³ but these are not discussed here. Ionic liquid nanostructure is solvophobic²⁴



Fig. 1 (A) SFA force-distance profile between a positive gold electrode ($\Delta U = +360$ mV vs. OCP) and a negative mica surface across $[\text{C}_4\text{mim}^+][\text{NTf}_2^-]$. Data points are equilibrium forces when bringing together the surfaces. Solid lines are fitted interaction potentials, where the total interaction (black) is a superposition of long range Poisson–Boltzmann (PB) double layer attraction (blue), short range mica–gold van der Waals attraction (red) and short range steric repulsion (green). The double layer force exhibits a decay length of $(\kappa^*)^{-1} = 10 \pm 2$ nm (eqn (3)). (B) SFA measurements between two negative mica surfaces across $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$ at room temperature ($T = 22$ °C) and at $T = 50$ °C. Data points are equilibrium forces on approach, and solid lines are fitted PB interaction potentials (eqn (2) and (3)). As the temperature is increased, $(\kappa^*)^{-1}$ decreases, consistent with the thermal disruption of electrostatic correlations and/or effective dissociation of ionic liquid ions. Figure panel (B) reprinted from M. A. Gebbie, H. A. Dobbs, M. Valtiner, and J. N. Israelachvili, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, 7432–7437.





The mechanism of structural forces in ionic liquids has the same origin as structural forces in molecular liquids, where

In contrast, protic ionic liquids can have cation radii that are less than 0.5 nm, and AFM imaging has suggested that protic ionic liquids with small ammonium cations, like ethylammonium nitrate (EAN) form surface-bound structures with a spacing that closely matches the charge sites of the underlying mica lattice.³⁷ This may be noteworthy, as studies of EAN using the SFA³⁸ and colloid probe AFM^{39,40} techniques report solely short range




Fig. 3 Force-distance profile for a Si_3N_4 AFM tip approaching (blue) and retracting from (red) a mica surface in (A) $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$, (B) $[\text{C}_4\text{mpyr}^+][\text{NTf}_2^-]$, (C) $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$, and (D) $[\text{C}_4\text{mim}^+][\text{PF}_6^-]$. Reproduced from ref. 24 with permission from the PCCP Owner Societies.

interaction forces at room temperature. Nonetheless, force measurements on aprotic ionic liquids in contact with mica surfaces consistently uncover the presence of long range interaction forces, where some portion of the mica surface potential necessarily propagates beyond the near surface nanostructure.

The situation is even less well resolved for surfaces other than mica. For example, Lauw *et al.*⁴¹ reported cation concentrations can be enhanced relative to anions in the layers of 1-butyl-1-methylpyrrolidinium ($[\text{C}_4\text{mpyr}^+]$) bis(trifluoromethylsulfonyl)imide ($[\text{NTf}_2^-]$) at gold electrode surfaces, even over a range of positive electrochemical potentials. Thus, knowledge of surface charge alone is insufficient to predict the major ionic species that will be present adjacent to charged surfaces, and one must also carefully consider the roles played by specific ion-surface interactions and entropic effects.

Trace water can significantly impact ionic liquid near-surface nanostructure. Experiments have shown that trace water can either enhance or disrupt bound ion layers,^{14,31,42} possibly depending on the exact nature of the ion-surface interactions. Trace amounts of water may enhance ion layering on mica by facilitating K^+ dissociation,⁴³ but above a certain water threshold, the IL layering can be disrupted due to both packing and capillary effects;^{14,42–44} the details are strongly ionic liquid-dependent. At saturation, water molecules can be in excess or depleted at surfaces depending on whether the surface is hydrophilic or hydrophobic respectively, and the affinity of the ions for the surface.⁴

Long range forces across ionic liquids

Gebbie *et al.*¹² first reported long range forces in ionic liquids in 2013 for $[\text{C}_4\text{mim}^+][\text{NTf}_2^-]$ (Fig. 1A). A long range, monotonic attraction across a gold surface held at positive bias and a mica surface was measured with a characteristic exponential decay length of 10 ± 2 nm (Fig. 1A). Mica becomes negatively charged when immersed in ionic liquids, due to desorption of surface-bound potassium ions, adding a negligible K^+ content to bulk ionic liquids.

A weaker monotonic attraction was measured when a negative potential was applied to the gold.¹² At first glance this is surprising, as a repulsion is expected between surfaces with the same charge. However, neutron reflectivity measurements⁴¹ for a structurally similar ionic liquid have shown that the first adsorbed layer can be cation rich, even at positive potentials, suggesting that $[\text{C}_4\text{mim}^+][\text{NTf}_2^-]$ could fully screen gold surfaces, which would account for the absence of a long range repulsion. The asymmetry in the ability of cations and anions to neutralize the surface potential highlights the fact that specific ion-surface interactions and the properties of bound ion layers are critical determinants of long range forces in ionic liquids.

The presence of long range surface forces has since been reported in several independent laboratories^{13–16,37} (Fig. 4), and it is clear that long range forces do operate for several combinations of ionic liquids and charged surfaces. The magnitudes of the long range surface forces measured in ionic liquids, normalized by the probe radius of curvature, (F/R), typically fall





Fig. 4 Comparison of SFA force-distance profiles between symmetric mica surfaces across differing ionic liquids and humidity conditions. Long range repulsive forces with a characteristic decay length of $(\kappa^*)^{-1} \sim 5\text{--}10$ nm are observed in dried ionic liquids and ionic liquids that have been exposed to relatively low humidity environments ($\text{RH} \leq 37\%$). Forces measured across (A) dried $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$ (blue) and dried $[\text{C}_3\text{mim}^+][\text{NTf}_2^-]$ (black),¹³ (B) dried (red), humid (blue) and semi-humid (purple) $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$,¹⁴ (C) dried (orange, red) and semi-humid (purple, blue, green), $[\text{C}_2\text{mim}^+][\text{FAP}]$,¹⁵ and (D) dried $[\text{C}_4\text{mpyr}^+][\text{NTf}_2^-]$.¹⁶ Figure panel (B) reprinted with permission from H. W. Cheng *et al.*, *Adv. Mater. Interfaces*, 2015, **2**, 1500159. Copyright 2015 Wiley and Sons. Figure panel (C) adapted with permission from R. M. Espinosa-Marzal, A. Arcifa, A. Rossi, and N. D. Spencer, *J. Phys. Chem. Lett.*, 2014, **5**, 179–184. Copyright 2014 American Chemical Society.

in the range of $0.1\text{--}1\text{ mN m}^{-1}$, and the minimum F/R magnitude that can be detected in typical force measuring experiments can be lower than 0.1 mN m^{-1} . Thus, long range forces in ionic liquids can routinely be measured *via* both SFA and AFM under carefully controlled conditions, as discussed in prior work.^{13–16,49} For comparison, long range diffuse electric double layer forces measured in mM concentration aqueous electrolyte solutions can exhibit F/R magnitudes of between $0.1\text{--}10\text{ mN m}^{-1}$, depending on the relative electrolyte concentrations and surface potentials.³⁸

Long range interaction forces in ionic liquids exhibit ionic liquid-dependent exponential decay lengths in the long range, asymptotic limit, with decay lengths ranging from between about $5\text{--}10$ nm for a diverse range of aprotic ionic liquids (Fig. 5).^{12–16} To date, the ionic liquids studied exhibit remarkably similar asymptotic interaction decay lengths, despite pronounced differences in short range ion structuring forces, implying a common underlying mechanism in the long range force regime.

It has also been shown that the long range exponentially decaying force is not unique to pure ionic liquids, or to ionic liquids at all: Smith *et al.*¹⁶ studied the interaction force between two (negative) mica sheets across a wide range of ionic liquids, ionic liquids mixed with a polar solvent in varying fractions, and NaCl in water at high concentration. In every case, the force law was qualitatively similar, with oscillations at short distances, giving way to a monotonic exponentially decaying force at longer range.

The decay length of the long range part was shown to depend on ion density and the dielectric constant. Decay lengths for these high-concentration electrolytes and diluted ionic liquids are presented in Fig. 5 alongside the pure ionic liquids for comparison. When the dielectric constant of the fluid is taken into account,¹⁶ all data, including previous literature data for aqueous salt solutions, such as the LiCl solutions studied by Baimpos *et al.*,¹⁷ fits onto a master curve. This means that finer molecular details of the ionic liquid ions,






Fig. 5 Ratio of the measured decay length of the long range component of the surface force, $(\kappa^*)_{\text{exp}}^{-1}$, to the theoretical Debye length, κ_D^{-1} , for a wide range of aprotic ionic liquids and electrolyte solutions. The ratio, $(\kappa^*)_{\text{exp}}^{-1}/\kappa_D^{-1}$, is plotted versus the dimensionless parameter d/κ_D^{-1} , where d is the mean ion diameter for each electrolyte. d/κ_D^{-1} scales as $(c/\epsilon)^{1/2}$ and can be considered as a dielectric-scaled ion concentration. The value of dielectric constant, ϵ , for each type of electrolyte also varies with concentration; here values of ϵ were either taken from the literature or calculated using effective medium theory. The pure ionic liquids are $[\text{C}_4\text{mpyr}^+][\text{NTf}_2^-]$, $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$, $[\text{C}_2\text{mim}^+][\text{BF}_4^-]$, $[\text{C}_2\text{mim}^+][\text{OTf}^-]$; the electrolyte solutions are aqueous NaCl, CsCl, LiCl, and $[\text{C}_4\text{mpyr}^+][\text{NTf}_2^-]$ in propylene carbonate solvent. For details of the determination of d , ϵ , and other discussion see Smith *et al.* *J. Phys. Chem. Lett.*, 2016. Adapted with permission from A. M. Smith, A. A. Lee, and S. Perkin, *J. Phys. Chem. Lett.*, 2016, 7, 2157–2163. Copyright 2016 American Chemical Society.

such as the size of cation hydrocarbon substituents, do not control the fundamental nature of the long range force. The long range force is of purely electrostatic origin.

Surface properties therefore play a key role in determining the presence and magnitude of long range forces in ionic liquids, while having little impact on the long-range interaction decay length. To date, pronounced long range surface interactions have only been observed when ionic liquids are confined between, or located directly adjacent to, charged surfaces.^{12–16,44} Experiments with *in situ* electrochemical control demonstrate that interaction forces between surfaces and/or probe tips, whether attractive¹² or repulsive,⁴⁵ systematically increase in magnitude as the surface potential difference is increased. In contrast, the interaction decay lengths appear to be independent of the surface potential. Both the dependence of interaction strengths on surface potential and the independence of the interaction decay lengths are consistent with the conclusion that the long range force is probing a bulk correlation length, determined by bulk properties such as solution ionic strength and dielectric permittivity.

Long range forces in ionic liquids are highly sensitive to temperature. Recent SFA measurements¹³ show that long range interaction decay lengths for two different $[\text{C}_R\text{mim}^+][\text{NTf}_2^-]$ ionic liquids systematically decrease as the temperature is increased from 22–75 °C. For all temperatures, the structures formed by the near-surface bound ion layers and mica surface

potential remain unchanged, while the interaction decay length decreases from 6.7 ± 0.1 nm (22 °C) to 3.4 ± 0.2 nm (60 °C) for $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$ and from 7.4 ± 0.1 nm (22 °C) to 3.5 ± 0.2 nm (74 °C) for $[\text{C}_3\text{mim}^+][\text{NTf}_2^-]$. It was proposed that increasing the temperature disrupts ionic correlations in bulk ionic liquids, enabling ionic liquid ions to more effectively screen surfaces over shorter distances. In contrast, forces that are purely entropic in nature would be expected to be repulsive and to grow in range as the temperature is increased.³⁵

The presence of trace amounts of water in ionic liquids also impacts long range interaction forces, however, there are few experimental works that systematically probe the specific impact of very small fractions of water on the long range interactions. Espinosa-Marzal and colleagues¹⁵ showed that equilibrating the ionic liquid $[\text{C}_2\text{mim}^+]$ tris(pentafluoroethyl)trifluorophosphate ($[\text{FAP}^-]$) at 37% relative humidity modestly decreases the magnitude of both the long and short range forces measured between mica surfaces, without qualitatively altering the range of long range forces (Fig. 4C).

In a separate paper⁴³ the decay length decreased with increasing humidity for a hydrophilic ionic liquid, in qualitative accordance with the dilution results of Smith *et al.*;¹⁶ the added water leads to an increased dielectric constant. In comparison, Cheng *et al.*¹⁴ show that saturating $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$ with water at 44% relative humidity gives rise to an unexpected attractive instability between 1.5 and 5 nm (Fig. 4D), attributed to capillary bridging of the mica surfaces by water films that adsorbed at each of the hydrophilic mica-liquid interfaces. Similar long range capillary forces in water saturated $[\text{C}_R\text{mim}^+][\text{NTf}_2^-]$ ionic liquids were recently observed by Wang *et al.*⁴⁶ using AFM. These results show that the presence of water in ionic liquids can result in complex, and at times unexpected, changes to surface forces.^{14,46,47}

While the presence of long range surface forces is increasingly measured for numerous combinations of ionic liquids and charged surfaces, long range interaction forces are not measured in every experiment on ionic liquids. In 1988, Horn and colleagues³⁸ observed solely short range ion binding forces in an SFA study of the protic ionic liquid EAN between mica surfaces. Similarly, in 2010, Bou-Malham *et al.*⁴⁸ performed an SFA study on aprotic ionic liquids where long range forces were not reported. Measurements were performed on two aprotic ionic liquids, $[\text{C}_4\text{mim}^+]$ tetrafluoroborate ($[\text{BF}_4^-]$) and $[\text{C}_4\text{mim}^+]$ hexafluorophosphate ($[\text{PF}_6^-]$), between mica surfaces and only short range oscillatory forces were observed. The apparent absence of long range forces in those experiments appears to be inconclusive, because the measurement of the long range force baselines required for the observation of weak long range forces is not necessary when addressing the near-surface structure that was the focus of those studies.

It remains to be shown whether an SFA experiment measuring surface forces across EAN, with high resolution and measured from very large distances would give rise to an additional weak monotonic component. Recent, AFM colloid probe measurements of EAN at room temperature yielded solely short range interaction forces,^{39,40} while AFM colloidal probe measurements of EAN at a significantly elevated






Fig. 6 Colloid probe AFM force–distance measurements between a silica probe and mica surface across the protic ionic liquid ethyl ammonium nitrate (EAN) at 80 °C and 120 °C. In EAN, no long range forces of electrostatic origin are observed between room temperature and 80 °C.^{38–40} However, at 120 °C, a long range force consistent with an electrostatic double layer force is manifested.⁴⁹ Reproduced from ref. 49 with permission from the Royal Society of Chemistry.

temperature of 120 °C clearly demonstrate the presence of long range monotonic forces (Fig. 6).⁴⁹ In this light, it appears unlikely that the absence of long range forces in the work of Horn *et al.* is due to technical difficulties.

The onset of a long range force at high T initially appears to contradict the temperature dependence of long range forces reported previously.¹³ However the temperature dependence of the long ranged force in EAN has not yet been examined. Thus, the current state of understanding for EAN is that significant increases in temperature from 22–120 °C leads to a transition from full surface charge screening at low temperatures to dilute electrolyte behaviour at 120 °C. At this stage, it is not possible to say whether this is due to changes in: (i) the ability of EAN to neutralise the surface charge at short range, the oscillatory forces are however indistinguishable from those at 80 °C as can be inferred from Fig. 6 or (ii) a change in bulk properties, for example temperature induced breakdown in the strongly hydrogen bonded, spongelike structure.

Of the other protic ILs studied, the only one to demonstrate evidence for a long ranged repulsion is dimethyl ethylammonium formate (DMEAF).³⁷ However since this is the protic ionic liquid system with both the largest cation, and the least hydrogen bonding, this observation provides little assistance in distinguishing between these two possibilities.

Open questions and plausible origins of long range forces

The classical mean-field approach to treating electrostatic screening in electrolytes provides a useful starting point for understanding prior proposals for explaining the origin of long range forces in ionic liquids. The (formal) Debye length is defined as:

$$(\kappa_D)^{-1} = \sqrt{\frac{\epsilon\epsilon_0 k_B T}{2n(q)^2}} \quad (1)$$

where $(\kappa_D)^{-1}$ (meters) is the Debye length, n (meters^{−3}) is the number density of ions, $q = ze$ (Coulombs) is the ion charge, z is the ion valence, e (Coulombs) is the elementary charge, ϵ is the dielectric permittivity of the electrolyte solution, ϵ_0 (farads per meter) is the vacuum permittivity, T (Kelvin) is the temperature, and k_B is the Boltzmann constant (J/K). The Debye length decreases as the ion concentration increases. At the limit of low ion density, when the assumptions leading to the Debye–Hückel theory hold, the Debye length of an electrolyte can be measured in force-distance experiments by fitting the measured force law to an appropriate form of the Poisson–Boltzmann equation. For example, eqn (2) shows a constant potential solution to the linearized Poisson–Boltzmann equation for two symmetric planar surfaces:

$$E_{EDL} = \frac{4e^2 n}{k_B T \kappa_D} \psi^2 \exp(-\kappa_D D) \quad (2)$$

where ψ (volts) is the surface potential.

Typical ionic liquids have a total bulk ion concentration of about 7 M, which yields a formal Debye length on the order of 0.1 nm. This is smaller than the size of an individual ionic liquid ion and is several orders of magnitude lower than the experimentally observed decay length; it is clear that a standard application of Debye–Hückel theory could not explain the observed long range forces.

The inapplicability of the standard Debye length, $(\kappa_D)^{-1}$, at such high electrolyte concentrations is unsurprising since strong ion–ion correlations exist at high ionic strengths.^{50,51} Poisson–Boltzmann equations, integral equation theories, and density functional theories have been developed to overcome the deficiencies in mean-field models, and these pioneering studies have been the subject of excellent reviews.⁵⁰ Nonetheless, to our knowledge, these theories do not quantitatively predict a decay length that is orders of magnitude greater than the formal Debye length.

Currently, the molecular origin of long range forces in ionic liquids remains an active topic of scientific discussion and theoretical exploration, and different ways of renormalizing the parameters present in eqn (1)–(3) appear to be useful for developing intuition on ionic liquids. Specifically, the interaction decay length in ionic liquids can be framed in terms of an effective Debye length with rescaled parameters, where the formal Debye length of eqn (1), $(\kappa_D)^{-1}$, is replaced by an effective Debye length in ionic liquids, $(\kappa^*)^{-1}$, shown in eqn (3).

$$(\kappa^*)^{-1} = \sqrt{\frac{\epsilon\epsilon_0 k_B T}{2n^*(q^*)^2}} \quad (3)$$

$(\kappa^*)^{-1}$ (meters) is the effective Debye length, n^* (meters^{−3}) is the number density of electrostatic monopoles (free dissociated ions, ion vacancies and/or higher order aggregates), $q^* = ze$ (Coulombs) is the carrier effective charge and z is the carrier effective valence. The effective concentration, effective charge and effective dielectric function appear to be multiple angles to view the same physical phenomenon; below, we organize our



discussion in terms of the differing approaches that have been proposed.

Renormalizing the effective ion concentration, n^*

Initially, Gebbie *et al.*¹² proposed that ionic liquids behave as very weakly dissociated, in other words dilute, ionic “solutions.” In this model, a small thermally excited population of “free ions” exist in equilibrium with a strongly correlated ionic network where each charge is neutralised by the sum of its neighbours charge, “dielectric solvent.” This conclusion was reached by interpreting the 10 nm interaction decay length reported in ref. 12 as the effective Debye length, $(\kappa^*)^{-1}$, of $[\text{C}_4\text{mim}^+][\text{NTf}_2^-]$ at room temperature. This would indicate that less than 0.1% of the total number of ions independently contribute to electrostatic screening which was a surprising conclusion at the time.

This effective Debye length was used in conjunction with the Boltzmann distribution to calculate the energetics for the effective dissociation for a virtual $[\text{C}_4\text{mim}^+][\text{NTf}_2^-]$ ion pair within the bulk ionic liquid. It was shown¹² that the effective dissociation energy determined from the measured interaction decay length was in agreement with a simple model that uses only the calculated vacuum ion pair interaction energy⁵² and measured low frequency dielectric permittivity⁵³ of $[\text{C}_4\text{mim}^+][\text{NTf}_2^-]$ (eqn (4) and (5)).

$$n^* = n_b \exp\left(-\frac{E_d}{k_B T}\right) \quad (4)$$

$$E_d = \frac{E_{\text{pair}}}{\epsilon} \quad (5)$$

where n^* (meters⁻³) is the effective dissociated ion density that determines the effective Debye length, n_b (meters⁻³) is the bulk total number density of ionic liquid ions, E_d (joules) is the dissociation energy of a virtual ion pair immersed in a surrounding ionic liquid network, E_{pair} (joules) is the ensemble average vacuum interaction energy of two ionic liquid ions, ϵ is the low frequency dielectric permittivity of the ionic liquid.

The agreement between this simple calculation and experimental observations¹² raises the intriguing possibility that the relevant energetics for cation–anion interactions for two arbitrary ions in bulk ionic liquids can be predicted from knowledge of the ensemble average ion pair potential and ionic liquid low frequency dielectric permittivity. The permittivity is a measured property that, to first order, appears to account for the influence of electrostatic correlations, for example, ion pairing. Hence, the picture is that the majority of ionic liquid ions are strongly associated and cannot move independently, nor obey the Poisson–Boltzmann equation, leaving the ions to effectively behave as a dielectric network. This dielectric network stabilizes the presence of “free” ions and/or charged thermal excitations.

Importantly, ion pairing and/or ion aggregation in ionic liquids can give rise to an effective low frequency dielectric behaviour in bulk ionic liquids, even when pairing and/or aggregation is highly dynamic and transient, as shown very recently theoretically by Kjellander.²⁰ This would explain why

NMR diffusivity measurements fail to find evidence of long lived ion pairing in bulk ionic liquids,⁵⁴ while surface forces measurements^{12–16} and dielectric spectroscopy measurements⁵⁵ are suggestive of ionic correlation-driven aggregation. We stress that ionic aggregation and/or pairing in ionic liquids is likely a highly dynamic process.

This effective dissociation framework¹² has several testable predictions and practical implications:

- First, any model of long range forces that invokes the thermal disruption of ionic correlations, an effective thermal “dissociation” of ionic liquid ions, necessitates that increases in temperature should systematically decrease the decay length of long range forces. Gebbie *et al.* confirmed this prediction using $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$ and $[\text{C}_3\text{mim}^+][\text{NTf}_2^-]$ (Fig. 1B).¹ For these ionic liquids, increasing the temperature decreases the long range interaction decay length, which was interpreted as an increase in the thermally excited free ion population.

- Second, the energetics of ion “dissociation” should exhibit a temperature dependence solely through the impact of temperature on the dielectric permittivity. An analysis using the theoretical framework outlined in ref. 12 and eqn (3) and (4) under varying temperatures was carried out,¹ and the dissociation energy was found to be independent of temperature for $[\text{C}_2\text{mim}^+][\text{NTf}_2^-]$ and $[\text{C}_3\text{mim}^+][\text{NTf}_2^-]$ up to 60 °C. This temperature independence appears to break down above 75 °C, perhaps due to changes in dielectric properties at high temperatures.

- Third, eqn (4) and (5) suggest that if one could identify two ionic liquids with similar ion pair vacuum interaction energies, E_{pair} , but significantly different low frequency dielectric permittivities, the ionic liquid with a higher dielectric permittivity should exhibit a significantly higher degree of ionic dissociation, and resultant smaller electrostatic screening length. This experiment could be realized by maintaining the cation core of a common aprotic ionic liquid, such as the $[\text{C}_R\text{mim}^+][\text{NTf}_2^-]$ class of ionic liquids, and comparing an alkyl-imidazolium substituted cation to an analogous imidazolium cation that incorporates a substituent with oxygen atoms or other polar groups. If confirmed, this would provide a new strategy for engineering the effective degree of dissociation of ionic liquid ions by controllably increasing (or decreasing) the ionic liquid permittivity. Hence, the cation substituents would provide functionalities that go beyond the crystal-suppressing symmetry breaking function of traditional alkyl substituents.

The ion aggregation model has been successful in describing the phenomenology and scaling of experiments on ionic liquids. Importantly, eqn (5) describes the energy required to separate charges in a pure dielectric medium with no ionic screening, thus corresponding to the limit of low free ion concentration. Models that explicitly account for screening by surrounding ions and ionic aggregates would prove beneficial for understanding ionic liquids. For multicomponent electrolyte solutions, theories must explicitly account for both dielectric properties and ionic screening. A simple mean-field model of a typical ionic liquid that explicitly considers ion-pairs as dipoles and the screened ion–ion, dipole–dipole, and dipole–ion interactions on the same footing showed that only 1/3 of ions are



More broadly, the observation of long range interactions in ionic liquids provides a series of experimental results that necessarily challenge the prevailing perspective⁴ on the relationship between electrostatic correlations and screening in ionic liquids and other highly concentrated electrolytes. At this point, there is solid evidence that the long range interactions measured in ionic liquids are truly equilibrium interactions, as opposed to complex time- and history-dependent dynamic effects.

The precise origin of long range forces in ionic liquids remains open, yet there are promising phenomenological theories^{12,13,16} as well as very recent rigorous statistical physical analyses²⁰ implicating strong electrostatic correlations between ions. As discussed above, these frameworks appear to be broadly compatible and suggest several potentially fruitful avenues for future research, with diverse implications for ionic liquids and other highly concentrated electrolytes, from designing ionic liquids for energy storage applications, to rationalizing electrostatic correlations in biological self-assembly.

Acknowledgements

This work was supported by the US Department of Energy, Office of Basic Energy Sciences, Division of Materials Science and Engineering Award DE-FG02-87ER-45331: to M. A. G. (analysis and manuscript writing), to H. A. D. (manuscript editing and discussion), and to J. N. I. (manuscript editing and discussion). M. A. G. acknowledges funding provided by the GLAM Postdoctoral Fellowship program in the Geballe Laboratory for Advanced Materials at Stanford University. S. P. is supported by The Leverhulme Trust (RPG-2015-328) and the ERC (Starting Grant 676861 LIQUISWITCH). A. A. L. is supported by a George F. Carrier Fellowship at Harvard University. AMS is supported by a Doctoral Prize from the EPSRC. R. A. and G. W. acknowledge funding support from the Australian Research Council. X. B. acknowledges the financial supports of the CRC funding program (Tiers 2) and NSERC Discovery program. MR acknowledges support from The Knut and Alice Wallenberg Foundation for ILEAP – Ionic Liquid Lubricants Enabling Advanced Performance.

References

- More broadly, the observation of long range interactions in ionic liquids provides a series of experimental results that necessarily challenge the prevailing perspective⁴ on the relationship between electrostatic correlations and screening in ionic liquids and other highly concentrated electrolytes. At this point, there is solid evidence that the long range interactions measured in ionic liquids are truly equilibrium interactions, as opposed to complex time- and history-dependent dynamic effects.
- The precise origin of long range forces in ionic liquids remains open, yet there are promising phenomenological theories^{12,13,16} as well as very recent rigorous statistical physical analyses²⁰ implicating strong electrostatic correlations between ions. As discussed above, these frameworks appear to be broadly compatible and suggest several potentially fruitful avenues for future research, with diverse implications for ionic liquids and other highly concentrated electrolytes, from designing ionic liquids for energy storage applications, to rationalizing electrostatic correlations in biological self-assembly.
- ## Acknowledgements
- This work was supported by the US Department of Energy, Office of Basic Energy Sciences, Division of Materials Science and Engineering Award DE-FG02-87ER-45331: to M. A. G. (analysis and manuscript writing), to H. A. D. (manuscript editing and discussion), and to J. N. I. (manuscript editing and discussion). M. A. G. acknowledges funding provided by the GLAM Postdoctoral Fellowship program in the Geballe Laboratory for Advanced Materials at Stanford University. S. P. is supported by The Leverhulme Trust (RPG-2015-328) and the ERC (Starting Grant 676861 LIQUISWITCH). A. A. L. is supported by a George F. Carrier Fellowship at Harvard University. AMS is supported by a Doctoral Prize from the EPSRC. R. A. and G. W. acknowledge funding support from the Australian Research Council. X. B. acknowledges the financial supports of the CRC funding program (Tiers 2) and NSERC Discovery program. MR acknowledges support from The Knut and Alice Wallenberg Foundation for ILEAP – Ionic Liquid Lubricants Enabling Advanced Performance.
- ## References
- 1 J. M. Tarascon and M. Armand, *Nature*, 2001, **414**, 359–367.
 - 2 P. Simon and Y. Gogotsi, *Nat. Mater.*, 2008, **7**, 845–854.
 - 3 B. Eisenberg, *Biophys. J.*, 2013, **104**, 1849–1866.
 - 4 M. V. Fedorov and A. A. Kornyshev, *Chem. Rev.*, 2014, **114**, 2978–3036.
 - 5 T. Welton, *Chem. Rev.*, 1999, **99**, 2071–2083.
 - 6 M. Galinski, A. Lewandowski and I. Stepniak, *Electrochim. Acta*, 2006, **51**, 5567–5580.
 - 7 C. F. Ye, W. M. Liu, Y. X. Chen and L. G. Yu, *Chem. Commun.*, 2001, 2244–2245, DOI: 10.1039/b106935g.
 - 8 M. Antonietti, D. B. Kuang, B. Smarsly and Z. Yong, *Angew. Chem., Int. Ed.*, 2004, **43**, 4988–4992.
 - 9 M. Armand, F. Endres, D. R. MacFarlane, H. Ohno and B. Scrosati, *Nat. Mater.*, 2009, **8**, 621–629.
 - 10 J. Jeong, N. Aetukuri, T. Graf, T. D. Schladt, M. G. Samant and S. S. Parkin, *Science*, 2013, **339**, 1402–1405.
 - 11 M. Z. Bazant, B. D. Storey and A. A. Kornyshev, *Phys. Rev. Lett.*, 2011, **106**, 046102.
 - 12 M. A. Gebbie, M. Valtiner, X. Banquy, E. T. Fox, W. A. Henderson and J. N. Israelachvili, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 9674–9679.
 - 13 M. A. Gebbie, H. A. Dobbs, M. Valtiner and J. N. Israelachvili, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, 7432–7437.
 - 14 H.-W. Cheng, P. Stock, B. Moeremans, T. Baimpos, X. Banquy, F. U. Renner and M. Valtiner, *Adv. Mater. Interfaces*, 2015, **2**, 1500159.
 - 15 R. M. Espinosa-Marzal, A. Arcifa, A. Rossi and N. D. Spencer, *J. Phys. Chem. Lett.*, 2014, **5**, 179–184.
 - 16 A. M. Smith, A. A. Lee and S. Perkin, *J. Phys. Chem. Lett.*, 2016, **7**, 2157–2163.
 - 17 T. Baimpos, B. R. Shrestha, S. Raman and M. Valtiner, *Langmuir*, 2014, **30**, 4322–4332.
 - 18 A. Yochelis, *Phys. Chem. Chem. Phys.*, 2014, **16**, 2836–2841.
 - 19 A. Yochelis, *J. Phys. Chem. C*, 2014, **118**, 5716–5724.
 - 20 R. Kjellander, *Phys. Chem. Chem. Phys.*, 2016, **18**, 18985–19000.
 - 21 J. N. A. C. Lopes and A. A. H. Padua, *J. Phys. Chem. B*, 2006, **110**, 3330–3335.
 - 22 A. Triolo, O. Ruzzina, H. J. Bleif and E. Di Cola, *J. Phys. Chem. B*, 2007, **111**, 4641–4644.
 - 23 J. D. Holbrey and K. R. Seddon, *J. Chem. Soc., Dalton Trans.*, 1999, 2133–2139, DOI: 10.1039/a902818h.
 - 24 R. Hayes, G. G. Warr and R. Atkin, *Phys. Chem. Chem. Phys.*, 2010, **12**, 1709–1723.
 - 25 R. G. Horn and J. N. Israelachvili, *J. Chem. Phys.*, 1981, **75**, 1400–1411.
 - 26 H. K. Christenson, *J. Chem. Phys.*, 1983, **78**, 6906–6913.
 - 27 J. N. Israelachvili and R. M. Pashley, *Nature*, 1983, **306**, 249–250.
 - 28 S. Perkin, T. Albrecht and J. Klein, *Phys. Chem. Chem. Phys.*, 2010, **12**, 1243–1247.
 - 29 L. J. D. Frink and F. van Swol, *J. Chem. Phys.*, 1998, **108**, 5588–5598.
 - 30 R. Atkin and G. G. Warr, *J. Phys. Chem. C*, 2007, **111**, 5162–5168.
 - 31 S. Perkin, L. Crowhurst, H. Niedermeyer, T. Welton, A. M. Smith and N. N. Gosvami, *Chem. Commun.*, 2011, **47**, 6572–6574.
 - 32 A. M. Smith, K. R. Lovelock, N. N. Gosvami, P. Licence, A. Dolan, T. Welton and S. Perkin, *J. Phys. Chem. Lett.*, 2013, **4**, 378–382.
 - 33 M. Mezger, H. Schroder, H. Reichert, S. Schramm, J. S. Okasinski, S. Schoder, V. Honkimaki, M. Deutsch, B. M. Ocko, J. Ralston, M. Rohwerder, M. Stratmann and H. Dosch, *Science*, 2008, **322**, 424–428.
 - 34 A. Sheehan, L. A. Jurado, S. N. Ramakrishna, A. Arcifa, A. Rossi, N. D. Spencer and R. M. Espinosa-Marzal, *Nanoscale*, 2016, **8**, 4094–4106.
 - 35 J. N. Israelachvili, *Intermolecular and surface forces*, Academic Press, Burlington, MA, 3rd edn, 2011.
 - 36 H. K. Christenson and N. H. Thomson, *Surf. Sci. Rep.*, 2016, **71**, 367–390.
 - 37 A. Elbourne, K. Voitchovsky, G. G. Warr and R. Atkin, *Chem. Sci.*, 2015, **6**, 527–536.
 - 38 R. G. Horn, D. F. Evans and B. W. Ninham, *J. Phys. Chem.*, 1988, **92**, 3531–3537.
 - 39 R. A. Asencio, E. D. Cranston, R. Atkin and M. W. Rutland, *Langmuir*, 2012, **28**, 9967–9976.
 - 40 N. Hjalmarsson, R. Atkin and M. W. Rutland, *Phys. Chem. Chem. Phys.*, 2016, **18**, 9232–9239.
 - 41 Y. Lauw, M. D. Horne, T. Rodopoulos, V. Lockett, B. Akgun, W. A. Hamilton and A. R. Nelson, *Langmuir*, 2012, **28**, 7374–7381.
 - 42 R. M. Espinosa-Marzal, A. Arcifa, A. Rossi and N. D. Spencer, *J. Phys. Chem. C*, 2014, **118**, 6491–6503.
 - 43 L. A. Jurado, H. Kim, A. Rossi, A. Arcifa, J. K. Schuh, N. D. Spencer, C. Leal, R. H. Ewoldt and R. M. Espinosa-Marzal, *Phys. Chem. Chem. Phys.*, 2016, **18**, 22719–22730.
 - 44 H. Li, R. J. Wood, F. Endres and R. Atkin, *J. Phys.: Condens. Matter*, 2014, **26**, 284115.
 - 45 R. Hayes, N. Borisenko, M. K. Tam, P. C. Howlett, F. Endres and R. Atkin, *J. Phys. Chem. C*, 2011, **115**, 6855–6863.
 - 46 Z. Wang, H. Li, R. Atkin and C. Priest, *Langmuir*, 2016, **32**, 8818–8825.
 - 47 H. W. Cheng, J. N. Dienemann, P. Stock, C. Merola, Y. J. Chen and M. Valtiner, *Sci. Rep.*, 2016, **6**, 30058.
 - 48 I. Bou-Malham and L. Bureau, *Soft Matter*, 2010, **6**, 4062.
 - 49 N. Hjalmarsson, R. Atkin and M. W. Rutland, *Chem. Commun.*, 2016, DOI: 10.1039/c6cc07396d.
 - 50 J. P. Hansen and H. Lowen, *Annu. Rev. Phys. Chem.*, 2000, **51**, 209–242.
 - 51 A. A. Lee, D. Vella, S. Perkin and A. Goriely, *J. Phys. Chem. Lett.*, 2015, **6**, 159–163.

- 52 S. Tsuzuki, H. Tokuda, K. Hayamizu and M. Watanabe, *J. Phys. Chem. B*, 2005, **109**, 16474–16481.
- 53 H. Weingaertner, *Angew. Chem., Int. Ed.*, 2008, **47**, 654–670.
- 54 H. Tokuda, S. Tsuzuki, M. A. B. H. Susan, K. Hayamizu and M. Watanabe, *J. Phys. Chem. B*, 2006, **110**, 19593–19600.
- 55 V. Lockett, M. Horne, R. Sedev, T. Rodopoulos and J. Ralston, *Phys. Chem. Chem. Phys.*, 2010, **12**, 12499–12512.
- 56 D. J. Bozym, B. Uralcan, D. T. Limmer, M. A. Pope, N. J. Szamreta, P. G. Debenedetti and I. A. Aksay, *J. Phys. Chem. Lett.*, 2015, **6**, 2644–2648.
- 57 S. Alexander, P. M. Chaikin, P. Grant, G. J. Morales, P. Pincus and D. Hone, *J. Chem. Phys.*, 1984, **80**, 5776–5781.
- 58 R. Kjellander and D. J. Mitchell, *Mol. Phys.*, 1997, **91**, 173–188.
- 59 H. J. Butt, *Biophys. J.*, 1991, **60**, 1438–1444.
- 60 U. Raviv, P. Laurat and J. Klein, *J. Chem. Phys.*, 2002, **116**, 5167–5172.
- 61 M. Todorova and J. Neugebauer, *Phys. Rev. Appl.*, 2014, **1**, 014001.
- 62 R. M. Espinosa-Marzal, T. Drobek, T. Balmer and M. P. Heuberger, *Phys. Chem. Chem. Phys.*, 2012, **14**, 6085–6093.
- 63 C. Daguenet, P. J. Dyson, I. Krossing, A. Oleinikova, J. Slattery, C. Wakai and H. Weingartner, *J. Phys. Chem. B*, 2006, **110**, 12682–12688.
- 64 V. Reus, L. Belloni, T. Zemb, N. Lutterbach and H. Versmold, *J. Phys. II*, 1997, **7**, 603–626.
- 65 D. Qiu, T. Cosgrove, A. M. Howe and C. A. Dreiss, *Langmuir*, 2006, **22**, 546–552.
- 66 J. Eastoe, B. M. H. Cazelles, D. C. Steytler, J. D. Holmes, A. R. Pitt, T. J. Wear and R. K. Heenan, *Langmuir*, 1997, **13**, 6980–6984.
- 67 E. W. Kaler, J. F. Billman, J. L. Fulton and R. D. Smith, *J. Phys. Chem.*, 1991, **95**, 458–462.
- 68 Z. He and P. Alexandridis, *Phys. Chem. Chem. Phys.*, 2015, **17**, 18238.

