

Cite this: *Chem. Sci.*, 2019, 10, 1219

All publication charges for this article have been paid for by the Royal Society of Chemistry

## Novel near-infrared II aggregation-induced emission dots for *in vivo* bioimaging†

Jiacheng Lin,<sup>‡ab</sup> Xiaodong Zeng,<sup>‡ab</sup> Yuling Xiao,<sup>‡ab</sup> Lin Tang,<sup>a</sup> Jinxia Nong,<sup>a</sup> Yufang Liu,<sup>a</sup> Hui Zhou,<sup>‡ab</sup> Bingbing Ding,<sup>a</sup> Fuchun Xu,<sup>c</sup> Hanxing Tong,<sup>d</sup> Zixin Deng<sup>a</sup> and Xuechuan Hong<sup>‡abc</sup>

Near-infrared II fluorescence imaging holds great promise for *in vivo* imaging and imaging-guided surgery with deep penetration and high spatiotemporal resolution. However, most NIR-II aromatic luminophores suffer from the notorious aggregation-caused quenching (ACQ) effect in the aqueous solution, which largely hinders their biomedical application *in vivo*. In this study, the first NIR-II organic aggregation-induced emission (AIE) fluorophore (HLZ-BTED), encapsulated as nanoparticles (HLZ-BTED dots) for *in vivo* biomedical imaging, was designed and synthesized. The NIR-II AIE HLZ-BTED dots showed high temporal resolution, high photostability, outstanding water-solubility and biocompatibility *in vitro* and *in vivo*. The HLZ-BTED dots were further used for long-term breast tumor imaging and visualizing tumor-feeding blood vessels, long-term hind limb vasculature and incomplete hind limb ischemia. More importantly, as a proof-of-concept, this is the first time that non-invasive and real-time NIR-II imaging of the gastrointestinal tract in health and disease has been performed, making the AIE dots a promising tool for gastrointestinal (GI) tract research, such as understanding the healthy status of GI peristalsis, diagnosing and evaluating intestinal motility dysfunction, and assessing drug effects on intestinal obstruction.

Received 2nd October 2018  
Accepted 9th November 2018

DOI: 10.1039/c8sc04363a

rsc.li/chemical-science

## Introduction

Optical fluorescence imaging in the near-infrared window (NIR, 700–1700 nm) is a well-established and powerful tool for biomedical application in scientific research and clinical practice, such as noninvasive *in vivo* vascular imaging,<sup>1</sup> lymphatic mapping,<sup>2</sup> cancer diagnosis,<sup>3</sup> and image-guided surgery.<sup>4</sup> Compared with the conventional first near-infrared window (NIR-I, 700–900 nm) fluorescence imaging, fluorescence imaging in the second near-infrared window (NIR-II, 1000–1700 nm) has deeper penetration and higher spatiotemporal resolution due to low photon absorption and scattering and low auto-fluorescence of normal biological tissues.<sup>5</sup> Numerous organic or inorganic nanostructured NIR-II contrast agents

such as rare earth nanoparticles (NPs),<sup>6</sup> quantum dots (QDs),<sup>7</sup> carbon nanotubes (CNTs),<sup>8</sup> and conjugated polymer NPs<sup>9</sup> have been reported for biological imaging. Recently, the first rapidly renally excreted NIR-II fluorophore **CH1055** has been developed based on a benzobisthiadiazole (BBTD)-based donor–acceptor–donor (D–A–D) structure for high-quality NIR-II biomedical imaging.<sup>10</sup> After that, a series of small-molecule organic NIR-II dyes were reported for use in bioimaging with or without the BBTD core.<sup>11</sup> However, most of these NIR-II aromatic luminophores suffer from the notorious aggregation-caused quenching (ACQ) effect, with strong emission as isolated molecules but poor emission efficiency in the aggregate state due to intense intermolecular  $\pi$ – $\pi$  stacking interactions, which largely hinders their biomedical application in NIR-II imaging.

Research efforts to solve the ACQ problem by hampering chromophore aggregation have ended with limited success until 2001.<sup>12</sup> A novel AIE phenomenon, an exactly opposite phenomenon to overcome the notorious ACQ effect was reported by Tang and his co-workers.<sup>12</sup> The luminescent molecules with AIE characteristics usually show strong fluorescence emission upon aggregation, and weak or non-emissive fluorescence in dilute solvents. So far, a variety of small fluorescent organic molecules have been developed as AIE fluorogens (AIEgens) with visible (Vis) or NIR-I fluorescence emission.<sup>13</sup> Recently, two short-wave infrared (SWIR, 900–1700 nm) AIEgens **TQ-BPN** and **TB1** were reported with the maximum fluorescence

<sup>a</sup>State Key Laboratory of Virology, Key Laboratory of Combinatorial Biosynthesis and Drug Discovery (MOE), Hubei Provincial Key Laboratory of Developmentally Originated Disease, Wuhan University School of Pharmaceutical Sciences, Wuhan 430071, China. E-mail: xhy78@whu.edu.cn

<sup>b</sup>Shenzhen Institute of Wuhan University, Shenzhen, 518057, China

<sup>c</sup>Innovation Center for Traditional Tibetan Medicine Modernization and Quality Control, Medical College, Tibet University, Lhasa, 850000, China

<sup>d</sup>Department of General Surgery, Zhongshan Hospital, Fudan University, Shanghai, 200032, China

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8sc04363a

‡ These authors contributed equally to this work.












Fig. 4 Biocompatibility study of the HLZ-BTED dots. (A) Cell viability of 4T1 and L929 cells after incubation with different concentrations of the HLZ-BTED dots for 24 h ( $n = 3$ ). (B) Body weight of normal mice on different days ( $n = 3$ ) treated with PBS,  $7.5 \text{ mg kg}^{-1}$ , and  $15 \text{ mg kg}^{-1}$  HLZ-BTED dots. (C) Representative hematoxylin and eosin stained images of major organs (heart, liver, spleen, lungs, and kidneys) from the control mice and HLZ-BTED dot injected mice at 31 days post-treatment. Scale bar:  $100 \mu\text{m}$ .




Fig. 5 Representative *in vivo* long-term NIR-II fluorescence images (808 nm excitation,  $90 \text{ mW cm}^{-2}$ ,  $1000 \text{ nm LP}$ ,  $100 \text{ ms}$ ,  $n = 3$ ) of the 4T1 breast tumor model at different time points after tail vein injection of the HLZ-BTED dots ( $0.2 \text{ mL}$ ,  $10 \text{ mg kg}^{-1}$ ). The white circles indicate the location of the 4T1 tumor. Scale bar:  $1 \text{ cm}$ .


organs, indicating that the HLZ-BTED dots have excellent biocompatibility and are more applicable for NIR-II bioimaging *in vivo*.

Breast cancer is the most common cancer in women worldwide.<sup>19</sup> Early diagnosis of breast cancer with the use of systemic therapy has an excellent prognosis and increases the long-term disease-free survival (DFS).<sup>20</sup> Firstly, the *in vitro* cellular endocytosis of the HLZ-BTED dots was studied using 4T1 breast cancer cells. As shown in Fig. S7,† the NIR-II fluorescence intensity of the 4T1 cells incubated with the HLZ-BTED dots showed explicit time dependence, suggesting that the AIE dots can permeate into the cells *via* the cellular membrane. To explore the ability of the HLZ-BTED dots to diagnose breast cancer early, a mouse model of breast cancer was established. 4T1 breast cancer cells (roughly  $2 \times 10^6$ ) were subcutaneously injected into the right leg of female BALB/c mice. NIR-II fluorescence imaging of 4T1 breast tumors was carried out by intravenously injecting the AIE dots ( $0.2 \text{ mL}$ ,  $10 \text{ mg kg}^{-1}$ ) into 4T1 tumor-bearing mice ( $n = 3$ ) through the tail vein when the tumors reached  $\sim 60 \text{ mm}^3$  in a week. The fluorescence intensity ( $1000 \text{ nm LP}$ ,  $100 \text{ ms}$ ) in tumors reached the highest level at 48 h post-injection under 808 nm excitation with a power density of  $90 \text{ mW cm}^{-2}$  (Fig. 5 and S8†). For up to 8 days post-injection, the NIR-II fluorescence signal of tumors was extremely strong and clearly delineated the normal tissues, indicating that the AIE HLZ-BTED dots are adequate for long-term tumor imaging and real-time image-based monitoring for solid tumor therapy. An *ex vivo* biodistribution study was performed for the HLZ-BTED dots to evaluate their distribution in major organs at 192 h (Fig. S9†). High accumulation was observed in the liver

and spleen, which suggested that the clearance routes of the HLZ-BTED dots were through the hepatobiliary system. In addition, the AIE dots showed high accumulation in 4T1 tumors for up to 8 days (Fig. S9†).

Transcatheter arterial embolization (TAE) is widely used in the clinic to increase survival for cancer patients who are not suitable for chemotherapy and surgical resection.<sup>21</sup> To estimate the potential feasibility of the HLZ-BTED dots as a NIR-II contrast agent for the *in vivo* imaging of the tumor-feeding blood vessels to implement TAE, the AIE HLZ-BTED dots were *i.v.* injected into 4T1 tumor bearing BALB/c mice ( $n = 3$ ). At 2 min post-injection, the tumor vasculature was visualized using an InGaAs camera with  $1000 \text{ nm}$  (Fig. 6A,  $200 \text{ ms}$  exposure time) and  $1250 \text{ nm}$  (Fig. 6B,  $800 \text{ ms}$  exposure time) long-pass filters under 808 nm laser excitation. The NIR-II fluorescence images of the 4T1 tumor blood vessels showed a more distinct vascular network with the  $1250 \text{ nm}$  long-pass filter than with the  $1000 \text{ nm}$  one, owing to both reduced biological auto-fluorescence and photon scattering in the longer wavelength NIR-II region. Meanwhile a similar image quality *via* different long-pass filters has been demonstrated in the hind limb vasculature of C57BL/6 mice (Fig. 6C and D). Moreover, we performed a surgery to induce incomplete hind limb ischemia to mimic the NIR-II fluorescence imaging-guided vascular embolization procedure. The occlusion site of the blood supply vessels was precisely monitored by NIR-II imaging using the HLZ-BTED dots (Fig. 6E). Then the capability of long-term vasculature imaging using the HLZ-BTED dots was investigated. As shown in Fig. 6F, at 4 h post-injection, the vasculature was still clearly visualized through NIR-II imaging which would be adequate for surgical operation. The resolution ( $497.6 \mu\text{m}$ ) of the hind limb





**Fig. 6** (A and B) *In vivo* visualization of tumor-feeding vessels. NIR-II tumor blood vessel fluorescence images of the 4T1 breast tumor-bearing mice were obtained at 2 min post tail vein injection of the HLZ-BTED dots (0.2 mL, 10 mg kg<sup>-1</sup>) with 1000 nm (A), 200 ms, 808 nm excitation, 90 mW cm<sup>-2</sup> and 1250 nm (B), 800 ms, 808 nm excitation, 180 mW cm<sup>-2</sup> long-pass filters, respectively. The red circles indicate the location of the tumor, and yellow arrows indicate the tumor-feeding arteries. (C and D) NIR-II fluorescence images of the hind limb vessels in C57BL/6 mice at 2 min after tail vein injection of the HLZ-BTED dots (0.2 mL, 15 mg kg<sup>-1</sup>) with 1000 nm (C), 60 ms, 808 nm excitation, 90 mW cm<sup>-2</sup> and 1250 nm (D), 500 ms, 808 nm excitation, 180 mW cm<sup>-2</sup> long-pass filters, respectively. (E) *In vivo* NIR-II fluorescence images (1250 nm LP, 400 ms, 808 nm excitation, 180 mW cm<sup>-2</sup>) of incomplete left hind limb ischemia pre-injection and 60 s, 120 s, and 180 s after HLZ-BTED dot injection, where the white arrow located the occlusion site in the femoral artery. (F) Intravital long-term hind limb vasculature NIR-II imaging (1250 nm LP, 400 ms, 808 nm excitation, 180 mW cm<sup>-2</sup>) from 0 min to 240 min after tail vein injection of the HLZ-BTED dots. Scale bar (A–F): 1 cm. (G) The vessel FWHM width based on the cross-sectional intensity profile measured along the white line in (F) (1 min) with the peak fitted to Gaussian functions (the red curve is the Gaussian fit to the profile).

vasculature was investigated *via* the Gaussian-fitted full width at half maximum (FWHM), which further confirmed the excellent ability of the HLZ-BTED dots to precisely map vascular vessels with NIR-II imaging (Fig. 6G).

The gastrointestinal (GI) tract is an organ system, which includes all organ structures between the mouth and the anus, such as the esophagus, stomach, small intestine, and large intestine. The GI tract is responsible for ingesting and digesting food, extracting and absorbing nutrients, and expelling the waste as feces. Recently, non-invasive imaging modalities such as positron emission tomography (PET),<sup>22</sup> magnetic resonance imaging (MRI),<sup>23</sup> X-ray computed tomography (CT),<sup>24</sup> and photoacoustic (PA)<sup>22</sup> imaging, have played an important role in GI function research and the diagnosis and prognosis of GI diseases. However, these imaging modalities are limited by their serious drawbacks, notably radiation risk, expensive instrument cost, long imaging times and limited spatial resolution for real-time gut function study such as physiological or

pathological intestinal motility. NIR-II fluorescence imaging may provide a reliable method for monitoring intestinal motility disorders or dysfunction and providing evaluation information of GI therapeutic agents *in vivo* due to superior temporal-spatial resolution.<sup>25</sup> Therefore, as a proof of concept, we demonstrated the feasibility of the AIE HLZ-BTED dots for *in vivo* NIR-II imaging of the GI tract by oral administration. First, the fluorescence intensity and hydrodynamic sizes of the HLZ-BTED dots in different pH buffers ranging from 1 to 9 or in gastric pH 1 buffer were investigated respectively. No significant difference was observed between different conditions, demonstrating the excellent stability and capability of the AIE dots as a GI tract contrast agent (Fig. S10†). Second, healthy BALB/c mice ( $n = 3$ ) were gavaged with the HLZ-BTED dots (0.1 mL, 5 mg kg<sup>-1</sup>) for real-time imaging of intestinal motility and monitoring the behavior of micelles within the GI tract. NIR-II fluorescence images at different time points (5, 10, 20, 30, 60, 90, 180, 480, 600 and 1440 min) were obtained under 808 nm laser irradiation (180 mW cm<sup>-2</sup>, within the safety limits (329 mW cm<sup>-2</sup>) determined by the International Commission on Non-ionizing Radiation Protection) through an InGaAs camera (200 ms exposure time) with a 1250 nm long-pass filter. As shown in Fig. 7A, from 5 min to 480 min after gavaging, the fluorescent signals were observed in the stomach, duodenum, jejunum, ileum, cecum, and colon. During the imaging procedure of the intestine, the contractile function of the intestine was demonstrated by the recorded Movie S1.† The fluorescence intensity movement in the small intestine indicated vigorous



**Fig. 7** (A and B) Representative non-invasive NIR-II fluorescence images (1250 nm LP, 200 ms, 808 nm excitation, 180 mW cm<sup>-2</sup>) of the gastrointestinal tract (GI tract) in BALB/c mice gavaged with the HLZ-BTED dots (0.1 mL, 5 mg kg<sup>-1</sup>) for real-time monitoring of gastrointestinal peristalsis of normal mice (A) and mice anesthetized (B) using pentobarbital sodium ( $n = 3$  mice). The white arrow indicates the feces of mice (Fig. 7A, 600 min). Scale bar: 1 cm.



contractility with peristalsis and segmental contractions of the intestine at 330 min after gavaging. By fluorescence intensity map analysis of regions of interest (ROIs),<sup>25f</sup> the intestine peristalsis frequency and propagation velocity was about 28–40 cycles  $\text{min}^{-1}$  and  $2.61 \pm 0.23 \text{ cm s}^{-1}$ , respectively. At 600 min after gavaging, the fluorescent signal was first detected in feces, indicating that the NIR-II dots moved out of the body from the rectum and anus. At 24 h after gavaging, no fluorescent signal of the abdomen was detected. These results illustrated that the **HLZ-BTED** dots can be totally excreted through the GI tract within 24 h and are not absorbed into the body through the intestine. The **HLZ-BTED** dots used in GI tract imaging are excreted quickly due to low absorption in the GI tract while the **HLZ-BTED** dots would interact with serum plasma and the immune system by i.v. injection, leading to the slow clearance routes of the **HLZ-BTED** dots through the hepatobiliary system. Moreover, in Fig. 7B and 8C, the experimental results showed that the use of an anesthetic, pentobarbital sodium, could suppress duodenum motility and gastric emptying, indicating that the NIR-II **HLZ-BTED** dots could be a powerful tool for assessing the effects of drugs on the GI tract.

To investigate the ability of the **HLZ-BTED** dots to diagnose intestinal diseases such as obstructions which impede or block peristaltic movement, we built a mouse model of intestinal partial obstruction according to the reported methods.<sup>26</sup> Then, we gavaged the mice with intestinal obstruction with the **HLZ-BTED** dots (0.1 mL,  $5 \text{ mg kg}^{-1}$ ). The NIR-II images shown in Fig. 8A and 7A exhibited differences in the intestinal distribution of dots between the mice with obstruction and healthy mice. At 5 h after gavaging, the lower small intestine was fully filled and distended after dissection, and the fluorescent signal was strong in the upper and medial small intestine (Fig. 8B), indicating that the **HLZ-BTED** dots were accumulated in these parts and are suitable for diagnosing intestinal obstruction.



Fig. 8 (A) Representative non-invasive NIR-II fluorescence images (1250 nm LP, 200 ms, 808 nm excitation,  $180 \text{ mW cm}^{-2}$ ) of the gastrointestinal tract in BALB/c mice gavaged with the **HLZ-BTED** dots ( $0.1 \text{ mL}$ ,  $5 \text{ mg kg}^{-1}$ ) for visualizing intestinal obstruction ( $n = 3$  mice). (B) *Ex vivo* NIR-II image (200 ms, 1250 nm LP) of the GI tract in normal mice or mice with intestinal obstruction at 5 h after gavage. (C) Representative *ex vivo* NIR-II fluorescence image (1250 nm LP, 200 ms) of the GI tract in BALB/c mice which are normal or anesthetized by pentobarbital sodium at 2 h after gavage. Scale bar: 1 cm.

## Experimental section

All animal experiments were performed according to the Chinese Regulations for the Administration of Affairs Concerning Experimental Animals and approved by the Institutional Animal Care and Use Committee (IACUC) of Wuhan University. And all experimental details are provided in the ESI.†

## Conclusions

In conclusion, we have successfully designed and synthesized a novel NIR-II organic fluorophore **HLZ-BTED** with AIE features. The organic fluorophore could be encapsulated as **HLZ-BTED** dots with good water-solubility and biocompatibility for *in vivo* NIR-II biomedical imaging. We further performed *in vivo* NIR-II imaging experiments that revealed the excellent passive targeted long-term NIR-II fluorescence imaging performance of the **HLZ-BTED** dots in breast tumor-bearing mice. Moreover, the **HLZ-BTED** dots were an excellent NIR-II imaging contrast agent for superior spatial-temporal resolution of tumor-feeding vessels imaged with a 1250 nm long-pass filter, precisely monitoring the occlusion site of incomplete hind limb ischemia and long-term hind limb vasculature, and have the potential capability for image-guided transcatheter arterial embolization. To the best of our knowledge, as a proof-of-concept, this is the first time that non-invasive and real-time NIR-II imaging of the gastrointestinal tract in health and in disease has been performed. Furthermore, dynamic NIR-II fluorescence imaging with gavage of the **HLZ-BTED** dots can be used to monitor intestinal motility in real-time, and thus provides a method for the diagnosis of intestinal motility disorders and information on the response of the intestine to therapeutic agents.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This work was partially supported by grants from the NSFC (81773674, 81573383, and 21390402), NSFHP (2017CFA024, 2017CFB711, and 2016ACA126), ABRPSTCS (SYG201521), NSFJP (BK20160387), Shenzhen Science and Technology Research Grant (JCYJ20170303170809222), Fundamental Research Funds for the Central Universities and Open Research Fund Program of the Hubei Province Engineering and Technology Research Center for Fluorinated Pharmaceuticals.

## Notes and references

- (a) X. D. Zhang, H. Wang, A. L. Antaris, L. Li, S. Diao, R. Ma, A. Nguyen, G. Hong, Z. Ma, J. Wang, S. Zhu, J. M. Castellano, T. Coray, Y. Liang, J. Luo and H. Dai, *Adv. Mater.*, 2016, **28**, 6872–6879; (b) H. Wan, J. Yue, S. Zhu, T. Uno, X. Zhang, Q. Yang, K. Yu, G. Hong, J. Wang, L. Li, Z. Ma, H. Gao,





- Y. Zhong, J. Su, A. L. Antaris, Y. Xia, J. Luo, Y. Liang and H. Dai, *Nat. Commun.*, 2018, **9**, 1171.
- 2 (a) E. C. Rossi, A. Ivanova and J. F. Boggess, *Gynecol. Oncol.*, 2012, **124**, 78–82; (b) G. Wishart, S.-W. Loh, L. Jones and J. Benson, *Eur. J. Surg. Oncol.*, 2012, **38**, 651–656; (c) C. Hirche, D. Murawa, Z. Mohr, S. Kneif and M. Hünerbein, *Breast Cancer Res. Treat.*, 2010, **121**, 373–378.
- 3 (a) M. Herranz and A. Ruibal, *J. Oncol.*, 2012, **2012**, 863747; (b) C. Zeng, W. Shang, X. Liang, X. Liang, Q. Chen, C. Chi, Y. Du, C. Fang and J. Tian, *ACS Appl. Mater. Interfaces*, 2016, **8**, 29232–29241; (c) S. Luo, E. Zhang, Y. Su, T. Cheng and C. Shi, *Biomaterials*, 2011, **32**, 7127–7138; (d) J. Yang, Q. Xie, H. Zhou, L. Chang, W. Wei, Y. Wang, H. Li, Z. Deng, Y. Xiao, J. Wu, P. Xu and X. Hong, *J. Proteome Res.*, 2018, **17**, 2428–2439.
- 4 (a) Y. Sun, X. Zeng, Y. Xiao, C. Liu, H. Zhu, H. Zhou, Z. Chen, F. Xu, J. Wang, M. Zhu, J. Wu, M. Tian, H. Zhang, Z. Deng, Z. Cheng and X. Hong, *Chem. Sci.*, 2018, **9**, 2092–2097; (b) Y. Sun, M. Ding, X. Zeng, Y. Xiao, H. Wu, H. Zhou, B. Ding, C. Qu, W. Hou, A. Erbu, Y. Zhang, Z. Cheng and X. Hong, *Chem. Sci.*, 2017, **8**, 3489–3493.
- 5 (a) F. Ding, Y. Zhan, X. Lu and Y. Sun, *Chem. Sci.*, 2018, **9**, 4370–4380; (b) G. Hong, A. L. Antaris and H. Dai, *Nat. Biomed. Eng.*, 2017, **1**, 0010; (c) F. Ding, S. Chen, W. Zhang, Y. Tu and Y. Sun, *Bioorg. Med. Chem.*, 2017, **25**, 5179–5184; (d) Y. Xu, M. Tian, H. Zhang, Y. Xiao, X. Hong and Y. Sun, *Chin. Chem. Lett.*, 2018, **29**, 1093–1097; (e) S. He, J. Song, J. Qu and Z. Cheng, *Chem. Soc. Rev.*, 2018, **47**, 4258–4278; (f) W. Zhu, *Sci. China: Chem.*, 2016, **59**, 203–204; (g) B. Li, L. Lu, M. Zhao, Z. Lei and F. Zhang, *Angew. Chem., Int. Ed.*, 2018, **57**, 7483–7487; (h) H. Zhou, Y. Xiao and X. Hong, *Chin. Chem. Lett.*, 2018, **29**, 1425–1428; (i) J. Yang and X. Hong, *Sci. China: Chem.*, 2018, DOI: 10.1007/s11426-018-9341-7.
- 6 (a) L. Liu, S. Wang, B. Zhao, P. Pei, Y. Fan, X. Li and F. Zhang, *Angew. Chem.*, 2018, **130**, 7640–7644; (b) Y. Fan, P. Wang, Y. Lu, R. Wang, L. Zhou, X. Zheng, X. Li, J. A. Piper and F. Zhang, *Nat. Nanotechnol.*, 2018, **13**, 941–946; (c) P. Wang, Y. Fan, L. Lu, L. Liu, L. Fan, M. Zhao, Y. Xie, C. Xu and F. Zhang, *Nat. Commun.*, 2018, **9**, 2898; (d) Y. Zhong, Z. Ma, S. Zhu, J. Yue, M. Zhang, A. L. Antaris, J. Yuan, R. Cui, H. Wan, Y. Zhou, W. Wang, N. F. Huang, J. Luo, Z. Hu and H. Dai, *Nat. Commun.*, 2017, **8**, 737; (e) X. Jiang, C. Cao, W. Feng and F. Li, *J. Mater. Chem. B*, 2016, **4**, 87–95; (f) D. Naczynski, M. Tan, M. Zevon, B. Wall, J. Kohl, A. Kulesa, S. Chen, C. Roth, R. Riman and P. Moghe, *Nat. Commun.*, 2013, **4**, 2199.
- 7 (a) M. Zhang, J. Yue, R. Cui, Z. Ma, H. Wan, F. Wang, S. Zhu, Y. Zhou, Y. Kuang, Y. Zhong, D.-W. Pang and H. Dai, *Proc. Natl. Acad. Sci. U. S. A.*, 2018, 201806153; (b) Z. Ma, M. Zhang, J. Yue, C. Alcazar, Y. Zhong, T. C. Doyle, H. Dai and N. F. Huang, *Adv. Funct. Mater.*, 2018, **28**, 1803417; (c) G. Chen, S. Lin, D. Huang, Y. Zhang, C. Li, M. Wang and Q. Wang, *Small*, 2018, **14**, 1702679; (d) C. Li, Y. Zhang, G. Chen, F. Hu, K. Zhao and Q. Wang, *Adv. Mater.*, 2017, **29**, 1605754; (e) G. Chen, F. Tian, C. Li, Y. Zhang, Z. Weng, Y. Zhang, R. Peng and Q. Wang, *Biomaterials*, 2015, **53**, 265–273; (f) G. Hong, J. T. Robinson, Y. Zhang, S. Diao, A. L. Antaris, Q. Wang and H. Dai, *Angew. Chem., Int. Ed.*, 2012, **51**, 9818–9821; (g) Y. Zhang, G. Hong, Y. Zhang, G. Chen, F. Li, H. Dai and Q. Wang, *ACS Nano*, 2012, **6**, 3695–3702.
- 8 (a) G. Hong, S. Diao, J. Chang, A. L. Antaris, C. Chen, Z. Bo, Z. Su, D. N. Atochin, P. L. Huang, K. I. Andreasson, C. J. Kuo and H. Dai, *Nat. Photonics*, 2014, **8**, 723–730; (b) J. T. Robinson, K. Welscher, S. M. Tabakman, S. P. Sherlock, H. Wang, R. Luong and H. Dai, *Nano Res.*, 2010, **3**, 779–793.
- 9 (a) K. Shou, Y. Tang, H. Chen, S. Chen, L. Zhang, A. Zhang, Q. Fan, A. Yu and Z. Cheng, *Chem. Sci.*, 2018, **9**, 3105–3110; (b) G. Hong, Y. Zou, A. L. Antaris, S. Diao, D. Wu, K. Cheng, X. Zhang, C. Chen, B. Liu, Y. He, J. Z. Wu, J. Yuan, B. Zhang, Z. Tao, C. Fukunaga and H. Dai, *Nat. Commun.*, 2014, **5**, 4206.
- 10 (a) A. L. Antaris, H. Chen, K. Cheng, Y. Sun, G. Hong, C. Qu, S. Diao, Z. Deng, X. Hu, B. Zhang, X. Zhang, O. K. Yaghi, Z. R. Alamparambil, X. Hong, Z. Cheng and H. Dai, *Nat. Mater.*, 2016, **15**, 235–242; (b) A. L. Antaris, H. Chen, S. Diao, Z. Ma, Z. Zhang, S. Zhu, J. Wang, A. X. Lozano, Q. Fan, L. Chew, M. Zhu, K. Cheng, X. Hong, H. Dai and Z. Cheng, *Nat. Commun.*, 2017, **8**, 15269; (c) Y. Feng, S. Zhu, A. L. Antaris, H. Chen, Y. Xiao, X. Lu, L. Jiang, S. Diao, K. Yu, Y. Wang, S. Herraiz, J. Yue, X. Hong, G. Hong, Z. Cheng, H. Dai and A. J. Hsueh, *Chem. Sci.*, 2017, **8**, 3703–3711.
- 11 (a) K. Shou, C. Qu, Y. Sun, H. Chen, S. Chen, L. Zhang, H. Xu, X. Hong, A. Yu and Z. Cheng, *Adv. Funct. Mater.*, 2017, **27**, 1700995; (b) S. Zhu, Q. Yang, A. L. Antaris, J. Yue, Z. Ma, H. Wang, W. Huang, H. Wan, J. Wang, S. Diao, B. Zhang, X. Li, Y. Zhong, K. Yu, G. Hong, J. Luo, Y. Liang and H. Dai, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, 962–967; (c) Q. Yang, Z. Ma, H. Wang, B. Zhou, S. Zhu, Y. Zhong, J. Wang, W. Hao, A. Antaris, M. Rui, X. Zhang, J. Yang, X. Zhang, H. Sun, W. Liu, Y. Liang and H. Dai, *Adv. Mater.*, 2017, **29**, 1605497; (d) Y. Sun, C. Qu, H. Chen, M. He, C. Tang, K. Shou, S. Hong, M. Yang, Y. Jiang, B. Ding, Y. Xiao, L. Xing, X. Hong and Z. Cheng, *Chem. Sci.*, 2016, **7**, 6203–6207; (e) Q. Yang, Z. Hu, S. Zhu, R. Ma, H. Ma, Z. Ma, H. Wan, T. Zhu, Z. Jiang, W. Liu, L. Jiao, H. Sun, Y. Liang and H. Dai, *J. Am. Chem. Soc.*, 2018, **140**, 1715–1724; (f) S. Zhu, S. Herraiz, J. Yue, M. Zhang, H. Wan, Q. Yang, Z. Ma, Y. Wang, J. He, A. L. Antaris, Y. Zhong, S. Diao, Y. Feng, Y. Zhou, K. Yu, G. Hong, Y. Liang, A. J. Hsueh and H. Dai, *Adv. Mater.*, 2018, **30**, 1705799.
- 12 J. Luo, Z. Xie, J. W. Lam, L. Cheng, H. Chen, C. Qiu, H. S. Kwok, X. Zhan, Y. Liu, D. Zhu and B. Z. Tang, *Chem. Commun.*, 2001, 1740–1741.
- 13 (a) Z. Chen, J. Zhang, M. Song, J. Yin, G.-A. Yu and S. H. Liu, *Chem. Commun.*, 2015, **51**, 326–329; (b) J. Mei, N. L. Leung, R. T. Kwok, J. W. Lam and B. Z. Tang, *Chem. Rev.*, 2015, **115**, 11718–11940; (c) G. Feng and B. Liu, *Acc. Chem. Res.*, 2018, **51**, 1404–1414; (d) J. Shi, Q. Deng, C. Wan, M. Zheng, F. Huang and B. Tang, *Chem. Sci.*, 2017, **8**, 6188–6195; (e) T. He, N. Niu, Z. Chen, S. Li, S. Liu and J. Li, *Adv. Funct. Mater.*, 2018, **28**, 1706196; (f) Q. Hu, M. Gao, G. Feng and



