

## REVIEW

View Article Online  
View Journal | View Issue



Cite this: *Org. Biomol. Chem.*, 2021, **19**, 9527

## Molecular cages for biological applications

Lucía Tapia, Ignacio Alfonso \* and Jordi Solà \*

Artificial receptors able to recognise biologically relevant molecules or ions have gained interest in the chemical community because they offer a plethora of possibilities. Molecular cage compounds are polycyclic compounds with a cavity designed for the encapsulation of guest species. Once inside the host cavity, the substrate can be transported through membranes and protected from the action of enzymes or other reactive species, thus offering the possibility of interfering with biological systems. Commonly, enzymes have been an inspiration for chemists in the search and design of defined cavities for different purposes. However, the chemical preparation of molecular cages has struggled with many synthetic challenges but this effort is worthwhile as they are a very promising tool for many applications ranging from sensing, delivery, purification or even promotion of/prevention from chemical modifications. Since the early reports at the end of the 60s, this field has experienced a growing interest; this review summarises the progress in the preparation and study of cage-like compounds highlighting their importance in biological applications.

Received 5th September 2021,  
Accepted 6th October 2021

DOI: 10.1039/d1ob01737c

rsc.li/obc

## Introduction

Non-covalent interactions are ubiquitous and govern almost every aspect of biological systems, from replication and transcription of genetic material to the structure and activity of biomacromolecules. In fact, the formation of host-guest

systems can control the activity of proteins through the binding of an effector thus promoting transport, enzyme activity or starting signalling cascades.<sup>1,2</sup> For years, supramolecular chemists have been focused on the design, preparation and study of small host-guest complexes, aiming to someday reach this complexity found in biological systems. However, mimicking the naturally occurring processes with artificial receptors remains a challenge. Inspired by nature, chemists have prepared compounds to specifically recognise relevant molecules in a biological context. The most conventional

Department of Biological Chemistry, Institute for Advanced Chemistry of Catalonia, IQAC-CSIC, Jordi Girona 18-26, 08034 Barcelona, Spain.  
E-mail: ignacio.alfonso@iqac.csic.es, jordi.sola@iqac.csic.es



Lucía Tapia

*Lucía Tapia studied Chemistry at the University of Barcelona (Spain). After completing her degree she took a master in Pharmaceutical Chemistry at IQS (Ramon Llull university, Spain). In 2017 she started her research in Supramolecular Chemistry in the Institute for Advanced Chemistry of Catalonia as a PhD student, obtaining the PhD degree in 2021. Her research interests are in the field of Supramolecular*

*Chemistry particularly in the synthesis and study of synthetic receptors for biomolecules with potential biological applications.*



Ignacio Alfonso

*Ignacio Alfonso studied Chemistry at the University of Oviedo (Spain), obtaining his PhD at the same University in 1999. After two post-doctoral studies at The Scripps Research Institute in La Jolla (USA, 2000–2002) and back to the University of Oviedo (2002–2004), he obtained the Ramón y Cajal contract and moved to Universitat Jaume I (Castellón). Since 2007 he holds a permanent position at*

*IQAC-CSIC in Barcelona, where he leads the Supramolecular Chemistry group. His research interests are all fields related to Supramolecular Chemistry, especially the molecular recognition of biomolecules and the study of dynamic complex systems.*



This review aims to present a broad vision of the field, from historical breakthroughs to recent developments in polymacrocyclic compounds and their application in the study or modulation of biological processes.

Metal ions are extensively present in biological systems. They participate in many processes involved in life such as catalysis, transport, metabolic regulation or cell signalling. Dysregulation of these ions can cause several diseases and an excessive accumulation also can result in toxicity. Thus, detection and imaging of metal ion concentrations in biological media is a hot topic in biochemical research. As mentioned above, cryptands were probably the first three-dimensional cavities designed for ion recognition. These molecules show a great affinity for alkali metals even in aqueous solutions. The binding constants can be up to 5 orders of magnitude higher than their monocyclic analogues. Cryptands also show selectivity depending on their size, as crown ethers do.<sup>28</sup> The nature of the cation–ligand interactions will also determine their affinity towards different cations. Ion–dipole interactions dominate in the case of oxygen atoms, favouring the recognition of alkali cations (hard electropositive cations). On the other hand, dative coordinative bonds dominate with nitrogen sites, being favourable for the binding of transition metal ions (soft cations).<sup>28</sup>

These important features combined make cryptands a suitable platform for the development of selective chemosensors. Conjugation of cryptands with different fluorophores such as anthracene allows the detection of cations thanks to PET mechanisms.<sup>29–31</sup> Thus, the detection of metal cations was one of the first examples of the use of cryptands in aqueous media. Not surprisingly, these compounds evolved to permit the monitoring of cations in real life samples.

Potassium is the major cation inside the cells and plays an important role in several processes like maintenance of electrolyte balance, acid–base homeostasis, hormone secretion or glucose and insulin metabolism to name a few. Detection and



## Jordi Solà

*Jordi Solà received a PhD in Chemistry from the University of Barcelona in 2006. After, he moved to UK where he performed postdoctoral studies with Prof. J. Clayden (University of Manchester) and later with Prof. D. A. Leigh (University of Edinburgh). In 2012 he received Ramon y Cajal Fellowship and joined the Institute for Advanced Chemistry of Catalonia as independent researcher. Since 2017 he is Científico Titular at*

*the same institute belonging to the CSIC. His research interest interests are Supramolecular Chemistry and Dynamic Chemical Systems.*

monitoring of potassium levels is critical to diagnose and understand some pathologies related to  $K^+$  levels. In a pioneer work, He and co-workers described a chemical sensor suitable for the measurement of extracellular potassium either in serum or blood samples based on a triazacryptand (TAC, 1) and a *o*-alkoxyaniline PET donor (compound 2, Fig. 1).<sup>32</sup> This cryptand overcomes the use of PBFI (potassium binding bensofurane isophthalate), which uses an aza-18-crown-6 ether derivative as the recognition unit. This molecule is sensitive to potassium concentrations and allows to measure  $K^+$  flux in cells but, despite being commercially available, it has a poor selectivity in front of  $Na^+$ .<sup>33</sup> Conversely, He's sensor responds rapidly and reversibly to changes in  $K^+$  concentrations typical of blood samples and without interference of other analytes or to variations of pH (Fig. 1). The importance of this invention is highlighted by its use in the Roche OPTI CCA analyser, a commercially available blood analyser.

Following this breakthrough, a number of TACs has been designed and synthesised for sensing and biological imaging.



Fig. 1 (A) Triaazacryptand[2,2,3]-1-(2-methoxyethoxy) benzene (TAC). (B) He's design for an optical sensor. (C) Excitation and emission spectra of a sensor disk exposed to different potassium chloride solutions in TRIS-HEPES buffer at pH 7.40 (left, excitation spectra, fixed emission at 540 nm; right, emission spectra, fixed excitation at 470 nm. Reproduced with permission ref. 32, copyright American Chemical Society.

Verkman group prepared a series of sensors based on this ligand. For example, they reported the preparation of a water-soluble,  $K^+$  sensor, consisting of triazacryptand coupled to 3,6-bis(dimethylamino)xanthylium (TAC-Red 3, Fig. 2). The sensor showed specificity for  $K^+$ ,  $Rb^+$  and  $Cs^+$  over  $Na^+$  and  $Li^+$ ; the titration experiments showed an estimated lower detection limit of 0.2 mM for  $K^+$ . Very importantly, divalent cations showed no fluorescence response. The long-wavelength excitation and bright fluorescence minimized interference from other blood components. As this compound was not cell-permeable they were able to stain the extracellular space of mouse brain by exposing the brain dura to a solution containing the sensor. Thus, after stimulation, the sensor was able to visualise  $K^+$  waves at the cortical surface *in vivo*.<sup>34</sup> Later on, the same group reported the preparation of a green-fluorescing dextran conjugate of TAC-bodipy 4 useful for measuring  $K^+$  transport across cell membranes and selective against  $Cl^-$  or  $Na^+$ .<sup>35</sup> Moreover, they developed a TAC-lime sensor (with green fluorescence) dextran conjugate, containing tetramethylrhodamine as reference chromophore with  $K^+$  red fluorescence. The dual probe allowed the determination of  $K^+$  in airway surface liquid (the thin fluid layer lining airway surface epithelial cells).<sup>36</sup> Interestingly, the group also reported modifications of the triazacryptand (TAC) moiety that facilitated the synthesis greatly improving the yield and without losing recognition properties by removing 2 methyl groups of the sensor (ddTAC-CHO, 6)<sup>37</sup>



Fig. 2 Different TAC-conjugates described by Verkman and Meldrum.




The same group also reported a polymerizable monomer formed by a TAC recognition moiety and a 4-amino-naphthalimide derived fluorophore containing a vinyl group. This monomer was able to penetrate human glioblastoma and oesophagus premalignant cell lines and be used to monitor  $K^+$  efflux. Moreover, thanks to the vinyl group the monomer was polymerised either with 2-hydroxyl methacrylate or acrylamide to give thin films useful for the measurement of extracellular  $K^+$  levels and thus measuring the efflux in *E. coli* and *B. subtilis* cells.<sup>40</sup> Later on, the same authors reported a clever modification of a TAC-bodipy sensor. Introducing the lipophilic triphenylphosphonium cation ( $TPP^+$ ) into the molecule, they were able to target mitochondria (because of the accumulation of  $TPP^+$  in these organelles) and, consequently, to monitor  $K^+$  ion dynamics in them. The sensor has a good response in con-

A few cryptands have been described for the recognition of cations other than alkali metals. Particularly, cages targeting transition metals have been described because of the relevance of these cations in environmental and biological processes. Kim and Bharadwaj described a cryptand-rhodamine conjugate sensor able to detect ppb of Hg(II) in aqueous media. Binding of Hg<sup>2+</sup> to the cryptand promotes the opening of the spiro-lactam ring, and the resulting change in fluorescence emission can be detected by naked eye. The sensor also gave a good result in cell samples allowing it to be used in live cells, in this case HEL 293 cell line.<sup>45</sup>

Carbohydrate recognition represents a formidable challenge for binding in aqueous media. Davis' group has long been working in a series of cages able to bind monosaccharides<sup>47,48</sup> and disaccharides<sup>49–51</sup> using polycyclic compounds. The general strategy that the group has developed consists in preparing hosts with the 'temple' design. This design incorporates a 'roof' and a 'floor' composed of aromatic moieties capable to establish hydrophobic contacts reinforced by CH- $\pi$  interactions (Fig. 6).<sup>52</sup> These building motifs are linked by pillars containing polar groups able to form hydrogen bonds with the substrate. Thus, carbohydrates with all-equatorial substitution, with axial C-H atoms are recognised preferentially.

Using this successful approach, they described a much easier to synthesize compound **9** (Fig. 7A) which can be prepared in just 5 steps compared to the long synthesis (up to 21



**Fig. 3** Time-dependent fluorescence of U87MG cells stimulated by nigericin and observed under confocal fluorescence microscope using TAC-TFC: (a)  $t = 0$  (before the addition of nigericin) (b–f)  $t = 1, 3, 5, 10$  and 15 min, respectively, after the addition of nigericin. Adapted with permission from ref. 39, copyright American Chemical Society.





**Fig. 4** (A) Cryptand-bodipy based sensor **7** for the detection of  $\text{Fe}^{3+}$ . (B) Naked-eye fluorescence responses of **7** ( $7 \mu\text{M}$ ) in  $\text{H}_2\text{O}-\text{CH}_3\text{CN}$  (9:1) upon addition of 100 equiv. of various metal cations ( $\lambda_{\text{ex}} = 365 \text{ nm}$ ). Reproduced with permission from ref. 29, copyright American Chemical Society.



**Fig. 5** Images of live HCT-116 cells (a–c) in the presence of  $30 \mu\text{M} \text{Fe}^{3+}$  solution in MEM medium and without  $\text{Fe}^{3+}$  (d–f) at (a and d) 0 and (b and e) 60 min after incubation with sensor **7** ( $20 \mu\text{M}$ ). c and f are the phase contrast images of b and e, respectively. Reproduced with permission from ref. 46, copyright American Chemical Society.

steps) of previously reported cages. Surprisingly, with a simpler design compound **9** binds glucose more strongly than other monosaccharides (for example 50:1 *versus* galactose). This selectivity allows **9** to act as a sensor for glucose in water in concentrations similar to those found in blood using fluorescence spectroscopy.<sup>53</sup> Arguably compound **9** is not a cage since although it is a macrocyclic compound composed by roof and floor and two pillars it lacks a third pillar required to



**Fig. 6** (A) Cartoon representation of the 'temple' design for recognition of all-equatorial carbohydrate recognition described by Davis.<sup>52</sup> (B) A tricyclic monosaccharide receptor prepared by Davis group for the recognition of  $\beta$ -glucose (ref. 47).

be considered a cage. Anyway, it follows the same principle for recognition and we found it relevant enough to be included in this section.

Going a step further in their research, the group optimised the recognition of glucose in complex media. Thus, they prepared the bicyclic cage **10** which provides a cavity that almost perfectly complements the all-equatorial  $\beta$ -pyranoside substrates.<sup>54</sup> The cage features several urea groups that provide an array of polar interactions and hydrogen bonding rationally designed by the authors. The receptor's affinity for glucose gives a  $K_a$  in the order of  $18\,000 \text{ M}^{-1}$ , which compares to the affinity of lectins and glucose transporters. In addition, the binding to other saccharides is about 100 times lower. These high levels of affinity and selectivity allowed the group to explore **10** as sensor in real-world samples. This was achieved through the reading of the CD spectra induced by the binding of glucose to the achiral host (Fig. 7B). As the affinity of the receptor for glucose was too high for practical determinations at usual concentrations in blood, a competition experiment was developed. The addition of L-glucose effectively doubles the signal range from the CD spectrometer and allows sensitivity to be adjusted over a wide concentration range. Thus, the system was able to analyse the presence of glucose in human serum (although filtrated to remove interfering proteins), cell culture and beer.<sup>55</sup>

More recently, the same group has described cage **11** (Fig. 8) able to bind polysaccharides such as cellulose through a threading geometry.<sup>56</sup> Polysaccharides, such as the mentioned cellulose, are not soluble in water, the formation of a threaded complex improved the water solubility of the bound polysaccharide. Solubilisation of these compounds could open the door to their use as feedstock chemicals through enzyme modification.





**Fig. 7** (A) Macrocyclic artificial lectins described by Davis for the sensing of glucose **9** (ref. 33) and **10** (ref. 34). (B) CD spectra from the titration of D-glucose into filtered, diluted, glucose-free human serum, in the presence of L-glucose (2 mM) and **10** (0.25 mM). Modified from ref. 55 with permission from the Royal Society of Chemistry.



**Fig. 8** Receptor able to recognise and solubilize polysaccharides prepared by Davis.<sup>56</sup>

## Ionophores

Molecular receptors that are able to cross membranes have a wide range of applications in biological research. Since they are able to transport chemical species in their interior, they

can be used to deliver drugs otherwise insoluble or less able to cross the cell membranes. These molecules can also disrupt cell homeostasis by allowing the flux of ions.<sup>57</sup> There are several natural cation selective ionophores which have indeed served as an inspiration for the design of artificial cationophores like for instance polycyclic ethers<sup>58</sup> or cyclodextrins used as antibiotics.<sup>59</sup> In the field of ionophores crown ethers also paved the way in the study of applications of supramolecular chemistry in biological environments.<sup>60</sup> Early on, Lehn's group already demonstrated the ability of the related cryptands to transport alkali metals through lipid membranes.<sup>61,62</sup> However, although a stronger binding may afford better selectivity this may be harmful in terms of transport when an ion flux is needed. Loiseau studied the effects of two cryptands: (221)C<sub>10</sub> cryptand, selective for Na<sup>+</sup>, and the (222)C<sub>10</sub> cryptand, selective for K<sup>+</sup> in opossum kidney cells (OK). It was found that the cryptands inhibited the pH-sensitive K<sup>+</sup> conductance of OK cells by inducing K<sup>+</sup>/H<sup>+</sup> exchange at the plasma membrane. Moreover, the Na/K ATP synthase was inhibited due to the entry of K<sup>+</sup> and Na<sup>+</sup> into the mitochondria.<sup>63</sup>


Anion transport across a membrane is more difficult because anions are usually more solvated than cations. In an early example Smith's group synthesised a ditopic macrobicycle able to bind NaCl or KCl and transport them through vesicle membranes.<sup>64</sup> The biological relevance of Cl<sup>−</sup> transport across cell membranes turned it into the perfect target for ion transporters as it is related to several diseases with especial relevance like cystic fibrosis<sup>65</sup> and cancer.<sup>66,67</sup> At the cellular level, electrolyte imbalance, particularly chloride imbalance between intra and extracellular environments, often leads to apoptosis.<sup>68</sup> A widely studied family of artificial compounds able to transport chloride are calix[4]pyrroles. These molecules offer many possibilities in the development of transmembrane ion transporters *via* different mechanisms. The pyrrolic N–H provides the capacity to bind anions and the aromatic cup to bind larger polarizable cations. Examples of calix[4]pyrroles with chloride binding properties are presented in Fig. 9. The first compound **12** acts as a CsCl co-transporter thus being electroneutral.<sup>69</sup> Fluorination of the pyrrole rings enhances the NH acidity, increasing the Cl<sup>−</sup> binding and the transmembrane transport capacity of the molecules. This is why the second compounds, **13**, is an efficient nitrate/chloride and bicarbonate/chloride exchanger thus capable of uniport across a membrane.<sup>70</sup> As we have explained before, these two compounds cannot be considered cages as they are simple macrocycles with a conic shape, however, their modification to add a staple that links two opposite sides of the molecule converts them in cage compounds.

Gale and co-workers studied strapped calix[4]pyrroles (**14** and **15** in Fig. 9) which showed Cl<sup>−</sup>/NO<sub>3</sub><sup>−</sup> exchange in the presence of Cs<sup>+</sup> but also as well as in the presence of Na<sup>+</sup>, K<sup>+</sup>, and Rb<sup>+</sup>.<sup>71,72</sup> This is a clear example of the effect of both co-transport and anion exchange to the global transport activity of this family of compounds.<sup>73</sup> Importantly, shorter strapped receptors showed certain selectivity for fluoride over chloride trans-









**Fig. 12** General chemical structure of chloride receptors **19** based on pseudopeptidic tripodal cages (top) and crystal structure of compound **19a** (R<sup>1</sup> = CH<sub>2</sub>Ph(4-OH), R<sup>2</sup> = Me, bottom).

played pH dependent activity, with a better transport occurring at acidic pH which implies a H<sup>+</sup>/Cl<sup>−</sup> symport mechanism. The NMR studies performed in DPC micelles demonstrated that the cages bind chloride within the lipid phase, and that both chloride binding and exchange are more efficient at acidic pH. We took advantage of the fact that the chloride affinity and transport improve at lower pH to selectively induce cell death of cancer cells because the cages display higher cytotoxicity in the presence of pH gradients resembling those characteristic of tumour microenvironments (Fig. 13).<sup>81</sup>

Transport of other anions different from chloride has been much less developed. Iodide selective transport using molecular containers (not cages) like β-cyclodextrins<sup>82</sup> and calix[4]arenes has been reported.<sup>83</sup> Using dynamic covalent chemistry Roh and Kim reported a porphyrin based organic cage able to transport iodide ions. The cage was constructed from square-



**Fig. 13** Schematic representation of the cytotoxicity experiments (A549 cell line) and plots of the percent cell viability versus the concentration of the cages **19b–d**, performed at different external pH<sub>e</sub> values.

shaped building units with different substitution (3- and 4-substituted) and had a large internal cavity, comparable to the thickness of bilayer membranes. The transport of a diverse range of anions was promoted by a dehydration-driven, channel mechanism in vesicles.<sup>84</sup> In addition, the organic cage seemed to be able to promote iodide transport across the membrane of a living cell. This was demonstrated using cells transfected with yellow fluorescent protein (YFP-H148Q/I152L) which is quenched by iodide, proving its potential usefulness in living systems.

Sulphate is one of the most common anions in plasma and it is involved in a number of biological processes.<sup>85</sup> However, sulphate transport through lipid bilayers is also very challenging because of its high hydrophilicity, being poorly investigated. Jolliffe, Gale and co-workers reported the transport abilities through membrane bilayers of a series of compounds based on tris-ureas and tris-thioureas. Among them the cyclopeptidic cage compounds **20** and **21** were studied (Fig. 14). Compounds **20** and **21** proved to be able to transport chloride in POPC vesicles using a selective electrode. Control experiments showed that these molecules act as molecular carriers rather than membrane spanning channels. Moreover, using <sup>33</sup>S NMR techniques they obtained evidence for transmembrane sulphate transport by receptor **21**, whereas receptor **20** was unable to transport this anion.

## Imaging

Positron emission tomography (PET) provides valuable information in the diagnosis and treatment of different diseases. PET relies on administering radioactive tracers containing positron-emitting isotopes and then detecting the radiation, <sup>18</sup>F being commonly used since the 60s. Metal-based PET radiotracers have become increasingly used in preclinical and clinical studies, particularly the emitting isotopes of copper, gallium and zirconium.<sup>87</sup> The development of these agents requires chelators with *in vivo* stability and able to be conjugated with targeting molecules (antibodies, peptides, etc.). A few examples of this application will be discussed in this section. The hexaazamacrobicyclic sarcophagine cage Sar **22** (Fig. 15) forms very stable complexes with Cu<sup>2+</sup> that do not dis-



**Fig. 14** Cryptand-like tris-thioureas described by Jolliffe and Gale for the transport of sulphate.<sup>86</sup>







**Fig. 15** (A) Sarcophagine cage and derivatives. (B) *In vivo* PET imaging of A431 xenograft tumor-bearing NSG mice following administration of [64Cu][CuMeCOSar]-cetuximab at different times. Reproduced from ref. 93 with permission from the Royal Society of Chemistry.

sociate under physiological conditions. Sargeson described the conjugation of a Sar derivative to monoclonal antibodies targeting disialogangliosides for tumour specific PET imaging agent using neuroblastoma and melanoma model systems.<sup>88</sup> The same chelator was used by Rogers to prepare a conjugate of bombesin, a peptide that binds to gastrin-releasing peptide receptor (GRPr) overexpressed in cancer cells. The conjugate was able thus to deliver Cu into PC-3 prostate cancer cells.<sup>89</sup> Similar cages with different handlers have been used to prepare conjugates with peptides for the PET imaging of tumours.<sup>90,91</sup> Other targets like active platelets have also been addressed with the appropriate antibody-chelator conjugates, thus delivery of <sup>64</sup>Cu allows the early diagnosis of thrombotic events.<sup>92</sup> More recently a Sar-conjugated antibody targeting epidermal growth factor receptor (EGFR) showed high-quality PET images in mouse models.<sup>93</sup> The sarcophagine chelator, MeCOSar (23, Fig. 15A), was attached to the antibody cetuximab by treating the antibody with the *N*-hydroxysuccinimide ester MeCOSar-NHS. On the other hand, the anti-EGFR murine monoclonal antibody, Fab528 was conjugated to the Sar derivative 24 by means of an enzymatic coupling. In Fig. 15B the *in vivo* PET images following administration of [64Cu][CuMeCOSar]-cetuximab in A431 xenograft tumour-bearing NSG mice at different times are shown. Tumour uptake was evident at 1 h post-administration but the best tumour: background ratio is obtained after 24 h.

Sarcophagine functionalised ligands with tumour targeting peptides have also been used with Ga<sup>3+</sup> allowing to use commercially available <sup>68</sup>Ge/<sup>68</sup>Ga generators for their use as radio-

pharmaceutical agents.<sup>94</sup> Finally, other kinds of chelators have been described for other metals. Wadas group prepared di-macrocytic terephthalamide ligands for the stabilization of <sup>89</sup>Zr which are highly stable *in vivo* allowing the use of this element in PET imaging.<sup>95</sup>

## Complexation of biomolecules

Complexation of bio-molecules has many potential biomedical applications: for instance, sensing, enzyme monitoring or drug design. Besides, it can provide insights into significant biomolecular interaction mechanisms. The complexation of natural products using artificial receptors is a major topic in supramolecular chemistry. Non covalent interactions with bio-molecules (amino acids, biopolymers, enzymes or nucleic acids), which are the fundamentals of many pharmacological activities, can be studied using supramolecular model complexes. Also in the biochemistry field, supramolecular complexes are being used for the protection, transport or targeting of bioactive molecules. Encapsulation in a suitable host can provide selectivity in the protection of natural drugs from metabolism or even the manipulation of the guest redox properties.

The disruption of metalloenzymes by metal coordination or sequestration offers a way to deactivate their activity. Following this idea, the group of García-España reported a series of pyrazole-containing macrocyclic and cryptand-like compounds active against the iron superoxide dismutase (Fe-SOD), an essential enzyme for the survival of the protozoan *Trypanosoma cruzi*, responsible of the Chagas disease. Among them, compound 25 (Fig. 16) showed great *in vitro* and *in vivo* activity against intra and extracellular forms of the parasite being the most effective compound, especially against the chronic phase. Importantly, the compound was highly selective for Fe-SOD when compared to human CuZn-SOD enzyme.<sup>96</sup> The close structural similarity of the Fe-SOD enzymes in *T. cruzi* and *Leishmania* spp. induced the group to study the antiproliferative activity of some compounds, including 25, against *Leishmania infantum* and *Leishmania braziliensis* species. The results showed again that compound 25 was highly leishmanicidal and induced extensive parasite cell damage.<sup>97</sup>

Pseudo-peptidic cages offer a suitable platform to work in biological media. On the one hand, a variety of amino acids is available which allow a broad structural and functional diversity. On the other hand, the non-peptidic part can be modulated to implement the desired functions. In the last years, we have been working on the design and applications of pseudo-peptidic cages. In 2012 we described the efficient synthesis of macrobicyclic pseudo-peptidic compounds with two strategies, using organic anion templates and driving the formation of the product by forcing the configuration of the building blocks.<sup>98</sup> We described the synthesis of a family of pseudo-peptidic macrocycles with relatively high yields using a 3 + 2 imine condensation to form the most thermodynamically stable product followed by the *in situ* reduction of the imine group





**Fig. 16** Macrobicyclic compound **25** described by García-España with anti-*Trypanosoma cruzi* and anti-leishmaniosis activity.

(Scheme 1). It is worth mentioning that this one-pot reaction involves the formation of 6 covalent bonds in two steps which means a total of 12 reactions. This reaction can be carried out using a template (5TBA) but the final products are also obtained without template if the mixture is allowed to reach the equilibrium for enough time (24 h).

The recognition properties of different cages prepared following the synthesis described in Scheme 1. Those compounds were investigated for the recognition N-protected dipeptides.<sup>99</sup> The interactions were measured through different techniques (NMR, MS and in some cases fluorescence emission spectroscopy) which allowed the determination of the association constant of each cage-receptor pair.



**Scheme 1** Synthesis of pseudopeptidic cages described by Alfonso.<sup>98</sup> (a) Equilibrium at RT either in the absence or the presence of 5TBA. (b)  $\text{BH}_3 \cdot \text{py}$  (c)  $\text{HCl}$ .


Very interesting conclusions were extracted from this work: cyclohexane gives a rigidity to the moiety that is beneficial for the recognition of most dipeptides. Moreover, the side chains (R substituents) of the cage also play an important role in the recognition. For instance, Ser and Val enhance the interaction whereas Phe residues reduce it. The highest binding constants were found to be with the dipeptides containing aromatic rings (Phe and Tyr). With all the data collected from the different interaction studies (NMR, MS and molecular modelling analysis) and considering the stereoselection observed with the Ac-EY-OH dipeptide, an interaction model was proposed, which implies the inclusion of the aromatic Tyr residue within the cage cavity, along with a number of ionic and H-bonding host-guest interactions (Fig. 17).<sup>100</sup>

Given the promising results, new cage-like compounds were synthesized with the (*R,R*)-cyclohexane-1,2-diamine and Lys and Orn residues (CyLys and CyOrn, respectively). Those showed better affinity for Tyr residues in short peptides and also in a polypeptide (PolyE4Y). This was relevant because EYE epitope is a known substrate for protein tyrosine kinases (PTKs). The affinity of both CyOrn and CyLys for this sequence was better than those of the other cages (CySer, CyThr) due to the positive charge of the Lys and Orn residues that allows secondary electrostatic interactions between glutamic residues of the polypeptide and the side chains of the cage. CyLys and CyOrn were proven to inhibit PTK activity by protecting the Tyr from phosphorylation using a commercial assay kit. The kinase inhibition was in agreement with the  $K_d$  found for CyLys and CyOrn and showed that CyLys is the most efficient cage for kinase inhibition.<sup>100,101</sup> Moreover, the selective encapsulation of Tyr was also demonstrated in polypeptides containing several tyrosines and in short peptide with different sequences. More recently, we have extended the research to other cages (CyHis, CyAsp and CyGlu, based on His, Asp or Glu residues, respectively) and examined their protective action in the Src mediated phosphorylation of a series of Tyr containing peptides (Fig. 18). Remarkably, the inhibition of



**Fig. 17** Two views of the proposed binding model for the recognition of the dipeptide Ac-L-Glu-L-Tyr-OH (C-atoms in green) by CySer receptor. Non-polar hydrogens have been omitted for clarity and possible H-bonds are shown in dashed lines.





Importantly, the binding affinities of the cages strongly correlate with the inhibitory efficiency of the cages. As general trend the Asp cage was the most efficient inhibitor with Lys, Orn, and His showing interesting activities that deeply depended on the nature of the substrate. Overall, it could be concluded that the inhibitory effect of the cages was due to the protective action resulting from Tyr encapsulation within the cage while the binding strongly depends on the additional interactions between the side chains of the pseudopeptidic cages and the amino acid sequences surrounding the Tyr residue in the peptides. Interestingly, we also demonstrated that the cage protects peptides from phosphorylation in a completely different tyrosine kinase system, highlighting the general application of the concept.

The preparation of cage-like compounds with a well-defined inner cavity is synthetically challenging. However, these com-

Financial support from the Spanish Ministry of Science and Innovation (RTI2018-096182-B-I00, RED2018-102331-T, MCI/AEI/FEDER, EU), AGAUR (2017 SGR 208) and personal support for L. Tapia (BES-2016-076863) are gratefully acknowledged. We acknowledge support of the publication fee by the CSIC



- 1 T. W. Traut, *Allosteric Regulatory Enzymes*, Springer US., New York, 2008.
- 2 T. D. Pollard, W. Earnshaw, J. Lippincott-Schwartz and G. Johnson., *Cell biology*, Elsevier, Philadelphia, 2017.
- 3 G. Crini, *Chem. Rev.*, 2014, **114**, 10940–10975.
- 4 W. Sliwa and T. Girek, *Cyclodextrins: Properties and Applications*, Wiley-VCH Verlag GmbH & Co. KGaA, 2017.
- 5 K. Kim, N. Selvapalam, Y. H. Ko, K. M. Park, D. Kim and J. Kim, *Chem. Soc. Rev.*, 2007, **36**, 267–279.
- 6 S. J. Barrow, S. Kasera, M. J. Rowland, J. del Barrio and O. A. Scherman, *Chem. Rev.*, 2015, **115**, 12320–12406.
- 7 D. Das, K. I. Assaf and W. M. Nau, *Front. Chem.*, 2019, **7**, 619.
- 8 F. N. Pur, *Mol. Divers.*, 2016, **20**, 781–787.
- 9 M. M. Naseer, M. Ahmed and S. Hameed, *Chem. Biol. Drug Des.*, 2017, **89**, 243–256.
- 10 R. Kumar, A. Sharma, H. Singh, P. Suating, H. S. Kim, K. Sunwoo, I. Shim, B. C. Gibb and J. S. Kim, *Chem. Rev.*, 2019, **119**, 9657–9721.
- 11 S. Kubik, in *Chemistry of Nanocontainers*, ed. M. Albrecht and E. Hahn, Springer Berlin Heidelberg, Berlin, Heidelberg, 2012, pp. 1–34.
- 12 B. Dietrich, in *Comprehensive Supramolecular Chemistry*, ed. G. W. Gokel, Elsevier, Oxford, 1996, pp. 153–211.
- 13 J. M. Lehn, *Pure Appl. Chem.*, 1977, **49**, 13.
- 14 B. Dietrich, J. M. Lehn and J. P. Sauvage, *Tetrahedron Lett.*, 1969, **10**, 2885–2888.
- 15 B. Dietrich, J. M. Lehn and J. P. Sauvage, *Tetrahedron Lett.*, 1969, **10**, 2889–2892.
- 16 C. H. Park and H. E. Simmons, *J. Am. Chem. Soc.*, 1968, **90**, 2431–2432.
- 17 R. Chakrabarty, P. S. Mukherjee and P. J. Stang, *Chem. Rev.*, 2011, **111**, 6810–6918.
- 18 T. R. Cook and P. J. Stang, *Chem. Rev.*, 2015, **115**, 7001–7045.
- 19 N. J. Young and B. P. Hay, *Chem. Commun.*, 2013, **49**, 1354–1379.
- 20 N. Ahmad, H. A. Younus, A. H. Chughtai and F. Verpoort, *Chem. Soc. Rev.*, 2015, **44**, 9–25.
- 21 H. Sepehrpour, W. Fu, Y. Sun and P. J. Stang, *J. Am. Chem. Soc.*, 2019, **141**, 14005–14020.
- 22 J. W. Steed, *Chem. Soc. Rev.*, 2009, **38**, 506–519.
- 23 T. R. Cook, V. Vajpayee, M. H. Lee, P. J. Stang and K.-W. Chi, *Acc. Chem. Res.*, 2013, **46**, 2464–2474.
- 24 E. G. Percástegui, T. K. Ronson and J. R. Nitschke, *Chem. Rev.*, 2020, **120**, 13480–13544.
- 25 G. Yu, M. Jiang, F. Huang and X. Chen, *Curr. Opin. Chem. Biol.*, 2021, **61**, 19–31.
- 26 M. Mastalerz, *Angew. Chem., Int. Ed.*, 2010, **49**, 5042–5053.

- 51 J. D. Howgego, C. P. Butts, M. P. Crump and A. P. Davis, *Chem. Commun.*, 2013, **49**, 3110–3112.
- 52 A. P. Davis, *Org. Biomol. Chem.*, 2009, **7**, 3629–3638.
- 53 C. Ke, H. Destecroix, M. P. Crump and A. P. Davis, *Nat. Chem.*, 2012, **4**, 718–723.
- 54 R. A. Tromans, T. S. Carter, L. Chabanne, M. P. Crump, H. Li, J. V. Matlock, M. G. Orchard and A. P. Davis, *Nat. Chem.*, 2019, **11**, 52–56.
- 55 R. A. Tromans, S. K. Samanta, A. M. Chapman and A. P. Davis, *Chem. Sci.*, 2020, **11**, 3223–3227.
- 56 T. J. Mooibroek, J. M. Casas-Solvas, R. L. Harniman, C. M. Renney, T. S. Carter, M. P. Crump and A. P. Davis, *Nat. Chem.*, 2016, **8**, 69–74.
- 57 I. Alfonso and R. Quesada, *Chem. Sci.*, 2013, **4**, 3009–3019.
- 58 K. C. Nicolaou, M. O. Frederick and R. J. Aversa, *Angew. Chem., Int. Ed.*, 2008, **47**, 7182–7225.
- 59 H. Yamamura, K. Suzuki, K. Uchibori, A. Miyagawa, M. Kawai, C. Ohmizo and T. Katsu, *Chem. Commun.*, 2012, **48**, 892–894.
- 60 M. Kralj, L. Tušek-Božić and L. Frkanec, *ChemMedChem*, 2008, **3**, 1478–1492.
- 61 M. Castaing and J. M. Lehn, *J. Membr. Biol.*, 1987, **97**, 79–95.
- 62 M. Castaing, F. Morel and J. M. Lehn, *J. Membr. Biol.*, 1986, **89**, 251–267.
- 63 A. Loiseau, C. Leroy and M. Castaing, *Biochim. Biophys. Acta, Biomembr.*, 1997, **1330**, 39–49.
- 64 A. V. Koulov, J. M. Mahoney and B. D. Smith, *Org. Biomol. Chem.*, 2003, **1**, 27–29.
- 65 R. Planells-Cases and T. J. Jentsch, *Biochim. Biophys. Acta, Mol. Basis Dis.*, 2009, **1792**, 173–189.
- 66 S. Hong, M. Bi, L. Wang, Z. Kang, L. Ling and C. Zhao, *Oncol. Rep.*, 2015, **33**, 507–514.
- 67 K. O. Alfarouk, S. B. M. Ahmed, A. Ahmed, R. L. Elliott, M. E. Ibrahim, H. S. Ali, C. C. Wales, I. Nourwali, A. N. Aljarbou, A. H. H. Bashir, S. T. S. Alhoufie, S. S. Alqahtani, R. A. Cardone, S. Fais, S. Harguindey and S. J. Reshkin, *Cancers*, 2020, **12**, 898.
- 68 S. P. Yu, L. M. T. Canzoniero and D. W. Choi, *Curr. Opin. Cell Biol.*, 2001, **13**, 405–411.
- 69 C. C. Tong, R. Quesada, J. L. Sessler and P. A. Gale, *Chem. Commun.*, 2008, 6321–6323, DOI: 10.1039/B814988G.
- 70 P. A. Gale, C. C. Tong, C. J. E. Haynes, O. Adeosun, D. E. Gross, E. Karnas, E. M. Sedenberg, R. Quesada and J. L. Sessler, *J. Am. Chem. Soc.*, 2010, **132**, 3240–3241.
- 71 M. Yano, C. C. Tong, M. E. Light, F. P. Schmidtchen and P. A. Gale, *Org. Biomol. Chem.*, 2010, **8**, 4356–4363.
- 72 M. G. Fisher, P. A. Gale, J. R. Hiscock, M. B. Hursthouse, M. E. Light, F. P. Schmidtchen and C. C. Tong, *Chem. Commun.*, 2009, 3017–3019.
- 73 S. Peng, Q. He, G. I. Vargas-Zúñiga, L. Qin, I. Hwang, S. K. Kim, N. J. Heo, C.-H. Lee, R. Dutta and J. L. Sessler, *Chem. Soc. Rev.*, 2020, **49**, 865–907.
- 74 H. J. Clarke, E. N. W. Howe, X. Wu, F. Sommer, M. Yano, M. E. Light, S. Kubik and P. A. Gale, *J. Am. Chem. Soc.*, 2016, **138**, 16515–16522.
- 75 S.-K. Ko, S. K. Kim, A. Share, V. M. Lynch, J. Park, W. Namkung, W. Van Rossom, N. Busschaert, P. A. Gale, J. L. Sessler and I. Shin, *Nat. Chem.*, 2014, **6**, 885–892.
- 76 I.-W. Park, J. Yoo, B. Kim, S. Adhikari, S. K. Kim, Y. Yeon, C. J. E. Haynes, J. L. Sutton, C. C. Tong, V. M. Lynch, J. L. Sessler, P. A. Gale and C.-H. Lee, *Chem. – Eur. J.*, 2012, **18**, 2514–2523.
- 77 S.-H. Park, S.-H. Park, E. N. W. Howe, J. Y. Hyun, L.-J. Chen, I. Hwang, G. Vargas-Zúñiga, N. Busschaert, P. A. Gale, J. L. Sessler and I. Shin, *Chem*, 2019, **5**, 2079–2098.
- 78 R. Quesada, *Chem*, 2019, **5**, 1924–1926.
- 79 I. Martí, M. Bolte, M. I. Burguete, C. Vicent, I. Alfonso and S. V. Luis, *Chem. – Eur. J.*, 2014, **20**, 7458–7464.
- 80 I. Martí, J. Rubio, M. Bolte, M. I. Burguete, C. Vicent, R. Quesada, I. Alfonso and S. V. Luis, *Chem. – Eur. J.*, 2012, **18**, 16728–16741.
- 81 L. Tapia, Y. Pérez, M. Bolte, J. Casas, J. Solà, R. Quesada and I. Alfonso, *Angew. Chem., Int. Ed.*, 2019, **58**, 12465–12468.
- 82 N. Madhavan, E. C. Robert and M. S. Gin, *Angew. Chem., Int. Ed.*, 2005, **44**, 7584–7587.
- 83 I. Izzo, S. Licen, N. Maulucci, G. Autore, S. Marzocco, P. Tecilla and F. De Riccardis, *Chem. Commun.*, 2008, 2986–2988.
- 84 B. P. Benke, P. Aich, Y. Kim, K. L. Kim, M. R. Rohman, S. Hong, I.-C. Hwang, E. H. Lee, J. H. Roh and K. Kim, *J. Am. Chem. Soc.*, 2017, **139**, 7432–7435.
- 85 D. Markovich, *Physiol. Rev.*, 2001, **81**, 1499.
- 86 N. Busschaert, L. E. Karagiannidis, M. Wenzel, C. J. E. Haynes, N. J. Wells, P. G. Young, D. Makuc, J. Plavec, K. A. Jolliffe and P. A. Gale, *Chem. Sci.*, 2014, **5**, 1118–1127.
- 87 L. E. McInnes, S. E. Rudd and P. S. Donnelly, *Coord. Chem. Rev.*, 2017, **352**, 499–516.
- 88 S. D. Voss, S. V. Smith, N. DiBartolo, L. J. McIntosh, E. M. Cyr, A. A. Bonab, J. L. J. Dearling, E. A. Carter, A. J. Fischman, S. T. Treves, S. D. Gillies, A. M. Sargeson, J. S. Huston and A. B. Packard, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 17489.
- 89 K. A. Lears, R. Ferdani, K. Liang, A. Zheleznyak, R. Andrews, C. D. Sherman, S. Achilefu, C. J. Anderson and B. E. Rogers, *J. Nucl. Med.*, 2011, **52**, 470.
- 90 E. Gourni, L. Del Pozzo, E. Kheirallah, C. Smerling, B. Waser, J.-C. Reubi, B. M. Paterson, P. S. Donnelly, P. T. Meyer and H. R. Maecke, *Mol. Pharm.*, 2015, **12**, 2781–2790.
- 91 B. M. Paterson, P. Roselt, D. Denoyer, C. Cullinane, D. Binns, W. Noonan, C. M. Jeffery, R. I. Price, J. M. White, R. J. Hicks and P. S. Donnelly, *Dalton Trans.*, 2014, **43**, 1386–1396.
- 92 B. M. Paterson, K. Alt, C. M. Jeffery, R. I. Price, S. Jagdale, S. Rigby, C. C. Williams, K. Peter, C. E. Hagemeyer and P. S. Donnelly, *Angew. Chem., Int. Ed.*, 2014, **53**, 6115–6119.



- 93 S. E. Rudd, J. K. Van Zuylekom, A. Raicevic, L. A. Pearce, C. Cullinane, C. C. Williams, T. E. Adams, R. J. Hicks and P. S. Donnelly, *Chem. Sci.*, 2021, **12**, 9004–9016.
- 94 M. T. Ma, O. C. Neels, D. Denoyer, P. Roselt, J. A. Karas, D. B. Scanlon, J. M. White, R. J. Hicks and P. S. Donnelly, *Bioconjugate Chem.*, 2011, **22**, 2093–2103.
- 95 D. N. Pandya, S. Pailloux, D. Tatum, D. Magda and T. J. Wadas, *Chem. Commun.*, 2015, **51**, 2301–2303.
- 96 M. Sánchez-Moreno, C. Marín, P. Navarro, L. Lamarque, E. García-España, C. Miranda, O. Huertas, F. Olmo, F. Gómez-Contreras, J. Pitarch and F. Arrebola, *J. Med. Chem.*, 2012, **55**, 4231–4243.
- 97 P. Navarro, M. SÁNchez-Moreno, C. MarÍN, E. García-España, I. Ramírez-Macías, F. Olmo, M. J. Rosales, F. Gómez-Contreras, M. J. R. Yunta and R. Gutierrez-Sánchez, *Parasitology*, 2014, **141**, 1031–1043.
- 98 A. Moure, S. V. Luis and I. Alfonso, *Chem. – Eur. J.*, 2012, **18**, 5496–5500.
- 99 E. Faggi, A. Moure, M. Bolte, C. Vicent, S. V. Luis and I. Alfonso, *J. Org. Chem.*, 2014, **79**, 4590–4601.
- 100 E. Faggi, C. Vicent, S. V. Luis and I. Alfonso, *Org. Biomol. Chem.*, 2015, **13**, 11721–11731.
- 101 E. Faggi, Y. Perez, S. V. Luis and I. Alfonso, *Chem. Commun.*, 2016, **52**, 8142–8145.

