

ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: X. Qi, Y. Wang and X. Wu, *Chem. Commun.*, 2024, DOI: 10.1039/D4CC05361C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

COMMUNICATION

CO Insertion Enabled γ -C(sp³)-H Heteroarylyative Carbonylation of Tertiary Alcohols via Heteroaryl MigrationXin Qi,^{a,b} Yuanrui Wang,^{a,b} Xiao-Feng Wu,^{a,b,c*}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


The direct functionalization of remote C(sp³)-H is valuable but challenging, which is even more difficult to achieve the γ -C(sp³)-H functionalization of alcohols. Among the strategies, hydrogen atom transfer (HAT) is one of the solutions for such transformations. Herein, we designed a migration reaction involving carbon monoxide, forming alkoxy radical by photocatalysis, and used carbon monoxide to extend the carbon chain to provide a site for the migration of heteroaryl groups, which makes 1,4-HAT more advantageous, and relied on this strategy to successfully achieve the synthesis of 1,4-dicarbonyl compounds by γ -C(sp³)-H functionalization of alcohols.

Direct functionalization of C(sp³)-H bonds is an ideal synthetic method because of its straightforward and atomically efficient nature in the construction of carbon-carbon or carbon-heteroatomic bonds, which has led to its use in modern chemistry for the synthesis of a wide variety of molecules in biologically active natural products, pharmaceuticals, and functional materials using inexpensive and accessible stocks.^[1-5] The α - and β -C(sp³)-H-functionalization can be readily accomplished through transition metal catalysis, which have also been extensively investigated for a range of aliphatic substrates.^[6-8] Nevertheless, functionalization reactions targeting more distant positions, such as the γ - and δ -C, present a greater challenge. Such functionalization reactions are more difficult to achieve due to the difficulty of forming kinetically unstable intermediates.

In recent years, as chemists have gained a deeper understanding of free radical chemistry, the use of free radicals in organic synthesis has become more prevalent. Attempts have also been made to utilize the free radical pathway to achieve a number of transformations that were previously challenging to attain. Among the radicals, it is well known that oxygen and nitrogen radical can be used for various synthetic transformations.^[9-16] The use of oxygen or nitrogen radicals for

1,5-hydrogen atom transfer (1,5-HAT) can figure out the direct functionalization of the remote C(sp³)-H bond (Figure 1, eq. a). The oxygen or nitrogen radical can initially be generated through photocatalysis, which initiates a 1,5-hydrogen atom transfer. This process involves the transfer of a hydrogen atom from the carbon to an oxygen or nitrogen radical, leading to the formation of a highly reactive carbon radical. Subsequently, the radical is captured by another radical partner, thereby installing a functional group at the δ -C. The implementation of this strategy results in the functionalization of the inactivated C(sp³)-H bond. The transition state formed by the transfer of the hydrogen atom of γ -C is less stable than the six-membered ring transition state formed by the H atom transfer of the δ -C, hence the 1,4-HAT is much less studied than 1,5-HAT.

Figure 1. Direct functionalization of remote C(sp³)-H and experimental design.



Since the experimental preparation of carbon monoxide in the 18th century, this gas has been used as a cheap and readily available source of C1 for a variety of carbonylation reactions. The coupling reactions using CO as a carbonyl source are one of the most efficient and direct methods for the synthesis of aldehydes, ketones, acids, and many other derivatives of carboxylic acids.^[17-20] Among the numerous industrial scale

^a Dalian National Laboratory for Clean Energy, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China, e-mail: xwu2020@dicp.ac.cn

^b University of Chinese Academy of Sciences, Beijing, 101408, China,

^c Leibniz-Institut für Katalyse e.V., Albert-Einstein-Straße 29a, 18059 Rostock, Germany

* Supplementary Information available: See DOI: 10.1039/x0xx00000x



COMMUNICATION

ChemComm

applied carbonylation processes, the carbonylation of methanol with CO is currently the main production method worldwide for acetic acid producing.^[21–23]

Previously we have developed radical migration reactions induced by visible light with carbon monoxide insertion (Figure 1, eq. b).^[24] We propose that the insertion of carbon monoxide serves as a hinge for the migration of the heteroaryl group and that migration promotes more efficient capture of CO, which is a key factor in the carbonylation reaction. On this basis, we envisage the use of CO to extend the carbon chain to provide a site for heteroaryl 1,4-migration, which promotes CO capture, leading to the synthesis of 1,4-dicarbonyl compounds by γ -C(sp³)-H functionalization of alcohols (Figure 1, eq. c).

We synthesized a tertiary alcohol containing butyl and benzothiazole groups (**1f**) as the model substrate for optimizing this migration reaction. Blue LEDs was chosen as the light source and high oxidation potential [Ir(dF(CF₃)ppy)₂(dtbbpy)]PF₆ as the photocatalyst. A persulfate-type oxidant is essential and cannot be substituted by using PIDA, and we finally chose K₂S₂O₈ as the oxidant (Table 1, entries 4, 5). Here, the compatibility between the photocatalyst and oxidant could be an explanation for this result. The introduction of alternative phase transfer catalysts did not supersede the efficacy of *n*-Bu₄NHSO₄ (Table 1, entries 6, 7), which is postulated to be the optimal phase transfer catalyst for this reaction. After comparing several solvents, PhCF₃ was identified as the most suitable option here (Table 1, entry 8). This might benefit from the weak interaction between fluorine atom and proton of OH group in the substrate. The addition of a small quantity of water is essential for the dissolution of the persulfate (Table 1, entry 9). Furthermore, no product was formed in the absence of light (Table 1, entry 10).

Table 1. Optimization of reaction conditions.^a


Entry	Variation of reaction conditions	Yield (%) ^c
1	none	50(48)
2	without photocatalyst	0
3	without CO	0
4	Na ₂ S ₂ O ₈ instead of K ₂ S ₂ O ₈	46
5 ^b	PIDA, PIFA instead of K ₂ S ₂ O ₈	0
6	18-Crown-6-ether instead of <i>n</i> -Bu ₄ NHSO ₄	10
7	<i>n</i> -Bu ₄ NCl instead of <i>n</i> -Bu ₄ NHSO ₄	45
8	acetone, toluene, CH ₃ CN instead of PhCF ₃	0
9	without H ₂ O	0
10	without light	0

[a] Reaction conditions: **1** (0.1 mmol), photocatalyst (3 mol%), K₂S₂O₈ (0.25 mmol), Bu₄NHSO₄ (0.05 mmol), PhCF₃ (1.5 mL), H₂O (150 μ l) at rt for 36 h under CO (50 bar). [b] PhCF₃ (2.0 mL), H₂O (200 μ l). [c] Yield was determined by GC the isolated yield is given in parentheses.

After determined the optimized reaction conditions, we subsequently studied the substrates scope of this

transformation (Scheme 1). To reduce the amount of the non-carbonylic migration products and increase the yield of the target product, we replaced the butyl group (**1f**) with a propyl group (**1a**), as this avoids 1,5-HAT. Changing the substituent on the aryl group, it can be seen that the reaction can still take place. The introduction of a strong electron-withdrawing group like trifluoromethyl did not affect the reaction (**2b**), and substrates containing methyl or phenyl groups on the benzene ring also underwent a smooth carbonyl migration reaction to produce the corresponding 1,4-dicarbonyl compounds (**2c–2e**) in moderate to good yields. However, no desired reaction occurred when 1,1-diphenylbutan-1-ol was tested as the substrate under our standard conditions.

Scheme 1. Scope of substrates.^a



[a] Reaction conditions: **1** (0.1 mmol), photocatalyst (3 mol%), K₂S₂O₈ (0.25 mmol), Bu₄NHSO₄ (0.05 mmol), PhCF₃ (1.5 mL), H₂O (150 μ l) at rt for 36 h under CO (50 bar). [b] The yield is calculated based on the isolated alcohol.




View Article Online
DOI: 10.1039/D4CC05361G

Combined with the understanding based on previous work,^[24] we proposed a possible mechanism (Scheme 2). We chose tertiary alcohol containing butyl and benzothiazole to illustrate the reaction pathways and explain the selectivity. Photocatalysis of tertiary alcohol generates alkoxy radical, which initiate hydrogen atom transfer. Generally, 1,5-HAT accounts for the majority of the product, but in the presence of carbon monoxide, 1,4-HAT accounts for the majority of the product as CO provides a carbon atom at the γ -C allowing benzothiazole to 1,4-migrate to form the more dominant

product. In more detail, acyl radical intermediate **II** will be produced after capture one molecule of CO once the radical intermediate **I** been generated. Then, intermediate **III** will be formed after the 1,4-migration of benzothiazole which then be oxidized and gave cation intermediate **IV**. Finally, the targeted product can be produced after deprotonation. Benzothiazoles migrate to form the five-membered transition state, so by-product (**2f'**) cannot be avoided, but the use of tertiary alcohols containing a propyl group instead of an *n*-butyl group can avoid the formation of the by-products.

Scheme 2. Proposed mechanism.



In summary, we used tertiary alcohols containing benzothiazole moieties as the substrates studied γ -C(sp³)-H heteroarylation carbonylation through heteroaryl migration. The reaction proceeds via the formation of alkoxy radical via photocatalysis, extended the carbon chain using carbon monoxide to provide carbon sites for the migration of benzothiazoles, which made 1,4-HAT more advantageous, and relied on this strategy to successfully achieve the synthesis of distal γ -functionalized 1,4-dicarbonyl compounds.

Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the Supplementary Information.

Acknowledgements

We thank the financial supports from National Key R&D Program of China (2023YFA1507500) and DICP.

Notes and references

- X. Chen, K.M. Engle, D.H. Wang, J.Q. Yu, *Angew. Chem. Int. Ed.* 2009, **48**, 5094-115.
- V. Ritleng, C. Sirlin, M. Pfeffer, *Chem. Rev.* 2002, **102**, 1731-1770.
- C.-L. Sun, B.-J. Li, Z.-J. Shi, *Chem. Rev.* 2011, **111**, 1293-1314.
- A.M. Messinis, J.C.A. Oliveira, A.C. Stückl, L. Ackermann, *ACS Catal.* 2022, **12**, 4947-4960.
- M. Albicker, N. Cramer, *Angew. Chem. Int. Ed.* 2009, **48**, 9139-9142.
- Z. Wu, C. Pi, X. Cui, J. Bai, Y. Wu, *Adv. Synth. Catal.*, 2013, **355**, 1971-1976.
- Z. Li, R. Yu, H. Li, *Angew. Chem. Int. Ed.*, 2008, **47**, 7497-7500.
- L. Yang, H. Xie, G. An, G. Li, *J. Org. Chem.* 2021, **86**, 7872-7880.
- M. Kapoor, A. Singh, K. Sharma, M.H. Hsu, *Adv. Synth. Catal.* 2020, **362**, 4513.
- Q. Qin, S. Yu, *Org. Lett.* 2015, **17**, 1894-1897
- L. Wang, Y. Xia, K. Bergander, A. Studer, *Org. Lett.* 2018, **20**, 5817-5820.
- B. Xu, U.K. Tambar, *ACS Catal.* 2019, **9**, 4627-4631.
- X. Wu, H. Zhang, N. Tang, Z. Wu, D. Wang, M. Ji, Y. Xu, M. Wang, C. Zhu, *Nat. Commun.* 2018, **9**, 3343.
- M. Wang, L. Huan, C. Zhu, *Org. Lett.* 2019, **21**, 821-825.
- Y. Zhu, K. Huang, J. Pan, X. Luo, Q. Qin, J. Wei, X. Wen, L. Zhang, N. Jiao, *Nat. Commun.* 2018, **9**, 2625.
- J.-J. Guo, A. Hu, Y. Chen, J. Sun, H. Tang, Z. Zuo, *Angew. Chem. Int. Ed.* 2016, **55**, 15319-15322.
- A. Schoenberg, I. Bartoletti, R.F. Heck, *J. Org. Chem.* 1974, **39**, 3318-3326.
- L.-C. Wang, Y. Yuan, Y. Zhang, X.-F. Wu, *Nat Commun.* 2023, **14**, 7439.
- X.-F. Wu, X. Fang, L. Wu, R. Jackstell, H. Neumann, M. Beller, *Acc. Chem. Res.* 2014, **47**, 1041-1053.



COMMUNICATION

ChemComm

- 20 Z.-P. Bao, N.-X. Sun, X.-F. Wu, *Chin. J. Catal.*, 2024, **60**, 171-177.
- 21 A. Seayad, M. Ahmed, H. Klein, R. Jackstell, T. Gross, M. Beller, *Science* 2002, **297**, 1676-1678.
- 22 R. Franke, D. Selent, A. Börner, *Chem Rev.* 2012, **112**, 5675-5732.
- 23 M.J. Howard, M.D. Jones, M.S. Roberts, S.A. Taylor, *Catal. Today* 1993, **18**, 325-354.
- 24 Y. Wang, H. Yang, Y. Zheng, M. Hu, J. Zhu, Z.-P. Bao, Y. Zhao, X.-F. Wu, *Nat. Catal.* 2024, **7**, 1065-1075.

View Article Online
DOI: 10.1039/D4CC05361C



The data supporting this article have been included as part of the Supplementary Information.

View Article Online
DOI: 10.1039/D4CC05361C

Open Access Article. Published on 04 November 2024. Downloaded on 05/11/2024 19:40:06.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

