



NPR

Traversing the fungal terpenome: biosynthetic enzymes and pathways.

| | |
|-------------------------------|--|
| Journal: | <i>Natural Product Reports</i> |
| Manuscript ID: | NP-HIG-06-2014-000075.R1 |
| Article Type: | Highlight |
| Date Submitted by the Author: | 15-Aug-2014 |
| Complete List of Authors: | Schmidt-Dannert, Claudia; University of Minnesota, Quinn, Maureen; University of Minnesota, Flynn, Christopher; University of Minnesota, |
| | |

SCHOLARONE™
Manuscripts

Traversing the fungal terpenome

Cite this: DOI: 10.1039/x0xx00000x

Maureen B. Quin*, Christopher M. Flynn* and Claudia Schmidt-Dannert†

University of Minnesota, Dept. of Biochemistry, Molecular Biology and Biophysics, 1479 Gortner Avenue, St. Paul, MN 55108.

Received 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

†To whom correspondence should be addressed

*These authors contributed equally to this work.

www.rsc.org/

Covering: 1965-2014

Fungi (Ascomycota and Basidiomycota) are prolific producers of structurally diverse terpenoid compounds. Classes of terpenoids identified in fungi include the sesqui-, di- and triterpenoids. Biosynthetic pathways and enzymes to terpenoids from each of these classes have been described. These typically involve the scaffold generating terpene synthases and cyclases, and scaffold tailoring enzymes such as e.g. cytochrome P450 monooxygenases, NAD(P)⁺ and flavin dependent oxidoreductases, and various group transferases that generate the final bioactive structures. The biosynthesis of several sesquiterpenoid mycotoxins and bioactive diterpenoids has been well-studied in Ascomycota (e.g. filamentous fungi). Little is known about the terpenoid biosynthetic pathways in Basidiomycota (e.g. mushroom forming fungi), although they produce a huge diversity of terpenoid natural products. Specifically, many *trans*-humulyl cation derived sesquiterpenoid natural products with potent bioactivities have been isolated. Biosynthetic gene clusters responsible for the production of *trans*-humulyl cation derived protoilludanes, and other sesquiterpenoids, can be rapidly identified by genome sequencing and bioinformatic methods. Genome mining combined with heterologous biosynthetic pathway refactoring has the potential to facilitate discovery and production of pharmaceutically relevant fungal terpenoids.

- 1 **Introduction**
- 2 **Fungal sesquiterpenoids**
 - 2.1 **Cyclization of FPP to produce the sesquiterpene scaffold**
 - 2.2 **The trans-humulyl cation derived sesquiterpenoids**
 - 2.3 **Modifying the sesquiterpene scaffold**
- 3 **Fungal diterpenoids**
 - 3.1 **Cyclization of GGPP to produce the diterpene scaffold**
 - 3.2 **Modifying the diterpene scaffold**
- 4 **Fungal triterpenoids**
 - 4.1 **Oxidosqualene/lanosterol cyclases**
 - 4.2 **Discovery of triterpenoid biosynthetic gene clusters**
- 5 **Mining the fungal (sesqui)terpene**
 - 5.1 **Terpene synthase identification**
 - 5.2 **Identification of terpenoid biosynthetic gene clusters**
 - 5.3 **Characterization of terpenoid biosynthetic gene clusters**
- 6 **Conclusions and outlook**
- 7 **Acknowledgements**
- 8 **References**

1 Introduction

Fungi are masters of evolution. With a history spanning at least 900 million years,¹ fungi have successfully adapted to almost every habitat on earth.² Conservative estimates place global fungal diversity beyond that of land plants by a ratio of 10:1.³ While there remains some uncertainty regarding the exact number of fungal species that exist,⁴ it is clear that fungi have been afforded with a unique evolutionary and environmental fitness. Fungi owe this inherent ability to survive to their extensive repertoire of natural product pathways. Many of the natural products from these pathways have antimicrobial, antifungal, immunosuppressive or cytotoxic effects.⁵⁻⁷ These bioactive properties enable fungi to successfully take hold of an ecological niche by conferring the ability to compete for nutrients, to deter predators, and to communicate with other organisms in the environment.⁸ These same fungal bioactive compounds can also be harnessed by humans to produce valuable medicines such as antimicrobial, anticancer and antiviral agents.⁹⁻¹⁶ Natural product pathway discovery and engineering from fungi therefore holds great promise for the pharmaceutical industry.

The importance of natural products for the survival of fungi is evidenced by the fact that they often have many different secondary metabolic pathways that are typically clustered within fungal genomes.¹⁷ One advantage that this feature presents for natural product discovery and engineering is that the physical clustering of genes facilitates genome mining approaches in the identification of biosynthetic pathways. If one gene in a secondary metabolic pathway can be identified, it is highly likely that the other pathway genes will be closely associated and therefore relatively easily identified by bioinformatic methods.¹⁸⁻²⁰ The recent decrease in the cost of next-generation DNA sequencing technologies²¹ has led to a large increase in the number of sequenced fungal genomes that are publically available through databases such as the Joint Genome Institute (JGI). While some of the fungal species represented in this database are of interest due to their lignin degrading capabilities,²² recent trends have seen a shift towards genome sequencing and mining strategies to uncover natural product pathways in species that produce medicinally relevant compounds.²³⁻²⁵ With such a unique toolbox, those interested in fungal natural product discovery and biosynthesis can tap into the vast, yet largely unexplored contingent of bioactive natural product pathways.

One of the largest groups of bioactive natural products that have been identified is the terpenoids. With over 55,000 initial terpene scaffolds so far described, terpenoids belong to an incredibly structurally diverse class of natural products.²⁶ Despite this diversity, all terpenoids are derived from the simple five carbon precursor molecules dimethylallyl diphosphate (DMAPP) **1** and isopentenyl diphosphate (IPP) **2**. In fungi, these two isomers are produced from acetyl-CoA via the mevalonate pathway.²⁷ Condensation of IPP **2** and DMAPP **1** monomers results in linear hydrocarbons of varying length: C10 geranyl pyrophosphate (GPP) **3**, C15 (*2E*, *6E*)-farnesyl pyrophosphate (*(2E, 6E)*-FPP, or FPP) **4**, and C20 geranylgeranyl pyrophosphate (GGPP) **5**. These linear hydrocarbons undergo a dephosphorylation and cyclization cascade to produce terpenes. This highly complex reaction is catalyzed by enzymes known as terpene synthases.²⁸ Two distinct classes of terpene synthase exist, defined according to substrate activation mechanism. Class I terpene synthases catalyze an ionization-dependent cyclization of substrate, while class II terpene synthases catalyze a protonation-dependent cascade.^{28, 29 30-32} Depending on the length of the precursor molecule, fungal terpene synthases are known to produce sesquiterpenes (C15), diterpenes (C20) and triterpenes (C30) (Scheme 1). Further biosynthetic pathway enzymes such as cytochrome P450 monooxygenases, oxidoreductases, and different group transferases modify this initial terpene scaffold, producing the final bioactive terpenoid natural product.

This review will focus on the biosynthesis of the major fungal terpenoid natural product classes that have been described in the last three decades: the sesquiterpenoids, the diterpenoids and the triterpenoids. To date no *bona fide* fungal monoterpene synthases have been described. To keep this contribution focused, natural products of mixed isoprenoid biosynthetic origin (e.g. indole-

diterpenoids, meroterpenoids) that involve different enzymatic steps to install the terpenoid moieties in these compounds will not be presented here.

Considering that terpenoids are the focus of a large body of literature, this article will discuss examples of the terpene synthases and cyclases responsible for the generation of the different fungal terpenoid scaffolds as well as examples of known terpenoid biosynthetic pathways. Comparisons will be drawn between the terpenomes of the two major fungal divisions: the Ascomycota which includes many well-known filamentous fungi like *Aspergillus*, *Penicillium* and *Fusarium*, and the Basidiomycota which includes the mushroom-forming fungi. Special emphasis will be placed upon the highly diverse and bioactive *trans*-humulyl cation derived sesquiterpenes produced by the Basidiomycota. Finally, to illustrate the promise of genome mining for the discovery of new bioactive terpenoids from fungi, a description of current bioinformatic approaches used in the identification of sesquiterpenoid biosynthetic pathways in Basidiomycota will be provided as example along with objectives for the development of tools to explore the largely untapped terpenome of this group of fungi.

2 Fungal sesquiterpenoids

2.1 Cyclization of FPP to produce the sesquiterpene scaffold

The sesquiterpenoids are a diverse group of cyclic hydrocarbons, with more than 300 sesquiterpene scaffolds described.³³ All sesquiterpenes share in common a C15 backbone derived from the linear precursor FPP **4**. Typically, FPP **4** is cyclized by class I terpene synthases known as sesquiterpene synthases, which are characterized by the signature active site motifs DDXXD and NSE.³⁴⁻³⁶ These amino acid residues play an important role in coordinating the catalytically essential divalent metal ions that stabilize the pyrophosphate group of FPP **4** within the active site cavity.^{35, 36} Cyclization is initiated by a metal ion induced ionization of the substrate and departure of inorganic pyrophosphate (PPi), which promotes structural displacements including closure of the active site lid.³⁵ The resulting highly reactive carbocation undergoes an initial ring closure at the 1,10 or the 1,11 position. Some enzymes catalyze first the *trans-cis* isomerization of the 2,3-double-bond of (*2E,6E*)-FPP **4** to (*3R*)-nerolidylpyrophosphate **6** prior to generating a *cisoid*, allylic nerolidyl-carbocation upon PPi cleavage.^{37, 38} This nerolidyl-carbocation results in a known initial ring closure at either the 1,6 or the 1,10 position; 1,7 ring closure may be possible but has not yet been mechanistically shown (Scheme 1). Subsequent proton shifts, methyl shifts, and complex ring rearrangements are stabilized by the aromatic residues that line the active site cavity of all sesquiterpene synthases.²⁸ The reaction cascade is quenched either by attack by a water molecule,³⁹⁻⁴¹ or by deprotonation which, based on the fact that no active site base has yet been confirmed,⁴²⁻⁴⁴ is believed to be mediated by the leaving PPi group.^{45, 46} The final terpene product and divalent metal ions are released by the enzyme upon opening of the active site lid. Comprehensive mutational analyses and structural studies have provided detailed insights into the catalytic mechanisms used by

fungal sesquiterpene synthases, therefore this class of terpene synthase is relatively well understood (reviewed in:^{28, 30, 33}).

Structures are known for several microbial and plant sesquiterpene synthases.^{34, 35, 39, 47-52} However, crystal structures have only been solved for two fungal sesquiterpene synthases; aristolochene **7** synthase from *Penicillium roqueforti* and *Aspergillus terreus*;^{46, 48, 52, 53} and trichodiene **8** synthase from *Fusarium sporotrichioides*.^{42, 51, 54, 55}

Trichodiene **8** synthase (TS) from the plant pathogens *Trichothecium roseum*³⁸ and *Fusarium sporotrichioides* was the first characterized fungal enzyme.⁵⁶ Production of trichodiene **8** (Scheme 2) is the first committed step in the pathway to the trichothecene mycotoxins (see section 2.3) that cause the major cereal crop disease *Fusarium* head blight (FHB).⁵⁷ The mechanistic details of TS have been deduced by site directed mutagenesis³⁶ and structure solution of recombinant wild type and D100E enzyme in complex with PPi resulting from incubation with the substrate analogue 2-fluorofarnesyl diphosphate, and in complex with benzyl triethylammonium.^{42, 51, 54, 58} Removal of PPi from FPP **4** is initiated by three Mg²⁺ ions, two of which are coordinated by D100, and a third which is coordinated by the triad N225, S229 and E233.^{51, 54} The resulting cation is isomerized to (*3R*)-nerolidyl pyrophosphate (NPP) **6** upon recapture of PPi.⁵⁹ Rotation about the C2-C3 bond allows the intermediate to adopt a *cisoid* conformation, and departure of PPi results in a 1,6 ring closure to yield a (*6R*)- β -bisabolyl cation **9**. Residues K232, R304 and Y305 act to stabilize PPi, which remains bound at the active site throughout product generation.⁵⁸ Subsequent hydride transfer and a 7,11 ring closure produces a secondary cuprenyl cation **10**.^{60, 61} Finally, two methyl shifts and deprotonation, possibly mediated by the remaining PPi, results in trichodiene **8** as the major product (Scheme 2), and several minor products including α -cuprenene **11**, α -barbatene **12**, α - and β -bisabolene **13** and **14**.^{38, 42, 54}

The structure of aristolochene **7** synthase (AS) from *Penicillium roqueforti* (PR-AS)⁴⁸ and *Aspergillus terreus* (AT-AS) is also known.^{46, 52, 53} AS belongs to the subclass of 1,10 cyclizing sesquiterpene synthases, catalyzing the ionization of FPP **4** and an initial ring closure to yield germacrene A **15** (Scheme 3). Protonation at C6 leads to the eudesmane carbocation **16**,⁶² which following a methyl migration, hydride transfer, and deprotonation, yields (+)-aristolochene **7**.^{45, 63} Aristolochene **7** is the sesquiterpenoid scaffold of several mycotoxins, including the PR-toxin **17**, sporogen-AO1 and phomenone.⁶⁴

Detailed characterization of AS from the two distinct fungal sources has provided a unique perspective upon the evolution of sesquiterpene synthases. Both PR-AS and AT-AS follow the same cyclization mechanisms and display very similar three dimensional folds, indicating that the two enzymes share a common evolutionary ancestor. However, the underlying amino acid sequence of the two enzymes differs significantly, and this divergence in sequence has resulted in a divergence in fidelity. AT-AS produces (+)-aristolochene **7** (Scheme 3) as its sole product, while PR-AS is product promiscuous, producing (+)-aristolochene **7** as its major product and (-)-valencene **18** and germacrene A **15** as minor products (at a ratio of 94:2:4).

Comparative site directed mutagenesis of the active site metal coordinating sites of both enzymes highlighted the importance of these conserved motifs in product fidelity, in particular the NSE motif. In PR-AS, the product ratio of mutants N244D, S248A and E252D shifted to 20:80 (+)-aristolochene **7**: germacrene A **15**, while E252Q produced only germacrene A **15**. However, none of the mutants reported for PR-AS produced solely (+)-aristolochene **7**. The product ratio of the corresponding AT-AS mutants N219D and E227D shifted from (+)-aristolochene **7** as the only product to 44:56 and 26:74 (+)-aristolochene **7**: germacrene A **15**, respectively.⁴³ It would therefore appear that while the NSE metal coordinating motif serves as the modulator of product fidelity in AT-AS, other amino acids beyond these conserved residues play a role in determining the specific production of (+)-aristolochene **7** by PR-AS.

Three sesquiterpene synthases have recently been described from *Fusarium* strains. One terpene synthase (longiborneol **19** synthase FgCLM1, Scheme 2) from *F. graminearum* is presumed to catalyze a 1,6 cyclization to synthesize longiborneol **19** as the precursor of the antifungal compound culmorin.⁶⁵ Gene deletion studies in the rice pathogen *F. fujikuroi* identified two terpene synthases involved in the production of α -acorenol **20** (acorenol synthase Ff_sc6, Scheme 2) and koraiol **21** (koraiol synthase Ff_sc4, Scheme 4) via a 1,6 and presumably a 1,11-cyclization reaction, respectively, followed by quenching of the final carbocation with water to yield the sesquiterpenoid alcohol products.⁶⁶ The botrydial **22** biosynthetic gene cluster (see section 2.3) identified in the grey mold *Botrytis cinerea* encodes yet another 1,11-cyclizing enzyme (BcBOT2) which generates the tricyclic alcohol presilphiperfol-8 β -ol **23** (Scheme 4) as precursor for this phytotoxin.^{67, 68}

Surprisingly, until a few years ago no sesquiterpene synthases were known from Basidiomycota, despite the fact that these fungi are prolific producers of bioactive sesquiterpenoids, many of which are derived from the *trans*-humulyl cation,⁹ which will be discussed in more detail in the next section. Mining of the first genome of a mushroom-forming Basidiomycota, *C. cinerea*, led to the cloning and biochemical characterization of six enzymes (Cop1-6) catalyzing 1,10 and 1,6 cyclization reactions of 3R-NPP **6** to α -cuprenene **11**, germacrene D **24**, γ -cadinene **25**, δ -cadienene **26**, cubebol **27** shown in Schemes 2 and 5, and 1,10 cyclization of FPP **4** to α -muurolene **28** and germacrene A **15** shown in Scheme 3. One of the enzymes, Cop6, catalyzes highly selective synthesis of α -cuprenene **11** as the precursor of the antimicrobial compound lagopodin.⁶⁹⁻⁷¹

Of particular interest are enzymes that generate the 1,11-cyclization product Δ 6-protoilludene **29** (Scheme 4), the precursor to many pharmaceutically relevant compounds,⁹ including the cytotoxic compounds illudin M **30** and S **31** (Scheme 6) that are being developed as anticancer therapeutics.^{15, 72, 73} With the objective of characterizing the biosynthesis of the anticancer illudins and their derivatives (compounds **30-33**, Scheme 6) made by the Jack-O-Lantern mushroom *Omphalotus olearius*,^{74, 75} we sequenced its genome to apply a similar genome mining approach to find the Δ 6-protoilludene **29** and presumably co-localized illudin biosynthetic genes. We discovered a surprisingly large complement of ten sesquiterpene synthases

(Omp1-10) in this fungus. The recombinant enzymes catalyze all cyclization reactions of FPP shown in Scheme 1 (Schemes 2-5 show the major products for each Omp sesquiterpene synthase). The products made by of Omp1-5a/b are similar to those obtained with the *C. cinerea* enzymes Cop1-4, although Omp5a/b also makes as major products *epi*-zonarene **34** and γ -cadinene **25** (Scheme 5). Omp6 and Omp7 cyclize FPP highly selective into Δ 6-protoilludene **29** (Scheme 4) while the other enzymes also display new cyclization activities, producing barbatenes **12**, **35** (Omp9) and daucenes **36**, **37** (Omp 10).^{23, 76} Another Δ 6-protoilludene synthase was cloned from the honey mushroom *Armillaria gallica* (ArmGa1) which makes antimicrobial melleolide I **38** (Scheme 6) sesquiterpenoids.⁷⁷

The biochemical characterization of the identified sesquiterpene synthases from *Coprinus cinereus* (Cop1-6)⁶⁹⁻⁷¹ and of *Omphalotus olearius* (Omp1-10)^{23, 76} guided the subsequent development of *in silico* approaches for the directed discovery of new sesquiterpene synthases and their associated biosynthetic genes based upon cyclization mechanism of choice.²⁰ Initiatives such as the 1000 Fungal Genomes Project⁷⁸ have led to a substantial increase in the number of sequenced genomes that are publically available.⁷⁹ The increase in the number of sequenced Basidiomycota genomes has been particularly steep – from less than a handful in 2008 to more than 100 genomes listed in the genome database of the Joint Genome Institute (JGI) at the time of writing. Our own BLAST searches for terpene synthases across Basidiomycota with sequenced genomes led to the identification of hundreds of putative sesquiterpene synthases, as well as associated biosynthetic gene clusters.^{20, 23, 69}

The genome sequence of the wood-rotting mushroom *Stereum hirsutum* offers a unique opportunity to explore the terpenoid structural diversity of a fungal genus that is well known to produce a number of bioactive sesquiterpenoid natural products.⁸⁰⁻⁹³ These include the sterostreins (e.g. **39-42**), sterpurene **43** and its derivatives (e.g. **44-47**), hirsutene **48** and its derivatives (e.g. **49-53**), cadinane (**25**, **26**) (e.g. **54-55**) and drimane (e.g. **56**, **57**) compounds of which representative examples are shown in Scheme 6. We identified and predicted the cyclization mechanisms of 16 sesquiterpene synthases in this fungus. Presently, five of the predicted enzymes were cloned and biochemically characterized. Steh1|25180, 64702, 73029 are highly product specific Δ 6-protoilludene **29** synthases (Scheme 4) that are located in large biosynthetic gene clusters presumably involved in sterostrein (Scheme 6, e.g. compounds **39-42**) biosynthesis.^{89, 91} Steh1|159379 and Steh1|128017 are 1,6- and 1,10-cyclizing enzymes, respectively (Schemes 2 and 5). Within the clade of not yet characterized 1,11-cyclizing sesquiterpene synthases sequences identified in *S. hirsutum*, there may be enzymes that synthesize hirsutene **48** and sterpurene **43** (Scheme 4) for the above mentioned modified sesquiterpenoid compounds isolated from this fungal genus.

Furthermore, pentalene **58** (Scheme 4) has been observed as a minor volatile product in the culture headspace of *S. hirsutum* and *O. olearius*, and as a major volatile product of *C. cinerea*.^{20, 23, 69} However, no sesquiterpene synthase responsible for the production of pentalene **58** has yet been identified from these fungi. Site directed mutagenesis and quantum-chemical modelling

studies with bacterial pentalenene synthase indicate that the pathway to pentalenene **58** occurs through rearrangement of a protoilludyl cation **59**.^{44, 94} It is therefore likely that one or more of the multiple Δ 6-protoilludene synthases characterized from *S. hirsutum* and *O. olearius* are responsible for the production of pentalenene **58**, under unidentified reaction conditions.^{20, 23} On the other hand, Cop5 is likely the sesquiterpene synthase responsible for the production of pentalenene **58** by *C. cinerea*, which does not produce Δ 6-protoilludene **29**. However, an active form of the predicted sesquiterpene synthase could not be cloned from *C. cinerea* cDNA,⁶⁹ presumably due to incorrect prediction of gene borders, and/or the presence of frame shifting introns in all amplification products (see section 5 for more information).

2.2 The *trans*-humulyl cation derived sesquiterpenoids

While it is difficult to pin down the exact number of sesquiterpenoid-derived compounds characterized from the two major fungal divisions, searches in SciFinderTM and Web of ScienceTM return ten times more reports (~500) on sesquiterpenoids isolated from Basidiomycota than from Ascomycota. A diverse range of highly bioactive sesquiterpenoids with cytotoxic, antibacterial, nematocidal, and antiviral activities have been isolated from Basidiomycota (for reviews see:^{9, 95}). Many of these compounds are thought to be made to protect fungal fruiting bodies from consumption.^{96, 97}

Basidiomycota are known to synthesize sesquiterpenoids derived from 1,6 and 1,10 cyclized cations (Scheme 1), including germacranes,⁹⁵ bisabolanes,⁹⁶ cadinanes^{86, 90, 92} and drimanes,^{84, 98} including the mTOR signaling pathway inhibitor antrocin **57** produced by *Antrodia camphorata*⁹⁰ (Scheme 6) However, the vast majority of sesquiterpenoids with medicinal or cytotoxic properties isolated from Basidiomycota are derived from the 1,11 cyclized *trans*-humulyl cation **60** as shown in Schemes 4 and 7, which will be the emphasis of this section.⁹ Of particular interest are derivatives of the tricyclic protoilludyl cation^{9, 97} which have potential applications as anticancer, antifungal and antibiotic agents.⁹

The *trans*-humulyl cation can be directly deprotonated to produce α -humulene **61** (Scheme 4), which has been observed in *F. fujikorii* and *S. hirsutum*.^{20, 66} Modified humulanes, such as the potential antitumor compound mitissimol B **62** (Scheme 6), have also been isolated from the fruiting bodies of several members of the genus *Lactarius*.⁹⁹

Transformations of the *trans*-humulyl cation include a 2,10 ring closure, creating the distinctive dimethylated cyclobutyl ring of the caryophyllane scaffold. The mildly antibacterial sesquiterpenoid naematolon **63** (Scheme 6) is an oxidized, acetylated derivative of (*E*)- β -caryophyllene **64** produced by several members of the *Naematoloma* (*Hypholoma*) genus.⁹⁶

In addition, the caryophyllane scaffold can be further cyclized to produce the tricyclic 4:7:4 collybial scaffold by a secondary 3,5 closure. The collybial scaffold is the precursor to the antibiotic collybial **65** (Scheme 6), produced by *Collybia confluens*.¹⁰⁰ Hydroxylation at the C7 of the collybial scaffold results in koraial **21**. Alternatively, the caryophyllane scaffold can be cyclized to produce the triquinane presilphiperfolane scaffold, either via a

stepwise cyclization,⁶⁸ or via a concerted carbocation rearrangement.³² The resulting presilphiperfolane scaffold may then proceed through a 1,2 hydride shift and deprotonation to yield presilphiperfolan-1-ene **66**, a minor volatile product of *S. hirsutum*.²⁰ The presilphiperfolane scaffold may also undergo a 1,3 hydride shift and hydroxylated to produce **23**,⁶⁸ or be rearranged further to produce the silphiperfolane and silphinane families of triquinane sesquiterpenoids.¹⁰¹

Several tremulanes have been characterized from *Phellinus tremulae*, including tremulenediol A **67** (Scheme 6).^{102, 103} The potential route to the tremulane scaffold was deduced by ¹³C NMR analysis of tremulenediol A **67** formation by *P. tremulae*, and follows an initial 2,9 closure of the *trans*-humulyl cation **60**. Following this, an additional cyclobutyl ring formation occurs at C6-C9. A secondary ring-opening rearrangement yields the fifth methyl group on C6 (Scheme 9).¹⁰³

The africananes are defined by a cyclopropyl group fused to a central cycloheptane ring, formed by a 7,9 cyclization of the *trans*-humulyl cation **60**. Examples of modified africanane alcohols are leptographiol **68** (Scheme 6), isoleptographiol, and isoafrikanol, isolated from *Leptographium lundbergii*.^{97, 104} Interestingly, unmodified africananes have been identified in the headspace of *S. hirsutum* and *O. olearius* liquid cultures.^{20, 23} However, the sesquiterpene synthase(s) responsible for their production have not yet been identified.

The largest and most diverse group of *trans*-humulyl cation **60** derived sesquiterpenoids produced by Basidiomycota are those derived from the 4:6:5 tricyclic protoilludyl cation **59**.⁹ This scaffold is a key intermediate in the production of a large number of sesquiterpenes including e.g. sterpurene **43**, hirsutene **48**,¹⁰⁵ Δ 6-protoilludene **29** and thereof derived scaffolds,²³ and of pentalenene **58**⁹⁴ as described below and shown in Scheme 7.

Δ 6-protoilludene **29** is the precursor to diverse bioactive sesquiterpenoids, including the marasmanes, fommanosanes, (seco)lactaranes, hyrogammane and the illudanes (reviewed in:⁹). These compounds are hypothesized to be produced by secondary ring opening and concurrent oxidations of Δ 6-protoilludene **29**, with the initial steps likely catalyzed by one or more cytochrome P450 monooxygenases and/or other oxygenases that are typically located in biosynthetic gene clusters of fungal sesquiterpenoid pathways.^{20, 23, 106, 107} In the case of the illudins **30** and **31**, a ring opening and subsequent contraction of the Δ 6-protoilludene **29** cyclobutyl ring yields the reactive cyclopropyl ring that defines the illudanes. Presently, no biosynthetic enzymes have been characterized that initiate these ring opening reactions.

Reopening of the protoilludyl cation **59** at C5-C6, and a subsequent 5,7 cyclization may afford the cerapicane scaffold which is the precursor to the rare cucumane and bullerane-type sesquiterpenes.⁹ The cerapicane scaffold is transformed to the sterpurane scaffold by an unusual ring rearrangement reaction that was deduced by ¹³C NMR analysis of 9,12-dihydroxysterpurene **46** synthesized by *Stereum purpureum*.⁸¹ Further ring opening modifications can then contract the cyclobutyl ring of sterpurene **43** into the defining dimethylated cyclopropyl ring of the *iso*-lactaranes, several of which have been identified from *Lactarius rufus*,¹⁰⁸ *S. purpureum*,⁸⁰ *Phlebia uda*,¹⁰⁹ *Flammulina velutipes*,¹¹⁰ and *Merulius tremillosus*.¹¹¹ The

iso-lactaranes are often co-produced with sterpurenones, supporting the hypothesis of a shared biosynthetic pathway.⁸⁰ Examples of *iso*-lactaranes include the cytotoxic flammulinolide A **69**,¹¹⁰ sterepolides (e.g. **45**),⁸⁰ and the sterelactones (e.g. **47**) (Scheme 6).⁸⁷

Sterostrein A **39** produced by *S. ostrea* is a Δ 6-protoilludene **29** dimer hypothesized to be formed by a Diels-Alder reaction that installs the central benzene ring.⁸⁹ Biosynthesis of the melleolides (e.g. **38** in Scheme 6) by *Armillaria gallica* requires the esterification Δ 6-protoilludene **29** with orsellinic acid.^{77, 112}

The bioactive hirsutanes (e.g. compounds **49-53**, Scheme 6) produced by *S. hirsutum* and other Basidiomycota^{20, 93, 88, 113} may be synthesized either through 3,7 rearrangement of the protoilludyl cation **59** or directly via 3,7 cyclization of a bicyclic precursor shown in Scheme 7. While the intermediacy of a protoilludyl cation **59** has not been confirmed, a ceratopicane intermediate is consistent with the ¹³C labelling pattern observed in hirsutene **48** biosynthesis by *Stereum complicatum*.¹⁰⁵ The ceratopicane intermediate undergoes multiple rearrangements, shifting the C3 (and its attached C14 methyl group) to its final position, and is then deprotonated to produce hirsutene **48**.¹⁰⁵ Hirsutene **48** may be converted by a Wagner-Meerwein rearrangement to produce the pleurotellane scaffold.⁹

2.3 Modifying the sesquiterpene scaffold

Compared to pathways responsible for the biosynthesis of diterpenoid natural products discussed in the next section, very few fungal sesquiterpenoid biosynthetic pathways have been characterized on a gene level. One of the most extensively characterized fungal sesquiterpenoid biosynthetic pathways is the route to the trichothecene mycotoxins (Scheme 8). Over 200 trichothecenes have been identified from diverse sources including the plant pathogens *Fusarium* and *Myrothecium*, the soil associated filamentous fungi *Trichoderma* and the mold *Stachybotrys*.^{114, 115} This particular pathway has been the focus of a great deal of research by the USDA during the past 30 years due to the significant economic impact that trichothecene related blight and ear rot has on global cereal crops each year, as well as the toxic effects of trichothecene contaminated feed on livestock.¹¹⁵⁻¹¹⁷ The trichothecene family of toxins is subclassified as Type A, B, C, and D depending on the chemical modifications to the common tricyclic core structure 12,13-epoxytrichothec-9-ene (EPT), exemplified by isotrichodermol **70** in Scheme 8. Of particular interest is deoxynivalenol (DON) **71**, the Type B trichothecene that is most commonly associated with *Fusarium* head blight in cereal crops. DON **71** plays a vital role in activating fungal pathogenicity and virulence as part of a complex feedback exchange between plant, pathogen and environment.¹¹⁸⁻¹²⁰

The initial, common trichothecene biosynthetic steps in the pathway to DON **71** were established in the model fungus *Fusarium sporotrichioides*.^{56, 121} Later steps specific to the production of DON **71** were studied in a *F. graminearum* strain that was isolated from infected crops.¹²² Activation of biosynthesis of DON **71** is controlled by the global transcriptional regulator *TRI6*, which autoregulates in response to differences in

nutrient levels.¹²³ The first step on the biosynthetic pathway to all trichothecenes is the cyclization of FPP **4** to trichodiene **8** by the *TRI5* encoded trichodiene **8** synthase,⁵⁶ as described in section 2.1 and in Scheme 2. Trichodiene **8** is subjected to multiple oxidations by the cytochrome P450 monooxygenase *TRI4*, which installs oxygens at C2, C3, C11 and the C12-C13 epoxy group to give the intermediate isotrichotriol **72**.¹²⁴ Isotrichotriol **72** undergoes a spontaneous dual cyclization to yield isotrichodermol **70**, a modified form of EPT.¹²⁵ An acetyltransferase *TRI101* acetylates the hydroxyl group at C3 to produce isotrichodermin **73**, reducing toxicity of the intermediate to the host.^{126, 127} C15 of **73** is hydroxylated by the cytochrome P450 monooxygenase *TRI11*¹²⁸ to form intermediate **74** and acetylated by *TRI3*¹²⁹ yielding calonectrin **75**, a key branch point on the pathway to biosynthesis of Type A and Type B trichothecenes by *F. sporotrichioides* and *F. graminearum*, respectively.¹¹⁵ For DON **71** production, calonectrin **75** is hydroxylated at positions C7 and C8 by *TRI1* to give 7,8-dihydroxycalonectrin **76**.¹³⁰ The C8 hydroxyl group of **76** is then converted to a keto group by an unidentified enzyme, leading to 3,15-acetyldeoxynivalenol **77**. Different variants of the deacetylase *TRI8* remove the acetyl group from either the C3 or the C15 position, resulting in 15-acetyldeoxynivalenol (15-ADON) or 3-ADON **78**, (the latter compound is shown in Scheme 8), respectively.¹³¹ The final biosynthetic step, deacetylation of ADON to DON **71**, has yet to be elucidated (Scheme 8).

The ability of the fungal host to tolerate the toxic effects of intermediates and products of its own secondary metabolic pathways is key to survival. *TRI12* is an integral membrane protein that acts as an efflux pump to export trichothecenes from hyphal cells, removing toxic products that could otherwise damage the cell.¹³² Subcellular localization of *TRI12* to large motile vesicles¹³³ indicated that *F. graminearum* could rely on an encapsulation mechanism to sequester trichothecene biosynthesis. The cytochrome P450 monooxygenases *TRI1* and *TRI4* co-localize with *TRI12*, suggesting that at least part of the trichothecene biosynthetic pathway is integrated within cellular targeting hubs, “toxisomes”, preventing toxic intermediates from causing unwanted effects.¹³⁴ Whether this compartmentalization strategy is unique to *F. graminearum*, or whether it is a global mechanism employed by fungi producing bioactive secondary metabolites, remains to be established. Similarly, vesicles have been shown to also play an important role aflatoxin biosynthesis by *Aspergillus parasiticus*.^{135, 136}

A much less complex gene cluster has been identified in the mold *Botrytis cinerea* that is responsible for the biosynthesis of the phytotoxin botrydial **22** (Scheme 8).^{67, 68} Presilphiperfolan-8 β -ol **23**, the product of terpene synthase *BcBOT2* (Scheme 4), is converted by three cytochrome P450 monooxygenases and an acetyltransferase to the final phytotoxin. Most recently, screening of a genomic phage library and comparison with orthologous genes identified in the sequenced genome of a related strain, *P. chysogenum*, led to the identification of a 10-gene cluster encoding the biosynthetic pathway for the aristolochene 7-derived PR-toxin **17** in the blue cheese mold *P. roqueforti* (Scheme 8).¹³⁷

As discussed in the previous section, multiple copies of Δ 6-protoilludene **29** synthase are encoded in the genomes of *O.*

olearius (Omp6 and Omp7) and *S. hirsutum* (Steh1|25180, 64702, 73029) which are each located in predicted biosynthetic gene clusters responsible for the biosynthesis of illudin derived compounds (e.g. compounds **30-33**) and potentially the sterostreins (e.g. compounds **39-42**) isolated from *O. olearius* and *Stereum ostrea* BCC22955, respectively (Scheme 6).^{20, 23} The clusters share in common several cytochrome P450 monooxygenases, as well as further scaffold decorating enzymes including different types of oxygenases, oxidoreductases, group transferases and membrane transporters.

In *O. olearius*, Omp7 is located in a mini-gene cluster, including one P450 and an FAD-type oxidoreductase, while Omp6 is part of a large gene cluster.²³ It appears that the two copies of protoilludene **29** synthase may have arisen from a gene duplication event, thereby boosting the production of illudin compounds.²³ In the case of *S. hirsutum*, the three Δ 6-protoilludene **29** synthases may have resulted from gene duplication, or they may have diverged from a common ancestor. Differences in catalytic efficiencies of the enzymes, as well as a diversity in product fidelity, could represent a spectrum of evolutionary time points on the course to a specific Δ -6 protoilludene **29** synthase.²⁰ Furthermore, each of the three Δ -6 protoilludene **29** synthases is located in a distinct biosynthetic gene cluster which appear to be expressed at the same time under standard growth conditions.²⁰ Heterologous refactoring of the entire gene clusters (see section 5) will be required to understand the role that each of these pathways plays in defining the natural product landscape of this fungus. It remains to be seen if all three biosynthetic gene clusters are responsible for the biosynthesis of the different Δ 6-protoilludene **29** derived sterostreins (Scheme 8) isolated from *Stereum*, make yet uncharacterized compounds, or may even carry out a sophisticated combinatorial scheme among the three different clusters that each act on the same protoilludene scaffold.

3 Fungal diterpenoids

3.1 Cyclization of GGPP to produce the diterpene scaffold

Diterpenoids are a diverse class of natural products derived from the C20 precursor GGPP **5**, with at least 12,000 compounds already described.¹³⁸ The first committed step in diterpenoid biosynthesis is the cyclization of GGPP **5** to produce the diterpene scaffold, which occurs via a carbocation cascade. Classically, activation of the carbocation cascade by terpene synthases corresponds to the removal of the pyrophosphate group from the linear substrate (as described for the sesquiterpene synthases in section 2.1). This ionization-dependent reaction is catalyzed by class I terpene synthases.²⁸

Notably, the initial generation of the reactive carbocation follows a different path in the biosynthesis of labdane-related diterpenes.¹³⁹ Here, the carbocation cascade is preceded by a protonation-dependent bicyclization reaction to install the labdane bicycle. This reaction is catalyzed by class II diterpene synthases, more akin to the triterpene synthases (described in section 4.1).¹⁴⁰ In this case, the pyrophosphate group of GGPP **5** remains intact and is later removed in a class I cyclization reaction (Scheme 9).¹³⁹ Thus, this type of diterpene synthase appears to have

evolved as an amalgamation of ancestral bacterial class I and class II terpene synthases.¹³⁹

The first bifunctional fungal diterpene synthase was isolated and characterized from the gibberellin producer *Phaeosphaeria* sp. L487.¹⁴¹ The putative gene encoding the diterpene synthase was isolated using degenerate primers based upon conserved sequence motifs of plant copalyl diphosphate synthases. Characterization of the gene product revealed a conglomerate enzyme: that is, a bifunctional copalyl diphosphate synthase/*ent*-kaurene synthase (CPS/KS), which catalyzed a sequential class II followed by class I cyclization of GGPP **5** to produce (-)-*ent*-CDP **79** as an intermediate on the path to *ent*-kaurene **80** (Scheme 9).¹⁴² The class I activity is located in the N-terminal region, while the C-terminal region contains the class II activity. This differed from the previously characterized plant diterpene synthases, which appeared to involve separate mono-functional CPS and KS enzymes.^{143, 144}

Several structures of plant diterpene synthases have recently been solved and shown to contain both an N-terminal α -helical domain (α -domain) typical of class I terpenoid synthases and a C-terminal α -barrel (or $\gamma\beta$ -) domain typical of class II terpenoid synthases.¹⁴⁵⁻¹⁴⁷ Many of these modular $\gamma\beta\alpha$ -domain plant diterpene synthases, however, are monofunctional in that only one of the cyclization activities (CPS or KS) is functional (reviewed in: ¹³⁹). However, some plant enzymes (e.g. abietadiene synthase¹⁴⁷) are bifunctional, like the fungal enzymes. The $\gamma\beta\alpha$ -domain taxadiene synthase, however, converts GGPP **5** at its class I domain directly into the tricyclic taxadiene scaffold.¹⁴⁶ Although protein sequences of diterpene synthases from plants and fungi share little similarity except for some conserved motifs, homology modelling suggests that they share the same domain organization.

A number of CPS/KS-type fungal diterpene synthases have been identified in addition to the one from *Phaeosphaeria* sp. These include the *ent*-kaurene **80** synthase from the gibberellic acid producers *Fusarium fujikuroi* (teleomorph: *Gibberella fujikuroi*),^{141, 142, 148, 149} the *ent*-primara-8(14),15-diene **81** synthase AN1495 from *Aspergillus nidulans*,¹⁵⁰ phyllocladen-16 α -ol **82** synthase (PaDC1) and a CPS/KS synthase (PaDC2) with only a functional CPS domain¹⁵¹ from *Phomopsis amygdali*, and aphidicolan-16 β -ol **83** synthase (PbACS) from *Phoma betae*.¹⁵² Only one labdane-type diterpene synthase has thus far been cloned from a Basidiomycota: pleuromutilin **84** synthase from *Clitopilus passeckerianus*.¹⁵³ (see Scheme 1 for cyclization reactions).

Typically, these bifunctional diterpene synthases and their associated biosynthetic gene clusters (see section 3.2) have been identified by a sequence analysis strategy that first involved the identification of a GGPP **5** synthase (GGS) via genomic amplification followed by genome walking to identify flanking regions. GGPP **5** utilizing biosynthetic pathways require a dedicated GGS that extends the common isoprene precursor FPP **4** by one IPP **2** unit.^{148, 154} The gene encoding GGPP **5** synthase, which is essential for production of the precursor molecule is commonly clustered with other pathway genes.¹⁵⁵ This gene finding strategy has led to the discovery two unusual monofunctional, non-labdane-type diterpene synthases from

Phoma betae.^{156, 157} PaFS and PaPS, which synthesize fusicocca-2,10(14)-diene **85** and phomopsene **86** (Scheme 1), respectively, deviate from other CPS/KS-type fungal diterpene synthases in that they display multiple DDXXD motifs, as opposed to the prototypical N-terminal DXDD and C-terminal DEXXE motifs common to the dual-function diterpene synthases that carry out sequential type II and type I cyclization of GGPP **5** (Scheme 9).¹⁵⁸ The DXDD motif is required for the class II type protonation dependent cyclization reaction, like that carried out by squalene hopene cyclase,¹⁵⁹ while the DDXXD motif is associated with the class I type ionization dependent reaction, which has been described for terpene synthases and prenylchain synthases like GGPP **5** and FPP **4** synthases.^{34, 160}

Both PaFS and PaPS appear to derive from a gene fusion, resulting in chimeric proteins with an N-terminal class I terpene synthase domain and another C-terminal class I domain that contains the GGPP **5** synthase activity. This physical co-localization of diterpene synthase and GGPP **5** synthase may have evolved to improve flux of precursor molecules towards downstream modifying enzymes. A chimeric sesterterpene (C25) synthase AcOS was recently identified in *Aspergillus clavatus*. Here a GFPP (geranylarnesyl diphosphate) synthase domain supplies the substrate for subsequent cyclization by the terpene synthase domain into ophiobolin F **87** (Scheme 1).¹⁶¹ Structural analyses of plant and bacterial diterpene synthases indicate that modularity in domain architecture and diversity in mechanisms are a common theme for diterpene synthases.¹⁴⁵⁻¹⁴⁷

3.2 Modifying the diterpene scaffold

The diversity of diterpenoid compounds stems from the large number of modifications to the diterpene scaffold, typically catalyzed by decorating enzymes such as P450 monooxygenases, different types of oxidases and oxidoreductases as well as transferases. Diterpenoids are widely produced by plants, functioning as signaling compounds, as photosynthetic pigments, and in defense against infection.¹³⁸ While a number of enzymes involved in the biosynthesis of diterpenoids in plants and bacteria have been characterized in detail, fewer diterpenoid biosynthetic enzymes have been characterized from fungi.^{138, 162} Except for pleuromutilin **84** biosynthesis by *Clitopilus passeckeranus*,¹⁵³ all other characterized fungal diterpene pathways are from Ascomycota. Yet, diverse bioactive diterpenoids have been isolated from Basidiomycota, although their biosynthetic pathways remain to be identified (Table 1, see Scheme 10 for modified diterpenoid structures discussed below).¹⁶³

One of the first fungal species identified as a diterpenoid producer was the rice plant pathogen *F. fujikuroi*. This fungus causes stems of plants to become hyperelongated and etiolated, leading to what is commonly known as “foolish seedling” disease. The causative agent of this phenotype was discovered to be the gibberellin (GA) diterpenoids (GA compounds **101-104** Scheme 11) produced by *F. fujikuroi*. The biosynthetic pathway to the production of GAs by *F. fujikuroi*, which is clustered on the genome, has since been fully elucidated by Tudzynski and co-workers¹⁶⁴ (Scheme 11). Detailed characterization of the enzymes involved in GA biosynthesis by *F. fujikuroi* has highlighted key

differences in sequence and function from that of plant GA pathways,¹⁶⁵ such that it has been suggested that GA biosynthesis in bacteria, plants and fungi evolved separately.¹⁵⁵

In GA biosynthesis, GGPP **5** is the substrate for a bifunctional diterpene synthase CPS/KS.¹⁴⁹ The resulting tetracyclic *ent*-kaurene **80** diterpene is converted to *ent*-kaurenoic acid **105** through oxidation by the cytochrome P450 monooxygenase P450-4.¹⁴⁸ Hydroxylation at the C7 position of *ent*-kaurenoic acid **105** is carried out by P450-1, resulting in aldehyde GA12 **101** (note GA numbering is based on the order the compounds were identified), with a characteristic 6-5-6-5 ring structure. A further hydroxylation at the C3 position and an oxidation by the multifunctional P450-1 results in GA14 **102**, a key intermediate in the pathway. A closely related cytochrome P450, P450-2, catalyzes the essential oxidation of GA14 **102** at C20 to yield the biologically active form, GA4 **103**.^{148, 166} Finally, P450-3 hydroxylates GA4 to GA1 **104**, another bioactive form of the growth hormone.¹⁶⁷ Each of the steps catalyzed by the cytochrome P450s depends on a cytochrome P450 oxidoreductase (CPR) to transfer electrons from the cofactor NADPH via FAD and FMN.¹⁶⁸

Subsequent identification and characterization of the GA biosynthetic pathway from *S. manihoticola* revealed that the organization of the gene cluster, the underlying gene sequences, and the function of the encoded GA enzymes, were similar to that of *F. fujikuroi*.¹⁶⁹ Despite the fact that *S. manihoticola* lacked enzymes responsible for GA1 **104**, **3**, and **7** production, complementation with its gene cluster was sufficient to restore GA biosynthesis in *F. fujikuroi* mutants. This indicates that the GA pathways of these two distantly related fungi may share a common evolutionary ancestor. By contrast, characterization of the GA biosynthetic pathway in *Phaeosphaeria* sp. L487 shows that it is more similar to that of plants.^{170, 171} Here, the GA pathway branch point is GA12 **101** rather than GA14 **102**, which is the key intermediate linking the GA pathways of *F. fujikuroi* and *S. manihoticola*. Additionally, the C3 hydroxylation takes place later in the GA pathway of *Phaeosphaeria* sp. L487 than that of the other fungi.

Diversification of ancestral GA-pathways likely led to new diterpene pathways that produce different bioactive compounds. Several dozens of such putative diterpene biosynthetic gene clusters can be readily identified in published Ascomycota genome sequences. However, only very few diterpene biosynthetic gene clusters have been characterized so far. The only labdane-type diterpene pathway from a Basidiomycota has been disclosed in a patent application for pleuromutilin antibiotic biosynthesis in *Clitopilus passeckeranus*.¹⁵³

Identification of the aphidicolan-16 β -ol **83** synthase¹⁵² from *Phoma betae* (Scheme 1) facilitated subsequent identification and heterologous expression in *Aspergillus oryzae* of the aphidicolin **106** biosynthetic pathway (Scheme 6).¹⁷² In addition to the bifunctional diterpene synthase, this cluster encompasses two P450 enzymes that carry out three hydroxylations of the diterpenoid alcohol scaffold, as well as a transporter and the aforementioned GGPP **5** synthase. The two CPS/KS homologs PaDC1 phyllocladen-16 α -ol **82** synthase) and PaDC2 ((+)-CDP synthase) characterized from *P. amygdali* are each part of

separate gene clusters that are involved in the biosynthesis of phyllocladan-11 α ,16 α ,18-triol **107**.¹⁵¹ Interestingly, in addition to different types of oxygenases, each of two clusters includes its own GGPP synthase.

Genome mining recently led to the identification of an *ent*-primara-8(14),15-diene **81** synthase AN1495 in *Aspergillus nidulans* which is part of a gene cluster that has yet to be characterized.¹⁵⁰ Encoded within this cluster are not only a GGPP synthase, but also a HMG-CoA synthase, a mevalonate pathway specific enzyme, which may boost precursor supply for diterpenoid biosynthesis by this pathway.

The monofunctional diterpene synthase, fusicocca-2,10(14)diene **85** synthase (PaFS), from *P. amygdali* is part of a five gene cluster,¹⁵⁷ including three enzymes that catalyze oxidative modifications to the tricyclic diterpene scaffold.¹⁷³ This cluster together with a later discovered second, nine-gene cluster are required to synthesize the bioactive fusicocin A **108** diterpenoid.¹⁷⁴ The presence of a prenyltransferase previously shown to catalyze the prenylation of glucose¹⁷⁵ which is a moiety also present in fusicocin A **108**, led to the identification of this second fusicocin cluster in the draft genome sequence of this fungus. Biochemical studies of recombinant enzymes together with gene deletions were carried to complete characterization of fusicocin biosynthesis.¹⁷⁴ The same group also identified and dissected another fusicocadiene-type pathway for brassicene C **109** biosynthesis in *A. brassicola*; as seen with the other diterpene pathways, a contingent of oxidizing enzymes (five cytochrome P450 enzymes, an oxidoreductase and a dioxygenase) are involved in modifying the terpene hydrocarbon scaffold.^{158, 173, 176}

4 Fungal triterpenoids

4.1 Oxidosqualene/lanosterol cyclases

The triterpenoids are C30 prenyl chain derived compounds that are found widely in nature as steroids and sterols.¹⁷⁷ The biosynthetic pathway to ergosterol **110**, the major component of the plasma membrane in fungi, begins with the cyclization of squalene **111** to lanosterol **112**.¹⁷⁸ Epoxidation of squalene **111** yields 2,3-oxidosqualene **113** which adopts a *chair-boat-chair* conformation.¹⁷⁹ Cyclization of the linear polyisoprene chain is catalyzed by a Class II type terpene synthase oxidosqualene/lanosterol cyclase (OSC) as part of a protonation-dependent reaction.^{159, 180} Activation of the epoxide in **113** is initiated by a conserved Asp, and a sequential ring-forming cascade results in a 6-6-6-5 tetracyclic protosteryl cation **114**. Two methyl migrations and hydride shifts, followed by a final deprotonation, results in lanosterol **112** (Scheme 12), the last common intermediate to the lanostane triterpenoids.^{140, 181, 182}

Prokaryotic and eukaryotic triterpene cyclases share a similar overall protein fold, with two α -helical domains linked by a membrane spanning channel. The large aromatic-lined active site cavity spans the two domains, and the protonating group that initiates cyclization is located at the polar region at the top of the central channel.^{159, 180} Triterpene cyclases are also defined by conserved 16 amino acid long tandem repeats known as QW sequence motifs. These Gln and Trp residues provide a bonding

network between the α -helices, which stabilizes the enzyme during the exergonic reaction, allowing stringent control over product formation.¹⁸³ Beyond these structural similarities, bacterial squalene hopene cyclases (SHC) and eukaryotic OSC cyclases have diverged both in sequence and in mechanism. Notably, SHC is characterized by the presence of the protonating Asp in a conserved DXDD motif, while the protonating Asp of OSC is located in the motif XXDCX.¹⁸² This branching in the evolutionary tree of SHC and OSC has resulted in the mechanistically defining feature of the enzymes: SHC accepts squalene **111** as a substrate, while OSC accepts 2,3-oxidosqualene **113** as a substrate.^{182, 184}

The first eukaryotic OSC to be cloned and characterized was the ergosterol **110** biosynthetic pathway enzyme ERG7 from *Saccharomyces cerevisiae*.^{185, 186} The catalytic mechanism of ERG7 has since been elucidated by mutational studies. Substrate activation is highly dependent upon the maintenance of a protonating environment, and aromatic residues play a key role in the stabilization of carbocation intermediates, which funnels the product trajectory of ERG7 specifically towards lanosterol **112**.¹⁸⁷⁻¹⁹⁰ Further fungal OSCs have been characterized from diverse species, including the helvolic acid **115** antibiotic producers *Cephalosporium caerulens*¹⁹¹ and *Aspergillus fumigatus*,¹⁹² as well as the cytotoxic ganoderic acid (e.g. **116**) producer *Ganoderma lucidum*.^{193, 194} Notably, characterization of an oxidosqualene-clavainone cyclase from *Hypholoma sublateritium* revealed the presence of two OSCs: one specifically for primary and one dedicated to secondary metabolism for clavatic acid **117** production.¹⁹⁵

More recently, bioinformatic approaches led to a wider scale identification of putative OSCs from Ascomycota and Basidiomycota.¹⁹⁶ This database represents a diversity of OSC sequences that could potentially be used to engineer a product promiscuous fungal OSC, offering biosynthetic routes to a greater diversity of fungal triterpenoids, like those produced by plant OSCs.¹⁹⁷⁻¹⁹⁹

4.2 Discovery of triterpenoid biosynthetic gene clusters

Triterpenoid biosynthesis is relatively well understood and characterized in plants.^{197, 200-202} Comparatively little, however, is known about fungal triterpenoid biosynthesis; despite the isolation of a large number of bioactive fungal triterpenoids mostly from Basidiomycota.^{203, 204}

Recent genome sequencing and transcriptome projects have provided the first insights into the biosynthetic pathways to the triterpenoids.^{24, 205-207} *G. lucidum* is a prolific producer of bioactive ganoderic acid **116** triterpenoids, correspondingly it has an OSC and a large complement of cytochrome P450s encoded within its genome. Almost half of the genes encoding P450s are upregulated during the transition from primordia to fruiting bodies, which correlates with lanosterol **112** expression and triterpenoid production. Therefore, developmental stage of fungal growth plays an important role in regulation of secondary metabolite genes. Some of these same genes are physically clustered on the genome, albeit not with the gene encoding OSC.²⁴ Likewise, transcriptome analysis of the medicinally

relevant *Wolfiporia cocos* identified several P450s that were upregulated simultaneously with its OSC.²⁰⁵ Contrastingly, the genome of *A. fumigatus* shows that it contains multiple genes encoding OSC as opposed to a single copy. Furthermore, the putative enzymes related to helvolic acid **115** production are located in a gene cluster encoding protostadienol **118** synthase, multiple P450s, a reductase, acyltransferases and a dehydrogenase. Together, these enzymes are predicted to represent the full helvolic acid **115** biosynthetic pathway.¹⁹²

5 Mining the fungal (sesqui)terpenome

5.1 Terpene synthase identification

Numerous databases exist to search the rapidly increasing number of fungal genomes for biosynthetic gene candidates, and for predicting putative coding sequences. For example, the Joint Genome Institute (JGI) listed 362 fungal genomes (117 Basidiomycota) at the time of writing, with a goal of 1000 sequenced genomes from all families in the fungal kingdom in the next few years.⁷⁸ Users of the JGI web interface (<http://genome.jgi-psf.org>)²⁰⁸ benefit from standardized EST and/or RNAseq data available for many of the fungal genomes that are that are displayed along with gene annotations.

Putative terpene synthases can be identified by performing a Basic Local Alignment Search Tool (BLAST)²⁰⁹ search of target genome(s) using characterized terpene synthases sequences as query sequences. We have used BLAST searches of fungal genomes, combined with biochemical knowledge of the initial FPP **4** cyclization reaction (see Scheme 1) catalyzed by the *O. olearius* enzymes (Omp1-10, section 2.1) to develop a predictive framework for further, targeted discovery of additional fungal sesquiterpene synthases and their associated biosynthetic gene clusters.^{20,23}

We found that sesquiterpene synthase sequences identified by BLAST analysis in the sequenced Basidiomycota genomes (presently more than 1000 sequences in ~100 genomes) cluster in distinct clades according to initial cyclization mechanism (Figure 1).²³ This method led to the focused discovery and characterization of several 1,11-cyclizing terpene synthases, which produce *trans*-humulyl cation **60** derived sesquiterpenes (section 2.1.).^{20,23} Notably, these characterized Δ 6-protoilludene synthases form a single branch within the 1,11 cyclizing clade (Figure 1). Many other predicted *trans*-humulyl cation **60** producing sesquiterpene synthases remain yet to be discovered, potentially producing the diverse scaffolds shown in Scheme 7.

5.2 Identification of terpenoid biosynthetic gene clusters

Discovery of fungal terpenoid biosynthetic pathways by genome mining approaches is facilitated by the fact that fungi tend to cluster genes encoding biosynthetic pathways within their genome. Clusters can be identified by manually scanning the genome sequence \pm 20 kb upstream and downstream of predicted terpene synthase sequences e.g. identified using the strategy described in the previous section. BLAST searches of the NCBI protein database and the NCBI Conserved Domain Database (CDD) allows annotation of predicted genes surrounding the

terpene synthase. Typical enzyme classes and proteins expected in such clusters are cytochrome P450 monooxygenases, oxidoreductases and oxygenases, group transferases, transporters, transcription factors and enzymes required for isoprenoid precursor biosynthesis. The likely boundaries of such biosynthetic clusters are predicted by the presence of large gaps between genes (>10 kb), or the presence of several consecutive genes that are not likely to be associated with natural product biosynthesis.¹⁸

However, manual identification of biosynthetic gene clusters is subject to user interpretation, is extremely time consuming, and as a result is a significant bottleneck in the identification of putative biosynthetic pathways. Therefore, automated search algorithms are required for the large scale identification of fungal natural product pathways and several web-based analysis tools have been developed for this purpose (reviewed in:^{210,211}). When using these platforms, however, one has to keep in mind that these tools have often been trained for the identification of specific natural products gene clusters (e.g. non-ribosomal peptide (NRP) and polyketide synthase (PKS) pathways) common in certain groups of organisms (e.g. Ascomycota, bacteria). From our own experience we have found that these tools are not particularly adept at identifying terpenoid clusters in Basidiomycota. For example, antiSMASH 2.0²¹² identified four of the eleven manually identified putative terpene synthases in the genome of *O. olearius*.²³ Similarly, five out of eighteen manually predicted terpene synthase sequences were identified in *S. hirsutum*.²⁰ This and other tools also significantly underestimated biosynthetic gene cluster size compared to our manual annotations, probably due to the high degree of complexity and variability of fungal gene clusters, as well that fact that few Basidiomycota clusters have been fully characterized.²¹³ As more natural product pathways from Basidiomycota are characterized, these tools can be trained for more accurate, automated cluster prediction in these types of fungi.

5.3 Characterization of terpenoid biosynthetic gene clusters

Characterization of fungal terpenoid gene clusters identified in Ascomycota described throughout this contribution has relied on a combined approach of biochemical characterization of recombinant enzymes and specific gene deletions in the native producer host, as well as transfer of biosynthetic cluster genes into closely related fungal hosts for which genetic tools are available.²¹⁴⁻²¹⁶ In order to access the incredible diversity of Basidiomycota terpenoid natural product pathways encoded in the large (and increasing) number of fungal genomes, different tools and methods need to be developed to go from genome mining to terpenoid biosynthesis. Some of the problems that need to be overcome for this to happen will be discussed briefly using the Basidiomycota as an example, but similar difficulties may also be encountered with other, less well-studied classes of fungi.

Several major challenges significantly impede efficient characterization of terpenoid biosynthetic gene clusters in Basidiomycota: 1. Correct cDNA prediction, 2. Therefore, necessity to clone genes from cDNA and thus the need to establish laboratory growth of strains under conditions where target genes/pathways are expressed, 3. Fungal strains are not

genetically tractable, and therefore require heterologous gene expression and pathway assembly.

The first step following the identification of putative terpene synthases and the associated gene cluster in a genome requires a refined prediction of the correct coding sequences for each gene. However, this is in our experience not trivial for Basidiomycota, which tend to have very intron-rich genomes.²¹⁷ We have found several very small and unpredicted exons (sometimes only 11-18 bp in length²¹⁸ and introns/exons 27 and 28 bp in length^{23, 69}) in genes cloned from cDNA. Genes synthesized based on cDNA predictions are therefore often non-functional. Figure 2 illustrates different structural gene predictions, alternative splice variants and the functional cDNAs of sesquiterpene synthases cloned from *S. hirsutum*. The gene prediction program Augustus (<http://bioinf.uni-greifswald.de/augustus/>)^{219, 220} can be used to identify the most likely cDNA sequences for a biosynthetic enzyme. Gene predictions created from the putative terpene synthase ± 10 kb can then be used to manually identify potential gene start/stop codons by alignment in MEGA6.²²¹ This allows the design of a small set of cloning primers to attempt amplification of the correct ORF from cDNA. In addition, the program FGENESH (<http://linux1.softberry.com/>) by SoftBerry (Mount Kisco, NY) offers multiple methods to predict gene models.²²² However, these programs have been trained to a small set of fungal strains and often do not correctly predict the less common splicing pattern present in Basidiomycota.

Collection of fungal tissue sufficient for cDNA isolation is often a significant barrier, as many fungi, and especially many Basidiomycota, are difficult or impossible to culture in the laboratory and may be not available from field collections. Also, the target genes for cloning must be expressed at the time of cDNA preparation. Fungal gene expression is highly complex, subject to RNAi silencing and influenced by trans-acting elements of the genome architecture.^{223, 224} Therefore, a large number of biosynthetic gene clusters remain silent.¹⁹ Strategies have been developed to activate fungal natural product pathways including modification of culture conditions, co-culture,²²⁵ and epigenetic manipulation, typically in high-throughput assays to produce novel fungal compounds for isolation.^{19, 226-229} However, the conditions required to express specific genes are generally unknown, and cannot be predicted with confidence.

Most Basidiomycota are not genetically tractable and no suitable tractable Basidiomycota model systems are available that could correctly splice and functionally express heterologous biosynthetic gene clusters cloned from other members of this fungal division. Hence, characterization of terpenoid natural product pathways identified in these fungi has to rely on the tedious assembly method of one gene at a time, in alternative hosts, such as yeast. Successful expression of pathways in these hosts can be challenging, as some enzymes e.g. cytochrome P450 monooxygenases, require a specific reductase for optimal activity²³⁰ and modified terpenoids may be toxic to the expression host, therefore complex cellular sequestration and secretion machinery could be required to prevent reduced cell growth.¹³⁴ In the future, additional development of genetic platforms specifically designed for natural product pathway expression in

Basidiomycota, such as *Ustilago maydis*, is needed,^{231, 232} similar to the advanced tools currently available for Ascomycota.²³³

6 Conclusions and outlook

Fungi have an enormous capacity for terpenoid natural product biosynthesis, and therefore represent an incredible resource for the discovery of new biosynthetic pathways. Identification and characterization of some of the pathways responsible for terpenoid production has revealed a diversity in structure, chemistry and function, including new classes of enzymes. The increasing availability of fungal genome sequences is streamlining the identification of novel pathways and enzymes to a wide range of bioactive terpenoids. The majority of such biosynthetic studies have focused on Ascomycota and only recently have we begun to look into the terpenome of Basidiomycota. Genome surveys clearly show that we have barely begun to unlock some of this biosynthetic potential encoded in the relatively small number of fungi with sequenced genomes that represent just a tiny fraction of the fungal diversity (Figure 1).

Our ability to characterize the fungal terpenome is dependent on, and currently severely limited by, our capability to rapidly identify and subsequently biochemically characterize terpenoid biosynthetic genes and their diverse functions. Current approaches of first finding potential pathways and then using molecular biology strategies to tease out the functions of individual enzymes in a biosynthetic pathway are slow, cumbersome and inadequate in the face of rapidly increasing genomic information. The development of better bioinformatic and genetic tools for the mining and heterologous assembly of large natural product pathways will be absolutely essential to enable high-throughput pathway discovery. For this to occur, it will be important for the natural products community to adopt synthetic biology and modular pathway assembly principles along with suitable platform organisms for process automation. The ultimate goal for the development of such discovery pipelines will be the ability to quickly move from *in silico* pathway prediction to high-throughput biosynthetic pathway assembly, heterologous expression, and analytical profiling of products followed by bioactivity screening. Revitalization of the natural products drug discovery pipeline will critically depend on the implementation of such strategies.

7 Acknowledgements

The authors apologize to any colleagues whose work was not discussed due to space limitations. The authors' research in fungal terpenoid biosynthesis has been supported by the National Institutes of Health Grant GM080299 (to C.S-D.).

8 References

1. J. E. Galagan, M. R. Henn, L. J. Ma, C. A. Cuomo and B. Birren, *Genome Res.*, 2005, **15**, 1620-1631.

2. L. Tedersoo, T. W. May and M. E. Smith, *Mycorrhiza*, 2010, **20**, 217-263.
3. M. Blackwell, *Am. J. Bot.*, 2011, **98**, 426-438.
4. D. Bass and T. A. Richards, *Fungal Biol. Rev.*, 2011, **25**, 159-164.
5. M. J. Alves, I. C. Ferreira, J. Dias, V. Teixeira, A. Martins and M. Pintado, *Planta Med.*, 2012, **78**, 1707-1718.
6. M. J. Alves, I. C. Ferreira, A. Martins and M. Pintado, *J. Appl. Microbiol.*, 2012, **113**, 466-475.
7. B. M. Fraga, *Nat. Prod. Rep.*, 2012, **29**, 1334-1366.
8. J. Gershenzon and N. Dudareva, *Nat. Chem. Biol.*, 2007, **3**, 408-414.
9. W. R. Abraham, *Curr. Med. Chem.*, 2001, **8**, 583-606.
10. V. Elisashvili, *Int. J. Med. Mushrooms*, 2012, **14**, 211-239.
11. A. Evidente, A. Kornienko, A. Cimmino, A. Andolfi, F. Lefranc, V. Mathieu and R. Kiss, *Nat. Prod. Rep.*, 2014, **31**, 617-627.
12. M. Misiek and D. Hoffmeister, *Planta Med.*, 2007, **73**, 103-115.
13. S. P. Wasser, *Appl. Microbiol. Biotechnol.*, 2011, **89**, 1323-1332.
14. B. Z. Zaidman, M. Yassin, J. Mahajna and S. P. Wasser, *Appl. Microbiol. Biotechnol.*, 2005, **67**, 453-468.
15. R. Schobert, S. Knauer, S. Seibt and B. Biersack, *Curr. Med. Chem.*, 2011, **18**, 790-807.
16. S. Brase, A. Encinas, J. Keck and C. F. Nising, *Chem. Rev.*, 2009, **109**, 3903-3990.
17. J. D. Walton, *Fungal Genet. Biol.*, 2000, **30**, 167-171.
18. G. T. Wawrzyn, S. E. Bloch and C. Schmidt-Dannert, *Methods Enzymol.*, 2012, **515**, 83-105.
19. P. Wiemann and N. P. Keller, *J. Ind. Microbiol. Biotechnol.*, 2014, **41**, 301-313.
20. M. B. Quin, C. M. Flynn, G. T. Wawrzyn, S. Choudhary and C. Schmidt-Dannert, *ChemBioChem*, 2013, **14**, 2480-2491.
21. J. Shendure and E. Lieberman Aiden, *Nat. Biotechnol.*, 2012, **30**, 1084-1094.
22. D. Floudas, M. Binder, R. Riley, K. Barry, R. A. Blanchette, B. Henrissat, A. T. Martinez, R. Otillar, J. W. Spatafora, J. S. Yadav, A. Aerts, I. Benoit, A. Boyd, A. Carlson, A. Copeland, P. M. Coutinho, R. P. de Vries, P. Ferreira, K. Findley, B. Foster, J. Gaskell, D. Glotzer, P. Gorecki, J. Heitman, C. Hesse, C. Hori, K. Igarashi, J. A. Jurgens, N. Kallen, P. Kersten, A. Kohler, U. Kues, T. K. Kumar, A. Kuo, K. LaButti, L. F. Larrondo, E. Lindquist, A. Ling, V. Lombard, S. Lucas, T. Lundell, R. Martin, D. J. McLaughlin, I. Morgenstern, E. Morin, C. Murat, L. G. Nagy, M. Nolan, R. A. Ohm, A. Patyshakuliyeva, A. Rokas, F. J. Ruiz-Duenas, G. Sabat, A. Salamov, M. Samejima, J. Schmutz, J. C. Slot, F. St John, J. Stenlid, H. Sun, S. Sun, K. Syed, A. Tsang, A. Wiebenga, D. Young, A. Pisabarro, D. C. Eastwood, F. Martin, D. Cullen, I. V. Grigoriev and D. S. Hibbett, *Science*, 2012, **336**, 1715-1719.
23. G. T. Wawrzyn, M. B. Quin, S. Choudhary, F. Lopez-Gallego and C. Schmidt-Dannert, *Chem. Biol.*, 2012, **19**, 772-783.
24. S. Chen, J. Xu, C. Liu, Y. Zhu, D. R. Nelson, S. Zhou, C. Li, L. Wang, X. Guo, Y. Sun, H. Luo, Y. Li, J. Song, B. Henrissat, A. Levasseur, J. Qian, J. Li, X. Luo, L. Shi, L. He, L. Xiang, X. Xu, Y. Niu, Q. Li, M. V. Han, H. Yan, J. Zhang, H. Chen, A. Lv, Z. Wang, M. Liu, D. C. Schwartz and C. Sun, *Nature Communications*, 2012, **3**, 913.
25. K. E. Bushley, R. Raja, P. Jaiswal, J. S. Cumbie, M. Nonogaki, A. E. Boyd, C. A. Owensby, B. J. Knaus, J. Elser, D. Miller, Y. Di, K. L. McPhail and J. W. Spatafora, *PLoS Genet.*, 2013, **9**, e1003496.
26. D. W. Christianson, *Curr. Opin. Chem. Biol.*, 2008, **12**, 141-150.
27. H. M. Miziorko, *Arch. Biochem. Biophys.*, 2011, **505**, 131-143.
28. D. W. Christianson, *Chem. Rev.*, 2006, **106**, 3412-3442.
29. K. U. Wendt and G. E. Schulz, *Structure*, 1998, **6**, 127-133.
30. E. Oldfield and F. Y. Lin, *Angew. Chem. Int. Ed. Engl.*, 2012, **51**, 1124-1137.
31. J. S. Dickschat, *Nat. Prod. Rep.*, 2011, **28**, 1917-1936.
32. D. J. Tantillo, *Nat. Prod. Rep.*, 2011, **28**, 1956-1956.
33. D. J. Miller and R. K. Allemann, *Nat. Prod. Rep.*, 2012, **29**, 60-71.
34. C. A. Lesburg, G. Zhai, D. E. Cane and D. W. Christianson, *Science*, 1997, **277**, 1820-1824.
35. C. M. Starks, K. W. Back, J. Chappell and J. P. Noel, *Science*, 1997, **277**, 1815-1820.
36. D. E. Cane, Q. Xue and B. C. Fitzsimmons, *Biochemistry*, 1996, **35**, 12369-12376.
37. D. E. Cane, H. J. Ha, C. Pargellis, F. Waldmeier, S. Swanson and P. P. N. Murthy, *Bioorg. Chem.*, 1985, **13**, 246-265.
38. D. E. Cane, S. Swanson and P. P. N. Murthy, *J. Am. Chem. Soc.*, 1981, **103**, 2136-2138.
39. P. Baer, P. Rabe, C. A. Citron, C. C. de Oliveira Mann, N. Kaufmann, M. Groll and J. S. Dickschat, *ChemBioChem*, 2014, **15**, 213-216.
40. P. Rabe, C. A. Citron and J. S. Dickschat, *ChemBioChem*, 2013, **14**, 2345-2354.
41. S. Garms, T. G. Kollner and W. Boland, *J. Org. Chem.*, 2010, **75**, 5590-5600.
42. L. S. Vedula, J. Jiang, T. Zakharian, D. E. Cane and D. W. Christianson, *Arch. Biochem. Biophys.*, 2008, **469**, 184-194.
43. B. Felicetti and D. E. Cane, *J. Am. Chem. Soc.*, 2004, **126**, 7212-7221.
44. M. Seemann, G. Zhai, J. W. de Kraker, C. M. Paschall, D. W. Christianson and D. E. Cane, *J. Am. Chem. Soc.*, 2002, **124**, 7681-7689.
45. J. A. Faraldos, V. Gonzalez and R. K. Allemann, *Chem. Commun.*, 2012, **48**, 3230-3232.
46. M. Chen, N. Al-lami, M. Janvier, E. L. D'Antonio, J. A. Faraldos, D. E. Cane, R. K. Allemann and D. W. Christianson, *Biochemistry*, 2013, **52**, 5441-5453.
47. J. A. Aaron, X. Lin, D. E. Cane and D. W. Christianson, *Biochemistry*, 2010, **49**, 1787-1797.
48. J. M. Caruthers, I. Kang, M. J. Rynkiewicz, D. E. Cane and D. W. Christianson, *J Biol Chem*, 2000, **275**, 25533-25539.

49. H. A. Gennadios, V. Gonzalez, L. Di Costanzo, A. Li, F. Yu, D. J. Miller, R. K. Allemann and D. W. Christianson, *Biochemistry*, 2009, **48**, 6175-6183.
50. R. P. McAndrew, P. P. Peralta-Yahya, A. DeGiovanni, J. H. Pereira, M. Z. Hadi, J. D. Keasling and P. D. Adams, *Structure*, 2011, **19**, 1876-1884.
51. M. J. Rynkiewicz, D. E. Cane and D. W. Christianson, *Proc. Natl. Acad. Sci. USA*, 2001, **98**, 13543-13548.
52. E. Y. Shishova, L. Di Costanzo, D. E. Cane and D. W. Christianson, *Biochemistry*, 2007, **46**, 1941-1951.
53. E. Y. Shishova, F. Yu, D. J. Miller, J. A. Faraldos, Y. Zhao, R. M. Coates, R. K. Allemann, D. E. Cane and D. W. Christianson, *J. Biol. Chem.*, 2008, **283**, 15431-15439.
54. M. J. Rynkiewicz, D. E. Cane and D. W. Christianson, *Biochemistry*, 2002, **41**, 1731-1741.
55. L. S. Vedula, D. E. Cane and D. W. Christianson, *Biochemistry*, 2005, **44**, 12719-12727.
56. T. M. Hohn and F. Vanmiddlesworth, *Arch. Biochem. Biophys.*, 1986, **251**, 756-761.
57. L. E. Osborne and J. M. Stein, *Int. J. Food Microbiol.*, 2007, **119**, 103-108.
58. L. S. Vedula, Y. Zhao, R. M. Coates, T. Koyama, D. E. Cane and D. W. Christianson, *Arch. Biochem. Biophys.*, 2007, **466**, 260-266.
59. D. E. Cane and H. J. Ha, *J. Am. Chem. Soc.*, 1988, **110**, 6865-6870.
60. J. S. Dickschat, N. L. Brock, C. A. Citron and B. Tudzynski, *ChemBioChem*, 2011, **12**, 2088-2095.
61. Y. J. Hong and D. J. Tantillo, *Org. Lett.*, 2006, **8**, 4601-4604.
62. D. J. Miller, J. Gao, D. G. Truhlar, N. J. Young, V. Gonzalez and R. K. Allemann, *Org. Biomol. Chem.*, 2008, **6**, 2346-2354.
63. J. A. Faraldos, V. Gonzalez, M. Senske and R. K. Allemann, *Org. Biomol. Chem.*, 2011, **9**, 6920-6923.
64. K. Motohashi, J. Hashimoto, S. Inaba, S. T. Khan, H. Komaki, A. Nagai, M. Takagi and K. Shin-ya, *J. Antibiot. (Tokyo)*, 2009, **62**, 247-250.
65. S. P. McCormick, N. J. Alexander and L. J. Harris, *Appl. Environ. Microbiol.*, 2010, **76**, 136-141.
66. N. L. Brock, K. Huss, B. Tudzynski and J. S. Dickschat, *ChemBioChem*, 2013, **14**, 311-315.
67. C. Pinedo, C. M. Wang, J. M. Pradier, B. Dalmais, M. Choquer, P. Le Pecheur, G. Morgant, I. G. Collado, D. E. Cane and M. Viaud, *ACS Chem. Biol.*, 2008, **3**, 791-801.
68. C. M. Wang, R. Hopson, X. Lin and D. E. Cane, *J. Am. Chem. Soc.*, 2009, **131**, 8360-8361.
69. S. Agger, F. Lopez-Gallego and C. Schmidt-Dannert, *Mol. Microbiol.*, 2009, **72**, 1181-1195.
70. F. Lopez-Gallego, S. A. Agger, D. Abate-Pella, M. D. Distefano and C. Schmidt-Dannert, *ChemBioChem*, 2010, **11**, 1093-1106.
71. F. Lopez-Gallego, G. T. Wawrzyn and C. Schmidt-Dannert, *Appl. Environ. Microbiol.*, 2010, **76**, 7723-7733.
72. R. Schobert, S. Seibt, K. Mahal, A. Ahmad, B. Biersack, K. Effenberger-Neidnicht, S. Padhye, F. H. Sarkar and T. Mueller, *J. Med. Chem.*, 2011, **54**, 6177-6182.
73. M. J. Kelner, T. C. McMorris, W. T. Beck, J. M. Zamora and R. Taetle, *Cancer Res.*, 1987, **47**, 3186-3189.
74. T. C. McMorris and M. Anchel, *J. Am. Chem. Soc.*, 1965, **87**, 1594-1600.
75. M. S. Nair and M. Anchel, *Tetrahedron Lett.*, 1972, **13**, 2753-2754.
76. M. B. Quin, G. Wawrzyn and C. Schmidt-Dannert, *Acta Crystallogr. Sect. F*, 2013, **69**, 574-577.
77. B. Engels, U. Heinig, T. Grothe, M. Stadler and S. Jennewein, *J. Biol. Chem.*, 2011, **286**, 6871-6878.
78. F. Martin, D. Cullen, D. Hibbett, A. Pisabarro, J. W. Spatafora, S. E. Baker and I. V. Grigoriev, *New Phytol.*, 2011, **190**, 818-821.
79. C. A. Cuomo and B. W. Birren, *Methods Enzymol.*, 2010, **470**, 833-855.
80. W. A. Ayer and M. H. Saeedighomi, *Tetrahedron Lett.*, 1981, **22**, 2071-2074.
81. C. Abell and A. P. Leech, *Tetrahedron Lett.*, 1988, **29**, 4337-4340.
82. A. M. Ainsworth, A. D. M. Rayner, S. J. Broxholme, J. R. Beeching, J. A. Pryke, P. R. Scard, J. Berriman, K. A. Powell, A. J. Floyd and S. K. Branch, *Mycol. Res.*, 1990, **94**, 799-809.
83. B. S. Yun, I. K. Lee, Y. Cho, S. M. Cho and I. D. Yoo, *J. Nat. Prod.*, 2002, **65**, 786-788.
84. Y. H. Kim, B. S. Yun, I. J. Ryoo, J. P. Kim, H. Koshino and I. D. Yoo, *J. Antibiot.*, 2006, **59**, 432-434.
85. N.-H. Yoo, J.-P. Kim, B.-S. Yun, I.-J. Ryoo, I.-K. Lee, E.-S. Yoon, H. Koshino and I.-D. Yoo, *J. Antibiot.*, 2006, **59**, 110-113.
86. G. H. Li, M. Duan, Z. F. Yu, L. Li, J. Y. Dong, X. B. Wang, J. W. Guo, R. Huang, M. Wang and K. Q. Zhang, *Phytochemistry*, 2008, **69**, 1439-1445.
87. T. Opatz, H. Kolshorn and H. Anke, *J. Antibiot.*, 2008, **61**, 563-567.
88. J. C. Liermann, A. Schuffler, B. Wollinsky, J. Birnbacher, H. Kolshorn, T. Anke and T. Opatz, *J. Org. Chem.*, 2010, **75**, 2955-2961.
89. M. Isaka, U. Srisanoh, W. Choowong and T. Boonpratuang, *Org. Lett.*, 2011, **13**, 4886-4889.
90. Y. K. Rao, A. T. Wu, M. Geethangili, M. T. Huang, W. J. Chao, C. H. Wu, W. P. Deng, C. T. Yeh and Y. M. Tzeng, *Chem. Res. Toxicol.*, 2011, **24**, 238-245.
91. M. Isaka, U. Srisanoh, M. Sappan, S. Supothina and T. Boonpratuang, *Phytochemistry*, 2012, **79**, 116-120.
92. X. Zheng, G. H. Li, M. J. Xie, X. Wang, R. Sun, H. Lu and K. Q. Zhang, *Phytochemistry*, 2013, **86**, 144-150.
93. K. Ma, L. Bao, J. Han, T. Jin, X. Yang, F. Zhao, S. Li, F. Song, M. Liu and H. Liu, *Food Chem.*, 2014, **143**, 239-245.
94. L. Zu, M. Xu, M. W. Lodewyk, D. E. Cane, R. J. Peters and D. J. Tantillo, *J. Am. Chem. Soc.*, 2012, **134**, 11369-11371.
95. B. M. Fraga, *Nat. Prod. Rep.*, 2013, **30**, 1226-1264.
96. G. Erkel and T. Anke, in *Biotechnology: Products of Secondary Metabolism*, eds. H.-J. Rehm and G. Reed, Wiley, 2 edn., 1997, pp. 490-533.

97. R. Kramer and W.-R. Abraham, *Phytochem. Rev.*, 2011, **11**, 15-37.
98. B. J. M. Jansen and A. de Groot, *Nat. Prod. Rep.*, 2004, **21**, 449-477.
99. D.-Q. Luo, Y. Gao, J.-M. Gao, F. Wang, X.-L. Yang and J.-K. Liu, *J. Nat. Prod.*, 2006, **69**, 1354-1357.
100. B. Simon, T. Anke, U. Anders, M. Neuhaus and E. Hansske, *Zeitschrift Fur Naturforschung C-a Journal of Biosciences*, 1995, **50**, 173-180.
101. R. M. Coates, J. Z. Ho, M. Klobus and L. J. Zhu, *J. Org. Chem.*, 1998, **63**, 9166-9176.
102. W. A. Ayer and E. R. Cruz, *J. Org. Chem.*, 1993, **58**, 7529-7534.
103. E. R. Cruz, *Can. J. Chem.*, 1997, **75**, 834-839.
104. W. R. Abraham, L. Ernst, L. Witte, H. P. Hanssen and E. Sprecher, *Tetrahedron*, 1986, **42**, 4475-4480.
105. T. C. Feline, G. Mellows, R. B. Jones and L. Phillips, *J. Chem. Soc., Chem. Commun.*, 1974, **2**, 63-64.
106. P. S. Covello, K. H. Teoh, D. R. Polichuk, D. W. Reed and G. Nowak, *Phytochemistry*, 2007, **68**, 1864-1871.
107. T. Tokai, H. Koshino, N. Takahashi-Ando, M. Sato, M. Fujimura and M. Kimura, *Biochem. Biophys. Res. Commun.*, 2007, **353**, 412-417.
108. W. M. Daniewski and G. Vidari, in *Constituents of Lactarius (mushrooms)*, Springer-Verlag Wien, 1999, pp. 69-171.
109. A. Schuffler, B. Wollinsky, T. Anke, J. C. Liermann and T. Opatz, *J. Nat. Prod.*, 2012, **75**, 1405-1408.
110. Y. Wang, L. Bao, D. Liu, X. Yang, S. Li, H. Gao, X. Yao, H. Wen and H. Liu, *Tetrahedron*, 2012, **68**, 3012-3018.
111. O. Sterner, T. Anke, W. S. Sheldrick and W. Steglich, *Tetrahedron*, 1990, **46**, 2389-2400.
112. G. Lackner, M. Bohnert, J. Wick and D. Hoffmeister, *Chem. Biol.*, 2013, **20**, 1101-1106.
113. T. Opatz, H. Kolshorn, J. Birnbacher, A. Schuffler, F. Deininger and T. Anke, *Eur. J. Org. Chem.*, 2007, 5546-5550.
114. J. F. Grove, *Progr. Chem. Org. Nat. Prod.*, 2007, **88**, 63-130.
115. S. P. McCormick, A. M. Stanley, N. A. Stover and N. J. Alexander, *Toxins (Basel)*, 2011, **3**, 802-814.
116. A. E. Desjardins, *J. Agric. Food Chem.*, 2009, **57**, 4478-4484.
117. M. Kimura, T. Tokai, N. Takahashi-Ando, S. Ohsato and M. Fujimura, *Biosci., Biotechnol., Biochem.*, 2007, **71**, 2105-2123.
118. K. Kazan, D. M. Gardiner and J. M. Manners, *Mol. Plant Pathol.*, 2012, **13**, 399-413.
119. K. Audenaert, A. Vanheule, M. Hofte and G. Haesaert, *Toxins (Basel)*, 2014, **6**, 1-19.
120. Q. Gu, C. Zhang, X. Liu and Z. Ma, *Mol. Plant Pathol.*, 2014.
121. T. M. Hohn and P. D. Beremand, *Gene*, 1989, **79**, 131-138.
122. D. W. Brown, S. P. McCormick, N. J. Alexander, R. H. Proctor and A. E. Desjardins, *Fungal Genet. Biol.*, 2002, **36**, 224-233.
123. C. G. Nasmith, S. Walkowiak, L. Wang, W. W. Leung, Y. Gong, A. Johnston, L. J. Harris, D. S. Guttman and R. Subramaniam, *PLoS Path.*, 2011, **7**, e1002266.
124. S. P. McCormick, N. J. Alexander and R. H. Proctor, *Can. J. Microbiol.*, 2006, **52**, 636-642.
125. A. R. Hesketh, B. W. Bycroft, P. M. Dewick and J. Gilbert, *Phytochemistry*, 1993, **32**, 105-116.
126. S. P. McCormick, N. J. Alexander, S. E. Trapp and T. M. Hohn, *Appl. Environ. Microbiol.*, 1999, **65**, 5252-5256.
127. S. Ohsato, T. Ochiai-Fukuda, T. Nishiuchi, N. Takahashi-Ando, S. Koizumi, H. Hamamoto, T. Kudo, I. Yamaguchi and M. Kimura, *Plant Cell Rep.*, 2007, **26**, 531-538.
128. N. J. Alexander, T. M. Hohn and S. P. McCormick, *Appl. Environ. Microbiol.*, 1998, **64**, 221-225.
129. S. P. McCormick, T. M. Hohn and A. E. Desjardins, *Appl. Environ. Microbiol.*, 1996, **62**, 353-359.
130. D. W. Brown, R. B. Dyer, S. P. McCormick, D. F. Kendra and R. D. Plattner, *Fungal Genet. Biol.*, 2004, **41**, 454-462.
131. N. J. Alexander, S. P. McCormick, C. Waalwijk, T. van der Lee and R. H. Proctor, *Fungal Genet. Biol.*, 2011, **48**, 485-495.
132. N. J. Alexander, S. P. McCormick and T. M. Hohn, *Mol. Genet. Genomics*, 1999, **261**, 977-984.
133. J. Menke, Y. Dong and H. C. Kistler, *Mol. Plant-Microbe Interact.*, 2012, **25**, 1408-1418.
134. J. Menke, J. Weber, K. Broz and H. C. Kistler, *PLoS One*, 2013, **8**, e63077.
135. A. Chanda, L. V. Roze, S. Kang, K. A. Artymovich, G. R. Hicks, N. V. Raikhel, A. M. Calvo and J. E. Linz, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 19533-19538.
136. A. Chanda, L. V. Roze and J. E. Linz, *Eukaryot. Cell*, 2010, **9**, 1724-1727.
137. P. I. Hidalgo, R. V. Ullan, S. M. Albillos, O. Montero, M. A. Fernandez-Bodega, C. Garcia-Estrada, M. Fernandez-Aguado and J. F. Martin, *Fungal Genet. Biol.*, 2014, **62**, 11-24.
138. J. Zi, S. Mafu and R. J. Peters, *Annu. Rev. Plant Biol.*, 2014, **65**.
139. R. J. Peters, *Nat. Prod. Rep.*, 2010, **27**, 1521-1530.
140. I. Abe, *Nat. Prod. Rep.*, 2007, **24**, 1311-1331.
141. H. Kawaide, R. Imai, T. Sassa and Y. Kamiya, *J. Biol. Chem.*, 1997, **272**, 21706-21712.
142. H. Kawaide, T. Sassa and Y. Kamiya, *J. Biol. Chem.*, 2000, **275**, 2276-2280.
143. T. P. Sun and Y. Kamiya, *Plant Cell*, 1994, **6**, 1509-1518.
144. S. Yamaguchi, T. Saito, H. Abe, H. Yamane, N. Murofushi and Y. Kamiya, *Plant J.*, 1996, **10**, 203-213.
145. M. Koksall, H. Hu, R. M. Coates, R. J. Peters and D. W. Christianson, *Nat. Chem. Biol.*, 2011, **7**, 431-433.

146. M. Koksai, Y. Jin, R. M. Coates, R. Croteau and D. W. Christianson, *Nature*, 2011, **469**, 116-120.
147. K. Zhou, Y. Gao, J. A. Hoy, F. M. Mann, R. B. Honzatko and R. J. Peters, *J. Biol. Chem.*, 2012, **287**, 6840-6850.
148. B. Tudzynski and K. Holter, *Fungal Genet. Biol.*, 1998, **25**, 157-170.
149. B. Tudzynski, H. Kawaide and Y. Kamiya, *Curr. Genet.*, 1998, **34**, 234-240.
150. K. Bromann, M. Toivari, K. Viljanen, A. Vuoristo, L. Ruohonen and T. Nakari-Setälä, *PLoS One*, 2012, **7**, e35450.
151. T. Toyomasu, R. Niida, H. Kenmoku, Y. Kanno, S. Miura, C. Nakano, Y. Shiono, W. Mitsuhashi, H. Toshima, H. Oikawa, T. Hoshino, T. Dairi, N. Kato and T. Sassa, *Biosci. Biotechnol. Biochem.*, 2008, **72**, 1038-1047.
152. H. Oikawa, T. Toyomasu, H. Toshima, S. Ohashi, H. Kawaide, Y. Kamiya, M. Ohtsuka, S. Shinoda, W. Mitsuhashi and T. Sassa, *J. Am. Chem. Soc.*, 2001, **123**, 5154-5155.
153. R. Mitterbauer and T. Specht, *WO/2011/110610*, 2011, **EP2011/053571**, WO2011110610A2011110611.
154. K. Mende, V. Homann and B. Tudzynski, *Mol. Genet. Genomics*, 1997, **255**, 96-105.
155. C. Bomke and B. Tudzynski, *Phytochemistry*, 2009, **70**, 1876-1893.
156. T. Toyomasu, A. Kaneko, T. Tokiwano, Y. Kanno, Y. Kanno, R. Niida, S. Miura, T. Nishioka, C. Ikeda, W. Mitsuhashi, T. Dairi, T. Kawano, H. Oikawa, N. Kato and T. Sassa, *J. Org. Chem.*, 2009, **74**, 1541-1548.
157. T. Toyomasu, M. Tsukahara, A. Kaneko, R. Niida, W. Mitsuhashi, T. Dairi, N. Kato and T. Sassa, *Proc. Natl. Acad. Sci. USA*, 2007, **104**, 3084-3088.
158. A. Minami, N. Tajima, Y. Higuchi, T. Toyomasu, T. Sassa, N. Kato and T. Dairi, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 870-874.
159. K. U. Wendt, K. Poralla and G. E. Schulz, *Science*, 1997, **277**, 1811-1815.
160. L. C. Tarshis, M. Yan, C. D. Poulter and J. C. Sacchettini, *Biochemistry*, 1994, **33**, 10871-10877.
161. R. Chiba, A. Minami, K. Gomi and H. Oikawa, *Org. Lett.*, 2013, **15**, 594-597.
162. M. J. Smanski, R. M. Peterson, S. X. Huang and B. Shen, *Curr. Opin. Chem. Biol.*, 2012, **16**, 132-141.
163. J. W. Shen, Y. Ruan and B. J. Ma, *J. Basic Microbiol.*, 2009, **49**, 242-255.
164. B. Tudzynski, *Appl. Microbiol. Biotechnol.*, 2005, **66**, 597-611.
165. P. Hedden, A. L. Phillips, M. C. Rojas, E. Carrera and B. Tudzynski, *J. Plant Growth Regul.*, 2001, **20**, 319-331.
166. B. Tudzynski, M. C. Rojas, P. Gaskin and P. Hedden, *J. Biol. Chem.*, 2002, **277**, 21246-21253.
167. B. Tudzynski, M. Mihlan, M. C. Rojas, P. Linnemannstons, P. Gaskin and P. Hedden, *J. Biol. Chem.*, 2003, **278**, 28635-28643.
168. S. Malonek, M. C. Rojas, P. Hedden, P. Gaskin, P. Hopkins and B. Tudzynski, *J. Biol. Chem.*, 2004, **279**, 25075-25084.
169. C. Bomke, M. C. Rojas, F. Gong, P. Hedden and B. Tudzynski, *Appl. Environ. Microbiol.*, 2008, **74**, 5325-5339.
170. H. Kawaide, *Biosci., Biotechnol., Biochem.*, 2006, **70**, 583-590.
171. H. Kawaide, T. Sassa and Y. Kamiya, *Phytochemistry*, 1995, **39**, 305-310.
172. R. Fujii, A. Minami, T. Tsukagoshi, N. Sato, T. Sahara, S. Ohgiya, K. Gomi and H. Oikawa, *Biosci. Biotechnol. Biochem.*, 2011, **75**, 1813-1817.
173. Y. Ono, A. Minami, M. Noike, Y. Higuchi, T. Toyomasu, T. Sassa, N. Kato and T. Dairi, *J. Am. Chem. Soc.*, 2011, **133**, 2548-2555.
174. M. Noike, Y. Ono, Y. Araki, R. Tanio, Y. Higuchi, H. Nitta, Y. Hamano, T. Toyomasu, T. Sassa, N. Kato and T. Dairi, *PLoS One*, 2012, **7**, e42090.
175. M. Noike, C. Liu, Y. Ono, Y. Hamano, T. Toyomasu, T. Sassa, N. Kato and T. Dairi, *ChemBioChem*, 2012, **13**, 566-573.
176. M. Hashimoto, Y. Higuchi, S. Takahashi, H. Osada, T. Sakaki, T. Toyomasu, T. Sassa, N. Kato and T. Dairi, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 5640-5643.
177. R. A. Hill and J. D. Connolly, *Nat. Prod. Rep.*, 2013, **30**, 1028-1065.
178. K. Kristan and T. L. Rizner, *J. Steroidal Biochem. Mol. Biol.*, 2012, **129**, 79-91.
179. M. Garaiova, V. Zambojova, Z. Simova, P. Griac and I. Hapala, *FEMS Yeast Res.*, 2014, **14**, 310-323.
180. R. Thoma, T. Schulz-Gaschg, B. D'Arcy, J. Benz, J. Aebi, H. Dehmlow, M. Hennig, M. Stihle and A. Ruf, *Nature*, 2004, **432**, 118-122.
181. E. J. Corey and S. C. Virgil, *J. Am. Chem. Soc.*, 1991, **113**, 4025-4026.
182. G. Siedenburg and D. Jendrossek, *Appl. Environ. Microbiol.*, 2011, **77**, 3905-3915.
183. K. Poralla, A. Hewelt, G. D. Prestwich, I. Abe, I. Reipen and G. Sprenger, *Trends Biochem. Sci.*, 1994, **19**, 157-158.
184. T. Frickey and E. Kannenberg, *Environ. Microbiol.*, 2009, **11**, 1224-1241.
185. E. J. Corey, S. P. T. Matsuda and B. Bartel, *Proc. Natl. Acad. Sci. USA*, 1994, **91**, 2211-2215.
186. Z. Shi, C. J. Buntel and J. H. Griffin, *Proc. Natl. Acad. Sci. USA*, 1994, **91**, 7370-7374.
187. E. J. Corey, H. Cheng, C. Hunter Baker, S. P. T. Matsuda, D. Li and X. D. Song, *J. Am. Chem. Soc.*, 1997, **119**, 1289-1296.
188. T. K. Wu, H. Y. Wen, C. H. Chang and Y. T. Liu, *Org. Lett.*, 2008, **10**, 2529-2532.
189. C. H. Chang, H. Y. Wen, W. S. Shie, C. T. Lu, M. E. Li, Y. T. Liu, W. H. Li and T. K. Wu, *Org. Biomol. Chem.*, 2013, **11**, 4214-4219.
190. T. K. Wu, M. T. Yu, Y. T. Liu, C. H. Chang, H. J. Wang and E. W. G. Diau, *Org. Lett.*, 2006, **8**, 1319-1322.


191. I. Abe, K. Naito, Y. Takagi and H. Noguchi, *Biochim. Biophys. Acta*, 2001, **1522**, 67-73.
192. H. Mitsuguchi, Y. Seshime, I. Fujii, M. Shibuya, Y. Ebizuka and T. Kushiro, *J. Am. Chem. Soc.*, 2009, **131**, 6402-6411.
193. C.-H. Shang, L. Shi, A. Ren, L. Qin and M.-W. Zhao, *Biosci., Biotechnol., Biochem.*, 2010, **74**, 974-978.
194. C. R. Cheng, Q. X. Yue, Z. Y. Wu, X. Y. Song, S. J. Tao, X. H. Wu, P. P. Xu, X. Liu, S. H. Guan and D. A. Guo, *Phytochemistry*, 2010, **71**, 1579-1585.
195. R. P. Godio and J. F. Martin, *Fungal Genet. Biol.*, 2009, **46**, 232-242.
196. S. Racolta, P. B. Juhl, D. Sirim and J. Pleiss, *Proteins*, 2012, **80**, 2009-2019.
197. D. R. Phillips, J. M. Rasbery, B. Bartel and S. P. Matsuda, *Curr. Opin. Plant Biol.*, 2006, **9**, 305-314.
198. Y. T. Liu, T. C. Hu, C. H. Chang, W. S. Shie and T. K. Wu, *Org. Lett.*, 2012, **14**, 5222-5225.
199. S. Lodeiro, Q. Xiong, W. K. Wilson, M. D. Kolesnikova, C. S. Onak and S. P. Matsuda, *J. Am. Chem. Soc.*, 2007, **129**, 11213-11222.
200. T. Moses, J. Pollier, J. M. Thevelein and A. Goossens, *New Phytol.*, 2013, **200**, 27-43.
201. S. Sawai and K. Saito, *Front Plant Sci*, 2011, **2**, 25.
202. Z. Xue, L. Duan, D. Liu, J. Guo, S. Ge, J. Dicks, O. M. P. A. Osbourn and X. Qi, *New Phytol.*, 2012, **193**, 1022-1038.
203. J. L. Rios, I. Andujar, M. C. Recio and R. M. Giner, *J. Nat. Prod.*, 2012, **75**, 2016-2044.
204. M. Zhao, T. Godecke, J. Gunn, J. A. Duan and C. T. Che, *Molecules*, 2013, **18**, 4054-4080.
205. S. Shu, B. Chen, M. Zhou, X. Zhao, H. Xia and M. Wang, *PLoS One*, 2013, **8**, e71350.
206. D. Liu, J. Gong, W. Dai, X. Kang, Z. Huang, H. M. Zhang, W. Liu, L. Liu, J. Ma, Z. Xia, Y. Chen, Y. Chen, D. Wang, P. Ni, A. Y. Guo and X. Xiong, *PLoS One*, 2012, **7**, e36146.
207. G. J. Yu, M. Wang, J. Huang, Y. L. Yin, Y. J. Chen, S. Jiang, Y. X. Jin, X. Q. Lan, B. H. Wong, Y. Liang and H. Sun, *PLoS One*, 2012, **7**, e44031.
208. H. Nordberg, M. Cantor, S. Dusheyko, S. Hua, A. Poliakov, I. Shabalov, T. Smirnova, I. V. Grigoriev and I. Dubchak, *Nucleic Acids Res.*, 2014, **42**, D26-31.
209. S. F. Altschul, W. Gish, W. Miller, E. W. Myers and D. J. Lipman, *J. Mol. Biol.*, 1990, **215**, 403-410.
210. N. Fedorova, V. Moktali and M. H. Medema, *Fungal Secondary Metabolism: Methods and Protocols*, Springer Science 2012.
211. T. Weber, *Int. J. Med. Microbiol.*, 2014, **304**, 230-235.
212. K. Blin, M. H. Medema, D. Kazempour, M. A. Fischbach, R. Breitling, E. Takano and T. Weber, *Nucleic Acids Res.*, 2013, **41**, W204-212.
213. P. Wiemann, C. J. Guo, J. M. Palmer, R. Sekonyela, C. C. Wang and N. P. Keller, *Proc. Natl. Acad. Sci. USA*, 2013, **110**, 17065-17070.
214. K. A. Pahirulzaman, K. Williams and C. M. Lazarus, *Methods Enzymol.*, 2012, **517**, 241-260.
215. K. Tagami, A. Minami, R. Fujii, C. Liu, M. Tanaka, K. Gomi, T. Dairi and H. Oikawa, *ChemBioChem*, 2014.
216. T. Itoh, K. Tokunaga, Y. Matsuda, I. Fujii, I. Abe, Y. Ebizuka and T. Kushiro, *Nat Chem*, 2010, **2**, 858-864.
217. J. L. Da Lage, M. Binder, A. Hua-Van, S. Janecek and D. Casane, *BMC Evol. Biol.*, 2013, **13**, 40.
218. C. M. Flynn, 2014.
219. K. J. Hoff and M. Stanke, *Nucleic Acids Res.*, 2013, **41**, W123-128.
220. M. Stanke, R. Steinkamp, S. Waack and B. Morgenstern, *Nucleic Acids Res.*, 2004, **32**, W309-312.
221. K. Tamura, G. Stecher, D. Peterson, A. Filipiński and S. Kumar, *Mol. Biol. Evol.*, 2013, **30**, 2725-2729.
222. V. Solovyev, P. Kosarev, I. Seledsov and D. Vorobyev, *Genome Biol.*, 2006, **7 Suppl 1**, S10.11-12.
223. H. Nakayashiki, *FEBS Lett.*, 2005, **579**, 5950-5957.
224. L. M. Noble and A. Andrianopoulos, *Genome Biol. Evol.*, 2013, **5**, 1336-1352.
225. S. Bertrand, N. Bohni, S. Schnee, O. Schumpp, K. Gindro and J. L. Wolfender, *Biotechnol. Adv.*, 2014, **epub ahead of print**.
226. Y. M. Chiang, S. L. Chang, B. R. Oakley and C. C. Wang, *Curr. Opin. Chem. Biol.*, 2011, **15**, 137-143.
227. R. H. Cichewicz, *Nat. Prod. Rep.*, 2010, **27**, 11-22.
228. L. Du, J. B. King, B. H. Morrow, J. K. Shen, A. N. Miller and R. H. Cichewicz, *J. Nat. Prod.*, 2012, **75**, 1819-1823.
229. A. A. Brakhage and V. Schroeckh, *Fungal Genet. Biol.*, 2011, **48**, 15-22.
230. C. J. Paddon, P. J. Westfall, D. J. Pitera, K. Benjamin, K. Fisher, D. McPhee, M. D. Leavell, A. Tai, A. Main, D. Eng, D. R. Polichuk, K. H. Teoh, D. W. Reed, T. Treynor, J. Lenihan, M. Fleck, S. Bajad, G. Dang, D. Dengrove, D. Diola, G. Dorin, K. W. Ellens, S. Fickes, J. Galazzo, S. P. Gaucher, T. Geistlinger, R. Henry, M. Hepp, T. Horning, T. Iqbal, H. Jiang, L. Kizer, B. Lieu, D. Melis, N. Moss, R. Regentin, S. Secrest, H. Tsuruta, R. Vazquez, L. F. Westblade, L. Xu, M. Yu, Y. Zhang, L. Zhao, J. Lievens, P. S. Covelio, J. D. Keasling, K. K. Reiling, N. S. Renninger and J. D. Newman, *Nature*, 2013, **496**, 528-532.
231. G. Steinberg and J. Perez-Martin, *Trends Cell Biol.*, 2008, **18**, 61-67.
232. M. Feldbrugge, R. Kellner and K. Schipper, *Appl. Microbiol. Biotechnol.*, 2013, **97**, 3253-3265.
233. J. Yaegashi, B. R. Oakley and C. C. Wang, *J. Ind. Microbiol. Biotechnol.*, 2014, **41**, 433-442.
234. K. Asahi, Y. Honma, K. Hazeki, T. Sassa, Y. Kubohara, A. Sakurai and N. Takahashi, *Biochem. Biophys. Res. Commun.*, 1997, **238**, 758-763.
235. T. Sassa, T. Ooi, M. Nukina, M. Ikeda and N. Kato, *Biosci., Biotechnol., Biochem.*, 1998, **62**, 1815-1818.
236. T. Sassa, M. Togashi and T. Kitaguchi, *Agric. Biol. Chem.*, 1975, **39**, 1735-1744.
237. T. Toyomasu, K. Nakaminami, H. Toshima, T. Mie, K. Watanabe, H. Ito, H. Matsui, W. Mitsuhashi, T. Sassa

- and H. Oikawa, *Biosci., Biotechnol., Biochem.*, 2004, **68**, 146-152.
238. Q.-X. Wang, Q.-Y. Qi, K. Wang, L. Li, L. Bao, J.-J. Han, M.-M. Liu, L.-X. Zhang, L. Cai and H.-W. Liu, *Org. Lett.*, 2013, **15**, 3982-3985.
239. S. Afiyatullo, T. A. Kuznetsova, V. V. Isakov, M. V. Pivkin, N. G. Prokofeva and G. B. Elyakov, *J. Nat. Prod.*, 2000, **63**, 848-850.
240. H. F. Sun, X. M. Li, L. Meng, C. M. Cui, S. S. Gao, C. S. Li, C. G. Huang and B. G. Wang, *J. Nat. Prod.*, 2012, **75**, 148-152.
241. Y. Feng, F. Ren, S. Niu, L. Wang, L. Li, X. Liu and Y. Che, *J. Nat. Prod.*, 2014, [Epub ahead of print].
242. E. M. Wijeratne, B. P. Bashyal, M. X. Liu, D. D. Rocha, G. M. Gunaherath, J. M. U'Ren, M. K. Gunatilaka, A. E. Arnold, L. Whitesell and A. A. Gunatilaka, *J. Nat. Prod.*, 2012, **75**, 361-369.
243. X. N. Wang, B. P. Bashyal, E. M. Wijeratne, J. M. U'Ren, M. X. Liu, M. K. Gunatilaka, A. E. Arnold and A. A. Gunatilaka, *J. Nat. Prod.*, 2011, **74**, 2052-2061.
244. A. J. Hartley, K. de Mattos-Shiple, C. M. Collins, S. Kilaru, G. D. Foster and A. M. Bailey, *FEMS Microbiol. Lett.*, 2009, **297**, 24-30.
245. H. Shibata, A. Irie and Y. Morita, *Biosci., Biotechnol., Biochem.*, 1998, **62**, 2450-2452.
246. C. Valdivia, M. Kettering, H. Anke, E. Thines and O. Sterner, *Tetrahedron*, 2005, **61**, 9527-9532.
247. Y. Liu, Y. Li, Y. Ou, S. Xiao, C. Lu, Z. Zheng and Y. Shen, *Bioorg. Med. Chem. Lett.*, 2012, **22**, 5059-5062.
248. Y. X. Ou, Y. Y. Li, X. M. Qian and Y. M. Shen, *Phytochemistry*, 2012, **78**, 190-196.
249. Y.-Y. Li and Y.-M. Shen, *Helv. Chim. Acta*, 2010, **93**, 2151-2157.
250. X. Mazur, U. Becker, T. Anke and O. Sterner, *Phytochemistry*, 1996, **43**, 405-407.


Table 1. Representative examples of fungal diterpenoids with characterized bioactivities. *Denotes an endophytic or endolichenic isolate. †Diterpenoid biosynthetic gene clusters described. Representative structures (in bold) of for each fungal species are shown in Schemes 5 and 6. For a comprehensive list of diterpenoids produced by Basidiomycota see review.¹⁶³

| Biological activity | Fungal species | Compound name(s) | Reference |
|----------------------|---|---|---------------|
| Ascomycota | | | |
| Cytotoxic | <i>Alternaria brassicola</i> | Brassicicene C (109) † | 158, 173, 176 |
| Cytotoxic | <i>Cladosporium sp.</i> | Cotylenin A-E (88) | 234-236 |
| Phytotoxin | <i>Phomopsis amygdali</i> | Fusicoccins A (108) †, F | 157 |
| DNA-Pol. inhibitor | <i>Phoma betae</i> | Aphidicolin (106) † | 152, 237 |
| Anti-inflammatory | <i>Bipolaris coicis</i> * | Coicenals A (89) -D | 238 |
| Cytotoxic | <i>Acremonium striatisporum</i> | Virescenosides A(90) , B, C, M, N | 239 |
| Cytotoxic | <i>Aspergillus wentii</i> EN-48* | Asperolides A (91) -C | 240 |
| Cytotoxic | <i>Cercospora sp.</i> * | Cercosporenes A (92) -F | 241 |
| Cytotoxic | <i>Geopyxis sp.</i> * | Geopyxins A (93) -D | 242 |
| Cytotoxic | <i>Smardaea sp.</i> * | Smardaesidins A (94) -G | 243 |
| Growth hormone | <i>Fusarium fujikuroi</i> | Gibberellins GA14 (102) , 4 (103) , 3, 1 (104) | 148 |
| Growth hormone | <i>Phaeosphaeria sp.</i> L487 | Gibberellins GA12 (101) , 4 (103) , 9, 20, 1 (104) | 141, 171 |
| Growth hormone | <i>Sphaceloma manihoticola</i> | Gibberellins GA14 (102) , 4 (103) | 169 |
| Basidiomycota | | | |
| Antibacterial | <i>Clitopilus sp.</i> & <i>C. passeckeranus</i> | Pleuromutilin (84) † | 153, 244 |
| Antibacterial | <i>Sarcodon scabrosus</i> | Sarcodonin L (95) , M | 245 |
| Antifungal | <i>Coprinus heptemerus</i> | Heptemerones A-G (96) | 246 |
| Cytotoxic | <i>Coprinus plicatilis</i> | Plicatilisins A (97) -D | 247 |
| Cytotoxic | <i>Coprinus radians</i> | Radianspene A (98) -M | 248 |
| Cytotoxic | <i>Crinipellis sp.</i> | Crinipellins A-C (99) , D | 249 |
| Cytotoxic | <i>Lepista sordida</i> | Lepistol (100) | 250 |


SCHEMES 1-12




Scheme 1: Overview of fungal terpene biosynthesis. The mevalonate pathway yields the C5 isoprene precursors dimethylallyl diphosphate (DMAPP) **1** and its isomer isopentenyl diphosphate (IPP) **2**. Head-to-tail condensation of DMAPP **1** with one, two or three IPP **2** units catalyzed by prenyldiphosphate synthases yields geranyl diphosphate (GPP) **3**, farnesyl diphosphate (FPP) **4** and geranylgeranyldiphosphate (GGPP) **5** that are cyclized by ionization-dependent class I or protonation-dependent class II terpene synthases into structurally diverse mono-, sesqui- or diterpenoids. Head-to-head condensation of two FPP **4** molecules yields squalene **111** as the precursor of triterpenoids. The cyclic scaffolds of diterpenoids are synthesized by either monofunctional terpene synthases that contain a class I catalytic domain fused to a prenyldiphosphate domain, or bifunctional terpene synthases that contain class I and class II catalytic domains (see also Scheme 9). Sesquiterpene synthases catalyze different ring-closures of 2E,6E-FPP **4** or upon isomerization, of (3R)-nerolidyl diphosphate (NPP) **6** to generate different sesquiterpenoids. At present, *bona fide* monoterpene synthases have not been isolated from fungi. CDP: Copolyldiphosphate. *: Ophiobolin F **87** is synthesized from the C25 diphosphate precursor farnesylgeranyl diphosphate.




Scheme 2. Examples of sesquiterpenoids generated by 1,6-cyclization of FPP upon isomerization to NPP. The cyclization of FPP 4 to trichodiene 8 by trichodiene synthase (TS) illustrates a well-studied cyclization mechanism. Other characterized fungal enzymes (see text) cyclize NPP into different major products.




Scheme 3. Examples of sesquiterpenoids generated by 1,10-cyclization of FPP. The cyclization of FPP to (+)-aristolochene 7 by aristolochene synthase (AS) illustrates a well-studied cyclization mechanism. Other characterized fungal enzymes (see text) cyclize FPP into different major products.




Scheme 4. Examples of sesquiterpenoids generated by 1,11-cyclization of FPP. The initial *trans*-humulyl cation 60 cyclization product can be cyclized into different sesquiterpenoid scaffolds. Known fungal enzymes (see text) that carry out these cyclization reactions are shown.




Scheme 5. Examples of sesquiterpenoids generated by 1,10-cyclization of FPP upon isomerization to NPP. The initial *Z,E*-germacradienyl cation cyclization product can be cyclized into different sesquiterpenoid scaffolds. Major products of known fungal enzymes (see text) are shown. Asterisk denotes that an alternative 1,10-cyclization pathway (Figure 10) is possible for δ -cadinene.




Scheme 6. Representative examples of modified sesquiterpenoids isolated from Basidiomycota. Compounds are organized based on the sesquiterpenoid scaffolds that they are derived from (see text for details).




Scheme 7. Major fungal transformations of the *trans*-humulyl cation. For consistency, carbon numbering is based on position in the linear FPP 4 precursor for all mechanism discussions. Where necessary for final products, published compound names are used. Enzymes that have been characterized as catalyzing transformations are listed beneath their major product. The key cation intermediates **59** and **60** are shown in blue, while minor intermediate cations are shown in red. Isolated final products are shown in black, and sesquiterpene scaffolds of known compound families are shown in orange. Rearrangements performed by sesquiterpene synthases are shown with black arrows, while reactions thought to be catalyzed by secondary enzymes are indicated by red arrows.




Scheme 8: Identified sesquiterpenoid biosynthetic pathways in Ascomycota. Biosynthesis of (A) the trichothecene DON 71, (B) PR-toxin 17 and (C) botrydial 22 by the fungal strains shown.




Scheme 9. Cyclization mechanism of the bifunctional CPS/KS type diterpene synthases. The precursor GGPP 5 undergoes an initial protonation dependent cyclization yielding (-)-ent-CDP 79 containing the typical bicyclic ring system of labdane-type diterpenoids. An ionization dependent dephosphorylation reaction and further cyclizations result in ent-kaurene 80.



Scheme 10. Examples of fungal diterpenoid natural products and their producer organisms. (A) Fusicoccadiene (top) and labdane-type (bottom) diterpenoid compounds with known biosynthetic gene clusters. **(B)** Bioactive diterpenoids isolated from Ascomycota and Basidiomycota for which pathways are not known (see Table 1 for details and references)* Cotylenin A **88** biosynthesis is assumed to proceed via a pathway similar to fusicoccin A **108**¹⁷⁴.



Scheme 11. Proposed biosynthetic pathway to gibberellic acids (GA) in *F. fujikuroi*. Production of the intermediate *ent*-kaurene **80** from GGPP **5** is catalyzed by a bifunctional CPS/KS type diterpene synthase (see Scheme 9). *Ent*-kaurene **80** is oxidized to the final product GA1 by four cytochrome P450s.



Scheme 12. Triterpenoid biosynthesis in fungi. Cyclization of 2,3-oxidosqualene **113** by an oxidosqualene cyclase (OSC) via a protonation dependent reaction results in a 6-6-6-5 tetracyclic protosteryl cation that is the precursor to the membrane sterol ergosterol **110** and the bioactive triterpenoid examples shown.




Figure 1. Putative Basidiomycota sesquiterpene synthases form clades consistent with their initial cyclization reaction. 117 Basidiomycota genomes listed in JGI were BLAST searched for gene catalogued protein homologues to Steh1|73029, identifying 777 proteins. Of these, 577 were assembled in an unrooted neighbor-joining dendrogram with the 22 previously characterized Basidiomycota sesquiterpene synthases, Cop1-6 (*Coprinus cinerea*),⁶⁹ Pro1 (*A. gallica*),⁷⁷ Omp1-10 (*O. olearius*),²³ Steh1|159379, 128017, 25180, 73029, and 64702 (*S. hirsutum*).²⁰ Enzymes cluster in clades according to initial cyclization mechanism. Characterized enzymes responsible for a: 1,10 cyclization of (*E,E*)-FPP are indicated by purple triangles; 1,10 cyclization of (*3R*)-NPP **6** are indicated by blue circles; 1,6 cyclization of (*3R*)-NPP **6** are indicated by green diamonds; 1,11 cyclization of (*E,E*)-FPP **4** are indicated by orange squares. The branch consisting of Δ 6-protoilludene **29** synthases is also highlighted.




Figure 2. Comparison of the intron architecture predicted by JGI and Augustus, and that found in cloned and characterized sesquiterpene synthases from *S. hirsutum*. Included are the transcripts encoding the active Steh1|25180, 73029, 64702, 128017 and 159379 enzymes. Stop codons are indicated with black arrowheads. Note that Steh1|64702-short contained an additional splice site compared to the active transcript, while Steh1|159379 isolate 6 was spliced to include a stop codon. An active form of Steh1|113028 could not be obtained.²⁰²⁰