




Cite this: *Org. Biomol. Chem.*, 2020, **18**, 1349

Received 21st January 2020,  
Accepted 4th February 2020

DOI: 10.1039/d0ob00137f

rsc.li/obc

## Automated access to well-defined ionic oligosaccharides†

Yuntao Zhu, <sup>a</sup> Theodore Tyrikos-Ergas, <sup>a,b</sup> Kevin Schiefelbein,<sup>a</sup>  
Andrea Grafmüller, <sup>c</sup> Peter H. Seeberger <sup>a,b</sup> and Martina Delbianco <sup>\*a</sup>

**Ionic polysaccharides are part of many biological events, but lack structural characterisation due to challenging purifications and complex synthesis. Four monosaccharides bearing modifications not found in nature are used for the automated synthesis of a collection of ionic oligosaccharides. Structural analysis reveals how the charge pattern affects glycan conformation.**

Polysaccharides bearing charged functional groups are ubiquitous in Nature, including chitosan, pectin, and glycosaminoglycans (GAGs).<sup>1</sup> These polysaccharides play important roles in plant and bacteria cell-wall formation,<sup>2,3</sup> cellular interactions,<sup>4</sup> and signaling.<sup>5</sup> Due to their unique gelation properties and tendency to bind metals<sup>6</sup> and/or proteins,<sup>5</sup> they are useful biomaterials for drug delivery and tissue engineering.<sup>2,7–10</sup> Little is known on how a particular charge pattern can affect the material properties,<sup>11–13</sup> partially due to the lack of pure samples with well-defined structures. Polysaccharides, generally extracted from natural sources as heterogeneous mixtures, limit definitive structure–property correlations. A better understanding of how the charge pattern affects carbohydrate conformation and aggregation is necessary for the design of functional carbohydrate materials, as already demonstrated for soft materials based on peptides and proteins.<sup>14,15</sup>

Automated Glycan Assembly (AGA) is a powerful method for the quick production of well-defined natural and unnatural oligosaccharides.<sup>16–18</sup> Many natural ionic oligosaccharides have been prepared by AGA,<sup>19–23</sup> however several complications make the synthesis of these glycans challenging.<sup>24</sup> Uronic acid building blocks (BBs) suffer from low reactivity during

glycosylation.<sup>19,20,23</sup> Glycosylation of natural 2-amino-2-deoxy-sugars often require tedious protecting group manipulations.<sup>20,25</sup> Regioselective sulfation is challenging and the poor stability of sulfated moieties can be troublesome due to migration and/or cleavage during further synthetic manipulations.<sup>20,26</sup> An alternative approach to access glycans with a well-defined charge pattern relies on the insertion of unnatural charged sugar units, that are much easier to handle during the glycan synthesis.

Here, we describe four unnatural BBs based on a carboxylic acid masked as methyl ester and an amine masked as azide. All BBs are prepared in gram scale from commercially available precursors and show good reactivity and stability upon storage. The AGA of a collection of oligosaccharides with well-defined charge patterns illustrates the utility of these BBs. Cellulose-like structures were prepared, due to the importance of cellulose and its ionic derivatives, such as carboxymethyl cellulose.<sup>27,28</sup> Deprotection methods on solid support, such as methanolysis and Staudinger reduction, were optimised to provide the desired compounds in good yields.<sup>29,30</sup> Molecular dynamics (MD) simulations indicate the importance of the charge pattern for the oligosaccharide conformation.

The insertion of unnatural functionalities in defined position of the cellulose backbone is a powerful approach to manipulate the hydrogen-bonding and modulate the aggregation of cellulose.<sup>29</sup> To gain a better understanding of the roles played by the charged moieties for oligosaccharide conformation, four glucose BBs were synthesised (**BB-1–4**, Scheme 1). Unnatural BBs bearing a methyl carboxylate or an azide were designed to reveal negative or a positive charge, upon global deprotection. The different position of the substituent on the glucose ring (position 3 vs. 6) permits to further manipulate the H-bond network.

The BB design follows AGA considerations.<sup>18</sup> Thioglycosides are stable to prolonged storage and show good reactivity during glycosylations.<sup>31</sup> Benzoyl groups (Bz) were used as participant protecting groups to direct the formation of the  $\beta$ -(1–4) linkage. The fluorenylmethoxycarbonyl (Fmoc) was selected as a temporary protecting group that can be easily

<sup>a</sup>Department of Biomolecular Systems, Max-Planck-Institute of Colloids and Interfaces, Am Mühlenberg 1, 14476 Potsdam, Germany.

E-mail: [martina.delbianco@mpikg.mpg.de](mailto:martina.delbianco@mpikg.mpg.de)

<sup>b</sup>Department of Chemistry and Biochemistry, Freie Universität Berlin, Arnimallee 22, 14195 Berlin, Germany

<sup>c</sup>Department of Theory, Max-Planck-Institute of Colloids and Interfaces, Am Mühlenberg 1, 14476 Potsdam, Germany

†Electronic supplementary information (ESI) available. See DOI: 10.1039/d0ob00137f






**Scheme 1** Strategies for the synthesis of four unnatural BBs bearing masked charged functionalities.

removed in the deprotection cycle, before chain elongation. The remaining positions were selectively functionalised with either methyl carboxylate, azides or benzyl ethers (Bn).

All BBs can be prepared following a similar strategy based on the selective protection of position 4 and 6 with cyclic acetals, regioselective functionalisation of positions 2 and 3, and selective acetal opening (Scheme 1). The azido-functionalised compound **1**, obtained in seven steps starting from glucose diacetate (see ESI†), served for the synthesis of **BB-1**. Peracetylated ethyl β-D-thioglucofuranoside (**2**) allowed for the synthesis of **BB2-4**. Methanolysis of the acetyl groups, followed by 4,6-*O*-benzylidene protection using benzaldehyde dimethyl acetal catalysed by camphorsulfonic acid, afforded compounds **3-4**.<sup>32</sup> Alternatively, the 4,6-*O*-naphthylmethylene acetal was introduced using 2-naphthaldehyde catalysed by camphorsulfonic acid and trimethyl orthoformate, yielding compound **5**.<sup>33</sup>

2-*O*-Benzoyl protected **6** was obtained readily upon treatment with benzoyl chloride and pyridine. Treatment with di-*n*-butyltin oxide, followed by addition of the desired alkyl bromide in the presence of cesium fluoride (CsF), permitted the regioselective functionalisation of compounds **4-5**.<sup>34,35</sup> The addition of CsF to the stannylidene acetal facilitated the nucleophilic substitution with the relatively unreactive, but sterically not very demanding, bromomethyl acetate. Benzoylation afforded compounds **7-9**.

Selective opening of the 4,6-*O*-acetal liberated the desired hydroxyl groups for further chemical modification.<sup>36</sup> Compound **8** was subjected to reductive ring opening conditions, in the presence of triethylsilane and trifluoroacetic acid (TFA), to reveal the 4-hydroxyl group. Similar conditions were not compatible with compound **6**, resulting in an additional azide reduction. Treatment with sodium cyanoborohydride and iodine afforded the target product (**BB-1** precursor).



**Scheme 2** AGA of tailor-made charged cellulose-like oligosaccharides. Yields of isolated products after AGA, deprotection, and purification are reported.



sor). Cleavage of the 4,6-O-benzylidene using TFA–water, followed by regioselective functionalisation of the primary alcohol with 4-toluenesulfonyl chloride (TsCl) permitted the introduction of the azide group in position 6 (**BB-2** precursor). Compound **9** was reacted with borane–tetrahydrofuran (BH<sub>3</sub>–THF) complex and trimethylsilyl trifluoromethanesulfonate (TMSOTf) to liberate the 6-hydroxyl group, that was subsequently subjected to etherification, using bromomethyl acetate and NaH (**BB-4** precursor). The introduction of Fmoc completed all BBs syntheses.

**BB1–4** were used to prepare a collection of charged oligocellulose analogues by AGA (Scheme 2). Merrifield resin bearing the photolabile linker (**S0**) was selected as solid support.<sup>37</sup> Unmodified glucose units were introduced using the commer-

cially available **BB-0**. A Staudinger reduction on solid phase was developed, adapting the conditions from a previously reported method.<sup>30</sup> The resin was suspended in THF–water and ammonia and triethylphosphine were added, liberating the desired amino group in 24 h. Methanolysis on solid phase<sup>29</sup> to hydrolyse the benzoyl groups was performed with trace amounts of water, permitting the simultaneous hydrolysis of the methyl esters. UV cleavage released the partially protected oligosaccharides that, upon hydrogenation catalysed by Pd/C, afforded the target compounds.

Twelve analogues with different charge patterns were prepared (Scheme 2). All four unnatural BBs showed good reactivity during glycosylation and stability during post AGA manipulation. The modified BBs were successfully positioned



**Fig. 1** Dihedral analysis ( $\omega$ ,  $\Psi$  and  $\Phi$ ) obtained by MD simulations and representative snapshots of the oligomers showing specific intramolecular interactions (highlighted with circles) due to the modifications. The effect of the modification on the  $\omega$  torsion angles is highlighted with arrows (top graphs). The most affected  $\Psi$  and  $\Phi$  dihedrals are marked by red or blue boxes. For (AdA)<sub>2</sub> the two blue lines serve as guide to the eye for comparing different minima. The residues are numbered from the non-reducing end (R1) to the reducing end (R6).





at the non-reducing end of the glycan (e.g. DA<sub>5</sub> and GA<sub>5</sub>) or internally (e.g. (AGA)<sub>2</sub>). Multiple substitutions were also tolerated without any significant loss in yield (e.g. (ADA)<sub>4</sub>). Zwitterionic compounds (e.g. AGA<sub>2</sub>DA) were prepared, suggesting the possibility of using such BBs for the preparation of GAG analogues.

These compounds are important substrates to evaluate the effect of the charge pattern on the oligosaccharide structure. MD simulations were performed to systematically examine the effect of each modification on glycan conformation (Fig. 1). Four analogues with the same sequence were compared (i.e. (AXA)<sub>2</sub>). All the modifications resulted in more flexible structures when compared to the unsubstituted analogue A<sub>6</sub> (see ESI†). Specific intramolecular hydrogen bonds stabilise particular conformations that can trigger the formation of different materials. The substitution at position 6 (g and d) produces a drastic effect on the  $\omega$  torsion angles, much less affected by the modification at position 3 (G and D). The interaction between the NH<sub>3</sub><sup>+</sup>(6) and the OH(3) of the adjacent sugar (R + 1) stabilises the *gt* rotamers (red and yellow plots, Fig. 1).

This also affects the glycosidic bond geometry with a high population at negative degrees (−172°) for  $\Psi_1$ ,  $\Psi_4$  ((AgA)<sub>2</sub> red boxes). The opposite trend is observed when position 6 is substituted with COO<sup>−</sup> (d). Due to sterics, the *gg* conformation is preferred, with a small percentage of *tg* stabilised by a COO<sup>−</sup>...OH(2) interaction (Fig. 1). This interaction also affects the dihedrals of the adjacent glycosidic bond ((AdA)<sub>2</sub> blue boxes). The substitutions in position 3 influence predominantly the glycosidic bond geometry. The interaction between the NH<sub>3</sub><sup>+</sup>(3) and the O(5) of the previous residue (R-1) preserved a cellulose-like character ((AGA)<sub>2</sub> red boxes). In contrast, the carboxylate at position 3 can engage in additional H-bonds, as observed between COO<sup>−</sup> and OH(2) of the same residue as well as the OH(6) of the previous sugar (R-1), resulting in remarkable changes of the  $\Phi$  and  $\Psi$  dihedrals ((ADA)<sub>2</sub> blue boxes).

## Conclusions

Four monosaccharide BBs bearing masked carboxylic acid or amino groups are synthesised. Their good reactivity and stability make them useful BBs for creation of ionic oligosaccharides. Twelve well-defined ionic cellulose analogues were prepared by AGA. Multiply charged oligosaccharides as well as zwitterionic compounds are accessible. MD simulations demonstrate how the nature and the position of the modification (3 vs. 6) plays a major role for the flexibility and conformation of such oligosaccharides. These differences could play a major role in the formation of supramolecular assembly based on charged polysaccharides.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We thank the Max-Planck Society, the Minerva Fast Track Program, and the MPG-FhG Cooperation Project Glyco3Dysplay, for generous financial support. Open Access funding provided by the Max Planck Society.

## References

- 1 A. Varki, R. Cummings, J. Esko, H. Freeze, P. Stanley, G. Hart and P. H. Seeberger, *Essentials of glycobiology*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 3rd edn, 2017.
- 2 B. R. Thakur, R. K. Singh, A. K. Handa and M. Rao, *Crit. Rev. Food Sci. Nutr.*, 1997, **37**, 47–73.
- 3 D.-J. Scheffers and M. G. Pinho, *Microbiol. Mol. Biol. Rev.*, 2005, **69**, 585–607.
- 4 A. Varki, *Glycobiology*, 2017, **27**, 3–49.
- 5 Y. Van Kooyk and G. A. Rabinovich, *Nat. Immunol.*, 2008, **9**, 593.
- 6 M. Krzesłowska, *Acta Physiol. Plant.*, 2011, **33**, 35–51.
- 7 S. Gim, Y. Zhu, P. H. Seeberger and M. Delbianco, *Wiley Interdiscip. Rev.: Nanomed. Nanobiotechnol.*, 2019, e1558.
- 8 M. George and T. E. Abraham, *J. Controlled Release*, 2006, **114**, 1–14.
- 9 M. Witvrouw and E. De Clercq, *Gen. Pharmacol.*, 1997, **29**, 497–511.
- 10 R. De Philippis and M. Vincenzini, *FEMS Microbiol. Rev.*, 1998, **22**, 151–175.
- 11 S. Zhang, Z. Zhang and B. Vardhanabhuti, *Food Funct.*, 2014, **5**, 1829–1838.
- 12 I. Alshanski, J. Blaszkiewicz, E. Mervinetsky, J. Rademann, S. Yitzchaik and M. Hurevich, *Chem. – Eur. J.*, 2019, **25**, 12083–12090.
- 13 L. Schefer, J. Adamcik and R. Mezzenga, *Angew. Chem., Int. Ed.*, 2014, **53**, 5376–5379.
- 14 R. Mezzenga, J.-M. Jung and J. Adamcik, *Langmuir*, 2010, **26**, 10401–10405.
- 15 V. Castelletto, I. W. Hamley, J. Adamcik, R. Mezzenga and J. Gummel, *Soft Matter*, 2012, **8**, 217–226.
- 16 O. J. Plante, E. R. Palmacci and P. H. Seeberger, *Science*, 2001, **291**, 1523–1527.
- 17 P. H. Seeberger, *Acc. Chem. Res.*, 2015, **48**, 1450–1463.
- 18 M. Guberman and P. H. Seeberger, *J. Am. Chem. Soc.*, 2019, **141**, 5581–5592.
- 19 M. T. Walvoort, H. van den Elst, O. J. Plante, L. Kröck, P. H. Seeberger, H. S. Overkleeft, G. A. van der Marel and J. D. Codée, *Angew. Chem., Int. Ed.*, 2012, **51**, 4393–4396.
- 20 S. Eller, M. Collot, J. Yin, H. S. Hahm and P. H. Seeberger, *Angew. Chem., Int. Ed.*, 2013, **52**, 5858–5861.
- 21 H. S. Hahm, F. Broecker, F. Kawasaki, M. Mietzsch, R. Heilbronn, M. Fukuda and P. H. Seeberger, *Chem*, 2017, **2**, 114–124.
- 22 C.-F. Liang, H. S. Hahm and P. H. Seeberger, in *Glycosaminoglycans*, Springer, 2015, pp. 3–10.



- 23 M. W. Weishaupt, S. Matthies, M. Hurevich, C. L. Pereira, H. S. Hahm and P. H. Seeberger, *Beilstein J. Org. Chem.*, 2016, **12**, 1440–1446.
- 24 A. Pardo-Vargas, M. Delbianco and P. H. Seeberger, *Curr. Opin. Chem. Biol.*, 2018, **46**, 48–55.
- 25 M. Guberman, M. Bräutigam and P. H. Seeberger, *Chem. Sci.*, 2019, **10**, 5634–5640.
- 26 R. S. Loka, E. T. Sletten, U. Barash, I. Vlodavsky and H. M. Nguyen, *ACS Appl. Mater. Interfaces*, 2018, **11**, 244–254.
- 27 B. Thomas, M. C. Raj, J. Joy, A. Moores, G. L. Drisko and C. M. Sanchez, *Chem. Rev.*, 2018, **118**, 11575–11625.
- 28 L. T. Mika, E. Csefalvay and A. Nemeth, *Chem. Rev.*, 2017, **118**, 505–613.
- 29 Y. Yu, T. Tyrikos-Ergas, Y. Zhu, G. Fittolani, V. Bordoni, A. Singhal, R. J. Fair, A. Grafmüller, P. H. Seeberger and M. Delbianco, *Angew. Chem., Int. Ed.*, 2019, **58**, 13127–13132.
- 30 D. Senf, C. Ruprecht, G. H. de Kruijff, S. O. Simonetti, F. Schuhmacher, P. H. Seeberger and F. Pfrenkle, *Chem. – Eur. J.*, 2017, **23**, 3197–3205.
- 31 G. Veeneman, S. Van Leeuwen and J. Van Boom, *Tetrahedron Lett.*, 1990, **31**, 1331–1334.
- 32 A. M. Vibhute, V. Muvvala and K. M. Sureshan, *Angew. Chem., Int. Ed.*, 2016, **55**, 7782–7785.
- 33 S. Traboni, E. Bedini, M. Giordano and A. Iadonisi, *Adv. Synth. Catal.*, 2015, **357**, 3562–3572.
- 34 T. W. Rising, T. D. Claridge, N. Davies, D. P. Gamblin, J. W. Moir and A. J. Fairbanks, *Carbohydr. Res.*, 2006, **341**, 1574–1596.
- 35 T. B. Grindley, *Adv. Carbohydr. Chem. Biochem.*, 1998, **53**, 17.
- 36 M. Ohlin, R. Johnsson and U. Ellervik, *Carbohydr. Res.*, 2011, **346**, 1358–1370.
- 37 K. Le Mai Hoang, A. Pardo-Vargas, Y. Zhu, Y. Yu, M. Loria, M. Delbianco and P. H. Seeberger, *J. Am. Chem. Soc.*, 2019, **141**, 9079–9086.

