


 Cite this: *RSC Adv.*, 2021, **11**, 12249

 Received 2nd March 2021  
 Accepted 19th March 2021

DOI: 10.1039/d1ra01646f

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

# Transmembrane anion transport promoted by thioamides†

 Robert Pomorski,<sup>a</sup> María García-Valverde,<sup>b</sup> Roberto Quesada<sup>\*b</sup> and Michał J. Chmielewski<sup>†\*a</sup>

Thioamide groups represent useful hydrogen-bonding motifs for the development of active transmembrane anion transporters. Using a 1,8-di(thioamido)carbazole scaffold the superior performance of thioamides compared with the parent amides has been demonstrated.

## Introduction

Small molecules capable of performing protein functions represent attractive minimalistic models to study cellular processes and are appealing for potential biological applications.<sup>1–3</sup> In recent years, small molecules capable of facilitating the selective permeation of anions through lipid bilayers have emerged as promising candidates for the development of anti-cancer and antimicrobial agents as well as therapeutics for the treatment of channelopathies caused by faulty anion transport, such as cystic fibrosis, Dent's disease, and others.<sup>4–6</sup>

Transmembrane ion transport is a complex phenomenon influenced by a number of factors including binding energies and lipophilicity.<sup>7–9</sup> Subtle changes in the molecular structure of anionophores often result in dramatic differences in their performance.<sup>10–12</sup> Nevertheless, important insights into the design of this type of molecules have been obtained in recent years.<sup>13</sup> Hydrogen bonding is the most frequently employed interaction in the design of synthetic anion transporters. Several hydrogen bond donors arranged in a convergent manner ensure effective binding between the transporter and the anion, thus forming a supramolecular complex capable of permeating the membrane. Ureas and thioureas are among the most successful anion binding motifs employed in this type of molecules.<sup>14,15</sup> Thioureas usually outperform ureas as anion carriers because replacement of oxygen by sulfur improves the hydrogen bonding ability of the N–H groups, reduces tendency to self-assemble, and increases lipophilicity of the molecule. This strategy of using urea and thiourea derivatives has been

extensively used to produce anionophores with increased activity.<sup>16</sup> On the other hand, the potential of thioamides has remained largely unexplored despite a similar effect would be expected and amide groups are frequently employed in the construction of active anion receptors and transporters.<sup>17</sup> In this paper, we explore the potential of the thioamide moiety in the development of active anionophores.

## Results and discussion

We have recently reported a family of 1,8-diamidocarbazoles as anion receptors and identified compound **4A** as a highly active anion carrier (Fig. 1).<sup>18</sup> We envisaged the potential of this motif to prepare their thioamide derivatives and to explore their anion transport properties.

Thioamides **1T–5T** as well as amido-thioamides **5AT** and **6AT** were synthesized from the corresponding 1,8-diamidocarbazoles by treatment with Lawesson's reagent and fully characterized (Fig. 1, see also ESI† for details).

<sup>1</sup>H NMR titrations of the thioamides with TBACl in DMSO-*d*<sub>6</sub> + 0.5% H<sub>2</sub>O resulted in large downfield shifts of the carbazole NH protons ( $\Delta\delta_{\max}$  from 1.83 to 2.55 ppm, except for the most weakly bonding **1T**, for which  $\Delta\delta_{\max} = 0.94$  ppm was observed) as well as in much smaller downfield shifts of the thioamide NHs ( $0.42 < \Delta\delta_{\max} < 0.65$  ppm, again with the exception of **1T**, where  $\Delta\delta_{\max} = 0.89$  ppm) (Fig. 2). In case of the amido-thioamides **5AT** and **6AT**, the CSNH protons shifted slightly upfield upon chloride binding (by 0.13–0.18 ppm), mirroring the behaviour of CONH protons of their parent diamide receptors.<sup>19</sup> All these observations suggest that the hydrogen bonds formed with the anion by the central carbazole NHs are much stronger than the bonds formed by the (thio)amide side arms.

This supposition was further corroborated by the X-ray crystal structure of the chloride complex of dithioamide **3T**, in which the carbazole NH–Cl distance is much shorter (2.248 Å) than the two thioamide NH–Cl distances (2.581 Å and 2.830 Å), see Fig. 1. The inability of the thioamide arms to form shorter bonds with chloride might be related to the steric bulk of the sulfur atom, which enforces significant torsion of the thioamide

<sup>a</sup>Faculty of Chemistry, Biological and Chemical Research Centre, University of Warsaw, Żwirki i Wigury 101, 02-089 Warszawa, Poland. E-mail: mchmielewski@chem.uw.edu.pl

<sup>b</sup>Departamento de Química, Universidad de Burgos, Burgos 09001, Spain. E-mail: rquesada@ubu.es

† Electronic supplementary information (ESI) available: Compound characterization data and anion transport assays. CCDC 2027226. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d1ra01646f






Fig. 1 Compounds included in this study and solid-state X-ray structure of compound **3T** complexing chloride (as TBACl; TBA cation omitted for clarity).

arms with respect to the carbazole plane (44–46°) as well as to geometric mismatch between the relatively wide binding cavity of the receptor and the relatively small chloride anion.

Quantitative analysis of the titration data revealed that the association constants with chloride turned out to be 2–3 times smaller for thioamides in comparison to their amide precursors (Table 1). This is in contrast to what might have been expected based on the much higher acidity of thioamides,<sup>20</sup> but in line with some literature precedents.<sup>21,22</sup> Very likely, the beneficial effect of enhanced hydrogen bond donating ability of the

thioamide moieties<sup>23</sup> is counterbalanced here by unfavourable conformational preferences of the thioamides.<sup>24,25</sup>

The anion transport properties of these compounds were investigated in POPC phospholipid vesicles using a chloride selective electrode.<sup>26</sup> In these assays, the facilitated chloride efflux induced by aliquots of the compounds was monitored over time. All receptors were first screened at 2% carrier to POPC molar ratio and compounds eliciting more than 50% chloride efflux were further analysed at different concentrations. For these latter derivatives, the concentration-dependent chloride efflux activity was then fitted using the Hill equation to obtain the EC<sub>50</sub> parameters representing the amount of compound needed to induce 50% efflux of the encapsulated chloride. All the experiments were repeated at least three times using different batches of vesicles (Fig. 3).

The positive effect of replacing the amide groups by the thioamide was evident with all 1,8-di(thioamido)carbazoles outperforming their parent 1,8-diamidocarbazoles. The least active derivative was found to be the 1,8-dibenzothioamido derivative **1T**. For this compound, no EC<sub>50</sub> could be calculated. The poor performance of transporters bearing aromatic substituents, such as **1T**, is likely related to their much lower chloride affinity.<sup>18</sup> Thioamides **2T–5T** were found to be highly active in the chloride efflux assay with nitrate in the external solution, with EC<sub>50</sub> values in the submicromolar range. The most active derivative was **4T**, with a calculated EC<sub>50</sub> of 67 nM. The positive effect of replacing even a single amide group by thioamide is demonstrated in the case of compounds **5AT** and **6AT**. Compound **5AT** displayed transport efficacy rivalling the best performers identified in this study. Compound **6AT** is an active transporter derived from a non-chlorinated carbazole core. None of the diamides derived from this core showed any chloride transport activity in our previous investigations.<sup>18</sup>

Experiments were also performed by suspending the vesicles in an external sulfate solution or an external mixture of sulfate and bicarbonate. These assays allow studying the impact of the anion composition of the external buffer on the observed chloride efflux. Bicarbonate is an anion of biological relevance and is significantly more hydrophilic than chloride and nitrate,

Table 1 Association constants  $K_a$  for 1 : 1 complexes of compounds **1T–5T**, **5AT** and **6AT** as well as their amide congeners (in brackets) with chloride (added as tetrabutylammonium salt) determined from <sup>1</sup>H NMR titration experiments in DMSO-*d*<sub>6</sub> + 0.5% H<sub>2</sub>O at 293 K and transport activities expressed as EC<sub>50</sub> (nM)

Compound	$K_a$ (M <sup>-1</sup> )	EC <sub>50</sub> (nM) NO <sub>3</sub> <sup>-</sup> /Cl <sup>-</sup>	EC <sub>50</sub> (% molar) NO <sub>3</sub> <sup>-</sup> /Cl <sup>-</sup>	EC <sub>50</sub> (nM) HCO <sub>3</sub> <sup>-</sup> /Cl <sup>-</sup>	EC <sub>50</sub> (% molar) HCO <sub>3</sub> <sup>-</sup> /Cl <sup>-</sup>
<b>1T</b>	<10 (14) <sup>a</sup>	— <sup>b</sup>	— <sup>b</sup>	— <sup>b</sup>	— <sup>b</sup>
<b>2T</b>	49 (104) <sup>a</sup>	297	0.059	4290	0.858
<b>3T</b>	50 (109) <sup>a</sup>	107	0.022	2910	0.582
<b>4A</b>	123 <sup>a</sup>	184	0.037	— <sup>b</sup>	— <sup>b</sup>
<b>4T</b>	46	66	0.013	385	0.077
<b>5A</b>	159 <sup>a</sup>	— <sup>b</sup>	— <sup>b</sup>	— <sup>b</sup>	— <sup>b</sup>
<b>5T</b>	52	93	0.019	465	0.093
<b>5AT</b>	67	61	0.012	1719	0.344
<b>6A</b>	48 <sup>a</sup>	— <sup>b</sup>	— <sup>b</sup>	— <sup>b</sup>	— <sup>b</sup>
<b>6AT</b>	23	353	0.071	— <sup>b</sup>	— <sup>b</sup>

<sup>a</sup> The association constants determined for parent bisamides were taken from ref. 18. <sup>b</sup> The compound did not display enough activity to calculate an EC<sub>50</sub> value.






Fig. 2  $^1\text{H}$  NMR titration of 5T (5 mM) with TBACl in  $\text{DMSO-}d_6$  + 0.5%  $\text{H}_2\text{O}$ .  $T = 293$  K.

thus more difficult to extract into the lipid bilayer. This was reflected in the higher calculated  $\text{EC}_{50}$  values. Only thioamides 4T and 5T displayed  $\text{EC}_{50}$  values in the submicromolar range under these conditions. No significant chloride efflux was detected in the presence of sulfate alone in the external milieu (see Fig. S56<sup>†</sup>). This result supports an exchange mechanism accounting for the transport activity displayed by these compounds, where the higher hydrophilicity of the doubly charged sulfate renders the compounds inactive as a result of their inability to facilitate the transport of this anion. This result also rules out any unspecific transport or membrane destabilization effect promoted by these compounds. The ability of some of these compounds to exchange chloride and bicarbonate bodes well for their potential biological applications.




Fig. 3 Chloride efflux promoted by 4T at different concentrations (1–0.005% carrier to POPC molar ratio, 5–0.025  $\mu\text{M}$ ) in unilamellar POPC vesicles. Vesicles loaded with 489 mM NaCl were buffered at pH 7.2 with 5 mM phosphate and dispersed in 489 mM  $\text{NaNO}_3$  buffered at pH 7.2. Each trace represents the average of at least three trials.

## Conclusions

In conclusion, we have demonstrated the usefulness of thioamides in the development of highly active transmembrane anion transporters.<sup>27</sup> These compounds display enhanced transmembrane transport activity compared to the parent amide derivatives. Taking into account the widespread use of amides as hydrogen-bonding motifs in successful anionophores, it is expected that this strategy will allow the identification of new candidates with improved anion transport efficiency.

## Experimental

### *N,N'*-(3,6-Dichloro-9H-carbazole-1,8-diyl)dibenzothioamide (1T)

A 50 ml, 2-neck, round-bottomed flask was dried for 10 min with a stream of hot air (*ca.* 500 °C) from heat-gun and cooled down to rt in a desiccator. Then the flask was charged with diamide 1A (238 mg, 0.502 mmol), Lawesson's reagent (444 mg, 1.10 mmol) and a stir bar. Then the flask was equipped with a reflux condenser connected to a Schlenk line and its side neck was closed with a stopper. Air was removed by three pump/thaw cycles and then dry THF (30 ml) was added in a stream of argon through the side neck. The reaction mixture was intensively stirred and refluxed overnight. After this time the mixture was cooled down and 2.0 g of silica gel was added. Solvents were evaporated on a rotary evaporator and the solid residue was dried under high vacuum. Then the solid residue was suspended in hexane (40 ml) and put on top of a chromatographic column made of 110 g of silica suspended in  $\text{CHCl}_3$ . The column was eluted with  $\text{CHCl}_3$  (*ca.* 500 ml),  $\text{CHCl}_3$  :  $\text{CH}_3\text{CO}_2\text{Et}$  = 40 : 1 (*ca.* 500 ml) and  $\text{CHCl}_3$  :  $\text{CH}_3\text{CO}_2\text{Et}$  = 30 : 1 until the desired product was completely washed out. Fractions containing pure product were combined and evaporated on a rotary evaporator yielding 293 mg (98.0%) of yellow solid. Mp: decomposition at 301 °C.  $^1\text{H}$  NMR (500 MHz,  $\text{DMSO-}d_6$ )  $\delta_{\text{DMSO}}$ : 11.69 (s, 2H, thioamide NHs), 11.23 (s, 1H, carbazole NH), 8.37 (bs, 2H, carbazole CH-4/5), 8.03 (d,  $J = 7.6$  Hz, 4H, phenyl CH-ortho), 7.77 (bs, 2H, carbazole CH-2/7), 7.56 (t,  $J = 7.3$  Hz, 2H, phenyl CH-para), 7.48 (t,  $J = 7.5$  Hz, 4H, phenyl CH-meta).  $^{13}\text{C}$  NMR (126 MHz,  $\text{DMSO-}d_6$ )  $\delta_{\text{DMSO}}$ : 199.19, 141.09, 133.99, 131.30, 127.98, 125.66, 124.62, 124.46, 122.98, 119.23. Elemental analysis: calcd for  $\text{C}_{26}\text{H}_{17}\text{Cl}_2\text{N}_3\text{S}_2$ : C, 61.66; H, 3.38; N, 8.30, found: C, 61.56; H, 3.52; N, 8.27. HR MS (TOF MS  $\text{ES}^-$ )  $m/z$  calcd for  $\text{C}_{26}\text{H}_{16}\text{Cl}_2\text{N}_3\text{S}_2^-$ : 504.0163 found: 504.0168.

Compounds 2T–5T, 5AT and 6AT were prepared using similar procedures. Complete information is provided in the ESI.<sup>†</sup>

### *N,N'*-(3,6-Dichloro-9H-carbazole-1,8-diyl)dipropanthioamide (2T)

Yellow solid, yield 79.7%. Mp: decomposition at 270 °C.  $^1\text{H}$  NMR (500 MHz,  $\text{DMSO-}d_6$ )  $\delta_{\text{DMSO}}$ : 11.53 (s, 2H, thioamide NH); 10.84 (s, 1H, carbazole NH); 8.31 (d,  $J = 2.0$  Hz, 2H, carbazole CH); 7.75 (d,  $J = 2.0$  Hz, 2H, carbazole CH); 2.88 (q,  $J = 7.5$  Hz, 4H, methylene  $\text{CH}_2$ ); 1.35 (t,  $J = 7.5$  Hz, 6H, methyl  $\text{CH}_3$ ).  $^{13}\text{C}$  NMR (126 MHz,  $\text{DMSO-}d_6$ )  $\delta_{\text{DMSO}}$ : 208.02, 133.48, 124.93,



124.47, 123.81, 122.99, 119.04, 38.84, 13.92. Elemental analysis: calcd for  $C_{18}H_{17}Cl_2N_3S_2$ : C, 52.68; H, 4.18; N, 10.24, found: C, 52.64; H, 4.35; N, 10.09. HR MS (TOF MS  $ES^-$ )  $m/z$  calculated for  $C_{18}H_{16}Cl_2N_3S_2^-$ : 408.0163 found: 408.0161.

#### ***N,N'*-(3,6-Dichloro-9*H*-carbazole-1,8-diyl)dibutanethioamide (3T)**

Yellow solid, yield 80.2%. M.p. decomposition at 282 °C.  $^1H$  NMR (500 MHz, DMSO- $d_6$ )  $\delta_{DMSO}$ : 11.56 (s, 2H, thioamide NH), 10.64 (s, 1H, carbazole NH), 8.31 (d,  $J = 1.9$  Hz, 2H, carbazole CH), 7.71 (d,  $J = 2.0$  Hz, 2H, carbazole CH), 2.85 (t, 4H,  $J = 6.0$  Hz,  $CH_2C=S$ ), 1.93–1.81 (m, 4H,  $CH_2$ ), 1.02 (t,  $J = 7.4$  Hz, 6H,  $CH_3$ ).  $^{13}C$  NMR (126 MHz, DMSO- $d_6$ )  $\delta_{DMSO}$ : 206.45, 133.45, 124.90, 124.54, 123.78, 123.06, 119.12, 47.69, 22.61, 13.27. Elemental analysis: calcd for  $C_{20}H_{21}Cl_2N_3S_2$ : C, 54.79; H, 4.83; N, 9.58, found: C, 54.88; H, 5.06; N, 9.47. HR MS (TOF MS  $ES^-$ )  $m/z$  calcd for  $C_{20}H_{20}Cl_2N_3S_2^-$ : 436.0476 found: 436.0472.

#### ***N,N'*-(3,6-Dichloro-9*H*-carbazole-1,8-diyl)bis(3-methylbutanethioamide) (4T)**

Yellow solid, yield 64.7%.  $^1H$  NMR (500 MHz, DMSO- $d_6$ )  $\delta_{DMSO}$ : 11.57 (s, 2H, thioamide NH), 10.46 (s, 1H, carbazole NH), 8.33 (d,  $J = 1.5$  Hz, 2H, carbazole CH-4), 7.66 (d,  $J = 1.5$  Hz, 2H, carbazole CH-2), 2.75 (d,  $J = 7.2$  Hz, 4H,  $CH_2$ ), 2.40–2.25 (m, 2H, CH), 1.03 (d,  $J = 6.6$  Hz, 12H,  $CH_3$ ).  $^{13}C$  NMR (126 MHz, DMSO- $d_6$ )  $\delta_{DMSO}$ : 205.56, 133.46, 124.89, 124.66, 123.79, 123.15, 119.22, 54.88, 39.50, 29.07, 21.96. Elemental analysis: calcd for  $C_{22}H_{25}Cl_2N_3S_2$ : C, 56.64; H, 5.40; N, 9.01, found: C, 56.75; H, 5.54; N, 8.94. HR MS (TOF MS  $ES^-$ )  $m/z$  calcd for  $C_{22}H_{24}Cl_2N_3S_2^-$ : 464.0789 found: 464.0786.

#### ***N,N'*-(3,6-Dichloro-9*H*-carbazole-1,8-diyl)bis(3,3-dimethylbutanethioamide) (5T)**

Yellow solid, yield 95.1%. M.p. decomposition at 278 °C.  $^1H$  NMR (500 MHz, DMSO- $d_6$ )  $\delta_{DMSO}$ : 11.43 (s, 2H; thioamide NH); 10.30 (s, 1H; carbazole NH); 8.33 (d;  $J = 2.0$  Hz; 2H; carbazole CH-4/5); 7.59 (d;  $J = 2.0$  Hz; 2H; carbazole CH-2/7); 2.85 (s; 4H; methylene  $CH_2$ ); 1.14 (s; 18H; *t*-butyl  $CH_3$ ).  $^{13}C$  NMR (126 MHz, DMSO- $d_6$ )  $\delta_{DMSO}$ : 203.50, 133.43, 125.04, 124.75, 123.89, 123.17, 119.18, 59.12, 32.02, 29.77. Elemental analysis: calcd for  $C_{24}H_{29}Cl_2N_3S_2$ : C, 58.29; H, 5.91; N, 8.50, found: C, 58.37; H, 6.06; N, 8.25. HR MS (TOF MS  $ES^-$ )  $m/z$  calcd for  $C_{24}H_{28}Cl_2N_3S_2^-$ : 492.1102; found: 492.1113.

#### ***N*-(3,6-Dichloro-8-(3,3-dimethylbutanethioamido)-9*H*-carbazol-1-yl)-3,3-dimethylbutanamide (5AT)**

Yellow solid, yield 16.7%.  $^1H$  NMR (DMSO- $d_6$ )  $\delta_{DMSO}$ : 11.55 (s, 1H, thioamides NH), 10.56 (s, 1H, carbazole NH), 10.00 (s, 1H, amide NH), 8.30 (d,  $J = 1.8$  Hz, 1H, carbazole CH-5), 8.12 (d,  $J = 1.8$  Hz, 1H, carbazole CH-4), 7.90 (d,  $J = 1.8$  Hz, 1H, carbazole CH-2), 7.41 (d,  $J = 1.6$  Hz, 1H, carbazole CH-7), 2.88 (s, 2H,  $CH_2C=S$ ), 2.35 (s, 2H,  $CH_2C=O$ ), 1.16 (s, 9H, *t*-Bu on the thioamide side), 1.08 (s, 9H, *t*-Bu on the amide side).  $^{13}C$  NMR (DMSO- $d_6$ )  $\delta_{DMSO}$ : 203.85, 170.57, 133.40, 129.89, 125.06, 124.64, 124.59, 124.24, 124.00, 123.73, 122.96, 119.26, 117.74,

115.92, 58.90, 49.20, 32.07, 30.89, 29.78, 29.59. HR MS (TOF MS  $ES^-$ ):  $m/z$  calcd for  $C_{24}H_{28}Cl_2N_3OS^-$ : 476.1330, found: 476.1331. Elemental analysis calcd for  $C_{24}H_{29}Cl_2N_3OS$ : C, 60.24; H, 6.11; N, 8.78; found: C, 60.61; H, 6.39; N, 8.39.

#### ***N*-(8-(3,3-Dimethylbutanethioamido)-9*H*-carbazol-1-yl)-3,3-dimethylbutanamide (6AT)**

Yellow solid, yield 40.3%, mp decomposition above 282 °C.  $^1H$  NMR (500 MHz, DMSO- $d_6$ )  $\delta_{DMSO}$ : 11.47 (s; 1H; thioamide NH); 10.06 (s; 1H; carbazole NH); 9.95 (s; 1H; amide NH); 8.08 (d,  $J = 7.5$  Hz, 1H; CH-5 on the thioamide side); 7.94 (d,  $J = 7.7$  Hz, 1H; CH-4 on the amide side); 7.61 (dd;  $J_1 = 7.8$ ,  $J_2 = 1.0$  Hz; 1H; CH-2 on the amide side); 7.34 (d,  $J = 7.7$ , 1H; CH-7 on the thioamide side); 7.22 (t;  $J = 7.7$  Hz; 1H; CH-6 on the thioamide side); 7.17 (t;  $J = 7.8$  Hz; 1H; CH-3 on the amide side); 2.87 (s; 2H; methylene  $CH_2$  from the thioamide arm); 2.34 (s; 2H; methylene  $CH_2$  from the amide arm); 1.17 (s; 9H; *t*-butyl  $CH_3$  from the thioamide arm); 1.10 (s; 9H; *t*-butyl from the amide arm).  $^{13}C$  NMR (126 MHz, DMSO- $d_6$ )  $\delta_{DMSO}$ : 202.76, 170.19, 134.29, 131.48, 124.60, 124.47, 124.10, 123.61, 123.40, 119.43, 119.07, 119.05, 118.28, 116.28, 58.89, 49.13, 31.91, 30.84, 29.85, 29.68. Elemental analysis: calcd for  $C_{24}H_{31}N_3OS$ : C, 70.38; H, 7.63; N, 10.26, found: C, 70.07; H, 7.61; N, 10.05. HR MS (TOF MS  $ES^-$ )  $m/z$  calcd for  $C_{24}H_{30}N_3OS^-$ : 408.2110 found: 408.2112.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We thank the Polish National Science Centre for grant OPUS (2011/01/B/ST5/03900) and Consejería de Educación de la Junta de Castilla y León (project BU067P20). Part of this study was carried out at the Biological and Chemical Research Centre, University of Warsaw, established within a project co-financed by the European Union through the European Regional Development Fund under the Operational Programme Innovative Economy 2007–2013. The authors gratefully acknowledge Andrea Sancho-Medina and Óscar Fernández-Ramos for their contributions to transmembrane anion transport experiments.

## Notes and references

- 1 A. S. Grillo, A. M. SantaMaria, M. D. Kafina, A. G. Cioffi, N. C. Huston, M. Han, Y. A. Seo, Y. Y. Yien, C. Nardone, A. V. Menon, J. Fan, D. C. Svoboda, J. B. Anderson, J. D. Hong, B. G. Nicolau, K. Subedi, A. A. Gewirth, M. Wessling-Resnick, J. Kim, B. H. Paw and M. D. Burke, *Science*, 2017, **356**, 608–616.
- 2 S. Zheng, L. Huang, Z. Sun and M. Barboiu, *Angew. Chem., Int. Ed.*, 2021, **60**, 566–597.
- 3 F.-F. Shen, S.-Y. Dai, N.-K. Wong, S. Deng, A. S.-T. Wong and D. Yang, *J. Am. Chem. Soc.*, 2020, **142**, 10769–10779.
- 4 V. Kaushik, J. Yakisich, A. Kumar, N. Azad and A. Iyer, *Cancers*, 2018, **10**, 360.



- 5 J. T. Davis, P. A. Gale and R. Quesada, *Chem. Soc. Rev.*, 2020, **49**, 6056–6068.
- 6 N. Akhtar, O. Biswas and D. Manna, *Chem. Commun.*, 2020, **56**, 14137–14153.
- 7 H. Li, H. Valkenier, A. G. Thorne, C. M. Dias, J. A. Cooper, M. Kieffer, N. Busschaert, P. A. Gale, D. N. Sheppard and A. P. Davis, *Chem. Sci.*, 2019, **10**, 9663–9672.
- 8 I. Marques, P. M. R. Costa, M. Q. Miranda, N. Busschaert, E. N. W. Howe, H. J. Clarke, C. J. E. Haynes, I. L. Kirby, A. M. Rodilla, R. Pérez-Tomás, P. A. Gale and V. Félix, *Phys. Chem. Chem. Phys.*, 2018, **20**, 20796–20811.
- 9 N. J. Knight, E. Hernando, C. J. E. Haynes, N. Busschaert, H. J. Clarke, K. Takimoto, M. García-Valverde, J. G. Frey, R. Quesada and P. A. Gale, *Chem. Sci.*, 2016, **7**, 1600–1608.
- 10 X. Wu, A. M. Gilchrist and P. A. Gale, *Chem*, 2020, **6**, 1296–1309.
- 11 L. Martínez-Crespo, J. L. Sun-Wang, P. Ferreira, C. F. M. Mirabella, G. Aragay and P. Ballester, *Chem.–Eur. J.*, 2019, **25**, 4775–4781.
- 12 H. Valkenier, O. Akrawi, P. Jurček, K. Sleziaková, T. Lízal, K. Bartik and V. Šindelář, *Chem*, 2019, **5**, 429–444.
- 13 X. Wu, E. N. W. Howe and P. A. Gale, *Acc. Chem. Res.*, 2018, **51**, 1870–1879.
- 14 H. Valkenier, C. M. Dias, K. L. Porter Goff, O. Jurček, R. Puttreddy, K. Rissanen and A. P. Davis, *Chem. Commun.*, 2015, **51**, 14235–14238.
- 15 M. J. Spooner, H. Li, I. Marques, P. M. R. Costa, X. Wu, E. N. W. Howe, N. Busschaert, S. J. Moore, M. E. Light, D. N. Sheppard, V. Félix and P. A. Gale, *Chem. Sci.*, 2019, **10**, 1976–1985.
- 16 P. A. Gale, J. T. Davis and R. Quesada, *Chem. Soc. Rev.*, 2017, **46**, 2497–2519.
- 17 N. J. Andrews, C. J. E. Haynes, M. E. Light, S. J. Moore, C. C. Tong, J. T. Davis, W. A. Harrell Jr and P. A. Gale, *Chem. Sci.*, 2011, **2**, 256–260.
- 18 K. M. Bąk, K. Chabuda, H. Montes, R. Quesada and M. J. Chmielewski, *Org. Biomol. Chem.*, 2018, **16**, 5188–5196.
- 19 The rather uncommon upfield shifts of the (thio)amide protons upon chloride binding might be rationalised in terms of their engagement in strong hydrogen bonding interactions with solvent molecules prior to the addition of chloride.
- 20 F. G. Bordwell, *J. Am. Chem. Soc.*, 1988, **110**, 5903–5904.
- 21 S. J. Coles, P. A. Gale, M. B. Hursthouse, M. E. Light and C. N. Warrimer, *Supramol. Chem.*, 2004, **16**, 469–486.
- 22 T. Zieliński and J. Jurczak, *Tetrahedron*, 2005, **61**, 4081–4089.
- 23 H.-J. Lee, Y.-S. Choi, K.-B. Lee, J. Park and C.-J. Yoon, *J. Phys. Chem. A*, 2002, **106**, 7010–7017.
- 24 T. Zieliński, M. Kędziorek and J. Jurczak, *Chem.–Eur. J.*, 2008, **14**, 838–846.
- 25 I. Sandler, F. A. Larik, N. Mallo, J. E. Beves and J. Ho, *J. Org. Chem.*, 2020, **85**, 8074–8084.
- 26 L. A. Jowett and P. A. Gale, *Supramol. Chem.*, 2019, **31**, 297–312.
- 27 K. M. Bąk, B. van Kolck, K. Masłowska-Jarżyna, P. Papadopoulou, A. Kros and M. J. Chmielewski, *Chem. Commun.*, 2020, **56**, 4910–4913.

