

**ROY Confined in Hydrogen-Bonded Frameworks: Coercing  
Conformation of a Chromophore**

Journal:	<i>Materials Chemistry Frontiers</i>
Manuscript ID	QM-RES-03-2020-000200.R2
Article Type:	Research Article
Date Submitted by the Author:	11-Jun-2020
Complete List of Authors:	Tang, Sishuang; New York University, Chemistry Yusov, Anna; New York University, Chemistry Li, Yuantao; New York University, Chemistry Tan, Melissa; New York University, Department Chemistry and Molecular Design Institute Hao, Yunhui; New York University Li, Zongzhe; New York University, Chemistry Chen, Yu-Sheng; University of Chicago Advanced Photon Source,, ChemMatCARS Hu, Chunhua; New York University, Department of Chemistry Kahr, Bart; New York University, Department Chemistry and Molecular Design Institute Ward, Michael; New York University, Chemistry

# ROY Confined in Hydrogen-Bonded Frameworks: Coercing Conformation of a Chromophore

Sishuang Tang<sup>a</sup>, Anna Yusov<sup>a</sup>, Yuantao Li<sup>a</sup>, Melissa Tan<sup>a</sup>, Yunhui Hao<sup>a</sup>, Zongzhe Li<sup>a</sup>, Yu-Sheng Chen<sup>b</sup>, Chunhua T. Hu<sup>a</sup>, Bart Kahr<sup>a</sup>, Michael D. Ward<sup>a\*</sup>

<sup>a</sup>Department of Chemistry and the Molecular Design Institute, New York University, 100 Washington Square East, New York, NY 10003. <sup>b</sup>ChemMatCARS, Center for Advanced Radiation Sources, The University of Chicago, Lemont, Illinois 60439, United States

---

**Abstract:** 5-Methyl-2-[(2-nitrophenyl)amino]-3-thiophenecarbonitrile is a crystalline compound rich in conformational polymorphs largely owing to the flexible torsion angle that leads to distinct colors, earning it the moniker ROY (Red-Orange-Yellow). Guanidinium organosulfonate hydrogen-bonded frameworks form six crystalline inclusion compounds with ROY, described here, in which the framework limits conformational twisting out of plane. Three of the six inclusion compounds enforce greater planarity and  $\pi$ -conjugation than any of nine ROY polymorphs that have been characterized by single crystal X-ray diffraction.

---

*E-mail: mdw3@nyu.edu*

## Introduction


Hydrogen bonding guides the assembly of molecular components into organized supramolecular structures, from natural systems such as DNA to the design and synthesis of new materials. This themed issue is dedicated to Jean-Marie Lehn, who demonstrated in the early 1990s how hydrogen bonding can generate assemblies by molecular design, including cocrystals with molecular “strands” and “ribbons”<sup>1,2</sup> and polymeric liquid crystals.<sup>3</sup> The concepts emerged in parallel with other reports that illustrated the utility of hydrogen bonds in materials design, from hydrogen-bonded supramolecular polymers<sup>4,5</sup> to hydrogen-bonded assemblies and frameworks,<sup>6, 7, 8, 9, 10</sup> the subject of this contribution. Nearly contemporaneous with the honoree, Etter and Ward reported a series of compounds based on a 2D network assembled through complementary hydrogen-bonding between guanidinium (G) and sulfonate (S) groups of organosulfonates.<sup>11</sup> Since that time, our laboratory has created a large family of GS frameworks built from this 2D network with cavities having sizes and shapes that can be regulated by the choice of organosulfonate and the framework architecture, enabling the encapsulation of a large variety of guest molecules,<sup>12,13</sup> including laser dyes and other luminophores.<sup>14, 15</sup> Supramolecular hosts have been reported to exert influence on the conformation of their flexible guest molecules, suggesting an opportunity to regulate crystal properties through manipulation of guest conformation using the diverse library of GS frameworks.<sup>16,17</sup> Herein, we describe the inclusion of 5-methyl-2-[(2-nitrophenyl)amino]-3-thiophenecarbonitrile – also known as ROY for its Red, Orange and Yellow crystal polymorphs – in

various GS frameworks (Fig. 1). ROY has become the most vivid example of *conformational polymorphism*, a term coined by Bernstein and Hagler in 1978,<sup>18</sup> wherein crystal packing forces and molecular conformation are strongly interdependent, sometimes resulting in differently colored crystal forms.<sup>19, 20, 21, 22</sup> ROY molecules sequestered in GS host frameworks focuses conformational analysis on the interactions of the chromophores with the framework, as opposed to interactions with one another.

## Results and Discussion

ROY is associated with eleven polymorphs, including nine with associated crystal structures.<sup>23,24,25,26,27,28,29</sup> ROY owes its rich polymorphic behavior to the conformational flexibility of the S-C-N-C linkage, characterized by the torsion angle  $\theta_{\text{thio}}$  (Fig. 1B). The ROY polymorphs are distinguished by their lattice energy, crystal entropy and conformational energy.<sup>24</sup> The  $\theta_{\text{thio}}$  values influence the conjugation in ROY, leading to the range of colors among the polymorphs, For example,  $\theta_{\text{thio}} = 21.7^\circ$  for the red form,  $34.0 < \theta_{\text{thio}} < 52.6$  for the orange forms, and  $104.1^\circ < \theta_{\text{thio}} < 112.8^\circ$  for the yellow forms.

GS frameworks have proven highly versatile for guest inclusion because of the unusual persistence of two-dimensional (2D) guanidinium sulfonate (GS) network, which usually adopts a quasi-hexagonal symmetry owing to complementary 3-fold symmetry and hydrogen bond donors and acceptors. The pendant organic substituents attached to the sulfonate moiety project from the GS network, serving as pillars (for disulfonates with  $-\text{SO}_3^-$  groups on opposite ends) that support lamellar stacking as well as inclusion cavities between the sheets. The resilience of the GS network to a wide range of pillars and guests can be attributed to the strength of the charge-assisted hydrogen bonds and a unique structural compliance through puckering (defined by  $\theta_{\text{IR}}$ , Figure 1) of the GS sheet about a hydrogen-bonded hinge connecting adjacent GS ribbons, which provides a “shrink-wrapping” pathway for achieving close packing with retention of the hydrogen bond connectivity in the GS network. Moreover, the 2D character of the GS network permits an indefinite number of “projection topologies” defined by the pattern of “up-down” orientations of the organosulfonate groups from opposite sides of each GS sheet. This enables the lamellar architectures to form inclusion cavities with various sizes and shapes – as a consequence of templating by the guest molecules during crystal assembly – thereby accommodating a wide range of guests. The frameworks alone typically are colorless (guanidinium azobenzenedisulfonate,  $\text{G}_2\text{ABDS}$ , is an exception here), suggesting that the GS frameworks can sequester ROY molecules and enable determination of its conformation and associated color in a sequestered environment.




**Fig. 1.** (A) The quasi-hexagonal GS hydrogen bonded sheet illustrating the hydrogen-bonded major ribbons (one of these is highlighted in grey). (B) Molecular structures of the components in the guanidinium organosulfonate frameworks used herein and the molecular structure of ROY. The S-C-N-C torsion angle is denoted as  $\theta_{\text{thio}}$ . The numbers beneath the organosulfonate pillars correspond to their respective ROY inclusion compounds.

Crystallizations of  $\text{GS} \supset (\text{ROY})$  inclusion compounds were performed at the microscale by slow evaporation of solvent from solutions containing the guanidinium organosulfonate apohost and ROY. ROY crystals often formed concomitant with the inclusion compound; therefore crystallization was performed with an excess of the apohost to favor inclusion compound formation (see Supporting Information). The concomitant formation of ROY crystals prevented confirmation of inclusion compound stoichiometry by NMR spectroscopy and complicated determination of inclusion compound phase purity by powder X-ray diffraction (PXRD). Although the possibility of

polymorphism can never be excluded, polymorphism has never been observed in guanidinium organosulfonate compounds, and there was no evidence of polymorphism by visual inspection and Raman spectroscopy among the single crystals of each inclusion compound.

For example, crystallization of ROY with guanidinium 4,4'-biphenyldisulfonate ( $G_2BPDS$ ) by slow evaporation of methanol:acetonitrile solutions containing the dissolved  $G_2BPDS$  apohost afforded single crystals of  $G_2BPDS \supset (ROY)_{2/3}$  (**1**) as orangish-red {010} plates (Fig. S1, Table 1, Table S2). Single crystal X-ray diffraction confirmed the bilayer architecture (Fig. 2, Fig. S2), in which the long axes of the ROY molecules are aligned along one-dimensional channels flanked by the BPDS pillars (Fig. 3). Despite the non-integral stoichiometry, the ROY molecules are commensurate with the channel axis, with two ROY molecules commensurate with three pillars along the GS ribbons. The torsion angle  $\theta_{thio}$  of the ROY molecules is  $10.5^\circ$ , smaller than that of the red form R of ROY ( $21.7^\circ$ ), signaling more  $\pi$  conjugation of the phenyl and thiophene rings. The single crystal structure suggests that the near-planar conformation of ROY is a consequence of enforcement due to confinement in the narrow 1D channel (*ca.* 6.5 Å).




**Fig. 2.** Schematic representation of (A) bilayer, (B) zigzag and (C) double brick architectures in GS frameworks reported here.

**Table 1.** Structural and spectroscopic data for  $GS \supset ROY$  inclusion compounds.

	$G_2BPDS$ (1)	$G_2BSPE$ (2)	$G_2ABDS$ (3)	$G_2SDS$ (4)	$G_2BSPOE$ (5)	$G_2ADS$ (6)
Color	Red	Red	Red	Red	Red	Yellow
Architecture	Bilayer	Bilayer	Bilayer	Bilayer	Double-brick	Zigzag
$\theta_{thio}$ (degrees)	10.53	14.32/14.74	16.15/17.33	24.13	32.56	56.38
$\nu_{CN}$ ( $cm^{-1}$ )	2208	2215	2205	2215	2224	2228

The bilayer architecture also was observed for the G<sub>2</sub>BSPE, G<sub>2</sub>ABDS and G<sub>2</sub>SDS inclusion compounds with ROY. In these cases, however, the longer pillars result in the long axis of ROY aligned nearly parallel to the long axis of the pillars, reminiscent of enforced alignment of oligothiophene guests in GS frameworks.<sup>30</sup> Crystallization of ROY with guanidinium 1,2-bis(4-sulfonatophenyl)ethane (G<sub>2</sub>BSPE) afforded single crystals of G<sub>2</sub>BSPE⊃ROY (**2**) as red {001} plates (Fig. S1), along with a mixture of several ROY polymorphs. Single crystal X-ray diffraction revealed the bilayer architecture (Fig. S3), with torsion angles for the two ROY molecules in each asymmetric unit of  $\theta_{\text{thio}} = 14.3^\circ$  and  $14.7^\circ$ . In this case, the long axes of the ROY guests are nearly parallel to the long axes of the pillars.



**Fig. 3.** Crystal structures of inclusion compounds **1 - 6** (top to bottom) and their respective views from the top (second column), their respective ROY conformations (third column).

Crystallization with guanidinium azobenzenedisulfonate ( $G_2ABDS$ ) produced single crystal slabs of  $G_2ABDS \supset (ROY)_{3/4}(\text{methanol})_{1/4}$  (**3**) as (101) plates (Fig. S1). Although these crystals were red, azobenzene alone is red masking the true color of ROY. Nonetheless, ROY in **3** adopted two conformations with torsion angles of  $16.2^\circ$  and  $17.3^\circ$ , which would be consistent with the “red” form of ROY. Although this compound adopts the bilayer architecture (Fig. 3, Fig. S4), the unit cell ( $a = 35.698(9)$  Å;  $b = 7.334(2)$  Å;  $c = 48.5788(13)$  Å) is unusually large and the ROY guest arrangement is unusual for GS inclusion compounds. The large lattice constant along the  $\langle 101 \rangle$  direction is a consequence of unusual ordering of the guest molecules, wherein each channel contains alternating ROY tetrads and ROY pairs. This is a consequence of an unusual hydrogen-bonding pattern in the GS sheets consisting of a repeating pattern of four GS ribbons with the customary quasi-hexagonal motif connected by hydrogen bonds through the so-called “shifted ribbon” motif, which is occasionally observed in GS compounds (Fig. 3, Fig. S5). The nitro groups of ROY molecules in the tetrad are oriented opposite to the nitro groups of the ROY pairs. Crystallization with guanidinium 4,4'-stilbenedisulfonate ( $G_2SDS$ ) afforded single crystal  $G_2SDS \supset ROY$  (**4**) as  $\langle 100 \rangle$  needles (Fig. S1). Although **4** crystallizes in a bilayer architecture (Fig. 3, Fig. S6), the torsion angle  $\theta_{\text{thio}}$  of  $24.1^\circ$  is considerably larger than observed in compounds **1** – **3**. The long axes of ROY, like in BSPE frameworks, are parallel with the long axes of pillars.


Crystallization with guanidinium 1,2-bis(4-sulfonatophenoxy)ethane ( $G_2BSPOE$ ) afforded single crystals of  $G_2BSPOE \supset ROY$  (**5**) as red  $\langle 100 \rangle$  needles (Fig. S1), but with the double-brick architecture (Fig. 2, Fig. 3, Fig. S7), in which the projection of pillars on pairs of adjacent GS ribbons alternate across each GS sheet. The formation of the double-brick framework may be attributed to the large volume of the BSPOE pillar, which can frustrate inclusion of ROY in the smaller inclusion cavities of the bilayer architecture. The flexibility of BSPOE appears important, however, as the GS sheet puckers significantly such that the framework conforms to the ROY guests, which are nestled in pockets surrounding by two pairs of BSPOE pillars. The torsion angle  $\theta_{\text{thio}} = 33.2^\circ$ , is significantly higher than those in bilayer architectures. Unlike the aforementioned compounds, the ROY thiophene ring is substantially offset from the phenyl ring of the BSPOE pillar, suggested negligible  $\pi$ - $\pi$  interactions and structure-directing influence.

Crystallization with guanidinium 2,6-anthracene disulfonate ( $G_2ADS$ ) afforded single crystals of  $G_2ADS \supset (ROY)_2$  (**6**) as yellow  $\{001\}$  plates (Fig. S1), with the orthorhombic zigzag brick architecture (Fig. 2, Fig. 3, Fig. S8), in which the projections of the ADS pillars alternates in a zigzag manner across each GS sheet. This architecture, which typically is associated with larger inclusion cavities compared with the bilayer architecture, is likely a manifestation of the larger volume and greater rigidity of the ADS pillar compared with BPDS ( $185 \text{ \AA}^3$  versus  $166 \text{ \AA}^3$ ). Pairs of ROY molecules were encapsulated in the pockets flanked by ADS pillars. The torsion angle is  $\theta_{\text{thio}} = 56.4^\circ$ . The ROY thiophene ring is nearly



perpendicular to its neighboring ADS pillar, indicating negligible  $\pi$ - $\pi$  interaction and structure-directing influence.

The  $\theta_{thio}$  values for the included ROY molecules are in the range  $10.5^\circ \leq \theta_{thio} \leq 56.4^\circ$ , similar to the  $\theta_{thio}$  values for the red and orange ROY polymorphs. The Raman spectra for **1** – **6** revealed  $\nu_{CN}$  stretching mode frequencies that decreased with decreasing torsion angle  $\theta_{thio}$ , aligned with the trend reported for the red and orange ROY polymorphs (Fig. 4) and attributed to increased  $\pi$ -conjugation as  $\theta_{thio}$  approaches zero.<sup>28</sup> The confinement of the inclusion cavities in the GS frameworks exerts packing forces that result in  $\theta_{thio}$  values far below those for the yellow forms or the recently reported “pumpkin orange” form.<sup>29</sup>  $\theta_{thio}$  is smaller in inclusion compounds **1** – **3** than in the red ROY polymorph, which has the lowest value of  $\theta_{thio}$  among the polymorphs. This molecular flattening of ROY can be attributed to the ability of GS inclusion compounds to “shrink wrap” around the guests and achieve close packing through variable pillar conformations, puckering of the GS sheet, and adopting different framework architectures, all on display in compounds **1** – **6**. This feature is evident from the nearly uniform packing fractions of the inclusion compounds, which average  $0.70 \pm 0.02$ . (Table S2). Moreover, close inspection of the crystal structures of compounds **1** - **4**, which have the smallest values of  $\theta_{thio}$ , reveals near-parallelism of (pillar)phenyl-ROY(phenyl) and (pillar)phenyl-ROY(thiophene) planes, with substantial ring-ring ( $\pi$ - $\pi$ ) overlap, and interplanar distances comparable or somewhat less than the sum of the van der Waals radii for these rings (Fig. S9). In contrast, only the (pillar)phenyl-ROY(phenyl) rings are parallel in **5** and **6**, enabling larger values of  $\theta_{thio}$ , decreased  $\pi$ -conjugation, and larger  $\nu_{CN}$  stretching frequencies. Vibrational Stark spectroscopy has demonstrated that the vibrational frequency of nitriles, especially aromatic nitriles, can be sensitive to the surrounding electronic field.<sup>31</sup> The  $\pi$ - $\pi$  interactions between the pillar phenyl rings and the ROY thiophene ring in compounds **1** – **4**, as well as the coerced planarity from confinement, likely contribute to the trend in  $\nu_{CN}$  stretching frequencies.



**Fig. 4.** Dependence of the  $\nu_{CN}$  stretching frequency on the ROY torsion angle,  $\theta_{thio}$ , for inclusion compounds **1** – **6** and for the ROY polymorphs (denoted by their reported labels).

## Conclusions

Molecular building blocks with hydrogen bonding substituents have now produced a tremendous number of supramolecular assemblies and molecular frameworks. The guanidinium organosulfonate frameworks continue to surprise with their versatility, encapsulating a wide range of guest molecules, here confining ROY in channels and pockets that prevent severe conformational twisting out of plane. Among the nine solved crystal structures of ROY, the torsion angle  $\theta_{thio}$  ranges from  $21.7^\circ$  to  $112.7^\circ$ . The GS frameworks here, however, constrain the twisting of ROY, with an upper limit of  $\theta_{thio} = 56.4^\circ$  in one of the inclusion compounds and three others with a value of  $\theta_{thio}$  less than the lowest value among the polymorphs. These observations suggest that the properties of chromophores can be regulated by design of suitable molecular frameworks, particularly those that are amenable to adjustments in framework metrics with retention of generic architecture.

## Electronic Supplementary Information (ESI) available

Experimental and characterization details and additional figures. Crystal Structure available in the CCDC 1992984-1992989. For ESI and crystallographic data in CIF see DOI: 10.1039/x0xx00000x

## Acknowledgements

The authors acknowledge the support of the Materials Research Science and Engineering Center (MRSEC) Program of the National Science Foundation under Award Number DMR-1420073, the National Science Foundation through DMR-1308677, and the NSF Chemistry Research Instrumentation and Facilities Program (CHE-0840277). Synchrotron X-ray data were collected at the ChemMatCARS Sector 15 at the Advanced Photon Source (APS). NSF's ChemMatCARS Sector 15 is principally supported by the Divisions of Chemistry (CHE) and Materials Research (DMR), National Science Foundation, under grant number NSF/CHE-1834750. Use of the Advanced Photon Source, an Office of Science User Facility operated for the U.S. Department of Energy (DOE) Office of Science by Argonne National Laboratory, was supported by the U.S. DOE under Contract No. DE-AC02-06CH11357. MT is grateful for a Herman and Margaret Sokol graduate fellowship.

## Conflicts of Interest

There are no conflicts to declare.

## Notes and References

1. J.-M. Lehn, M. Mascal, A. DeCian, J. Fischer, Molecular recognition directed self-assembly of ordered supramolecular strands by cocrystallization of complementary molecular components, *J. Chem. Soc. Chem. Commun.*, 1990, 479.
2. J.-M. Lehn, M. Mascal, A. DeCian, J. Fischer, Molecular ribbons from molecular recognition directed self-assembly of self-complementary molecular components, *J. Chem. Soc. Perkin Trans. 2*, 1992, 461.
3. C. Fouquey, J.-M. Lehn, A.-M. Levelut, Molecular recognition directed self-assembly of supramolecular liquid crystalline polymers from complementary chiral components, *Adv. Mater.*, 1990, **2**, 254.
4. L. Brunsveld, B. J. B. Folmer, E. W. Meijer, R. P. Sijbesma, Supramolecular polymers, *Chem. Rev.*, 2001, **101**, 4071.
5. R. P. Sijbesma, F. H. Beijer, L. Brunsveld, B. J. B. Folmer, J. H. K. Ky Hirschberg, R. F. M. Lange, J. K. L. Lowe, E. W. Meijer, Reversible polymers formed from self-complementary monomers using quadruple hydrogen bonding, *Science*, 1977, **278**, 1601.
6. P. Brunet, M. Simard, J. D. Wuest, Molecular tectonics. Porous hydrogen-bonded networks with unprecedented structural integrity, *J. Am. Chem. Soc.*, 1997, **119**, 2737.
7. M. Simard, D. Su, J. D. Wuest, Use of hydrogen bonds to control molecular aggregation. Self-assembly of three-dimensional networks with large chambers, *J. Am. Chem. Soc.*, 1991, **113**, 4696.

8. M. W. Hosseini, Molecular Tectonics: From simple tectons to complex molecular networks, *Acc. Chem. Res.*, 2005, **38**, 313.
9. F. Garcia-Tellado, S. J. Geib, S. Goswami, A. D. Hamilton, Molecular recognition in the solid state: Controlled assembly of hydrogen-bonded molecular sheets, *J. Am. Chem. Soc.*, 1991, **113**, 9265.
10. O. Felix, M. W. Hosseini, A. De Cian, Design of 2-D hydrogen bonded molecular networks using pyromellitate dianion, cyclic bisamidinium dication as complementary tectons, *Solid State Sci.*, 2001, **3**, 789.
11. V. A. Russell, M. C. Etter, M. D. Ward, Layered materials by molecular design: Structural enforcement by hydrogen bonding in guanidinium alkane- and arenesulfonates, *J. Am. Chem. Soc.*, 1994, **116**, 1941.
12. K. T. Holman, A. M. Pivovar, J. A. Swift, M. D. Ward, Metric Engineering of soft molecular host frameworks, *Acc. Chem. Res.*, 2001, **34**, 107.
13. T. Adachi, M. D. Ward, Versatile and hydrogen-bonded host frameworks, *Acc. Chem. Res.*, 2016, **49**, 2669.
14. M. Handke, T. Adachi, C. Hu, M. D. Ward, Encapsulation of isolated luminophores within supramolecular cages, *Angew. Chem. Int. Ed. Engl.*, 2017, **56**, 13901.
15. A. C. Soegiarto, M. D. Ward, Directed organization of dye aggregates in hydrogen-bonded host frameworks, *Cryst. Growth Des.*, 2009, **9**, 3803.
16. B.-Q. Ma, Y. Zhang, P. Coppens, Multiple conformations of benzil in resorcinarene-based supramolecular host matrixes, *J. Org. Chem.*, 2003, **68**, 9467.
17. B.-Q. Ma, P. Coppens, Variable conformation of benzophenone in a series of resorcinarene-Based supramolecular frameworks, *Cryst. Growth Des.*, 2004, **4**, 1377.
18. J. Bernstein, A. T. Hagler, Conformational Polymorphism. The influence of crystal structure on molecular conformation, *J. Am. Chem. Soc.* 1978, **100**, 673.
19. P. H. Toma, M. P. Kelley, T. B. Borchardt, S. R. Byrn, B. Kahr, Chromoisomers and Polymorphs of 9-Phenylacridinium Hydrogen Sulfate, *Chem. Mater.*, 1994, **6**, 1317.
20. B. A. Nogueira, C. Castiglioni, R. Fausto, Color polymorphism in organic crystals, *Commun. Chem.*, 2020, **3**, 34.
21. S. R. Byrn, D. Curtin, I. Paul, X-ray crystal structures of the yellow and white forms of dimethyl 3,6-dichloro-2,5-dihydroxyterephthalate and a study of the conversion of the yellow form to the white form in the solid state, *J. Am. Chem. Soc.*, 1972, **94**, 890.
22. M. Tristani-Kendra, C. J. Eckhardt, J. Bernstein, E. Goldstein, Strong coupling in the optical spectra of polymorphs of a squarylium dye, *Chem. Phys. Lett.*, 1983, **98**, 57.
23. G. Stephenson, T. Borchardt, S. Byrn, J. Boyer, S. Snorek, C. Bunnell, L. Yu, Conformational and color polymorphism of 5-methyl-2-[(2-nitrophenyl) amino]-3-thiophenecarbonitrile, *J. Pharm. Sci.*, 1995, **84**, 1385.
24. L. Yu, A. G. Stephenson, C. A. Mitchell, C. A. Bunnell, S. V. Snorek, J. J. Bowyer, T. B. Borchardt, J. G. Stowell, S. R. Byrn, Thermochemistry and conformational polymorphism of a hexamorphic crystal system, *J. Am. Chem. Soc.*, 2000, **122**, 585.
25. S. Chen, H. Xi, L. Yu, Cross-nucleation between ROY polymorphs, *J. Am. Chem. Soc.*, 2005, **127**, 17439.
26. S. Chen, I. Guzei, L. Yu, New polymorphs of ROY and new record of coexisting polymorphs of solved structures, *J. Am. Chem. Soc.*, 2005, **127**, 9881.
27. M. Tan, A. G. Shtukenberg, S. Zhu, W. Xu, E. Dooryhee, S. M. Nichols, M. D. Ward, B. Kahr, Q. Zhu, ROY revisited, again: the eighth solved structure, *Faraday Discuss.*, 2018, **211**, 477.
28. L. Yu, Polymorphism in molecular solids: an extraordinary system of red, orange, and yellow crystals, *Acc. Chem. Res.*, 2010, **43**, 1257.
29. K. S. Gunshurst, J. Nyman, S. X. M. Boerrigter, The PO13 crystal structure of ROY, *CrystEngComm*, 2019, **21**, 1363.

- 
30. A. C. Soegiarto, A. Comotti, M. D. Ward, Controlled orientation of polyconjugated guest molecules in tunable host cavities, *J. Am. Chem. Soc.*, 2010, **132**, 14603.
31. S. S. Andrews, S. G. Boxer, Vibrational Stark effects of nitriles II. Physical origins of Stark effects from experiment and perturbation models, *J. Phys. Chem. A*, 2002, **106**, 469.

### Table of Contents Graphic and Description

The conformation of 5-Methyl-2-[(2-nitrophenyl)amino]-3-thiophenecarbonitrile, known as ROY for the Red-Orange-Yellow of its crystal polymorphs, is restricted by confinement in guanidinium organosulfonate hydrogen-bonded frameworks.

