

Cite this: *Chem. Sci.*, 2019, 10, 10733

All publication charges for this article have been paid for by the Royal Society of Chemistry

# The butterfly effect in bisfluorenylidene-based dihydroacenes: aggregation induced emission and spin switching†

Xiaodong Yin,<sup>ab</sup> Jonathan Z. Low,<sup>b</sup> Kealan J. Fallon,<sup>b</sup> Daniel W. Paley<sup>b</sup> and Luis M. Campos<sup>\*b</sup>

Linear acenes are a well-studied class of polycyclic aromatic hydrocarbons and their established physical properties have led to their widespread application across the field of organic electronics. However, their quinoidal forms – dihydroacenes – are much less explored and exhibit vastly different photophysical and electronic properties due to their non-planar, cross-conjugated nature. In this work, we present a series of difluorenylidene dihydroacenes which exhibit a butterfly-like structure with a quinoidal skeleton, resulting in comparatively higher optical gaps and lower redox activities than those of their planar analogs. We found that these compounds exhibit aggregation induced emission (AIE), activated through restriction of the “flapping” vibrational mode of the molecules in the solid state. Furthermore, anthracene-containing dihydroacenes exhibit thermally activated ground-state spin switching as evidenced by planarization of the acene core and diradical activity recorded by EPR. These two characteristics in this relatively unexplored class of materials provide new insights for the design of multifunctional materials.

Received 16th August 2019  
Accepted 4th October 2019

DOI: 10.1039/c9sc04096j

rsc.li/chemical-science

## Introduction

Polycyclic aromatic hydrocarbons (PAHs) are ubiquitous materials – they have been found in space<sup>1</sup> and have made their way into a variety of optoelectronic devices<sup>2,3</sup> and strong composite materials.<sup>4,5</sup> Oligoacenes are a subclass of PAHs that have sparked interest in organic electronics<sup>6,7</sup> due to the realization of their potential as active materials in organic field effect transistors (OFETs),<sup>8,9</sup> organic light-emitting diodes (OLEDs),<sup>10,11</sup> and third-generation photovoltaics.<sup>12–16</sup> In most of these systems, fine-tuning the chemistry of the core structure and intermolecular interactions is essential for stability and the overall performance of the materials. These molecules are generally flat and exhibit strong pi-stacking interactions that can be beneficial or detrimental for particular types of applications. For example, aggregation can impair optical emission, but it can be beneficial for directional charge transport.<sup>17</sup> In this vein, contorted PAHs can be tuned for practical applications. Additionally, introducing substituents with steric hindrance and conformational degrees of freedom can drastically alter

their physical properties. For example, acenes with a butterfly structure were originally studied a couple of decades ago, and the butterfly structure can be changed to a flat structure by electrochemical oxidation.<sup>18–21</sup>

Through precise molecular engineering of the core structure of oligoacenes, Tang and co-workers have shown that quinoidal derivatives of oligoacenes can exhibit aggregation induced restricted intramolecular vibration (RIV).<sup>22,23</sup> AIE is counterintuitive since aggregation tends to quench photoluminescence of organic chromophores.<sup>24–26</sup> For example, bis(diphenylmethylene)dihydroacenes are non-emissive in solution, but upon aggregation, they fluoresce in the solid state (Fig. 1A). This property does not translate to the fluoreno-quinoidal oligomers. In fact, fluoreno adducts are prone to stabilizing the radical character, as elegantly demonstrated by Wudl,<sup>27</sup> Wu,<sup>28</sup> Haley,<sup>29</sup> and others.<sup>30,31</sup> The fluoreno-derivative of Chichibabin's radical (Fig. 1B) is non-emissive; however, in its most stable structure, the diradical is readily characterized. This diradical character is not surprising, given that extending anthracene oligomers along the 9,10-positions disrupts pi-orbital overlap, thus forcing aromatization of the central ring and yielding the stable open-shell fluorenyl groups shown in Fig. 1B as the most favorable resonance structure. Such phenomena motivated our interest in investigating the structure–property relationship, extending the oligoacene core with bisfluoreno adducts. Interestingly, we found that the bisfluorenoanthracene in Fig. 1C exhibits both AIE in the solid state and the thermally accessible open-shell diradicaloid character.

<sup>a</sup>Beijing Key Laboratory of Photoelectric/Electrophotonic Conversion Materials, School of Chemistry and Chemical Engineering, Beijing Institute of Technology, Beijing 102488, P. R. China

<sup>b</sup>Department of Chemistry, Columbia University, New York, New York 10027, USA. E-mail: lcampos@columbia.edu

† Electronic supplementary information (ESI) available. CCDC 1947366–1947368. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc04096j

This journal is © The Royal Society of Chemistry 2019

Table 1 Summary of photophysical properties of difluorenylidene dihydroacenes

Entry	$\lambda_{\text{abs}}(\text{sol})^a/(\text{solid})^b/\text{nm}$	$\lambda_{\text{em}}(\text{sol})^a$	$\lambda_{\text{em}}(\text{solid})^b$	$\Phi_{\text{sol}}^a$	$\Phi_{\text{agg}}^b$	$\tau/\text{ns}(\text{sol})$	$\tau/\text{ns}(\text{solid})$
<b>bFA</b>	418/441	409, 431	512	0.8%	11.2%	0.30	8.84
<b>bFT</b>	413/440	513	511, 547	0.6%	24.9%	0.29	8.49
<b>bFP</b>	413/438	474	513	0.1%	18.5%	1.11	6.00

<sup>a</sup> In DCM solution; absolute quantum yields of these compounds were determined with an integrating sphere (Horiba Fluorolog-3). <sup>b</sup> Thin film on a quartz slide; absolute quantum yields of these compounds were determined with an integrating sphere (Horiba Fluorolog-3).

introduced into THF solutions, fluorescence from the samples increases dramatically due to the formation of molecular aggregates. Fig. 2A shows a series of photos of **bFA**, **bFT**, and **bFP** in different THF/water mixtures, clearly demonstrating the increase in fluorescence as the percentage of water increases. Fig. 2B shows the UV-Vis and fluorescence spectra of **bFT** as an example, which indicate the dramatic increase of the fluorescence of **bFT** in THF/water = 1/9 in comparison with that in pure THF. The UV-Vis and fluorescence spectra (Fig. S4†) for all three compounds in solutions containing different percentages of water were recorded. In mixtures containing 10–50% water by volume, the emission intensity remains weak for all three compounds, indicating that they were well solvated. When the percentage of water reaches 60%, the emission intensity shows a sharp increase that is greater than two orders of magnitude. Furthermore, the fluorescence lifetime ( $\tau$ ) in the solid state is longer than that in solution for all molecules, and in **bFA** and **bTA** dramatically so (Table 1 and Fig. S5†).

Combining the UV-Vis and fluorescence spectra, the quantum yield of the three compounds can be plotted against the percentage of water (Fig. 2c). The extent of the enhancement arising from aggregation-induced emission ( $\alpha_{\text{AIE}}$ ) can be defined as the quantum yield ratio of the aggregates to the solution ( $\Phi_{\text{agg}}/\Phi_{\text{sol}}$ ). As a guide,  $\alpha_{\text{AIE}} \sim 344$  is seen in tetraphenyl ethylene (TPE), a typical AIE chromophore.<sup>41</sup> In **bFA**, **bFT** and **bFP**,  $\alpha_{\text{AIE}}$  is 9.4, 40 and 185, respectively (Fig. 2c). The observed AIE in all three compounds is remarkable given that there are no rotational degrees of freedom, which is generally the key mechanism by which AIE occurs in conventional systems.

Here, the widely invoked restricted intramolecular rotation (RIR) model in AIE<sup>22</sup> cannot explain our observations. Thus, we turn to density functional theory (DFT) calculations to investigate the origin of the observed AIE. We first calculated the ground and excited state structures of the compounds in the gas phase using the Becke3–Lee–Yang–Parr level of theory with a 6-31G\*\* basis set (B3LYP/6-31G\*\*). Fig. 3a shows that for **bFP**, the bending angles of both the acene core ( $\theta_{\text{acene}}$ ) and the fluorenes ( $\theta_{\text{fluorene}}$ ) are more relaxed toward planarity in the excited state ( $S_1$ , blue) than in the ground state (red) by 6° and 17°, respectively. We compare these results to calculations confined within the crystal structure as shown in Fig. 3b, obtained using a combined quantum mechanics/molecular mechanics (QM/MM) method, ONIOM (B3LYP/6-31G\*\*: UFF), which is a reliable method of obtaining molecular structure information, embedded within a crystalline lattice (see details in the ESI†).<sup>23</sup> The result shows that the ground state and excited state

structures of **bFP** in the crystal are practically identical. Table S2† summarizes the computational results arising from structural changes for all three compounds under investigation. In all cases, a similar trend is observed where the ground state is more bent than the excited state in the gas phase, while both conformations are similar in a crystalline solid. The DFT results indicate that the AIE may result from restricted intramolecular vibrations (RIVs) only. This type of AIE has been observed before, but with only a slight enhancement in  $\alpha_{\text{AIE}}$ , e.g., ca. 60 for **BDBA**<sup>41</sup> and ca. 30 for **COTh-TMS**<sup>42</sup> with all carbon atoms  $sp^2$  hybridized. Our results are comparable to changes in the quantum yield reported for AIE in chromophores that exhibit marked conformational changes, such as RIR.<sup>43</sup> These results suggest that AIE may be common for other chromophores where RIR is absent, but RIV may result in unprecedented PL in the solid state.

Considering that these chromophores have resonance structures where the quinoidal structures can flatten and rehybridize into the open-shell diradicals, we sought to investigate the temperature-dependent structure–property relationship between the closed-shell and open-shell ground state structures (Fig. 1C).<sup>28,44</sup> Fig. 4a shows the <sup>1</sup>H-NMR spectrum of **bFA** as a function of temperature varying from 300 K to 385 K. We observe that the peaks of the  $H_a$  doublet (8.1 ppm) and  $H_b$  triplet (7.2 ppm, which overlaps with the solvent peak and a quartet) broaden considerably when the solution is heated beyond 373 K. Furthermore, electron paramagnetic resonance (EPR) spectroscopy performed at 430 K in dichlorobenzene shows a signal that matches well with previous reports of radicals localized on fluorene moieties (Fig. 4b).<sup>28,45</sup> EPR of a solid sample of **bFA** was also obtained, but show much weaker signals even at a high temperature (450 K). Taken together, the NMR and EPR results indicate that **bFA** adopts an open shell configuration at high temperatures, forming a triplet diradical.

To gain insight into the structural changes of the **bFA** backbone upon conversion to the open diradical, we synthesized an asymmetric analog, **bFA-t**, so that the signal from  $H_a$  and  $H_{a'}$  protons would be split into two distinct doublets, as shown in Fig. 4c. These doublets appear at 8.0–8.2 ppm. Fig. 4c also shows that as the temperature is increased, the signals from  $H_a$  and  $H_{a'}$  coalesce, indicating that the fluorene moiety is able to rotate freely. The free rotation of the fluorene group supports the formation of an open-shell diradical, since the double bond between fluorene and acene gains significant single bond character during this transformation. From the NMR data shown in Fig. 4c, we calculate that the energy barrier







Fig. 2 (A) Photographic image of quartz vials containing difluorenylidene acenequinoids displaying fluorescence in THF/H<sub>2</sub>O solutions with different water fractions irradiated using a 365 nm UV lamp; (B) UV-Vis and photoluminescence spectra of bFT in THF and THF/water = 1/9; (C) plots of quantum yield as a function of water fraction with excitation at 380 nm; the inset is the aggregation induced emission efficiency ( $\Phi_{\text{aggr}}/\Phi_{\text{sol}}$ ); absolute quantum yields of these samples were determined with an integrating sphere (Horiba Fluorolog-3).

of the rotation ( $\Delta G^\ddagger$ ) in bFA-t is 17.3 kcal mol<sup>-1</sup> (Fig. S11†). In contrast to bFA, neither bFT nor bFP show line broadening of the proton signals as the temperature is increased; rather, the




Fig. 3 (a) Optimized structures of the ground state (S<sub>0</sub>, red) and excited state (S<sub>1</sub>, blue) conformations of bFP in the gas and crystal state with the B3LYP/6-31G\*\* and ONIOM (B3LYP/6-31G\*\* : UFF) methods, respectively; (b) the ONIOM model, with the high layer presented as ball-and-stick molecules and the low layer presented as wireframe molecules.




Fig. 4 (a) VT-<sup>1</sup>H NMR of bFA in tetrachloroethane-d<sub>2</sub>; (b) EPR data of bFA in the solid state at 300 K (black) and 450 K (red) and in dichlorobenzene ( $4 \times 10^{-2}$  M) at 430 K (blue); inset: spin density illustration of the T<sub>1</sub> state of bFA<sub>twist</sub> (twisted configuration) at the UMO6L/6-311G\*\* level of theory; (c) VT-<sup>1</sup>H NMR of bFA-t and the molecular structure of bFA-t.

signals are sharpened, and no signal of EPR was observed (see Fig. S12†), indicating that these molecules do not have a thermally accessible open-shell ground-state configuration. Besides, the barrier to rotation ( $\Delta G^\ddagger$ ) for bFT is estimated to be ca. 21.7 kcal mol<sup>-1</sup> by the NMR data (Fig. S13†), which is much higher than that of bFA-t.

It is noteworthy to report our observation of a diprotonated derivative of bFA-t (bFAH-t) with strong blue fluorescence as the main product of the microwave-assisted Stille coupling reaction at ~180 °C (see Fig. S14†). This is strong evidence of H-abstraction of a diradical intermediate of bFA-t at high temperature.<sup>46,47</sup> The UV-Vis absorbance of bFAH-t indicates that the core of this molecule has an aromatic anthracene structure with a finely split peak at ca. 400 nm. Moreover, this



Fig. 5 (a) Energy diagram of **bFA** with rotation of the fluorene moiety; (b) HOMOs of the fluorene twisted structure of **bFA** (**bFA**<sub>twist</sub>), **bFT** (**bFT**<sub>twist</sub>), and **bFP** (**bFP**<sub>twist</sub>), with the  $y_0$  value and bond length (\*) of the C–C bond between fluorene and acenes.

diprotinated compound can be converted back to the quinoid structure by oxidation with DDQ, as shown in Fig. S15.†

In order to understand the spin switching process, we computed the stationary points and transition state for **bFA**, as it changed into the open-shell structure from the closed-shell ground state (DFT, UB3LYP/6-31G\*\*, Fig. 5A). As the dihedral angle joining fluorene to anthracene was planarized, an energy saddle point was observed in the conformation where one fluorene is twisted relative to anthracene whilst the other one is not. The theoretical energy barrier of 18 kcal mol<sup>−1</sup> compares well with the experimental result of deuterium-NMR of **bFA-t** with the same aromatic core (see Fig. S11†). When both fluorenes are twisted, a relatively stable state is formed with a comparable energy level (+3 kcal mol<sup>−1</sup>) to that of the original state. Single point energy calculations of both the single twisted structure and double twisted structure were conducted at the UM06L/6-311G\*\* level of theory.

The biradical character of organic materials can be assessed by the biradical character index ( $y_0$ ) and indicates the

contribution of the open-shell resonance form to the overall ground state structure, ranging from a pure biradical ( $y_0 = 1$ ) to a purely closed shell ( $y_0 = 0$ ).<sup>48–51</sup> For **bFA**,  $y_0$  values of 0 and 0.16 were obtained for the single twisted structure and double twisted structure, respectively, indicating the open-shell character of the double twisted structure (**bFA**<sub>twist</sub>) as shown in Fig. 5b. In addition, the theoretical C–C double bond length in **bFA**<sub>twist</sub> between the fluorene and anthracene moieties shows a remarkable extension (1.44 Å) compared to that in the butterfly configuration (1.36 Å), indicating the open shell character from a structural point of view. Crucially, the triplet state of **bFA**<sub>twist</sub> is calculated to be lower in energy than the diradical singlet state (as shown in Fig. 5), indicating that intersystem crossing is thermodynamically favorable in this compound. The spin density of the T<sub>1</sub> state of **bFA**<sub>twist</sub> is mainly distributed on the fluorene moiety, especially on carbons a and b, which match very well with the signal broadening observed by VT-<sup>1</sup>H NMR (see Fig. 4b). Theoretical analysis was also conducted for the twisted structures of **bFT** (**bFT**<sub>twist</sub>) and **bFP** (**bFP**<sub>twist</sub>), and both of them exhibit a closed shell ground state electronic structure ( $y_0 = 0$ , see Fig. 5b). These calculations match well with our experimental observations. Furthermore, the carbon–carbon bond length between fluorene and acenes of **bFT**<sub>twist</sub> and **bFP**<sub>twist</sub> is ca. 1.41 Å which is more comparable to a typical double bond, indicating closed-shell quinoidal structures.

From this analysis we conclude that, at elevated temperatures, aromatization is a strong enough driving force to access the open shell ground state in the anthraquinoid **bFA**, whereas the longer acenequinoid structures in **bFT** and **bFP** do not possess sufficient aromatic character in their aromatic conformation to exhibit this trait. This finding is supported by Clar's theory of aromatic sextets, where higher order acene molecules exhibit greater stability as two small acenes *versus* one large acene system (e.g. 2 × naphthalene *vs.* 1 × pentacene for **bFP**).<sup>52,53</sup> Besides, since the open-shell character of **bFA** is strongly related to the rotation of fluorene moieties, the weak EPR signal in the solid state even at a high temperature can be assigned to the constriction of intramolecular motion.

## Conclusions

In summary, we synthesized a series of difluorenylidene dihydroacene compounds using a facile and efficient method. These compounds exhibit wider optical gaps than expected due to their bent, butterfly-like quinoid structures. All compounds exhibit aggregation induced emission, which is rarely seen in such a kind of structure without any freely rotatable functionalities. A flapping intramolecular vibration was identified as the mechanism based on theoretical analysis (QM/MM calculations). Furthermore, a transformation between the bent quinoid butterfly configuration and the aromatic planar configuration is thermally accessible in the anthracene-containing material. As a result, the electronic structure is also switchable between the closed shell at a low temperature and open shell at a high temperature, which is confirmed by VT-<sup>1</sup>H NMR and EPR. This phenomenon can be assigned to



rotation of the fluorene moieties at a high temperature, leading to an intermediate structure with a flattened anthracene skeleton and moderate diradical character. Conversely, this transition is not thermally accessible in longer acenequinoids, due to the lower driving force as a function of the lower aromaticity of the aromatic conformation of these acenes. We envisage that the structure–property relationship reported herein will create new pathways for designing novel multi-functional materials.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This work was funded by the National Science Foundation (DMR-1807580). K. J. F. acknowledges the funding by the US-UK Fulbright Commission and Lloyd's Tercentenary Research Foundation. Single crystal X-ray diffraction was performed at the Shared Materials Characterization Laboratory at Columbia University. Use of the SMCL was made possible by funding from Columbia University.

## Notes and references

- 1 F. J. Lovas, R. J. McMahon, J. U. Grabow, M. Schnell, J. Mack, L. T. Scott and R. L. Kuczkowski, *J. Am. Chem. Soc.*, 2005, **127**, 4345–4349.
- 2 J. Wu, W. Pisula and K. Müllen, *Chem. Rev.*, 2007, **107**, 718–747.
- 3 A. Narita, X. Y. Wang, X. Feng and K. Müllen, *Chem. Soc. Rev.*, 2015, **44**, 6616–6643.
- 4 Z. Qin, G. S. Jung, M. J. Kang and M. J. Buehler, *Sci. Adv.*, 2017, **3**, e1601536.
- 5 C. Lee, X. Wei, J. W. Kysar and J. Hone, *Science*, 2008, **321**, 385–388.
- 6 J. E. Anthony, *Chem. Rev.*, 2006, **106**, 5028–5048.
- 7 J. E. Anthony, *Angew. Chem., Int. Ed.*, 2008, **47**, 452–483.
- 8 K. Ito, T. Suzuki, Y. Sakamoto, D. Kubota, Y. Inoue, F. Sato and S. Tokito, *Angew. Chem., Int. Ed.*, 2003, **115**, 1191–1194.
- 9 A. Naibi Lakshminarayana, A. Ong and C. Chi, *J. Mater. Chem. C*, 2018, **6**, 3551–3563.
- 10 L. C. Picciolo, H. Murata and Z. H. Kafafi, *Appl. Phys. Lett.*, 2001, **78**, 2378–2380.
- 11 S. A. Odom, S. R. Parkin and J. E. Anthony, *Org. Lett.*, 2003, **5**, 4245–4248.
- 12 M. B. Smith and J. Michl, *Chem. Rev.*, 2010, **110**, 6891–6936.
- 13 S. N. Sanders, E. Kumarasamy, A. B. Pun, M. T. Trinh, B. Choi, J. Xia, E. J. Taffet, J. Z. Low, J. R. Miller, X. Roy, X. Y. Zhu, M. L. Steigerwald, M. Y. Sfeir and L. M. Campos, *J. Am. Chem. Soc.*, 2015, **137**, 8965–8972.
- 14 E. Kumarasamy, S. N. Sanders, A. B. Pun, S. A. Vaselabadi, J. Z. Low, M. Y. Sfeir, M. L. Steigerwald, G. E. Stein and L. M. Campos, *Macromolecules*, 2016, **49**, 1279–1285.
- 15 S. N. Sanders, E. Kumarasamy, A. B. Pun, K. Appavoo, M. L. Steigerwald, L. M. Campos and M. Y. Sfeir, *J. Am. Chem. Soc.*, 2016, **138**, 7289–7297.
- 16 M. Einzinger, T. Wu, J. F. Kompalla, H. L. Smith, C. F. Perkinson, L. Nienhaus, S. Wieghold, D. N. Congreve, A. Kahn, M. G. Bawendi and M. A. Baldo, *Nature*, 2019, **571**, 90–94.
- 17 V. C. Sundar, J. Zaumseil, V. Podzorov, E. Menard, R. L. Willett, T. Someya, M. E. Gershenson and J. A. Rogers, *Science*, 2004, **303**, 1644–1646.
- 18 N. Martin, L. Sánchez, C. Seoane, E. Ortí, P. M. Viruela and R. Viruela, *J. Org. Chem.*, 1998, **63**, 1268–1279.
- 19 M. C. Díaz, B. M. Illescas, N. Martín, R. Viruela, P. M. Viruela, E. Ortí, O. Brede, I. Zilbermann and D. M. Guldi, *Chem.–Eur. J.*, 2004, **10**, 2067–2077.
- 20 N. E. Gruhn, N. A. Macías-Ruvalcaba and D. H. Evans, *Langmuir*, 2006, **22**, 10683–10688.
- 21 S. L. Broman, C. L. Andersen, T. Jousselein-Oba, M. Mansø, O. Hammerich, M. Frigoli and M. B. Nielsen, *Org. Biomol. Chem.*, 2017, **15**, 807–811.
- 22 Z. He, L. Shan, J. Mei, H. Wang, J. W. Y. Lam, H. H.-Y. Sung, I. D. Williams, X. Gu, Q. Miao and B. Z. Tang, *Chem. Sci.*, 2015, **6**, 3538–3543.
- 23 Z. He, L. Zhang, J. Mei, T. Zhang, J. W. Y. Lam, Z. Shuai, Y. Q. Dong and B. Z. Tang, *Chem. Mater.*, 2015, **27**, 6601–6607.
- 24 J. Mei, N. L. C. Leung, R. T. K. Kwok, J. W. Y. Lam and B. Z. Tang, *Chem. Rev.*, 2015, **115**, 11718–11940.
- 25 Z. Zhao, H. Nie, C. Ge, Y. Cai, Y. Xiong, J. Qi, W. Wu, R. T. K. Kwok, X. Gao, A. Qin, J. W. Y. Lam and B. Z. Tang, *Adv. Sci.*, 2017, **4**, 1700005.
- 26 D. Wang, M. M. S. Lee, G. Shan, R. T. K. Kwok, J. W. Y. Lam, H. Su, Y. Cai and B. Z. Tang, *Adv. Mater.*, 2018, **30**, 1802105.
- 27 X. Gong, M. Tong, F. G. Brunetti, J. Seo, Y. Sun, D. Moses, F. Wudl and A. J. Heeger, *Adv. Mater.*, 2011, **23**, 2272–2277.
- 28 Z. Zeng, Y. M. Sung, N. Bao, D. Tan, R. Lee, J. L. Zafra, B. S. Lee, M. Ishida, J. Ding, J. T. López Navarrete, Y. Li, W. Zeng, D. Kim, K. W. Huang, R. D. Webster, J. Casado and J. Wu, *J. Am. Chem. Soc.*, 2012, **134**, 14513–14525.
- 29 G. E. Rudebusch, J. L. Zafra, K. Jorner, K. Fukuda, J. L. Marshall, I. Arrechea-Marcos, G. L. Espejo, R. Ponce Ortiz, C. J. Gómez-García, L. N. Zakharov, M. Nakano, H. Ottosson, J. Casado and M. M. Haley, *Nat. Chem.*, 2016, **8**, 753–759.
- 30 D. Beaudoin, O. Levasseur-Grenon, T. Maris and J. D. Wuest, *Angew. Chem., Int. Ed.*, 2016, **55**, 894–898.
- 31 K. Cocq, V. Maraval, N. Saffon-Merceron, A. Saquet, C. Poidevin, C. Lepetit and R. Chauvin, *Angew. Chem., Int. Ed.*, 2015, **54**, 2703–2706.
- 32 M. E. Howe and M. A. Garcia-Garibay, *J. Org. Chem.*, 2019, **84**, 9570–9576.
- 33 F. Schwarz, G. Kastlunger, F. Lissel, C. Egler-Lucas, S. N. Semenov, K. Venkatesan, H. Berke, R. Stadler and E. Lörtscher, *Nat. Nanotechnol.*, 2015, **11**, 170–176.
- 34 O. Kahn and C. Jay Martinez, *Science*, 1998, **279**, 44–48.
- 35 M. Mannini, F. Pineider, P. Saintavitt, C. Danieli, E. Otero, C. Sciancalepore, A. M. Talarico, M. Arrio, A. Cornia, D. Gatteschi and R. Sessoli, *Nat. Mater.*, 2009, **8**, 194–197.
- 36 T. Matsumoto, G. N. Newton, T. Shiga, S. Hayami, Y. Matsui, H. Okamoto, R. Kumai, Y. Murakami and H. Oshio, *Nat. Commun.*, 2014, **5**, 3865.



- 37 A. Bouseksou, G. Molnár and G. Matouzenko, *Eur. J. Inorg. Chem.*, 2004, 4353–4369.
- 38 P. Gütllich, A. Hauser and H. Spiering, *Angew. Chem., Int. Ed.*, 1994, **33**, 2024–2054.
- 39 M. R. Geraskina, A. T. Buck and A. H. Winter, *J. Org. Chem.*, 2014, **79**, 7723–7727.
- 40 C. Y. Chiu, B. Kim, A. A. Gorodetsky, W. Sattler, S. Wei, A. Sattler, M. Steigerwald and C. Nuckolls, *Chem. Sci.*, 2011, **2**, 1480–1486.
- 41 N. L. C. Leung, N. Xie, W. Yuan, Y. Liu, Q. Wu, Q. Peng, Q. Miao, J. W. Y. Lam and B. Z. Tang, *Chem.–Eur. J.*, 2014, **20**, 15349–15353.
- 42 Z. Zhao, X. Zheng, L. Du, Y. Xiong, W. He, X. Gao, C. Li, Y. Liu, B. Xu, J. Zhang, F. Song, Y. Yu, X. Zhao, Y. Cai, X. He, R. T. K. Kwok, J. W. Y. Lam, X. Huang, D. Lee Phillips, H. Wang and B. Z. Tang, *Nat. Commun.*, 2019, **10**, 2952.
- 43 Z. Zhao, S. Chen, X. Shen, F. Mahtab, Y. Yu, P. Lu, J. W. Y. Lam, H. S. Kwok and B. Z. Tang, *Chem. Commun.*, 2010, **46**, 686–688.
- 44 M. Desroches, P. M. Burrezo, J. Boismenu-lavoie, M. Peça, J. G. Carlos, J. M. Matxain, D. Casanova, J. Morin and J. Casado, *Angew. Chem., Int. Ed.*, 2017, **56**, 16212–16217.
- 45 X. Lu, S. Lee, J. O. Kim, T. Y. Gopalakrishna, H. Phan, T. S. Herng, Z. Lim, Z. Zeng, J. Ding, D. Kim and J. Wu, *J. Am. Chem. Soc.*, 2016, **138**, 13048–13058.
- 46 H. Sakurai, H. Tobita, M. Kira and Y. Nakadaira, *Angew. Chem., Int. Ed.*, 1980, **19**, 620.
- 47 C. Wentrup, M. J. Regimbald-Krnel, D. Müller and P. Comba, *Angew. Chem., Int. Ed.*, 2016, **55**, 14600–14605.
- 48 G. E. Rudebusch, J. L. Zafra, K. Jorner, K. Fukuda, J. L. Marshall, I. Arrechea-Marcos, G. L. Espejo, R. Ponce Ortiz, C. J. Gómez-García, L. N. Zakharov, M. Nakano, H. Ottosson, J. Casado and M. M. Haley, *Nat. Chem.*, 2016, **8**, 753–759.
- 49 L. K. Montgomery, J. C. Huffman, E. A. Jurczak and M. P. Grendze, *J. Am. Chem. Soc.*, 1986, **108**, 6004–6011.
- 50 M. Abe, *Chem. Rev.*, 2013, **113**, 7011–7088.
- 51 W. Zeng, Z. Sun, T. S. Herng, T. P. Gonçalves, T. Y. Gopalakrishna, K.-W. Huang, J. Ding and J. Wu, *Angew. Chem., Int. Ed.*, 2016, **55**, 8615–8619.
- 52 E. Clar, in *Mobile Source Emissions Including Polycyclic Organic Species*, ed. D. Rondia, M. Cooke and R. K. Haroz, Springer Netherlands, Dordrecht, 1983, pp. 49–58.
- 53 M. Solà, *Front. Chem.*, 2013, **1**, 22.

