

Cite this: *Chem. Sci.*, 2020, 11, 2999

All publication charges for this article have been paid for by the Royal Society of Chemistry

Transient cavity dynamics and divergence from the Stokes–Einstein equation in organic aerosol†

Young-Chul Song,^a Stephen Ingram,^{ab} Robert E. Arbon,^{ab} David O. Topping,^c David R. Glowacki^{id}*^{abd} and Jonathan P. Reid^{id}*^a

The diffusion of small molecules through viscous matrices formed by large organic molecules is important across a range of domains, including pharmaceutical science, materials chemistry, and atmospheric science, impacting on, for example, the formation of amorphous and crystalline phases. Here we report significant breakdowns in the Stokes–Einstein (SE) equation from measurements of the diffusion of water (spanning 5 decades) and viscosity (spanning 12 decades) in saccharide aerosol droplets. Molecular dynamics simulations show water diffusion is not continuous, but proceeds by discrete hops between transient cavities that arise and dissipate as a result of dynamical fluctuations within the saccharide lattice. The ratio of transient cavity volume to solvent volume increases with size of molecules making up the lattice, increasing divergence from SE predictions. This improved mechanistic understanding of diffusion in viscous matrices explains, for example, why organic compounds equilibrate according to SE predictions and water equilibrates more rapidly in aerosols.

Received 9th December 2019

Accepted 15th February 2020

DOI: 10.1039/c9sc06228a

rsc.li/chemical-science

1. Introduction

Examining the relationship between the diffusion rates of small molecules and the viscosity of the surrounding molecular matrix is important for exploring problems as diverse as the molecular mechanisms of crystallization and the formation of amorphous phases in drying droplets,^{1–3} the controlled-release of active ingredients from structured micro-particles in pharmaceutical and consumer products,^{4–7} and the mass concentration of secondary organic aerosol particles in a polluted urban environment.^{8–10} The simplest relationship, the Stokes–Einstein (S–E) equation, expresses the inverse correlation between the translational diffusion coefficient, D , of a large spherical solute molecule of radius a , moving within a solvent continuum with a dynamic viscosity, η :^{11,12}

$$D = \frac{k_B T}{C \pi \eta a} \quad (1)$$

where C is a constant. However, in many important cases the “solvent” (*i.e.* the dominant component by mole fraction) may be a large organic molecule and the “solute” (*i.e.* the minor

component) may be a small molecule, *e.g.* water.^{11,13,14} For example, in the drying of aqueous-organic solution droplets, the evaporation of water can lead to an involatile solute surpassing its solubility limit, thereby becoming the major component with a mole fraction that can approach 1. The sudden removal of water can lead to a “frozen” organic-rich matrix with a sufficiently high viscosity such that nucleation and crystallization are delayed, unable to occur on an experimentally realisable timescale, with the solution composition crossing the threshold for a moisture-induced glass transition.¹⁵ Even then, the residual moisture content can impact product lifetime and particle morphology. Under these conditions, it is most appropriate to consider the diffusion of water within an organic matrix at infinite dilution of water; however, it is typical that a significant divergence from the S–E equation is observed in this limit.^{16–18} Modifications to the S–E equation have been suggested, including the use of a fractional exponent (*i.e.* $D \propto \eta^{-\alpha}$), that account for different relationships between the diffusion coefficient and viscosity.^{19–21}

Independent measurements of diffusion coefficients and viscosities over the appropriately wide ranges needed to observe the failure of the S–E equation are challenging. Most measurements report the *temperature-dependence* of viscosities and diffusion coefficients for super-cooled liquids or solutions of fixed composition, and can approach close to the glass transition temperature.^{12,22–24} By contrast, there are many fewer studies of the *compositional dependence* of the divergence from the S–E equation, for example with diminishing moisture content as the glass transition relative humidity (RH) is approached.^{11,13,17,18,20,25} Moisture acts as a plasticizer in

^aSchool of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, UK. E-mail: j.p.reid@bristol.ac.uk; glowacki@bristol.ac.uk

^bCentre for Computational Chemistry, University of Bristol, Cantock's Close BS8 1TS, UK

^cSchool of Earth and Environmental Science, University of Manchester, Manchester M13 9PL, UK

^dDepartment of Computer Science, University of Bristol, UK

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9sc06228a



atmospheric aerosol particles, regulates the viscosity and, thus, shelf-life of amorphous particles used in formulations, and could play a critical influence in governing crystal formation in drying droplets and films as opposed to the formation of an amorphous solid. Examining the compositionally dependent divergence of an organic solute–water mixture from S–E behaviour not only requires accurate measurements of diffusion coefficients and viscosities over as much as 15 orders of magnitude but requires accurate measurements of composition, recognising that both viscosity and diffusion coefficients are highly dependent on the identity for the functional groups forming the organic solute.²⁶ To access the full viscosity range, moisture must be removed from metastable supersaturated solution droplets without crystallization. We report here a systematic study of the failure of the S–E equation for a range of aqueous-saccharide solutions, varying the molecular size of the organic molecule forming the viscous matrix relative to water and exploring the detailed mechanism of water transport in the limit of a pure saccharide particle.

2. Measurements of diffusion coefficients of water in aqueous-saccharide aerosol particles

Not only are saccharides used widely as excipients for drug delivery^{27–29} and excipient particles are often prepared by spray drying,^{30–32} they find widespread application in the food industry and are commonly used as laboratory surrogates for high oxidized viscous secondary organic atmospheric aerosol.^{3,15,33–36} Using aerosol particles levitated in optical tweezers, we have carried out measurements which avoid the process of heterogeneous nucleation that occurs in the presence of a substrate, allowing access to particle viscosities spanning dilute aqueous solutions (10^{-3} Pa s) to an amorphous solid (10^{12} Pa s). The moisture content is readily altered by varying the relative humidity of the gas phase. Specifically, we considered five binary aqueous-saccharide solution aerosols: glucose (a mono-saccharide); sucrose, trehalose and maltose (all disaccharides); and raffinose (a tri-saccharide). We also consider aqueous aerosol droplets containing levoglucosan, a representative oxygenated compound of biomass burning aerosol particles in the atmosphere.³⁷

Fig. 1a shows examples of the time-dependence in the size response functions for aqueous-raffinose particles following transitions in RH. The significant changes in the equilibration time reflect the significant changes in particle viscosity that are observed over this range in RH/moisture content: equilibration at RHs above the glass-transition RH occurs on timescales $\ll 1$ hour; at low RH, the release of moisture from an amorphous glass occurs over many hours and indeed is not complete over the experimental timescales. The time-constant, τ , and “stretch factor” β of the multiexponential decay observed in both evaporation and condensation events show a path dependence, varying with both the initial and final RH, the initial particle size and the wait-time at intermediate RHs (see Fig. S4†). Tabulated values of both parameters observed in each of the




Fig. 1 Examples of each step in the workflow required to extract the compositional dependencies of the diffusion coefficients from a time-dependence in particle size. The panels show: (a) a collection of response functions for size changes of aqueous-raffinose particles following a step change in RH; (b) a single response function following a change in RH from 30 to 5% RH for a sucrose particle and the best-fit produced by the Fickian diffusion model. (c) The estimated compositional dependencies of the diffusion of water in the six binary aqueous-organic aerosol systems studied. The estimate of the diffusion coefficient for water in sucrose from the MD simulations also presented (yellow diamond).

new systems may be found in Table S1.† To fit the compositional/water activity dependence of the diffusion coefficient of water for binary solution aerosol droplets requires measurements at many RH transitions.^{38,39}

Measurements were performed over 6 RH transitions for 96 droplets for the six binary aqueous-organic aerosol systems studied (glucose, sucrose, trehalose, maltose, raffinose and levoglucosan). Transitions in size were slowest for maltose droplets at the lowest RHs. Moreover, the characteristic time-scale increases with increasing particle size for every binary organic system studied (see Fig. S3 and Table S1†). Time-constants for all particle sizes in the range 3–6 μm show the same ordering: maltose > raffinose > trehalose > sucrose > glucose \geq levoglucosan. In other words, the chain length of the organic appears to be important to the internal mixing dynamics but is not the only controlling factor.

The compositional dependencies of the diffusion coefficients of water estimated for these binary aqueous-organic systems are summarized in panel (c). For reference, the moisture driven glass transition RH has been reported as 53% for raffinose^{36,40} and 32% for maltose;³⁶ the majority of evaporation measurements for these two systems have been made with ultra-viscous and even glassy particles. A value of 23% RH has been reported for sucrose^{36,40} while glucose and levoglucosan are not expected to become glassy at any moisture content at this temperature;³⁴ indeed, levoglucosan crystallizes at an RH of 30% and diffusion coefficients cannot be measured below this. The trend in D_w is not monotonic with molecular weight: levoglucosan (162.1 g mol^{-1}) > glucose (180.2 g mol^{-1}) > raffinose (504.4 g mol^{-1}) > trehalose (342.3 g mol^{-1}) > sucrose (342.3 g mol^{-1}) > maltose (342.3 g mol^{-1}) at the same water activity. Water in the monosaccharide shows the fastest



diffusivity, and diffusion in the trisaccharide is faster than in the disaccharides when a fixed RH/water activity is considered. Indeed, this trend in the diffusion coefficient of water in the limit of a pure dry organic matrix is consistent with a previous assessment of the diffusion coefficients at the glass transition temperature for a subset of the compounds studied here.⁴¹

3. The relationship between diffusion and viscosity in mono-, di- and tri-saccharide particles

The diffusion coefficient measurements presented in Fig. 1 and our measurements of solution droplet viscosities³⁶ allow us to examine their correlation over wide ranges spanning more than 12 orders of magnitude in viscosity and 7 orders of magnitude in diffusion coefficient. The correlations for these systems are compared with predictions from the S-E equation in Fig. 2, assuming a molecular diameter for water of 0.2 nm. Typical error estimates in diffusion coefficient and viscosity are indicated by the representative error bars for each system. The diffusion coefficients for water in all organic-aqueous solutions increasingly deviate from the S-E equation with decreasing water activity and increasing viscosity. Even at the threshold of semi-solid behaviour (10^4 Pa s), the diffusion coefficient of water in aqueous-raffinose aerosol droplets is ~ 5 orders of magnitude larger than estimated by S-E. This is a consequence of the inapplicability of the S-E assumptions to estimations of the diffusion coefficient of a small molecule moving within

a matrix of large molecules, *i.e.* the translation of water is not characterized by simple Brownian motion.^{23,42}

Comparing the relative divergence of water diffusion coefficients from S-E predictions for the mono-, di- and tri-saccharides, the discrepancy increases systematically across this series. Water transport is fastest in solutions with the trisaccharide raffinose and slowest in solutions with the monosaccharide glucose at a certain solution viscosity; the disaccharides (sucrose, trehalose and maltose) fall in the intermediate range. These results suggest that the disparity in size between water and the organic molecule forming the matrix is key to determining the diffusion rate of water. It also explains why the particle size relaxation times and limiting D values (in dry air) did not directly scale with molecular weight: the particles exhibit different viscosities at the same water activity. Therefore, an independent viscosity axis, in this case produced using the aerosol particle coalescence technique, is crucial to separating the two effects.

It can be postulated that when forming a matrix from raffinose, a much larger molecule than water, the packing density of raffinose leaves sufficient free volume for water to move more readily through the network of organic molecules. When the organic molecule is closer in size to water, as in the case of the monosaccharide glucose, the tighter relative packing of glucose leads to a fewer adequately sized cavities. In this sense, the mechanism of impaired water transport more closely resembles percolation rather than diffusion,^{43,44} a process that is sensitive to the free volume of the medium.⁴⁵

Fig. 2 is instructive when considering the diffusion of water through the complex organic matrices found in atmospheric secondary organic aerosol (SOA), one particular motivation for the current study. For example, water transport in α -pinene SOA (orange bars) is more rapid than would be expected based on measurements of viscosity and estimates from the S-E equation,^{11,14,17} most closely resembling the di- and tri-saccharides. However, it should be recognised that the properties of SOA constituent molecules are considerably different. The average molecular weight of organic components identified in α -pinene SOA is $150\text{--}200\text{ g mol}^{-1}$,⁴⁷ albeit with a lower degree of oxygenation: typically the O : C ratio has been reported as 0.45–0.55.⁴⁸ The O : C ratios for trehalose and raffinose are 0.92 and 0.89, respectively. The faster diffusion of water in SOA than expected from the S-E equation may be attributed to the heterogeneity in composition at the molecular scale, leading to a porous network of channels through which water transport is more facile than expected.

4. The microscopic mechanism from molecular dynamics simulations

To better understand the microscopic mechanism of water transport, we carried out atomistic molecular dynamics (MD) simulations of water in sucrose (with 'concentrations' of one water molecule per 35 sucrose molecules) designed to mimic experimental water activities close to zero. See ESI† for further information on the MD simulations. The initial placement of




Fig. 2 Correlation of the diffusion coefficient of water with the viscosity of the aqueous-organic matrix. A prediction from the S-E equation is shown by the grey line. The relationship between the diffusion constant and viscosity of α -pinene SOA (orange markers) has been inferred by us from the literature, and has previously been discussed.^{17,46} The colour scale is the same as in Fig. 1.






Fig. 3 (a) Snapshots in the yz plane of the trajectory of a water molecule 'jumping' between interstices in an amorphous sucrose lattice. The red arrow indicated the direction of the observed hop. The periodic box is shown in grey. Sequential calculated potentials of mean force that the water experiences are shown in panels (b) to (d), separated in time by 1 ns.

the organic molecules is intended to replicate the amorphous packing structure that is expected to occur near the surface of a glassy sucrose droplet.⁴⁹ These MD simulations were

analysed to provide an independent estimate for the value of the intercept $D_{w,org}$. Fig. 1c shows the MD-derived value of $4.64 \times 10^{-17} \text{ m}^2 \text{ s}^{-1}$ is in good agreement with the experimental




Fig. 4 (a and b) The water hopping mechanism showing 2 and 3 metastable states (panels a & b respectively) arising from one of the nine trajectories. Subplots (i) show zy , $(-z)x$ projections of water molecule's position throughout the trajectory with different colours indicating different metastable states. Subplots (ii) shows a hidden Markov state model representation of the hopping behaviour. Each circle represents a metastable state, with the size related to its stability, the arrows show the hopping timescale in picoseconds from one state to another (same colour scheme as (i) subplots). Subplots (iii) show Bayesian estimates of the relaxation timescales associated with hopping between the states (thick line is mean, coloured region is a 95% credibility interval).



measurements, and indicates that our computational approach captures the physics of water diffusion in sucrose at low activities.

Inspection of our MD results reveals that the mechanism of water diffusion through the sucrose matrix proceeds by a hopping between cavities (Fig. 3). In general, a 'cavity' is defined as a sucrose interstitial domain where water has a significant lifetime based on Markov analysis. Our analysis

has enabled us to identify both reversible and irreversible examples of intercavity dynamics (see ESI†). Fig. 4 and much of the analysis described in this article focusses on clusters of 'cavities' between which water molecules make reversible kinetic hops, because this local equilibrium is amenable to analysis using standard tools in statistical mechanics.

Our analysis shows that water remains in a cavity (or cluster of cavities) until either (1) it achieves sufficient kinetic energy to




Fig. 5 Occupied volumes of (a) glucose (67.1%), (c) sucrose (66.0%) and (e) raffinose (62.4%) matrices are shown (same colour scheme as Fig. 1 again) showing the van der Waals radii of the saccharides within one snapshot of the short simulations. Fractional free volume are shown for (b) glucose, (d) sucrose, and (f) raffinose calculated by GROMACS throughout 10 ns simulations.



References

- M. L. F. Nascimento and E. Dutra Zanotto, *J. Chem. Phys.*, 2010, **133**, 174701.
- E. Mikhailov, S. Vlasenko, S. T. Martin, T. Koop and U. Pöschl, *Atmos. Chem. Phys.*, 2009, **9**, 9491–9522.
- T. Koop, J. Bookhold, M. Shiraiwa and U. Pöschl, *Phys. Chem. Chem. Phys.*, 2011, **13**, 19238–19255.
- G. M. Hidy, *Aerosols: Industrial and Environmental Science*, Elsevier Science, 1984.
- B. C. Hancock and G. Zograf, *J. Pharm. Sci.*, 1997, **86**, 1–12.
- C. Lorenz, H. Hagedorfer, N. Von Goetz, R. Kaegi, R. Gehrig, A. Ulrich, M. Scheringer and K. Hungerbühler, *J. Nanopart. Res.*, 2011, **13**, 3377–3391.
- A. E. Haddrell, J. F. Davies, R. E. H. Miles, J. P. Reid, L. A. Dailey and D. Murnane, *Int. J. Pharm.*, 2014, **463**, 50–61.
- M. Shiraiwa and J. H. Seinfeld, *Geophys. Res. Lett.*, 2012, **39**, L24801.
- H. Mai, M. Shiraiwa, R. C. Flagan and J. H. Seinfeld, *Environ. Sci. Technol.*, 2015, **49**, 11485–11491.
- A. M. Maclean, C. L. Butenhoff, J. W. Grayson, K. Barsanti, J. L. Jimenez and A. K. Bertram, *Atmos. Chem. Phys. Discuss.*, 2017, 1–17.
- R. M. Power, S. H. Simpson, J. P. Reid and A. J. Hudson, *Chem. Sci.*, 2013, **4**, 2597–2604.
- B. Chen, E. E. Sigmund and W. P. Halperin, *Phys. Rev. Lett.*, 2006, **96**, 1–4.
- H. C. Price, B. J. Murray, J. Mattsson, D. O'Sullivan, T. W. Wilson, K. J. Baustian and L. G. Benning, *Atmos. Chem. Phys.*, 2014, **14**, 3817–3830.
- V. Molinero and W. Goddard, *Phys. Rev. Lett.*, 2005, **95**, 045701.
- D. L. Bones, J. P. Reid, D. M. Lienhard and U. K. Krieger, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 11613–11618.
- R. M. Power, S. H. Simpson, J. P. Reid and A. J. Hudson, *Chem. Sci.*, 2013, **4**, 2597.
- H. C. Price, J. Mattsson, Y. Zhang, A. K. Bertram, J. F. Davies, J. W. Grayson, S. T. Martin, D. O'Sullivan, J. P. Reid, A. M. J. Rickards and B. J. Murray, *Chem. Sci.*, 2015, **6**, 4876–4883.
- Y. Chenyakin, D. A. Ullmann, E. Evoy, L. Renbaum-Wolff, S. Kamal and A. K. Bertram, *Atmos. Chem. Phys.*, 2017, **17**, 2423–2435.
- K. R. Harris, *J. Chem. Phys.*, 2009, **131**, 054503.
- H. C. Price, J. Mattsson and B. J. Murray, *Phys. Chem. Chem. Phys.*, 2016, **18**, 19207–19216.
- F. Fernandez-Alonso, F. J. Bermejo, S. E. McLain, J. F. C. Turner, J. J. Molaison and K. W. Herwig, *Phys. Rev. Lett.*, 2007, **98**, 1–4.
- M. Shrestha, Y. Zhang, M. A. Upshur, P. Liu, S. L. Blair, H. Wang, S. A. Nizkorodov, R. J. Thomson, S. T. Martin and F. M. Geiger, *J. Phys. Chem. A*, 2015, **119**, 4609–4617.
- S.-H. Chen, F. Mallamace, C.-Y. Mou, M. Broccio, C. Corsaro, A. Faraone and L. Liu, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 12974–12978.
- J. A. T. González, M. P. Longinotti and H. R. Corti, *J. Phys. Chem. B*, 2015, **119**, 257–262.
- S. Bastelberger, U. K. Krieger, B. Luo and T. Peter, *Atmos. Chem. Phys.*, 2017, **17**, 8453–8471.
- N. E. Rothfuss and M. D. Petters, *Environ. Sci. Technol.*, 2017, **51**, 271–279.
- Y. Zhao, B. G. Trewyn, I. I. Slowing and V. S. Y. Lin, *J. Am. Chem. Soc.*, 2009, **131**, 8398–8400.
- V. R. Sinha and R. Kumria, *Int. J. Pharm.*, 2001, **224**, 19–38.
- S. Sastry, J. Nyshadham and J. Fix, *Pharm. Sci. Technol. Today*, 2000, **3**, 138–145.
- J. D. Andya, Y. F. Maa, H. R. Costantino, P. A. Nguyen, N. Dasovich, T. D. Sweeney, C. C. Hsu and S. J. Shire, *Pharm. Res.*, 1999, **16**, 350–358.
- R. Vehring, *Pharm. Res.*, 2008, **25**, 999–1022.
- K. Mosén, K. Bäckström, K. Thalberg, T. Schaefer, H. G. Kristensen and A. Axelsson, *Pharm. Dev. Technol.*, 2004, **9**, 409–417.
- Y. Liu, B. Bhandari and W. Zhou, *J. Agric. Food Chem.*, 2006, **54**, 5701–5717.
- B. Zobrist, C. Marcolli, D. A. Pedernera and T. Koop, *Atmos. Chem. Phys.*, 2008, **8**, 5221–5244.
- J. P. Reid, A. K. Bertram, D. O. Topping, A. Laskin, S. T. Martin, M. D. Petters, F. D. Pope and G. Rovelli, *Nat. Commun.*, 2018, **9**, 956.
- Y. C. Song, A. E. Haddrell, B. R. Bzdek, J. P. Reid, T. Bannan, D. O. Topping, C. Percival and C. Cai, *J. Phys. Chem. A*, 2016, **120**, 8123–8137.
- S. Decesari, S. Fuzzi, M. C. Facchini, M. Mircea, L. Emblico, F. Cavalli, W. Maenhaut, X. Chi, G. Schkolnik, A. Falkovich, Y. Rudich, M. Claeys, V. Pashynska, G. Vas, I. Kourtchev, R. Vermeylen, A. Hoffer, M. O. Andreae, E. Tagliavini, F. Moretti and P. Artaxo, *Atmos. Chem. Phys.*, 2006, **6**, 375–402.
- S. Ingram, C. Cai, Y. C. Song, D. R. Glowacki, D. O. Topping, S. O'Meara and J. P. Reid, *Phys. Chem. Chem. Phys.*, 2017, **19**, 31634–31646.
- A. M. J. Rickards, Y. C. Song, R. E. H. Miles, T. C. Preston and J. P. Reid, *Phys. Chem. Chem. Phys.*, 2015, **17**, 10059–10073.
- H.-J. Tong, J. P. Reid, D. L. Bones, B. P. Luo and U. K. Krieger, *Atmos. Chem. Phys.*, 2011, **11**, 4739–4754.
- D. M. Lienhard, A. J. Huisman, U. K. Krieger, Y. Rudich, C. Marcolli, B. P. Luo, D. L. Bones, J. P. Reid, A. T. Lambe, M. R. Canagaratna, P. Davidovits, T. B. Onasch, D. R. Worsnop, S. S. Steimer, T. Koop and T. Peter, *Atmos. Chem. Phys.*, 2015, **15**, 13599–13613.
- Q. Liu, S. Huang and Z. Suo, *Phys. Rev. Lett.*, 2015, **114**, 1–5.
- V. K. S. Shante and S. Kirkpatrick, *Adv. Phys.*, 1971, **20**, 325–357.
- M. Shiraiwa, M. Ammann, T. Koop and U. Pöschl, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 11003–11008.
- R. P. White and J. E. G. Lipson, *Macromolecules*, 2016, **49**, 3987–4007.
- F. H. Marshall, R. E. H. Miles, Y.-C. Song, P. B. Ohm, R. M. Power, J. P. Reid and C. S. Dutcher, *Chem. Sci.*, 2016, **7**, 1298–1308.



- 47 S. Chen, W. H. Brune, A. T. Lambe, P. Davidovits and T. B. Onasch, *Atmos. Chem. Phys.*, 2013, **13**, 5017–5031.
- 48 X. Zhang, R. C. McVay, D. D. Huang, N. F. Dalleska, B. Aumont, R. C. Flagan and J. H. Seinfeld, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, 14168–14173.
- 49 E. Mikhailov, S. Vlasenko, S. T. Martin, T. Koop and U. Pöschl, *Atmos. Chem. Phys.*, 2009, **9**, 9491–9522.
- 50 R. Krishna and J. A. Wesselingh, *Chem. Eng. Sci.*, 1997, **52**, 861–911.
- 51 L. Jia and Y. Xu, *Atmos. Chem. Phys.*, 2018, **18**, 8137–8154.
- 52 W. Huang, H. Saathoff, A. Pajunoja, X. Shen, K.-H. Naumann, R. Wagner, A. Virtanen, T. Leisner and C. Mohr, *Atmos. Chem. Phys. Discuss.*, 2017, 1–23.
- 53 I. Kourtchev, C. Giorio, A. Manninen, E. Wilson, B. Mahon, J. Aalto, M. Kajos, D. Venables, T. Ruuskanen, J. Levula, M. Loponen, S. Connors, N. Harris, D. Zhao, A. Kiendler-Scharr, T. Mentel, Y. Rudich, M. Hallquist, J.-F. Doussin, W. Maenhaut, J. Bäck, T. Petäjä, J. Wenger, M. Kulmala and M. Kalberer, *Sci. Rep.*, 2016, **6**, 35038.

