

Cite this: *Chem. Sci.*, 2020, **11**, 11293

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 17th June 2020
Accepted 18th September 2020

DOI: 10.1039/d0sc03359f

rsc.li/chemical-science

Evidence for an enolate mechanism in the asymmetric Michael reaction of α,β -unsaturated aldehydes and ketones *via* a hybrid system of two secondary amine catalysts†

Nariyoshi Umekubo,^a Takahiro Terunuma,^a Eunsang Kwon^b and Yujiro Hayashi[✉] [✉]

The key nucleophile was found to be neither an enamine nor an enol, but an enolate in the direct Michael reaction of α,β -unsaturated aldehydes and non-activated ketones catalyzed by two amine catalysts namely diphenylprolinol silyl ether and pyrrolidine. This is a rare example of an enolate from a ketone serving as a key intermediate in the asymmetric organocatalytic reaction involving secondary amine catalysts because the ketone enolates are generally generated using a strong base, and the enamine is a common nucleophile in this type of reaction.

Introduction

Enolates are versatile and important synthetic intermediates for the formation of α -substituted carbonyl compounds, which are widely used in the synthesis of organic molecules. In order to retard nucleophilic attack on the carbonyl group, and to match the weak acidity of the proton, sterically hindered strong amide bases such as lithium diisopropylamide (LDA) and lithium hexamethyldisilyl-amide (LHMDS) are generally employed for the generation of enolates from ketones.^{1,2} In the field of organocatalysis³ organic super bases such as phosphazene⁴ and proazaphosphatrane⁵ have been developed, which have similar or much stronger basicity than LDA and LHMDS.

Contrary to organic strong bases, weak secondary and primary amines can be successfully employed for the α -functionalization of non-activated ketones in which the key intermediate is not an enolate but an enamine. Stork was a pioneer in this field of enamine chemistry where an equimolar amount of the amine was employed.⁶ However, in recent asymmetric reactions, catalytically generated enamines are essentially the key intermediates. In this context, chiral secondary and primary amines act as catalysts for the α -functionalization of ketones in bond forming reactions such as the aldol reaction,^{3,7} Mannich reaction,^{3,8} Michael reaction,³ α -aminooxylation³ and α -aminations.³ In the Mannich reaction of acetone catalysed by a primary amine-thiourea organocatalyst, an enol mechanism

was proposed,⁹ while an enamine mechanism is proved by calculation and ¹³C kinetic isotope effects in the Michael reaction of acetone catalysed by a similar catalyst.¹⁰ In the Michael reaction of propanal and methyl vinyl ketone catalysed by pyrrolidine, both enamine and enol mechanisms were computationally investigated, and an enamine is concluded to be a key nucleophile.¹¹ The discrimination of the reaction mechanism of enamine and enol is difficult, and the asymmetric reaction involving an enol is rare. Moreover, the reaction is very rare in which an enolate (not enol) is a key intermediate in the secondary amine mediated asymmetric catalytic reactions of non-activated ketones with excellent enantioselectivity. Although there is one report involving an enolate as an intermediate, as far as we are aware, the enantioselectivity is not sufficient.¹² In this study, we will demonstrate that the key nucleophile in the Michael reaction of cyclohexanone and α,β -unsaturated aldehyde catalyzed by a secondary amine was neither an enamine nor an enol but a rather unexpected enolate.


Results and discussion

Recently, we reported the first direct asymmetric Michael reaction of non-activated ketones and α,β -unsaturated aldehydes catalyzed by a combination of two organocatalysts namely diphenylprolinol silyl ether **1**¹³ and pyrrolidine or 4-hydroxyproline (**2**) (eqn (1), Table 1, entries 1, 2 and 3).¹⁴ Although two similar pyrrolidine-type catalysts were involved in the reaction, an iminium ion generated from the α,β -unsaturated aldehyde was the reactive Michael acceptor based on Mayr's electrophilicity principle.¹⁵ This implies that the enantio face-selectivity of the α,β -unsaturated aldehyde was controlled by diphenylprolinol silyl ether. We hypothesized that the nucleophile would be

^aDepartment of Chemistry, Graduate School of Science, Tohoku University, 6-3 Aramaki Aza Aoba, Aoba-ku, Sendai 980-8578, Japan. E-mail: yujiro.hayashi.b7@tohoku.ac.jp

^bResearch and Analytical Center for Giant Molecules, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0sc03359f



Table 1 The effect of the amines in the asymmetric Michael reaction of diphenylprolinol silyl ether^a


| Entry | Catalyst A | Catalyst B | Time [h] | dr ^b | Yield ^c [%] | ee ^d [%] |
|-------|------------|------------|----------|-----------------|------------------------|---------------------|
| 1 | (S)-1 | (S)-1 | 24 | 5 : 1 | 74 | 91 |
| 2 | (S)-1 | 2 | 40 | 15 : 1 | 74 | 96 |
| 3 | (R)-1 | 2 | 40 | 10 : 1 | 70 | −95 |
| 4 | (S)-1 | 3 | 30 | 5 : 1 | 68 | 93 |
| 5 | (R)-1 | 3 | 30 | 9 : 1 | 70 | −94 |

^a Unless otherwise shown, the reaction was performed by employing cinnamaldehyde (0.5 mmol), cyclohexanone (1.5 mmol), catalyst A (0.075 mmol), catalyst B (0.0375 mmol), *p*-nitrophenol (0.15 mmol), and water (1.5 mmol), in EtOH (0.4 mL) and toluene (0.1 mL) at room temperature. After the reaction, Wittig reagent (0.75 mmol) was added. See the ESI for details. ^b dr ratio (*syn* : *anti*) was determined by ¹H-NMR. ^c Isolated yield of the diastereomer mixture. ^d Determined by HPLC analysis on a chiral column material.

an enamine generated from the reaction of the ketone and either pyrrolidine or 4-hydroxyproline. It has been well established that the enamine generated from the ketone and proline is a reactive nucleophile.^{3,16} We reasoned that if the second amine can control the enantio-face selectivity of the ketone, both *syn* and *anti*-isomers can be selectively synthesized.¹⁷ This hypothesis led us to study the effect of the second chiral amine catalyst in more detail with the intention of enhancing the selective control of relative stereochemistry.

In our previous paper, we reported the results of (*S*)-diphenylprolinol silyl ether (*S*)-1 with (2*S*,4*R*)-4-hydroxyproline 2 and (*R*)-1 with (2*S*,4*R*)-2 in the Michael reaction of cinnamaldehyde and cyclohexanone (Table 1, entries 2 and 3).¹⁴ Although the enantio face-selectivity of an enamine originating from (2*S*,4*R*)-4-hydroxyproline 2 is not known, that of an (*R*)-2-(methoxymethyl)pyrrolidine 3 has been well investigated.¹⁸ Thus, the effect of this amine 3 with a combination of (*S*)-1 and (*R*)-1 was examined (entries 4 and 5). Although the diastereoselectivity was slightly different according to the relative fractions of the two catalysts, the *syn*-isomer was predominantly obtained in all cases. As for the enantioselectivity, the absolute configuration was controlled by the chirality of the diphenylprolinol silyl ether 1 (ref. 19) regardless of the chirality of the second amine catalyst 3. These results cast a doubt about the involvement of any enamine as an intermediate in the reaction.


**Fig. 1** Enamine, enol and enolate.**Scheme 1** The reaction of enamine 7 and iminium ion (*S*)-8.

The possible nucleophiles would be the enamine 4, the enol 5 and the enolate 6 (Fig. 1). Hence, we investigated the reactivity of these species. The reactivity of the enamine 7,¹⁸ which was generated from (*R*)-2-(methoxymethyl)pyrrolidine 3, was examined (Scheme 1). The equimolar reaction of enamine 7 and iminium ion (*S*)-8 (ref. 20) proceeds in the presence of 2,6-lutidine and MS4A to afford the bicyclic compound 11 (ref. 21) in 57% yield. 11 was formed by the Michael reaction, followed by an exchange of the enamine and iminium ion and cyclization.²² The stereochemistry at the α -position of the obtained cyclohexanone was *R* whereas the observed stereochemistry from the catalytic Michael reaction was the opposite *S* (Table 1, entry 4).²³ This result implies that the enamine 7 was not an intermediate in the catalytic reaction. Moreover, apart from the secondary amine, tertiary amines such as *i*-Pr₂NEt can successfully act as a co-catalyst to afford the Michael product with excellent enantioselectivity (71%, *syn* : *anti* = 5 : 1, 97% ee). This result was also a piece of evidence to support the claim that the enamine was not a key nucleophile in the reaction.

The equimolar reactions of the iminium ion with 1-methoxycyclohex-1-ene and 1-(trimethylsiloxy)cyclohex-1-ene, and a combination of 1-(trimethylsiloxy)cyclohex-1-ene and tris(dimethylamino)sulfonium difluorotrimethylsilicate (TASF) were investigated. The two former nucleophiles would possess a similar reactivity to 1-hydroxycyclohex-1-ene 5, but they did not react at all even after a longer reaction time (eqn (2)). In the reaction of a combination of silyl enol ether and TASF, which is known to generate an enolate,²⁴ the Michael product was obtained at −30 °C after 1 h with a similar diastereo- and enantio-





selectivity as observed in the catalytic reaction (eqn (3)). This result indicates that an enolate was the likely intermediate.



If the enolate was the actual nucleophile, a question arose about how the enolate would be generated from the ketone and pyrrolidine or *i*-Pr₂NEt. The direct deprotonation of the ketone by pyrrolidine would be very difficult considering their relative *pK_a* values. That is to say that the *pK_a* value of an α -proton on cyclohexanone in DMSO is 26.4 (ref. 25) while that of an ammonium ion of Et₃N is only 9.00.²⁶ This is why a strong base such as LDA is usually employed. But the enolate could be generated considering an equilibrium between the keto and enol forms, and that the O–H proton of the enol form is rather acidic, although the content of enol is very low.²⁷

Next, we investigated the generation speed of the enolate. The combined generation speed of both the enamine, the enol and the enolate can be monitored by the H/D exchange of the α -proton of the carbonyl group in the reaction with D₂O (eqn (4)). The generation speed of the enamine can also be monitored by the ¹⁶O/¹⁸O exchange in the reaction with H₂¹⁸O (eqn (5)).²⁸ Moreover, as a tertiary amine can also act as a co-catalyst in which the enamine would not be involved, the generation of only the enol and the enolate can be monitored by the H/D exchange reaction with D₂O in the case of the tertiary amine (eqn (4)).



We examined the H/D exchange experiments in the presence of (1) pyrrolidine + *p*-nitrophenol (red), (2) *p*-nitrophenol (blue),




Fig. 2 Generation of deuterated substrates in eqn (4), green: *i*-Pr₂NEt and *p*-nitrophenol, red: pyrrolidine and *p*-nitrophenol, blue: *p*-nitrophenol, yellow: *i*-Pr₂NEt, and generation of ¹⁸O labelled substrate in eqn (5), purple.

(3) *i*-Pr₂NEt + *p*-nitrophenol (green), and (4) *i*-Pr₂NEt (yellow) (Fig. 2). A reaction using H₂¹⁸O was conducted using pyrrolidine and *p*-nitrophenol (purple) (Fig. 2). Fig. 2 indicates that the generation speed of the enol and the enolate under the reaction conditions is rather fast compared to the Michael reaction. This indicates that the rapid kinetic of deprotonation is fast, although the concentration of the enol and enolate is quite low. The speed of generation of the enolate was faster in the case of *i*-Pr₂NEt and *p*-nitrophenol compared to pyrrolidine and *p*-nitrophenol. Although each acid and base are known to accelerate the keto/enol equilibrium,²⁹ *p*-nitrophenol did not promote the H/D exchange under the reaction conditions. Moreover, the tertiary amine did not promote the exchange either. However, it should be noted that a combination of both acid and base facilitates the generation of the enolate. Their cooperative role would be explained as follows (Scheme 2):³⁰ *p*-nitrophenol and *i*-Pr₂NEt partially form an ammonium ion, and all these species such as acids, bases and ammonium ions are present in a mixture under equilibrium based on their *pK_a* values.^{26,31} A protonation would occur at the carbonyl oxygen, which would increase the acidity of an α -proton of a carbonyl of cyclohexanone. Then, this α -proton of a carbonyl can be deprotonated by *i*-Pr₂NEt. From the generated vinyl alcohol, deprotonation of the O–H proton³² would proceed very fast to afford the enolate.³³ If so, the combination of acidity of the acid and basicity of the base should be important. In fact, when we used phenol or *p*-methoxyphenol³⁴ instead of *p*-nitrophenol, the reaction became slow (see the ESI†). Moreover, the reaction did not proceed at all in the presence of CF₃CO₂H³⁵ with a combination of *i*-Pr₂NEt (see the ESI†).

One of the roles of the acid is to facilitate the generation of iminium ions from the α,β -unsaturated aldehyde and diphenylprolinol silyl ether, but this result indicates that another role of the acid is to accelerate the generation of the enolate in combination with the amine.

Based on the above investigations, the reaction was considered to proceed as follows (Fig. 3): diphenylprolinol silyl ether reacts with cinnamaldehyde to generate an iminium salt. On the other hand, cyclohexanone is in equilibrium with the enol form in the presence of *p*-nitrophenol and pyrrolidine, and the





This journal is © The Royal Society of Chemistry 2020

- edn, 2013; (b) G. Stork and S. R. Dowd, *J. Am. Chem. Soc.*, 1963, **85**, 2178.
- 7 (a) *Modern Aldol Reactions*, ed. R. Mahrwald, Wiley-VCH, Weinheim, vol. 1–2, 2004; (b) N. Mase and Y. Hayashi, *Comprehensive Organic Synthesis*, Elsevier, Amsterdam, 2nd edn, 2014, vol. 2, pp. 273–339.
- 8 Y. Hayashi, *J. Synth. Org. Chem., Jpn.*, 2014, **72**, 1228.
- 9 D. A. Yalalov, S. B. Tsogoeva, T. E. Shubina, I. M. Martynova and T. Clark, *Angew. Chem., Int. Ed.*, 2008, **47**, 6624.
- 10 V. A. Roytman, R. W. Karugu, Y. Hong, J. S. Hirschi and M. J. Vetticatt, *Chem.–Eur. J.*, 2018, **24**, 8098.
- 11 M. P. Patil and R. B. Sunoj, *Chem.–Asian J.*, 2009, **4**, 714.
- 12 J. Muzart, *Tetrahedron: Asymmetry*, 2014, **25**, 697.
- 13 (a) Y. Hayashi, H. Gotoh, T. Hayashi and M. Shoji, *Angew. Chem., Int. Ed.*, 2005, **44**, 4212; (b) M. Marigo, T. C. Wabnitz, D. Fielenbach and K. A. Jørgensen, *Angew. Chem., Int. Ed.*, 2005, **44**, 794. For reviews, see; (c) C. Palomo and A. Mielgo, *Angew. Chem., Int. Ed.*, 2006, **45**, 7876; (d) A. Mielgo and C. Palomo, *Chem.–Asian J.*, 2008, **3**, 922; (e) L. W. Xu, L. Li and Z. H. Shi, *Adv. Synth. Catal.*, 2010, **352**, 243; (f) K. L. Jensen, G. Dickmeiss, H. Jiang, L. Albrecht and K. A. Jørgensen, *Acc. Chem. Res.*, 2012, **45**, 248; (g) H. Gotoh and Y. Hayashi, *Sustainable Catalysis*, Wiley, Hoboken, 2013, pp. 287–316; (h) B. S. Donslund, T. K. Johansen, P. H. Poulsen, K. S. Halskov and K. A. Jørgensen, *Angew. Chem. Int. Ed.*, 2015, **54**, 13860; (i) G. J. Reyes-Rodriguez, N. M. Rezayee, A. Vidal-Albalat and K. A. Jørgensen, *Chem. Rev.*, 2019, **119**, 4221.
- 14 Y. Hayashi and N. Umekubo, *Angew. Chem., Int. Ed.*, 2018, **57**, 1958.
- 15 S. Lakhdar, T. Tokuyasu and H. Mayr, *Angew. Chem., Int. Ed.*, 2008, **47**, 8723.
- 16 (a) Z. G. Hajos and D. R. Parrish, *J. Org. Chem.*, 1974, **39**, 1615; (b) U. Eder, G. Sauer and R. Wiechert, *Angew. Chem., Int. Ed. Engl.*, 1971, **10**, 496; (c) B. List, R. A. Lerner and C. F. Barbas, *J. Am. Chem. Soc.*, 2000, **122**, 2395.
- 17 For the reactions to control relative and absolute configurations *via* two catalysts, see; (a) S. Krautwald, D. Sarlah, M. A. Schafroth and E. M. Carreira, *Science*, 2013, **340**, 1065; (b) S. Krautwald, M. A. Schafroth, D. Sarlah and E. M. Carreira, *J. Am. Chem. Soc.*, 2014, **136**, 3020; (c) X. Huo, R. He, X. Zhang and W. Zhang, *J. Am. Chem. Soc.*, 2016, **138**, 11093; (d) X. Jiang, J. J. Beiger and J. F. Hartwig, *J. Am. Chem. Soc.*, 2017, **139**, 87; (e) F. A. Cruz and V. M. Dong, *J. Am. Chem. Soc.*, 2017, **139**, 1029; (f) L. Wei, Q. Zhu, S.-M. Xu, X. Chang and C.-J. Wang, *J. Am. Chem. Soc.*, 2018, **140**, 1508.
- 18 (a) S. J. Blarer, W. B. Schweizer and D. Seebach, *Helv. Chim. Acta*, 1982, **65**, 1637; (b) S. J. Blarer and D. Seebach, *Chem. Ber.*, 1983, **116**, 2250; (c) D. Seebach, M. Missbach, G. Calderari and M. Eberle, *J. Am. Chem. Soc.*, 1990, **112**, 7625; (d) E. Butkus and A. Stončius, *Synlett*, 1999, 234; (e) T. Husch, D. Seebach, A. K. Beck and M. Reiher, *Helv. Chim. Acta*, 2017, **100**, e1700182.
- 19 Y. Hayashi, D. Okamura, T. Yamazaki, Y. Ameda, H. Gotoh, S. Tsuzuki, T. Uchimarui and D. Seebach, *Chem.–Eur. J.*, 2014, **20**, 17077.
- 20 H. Gotoh, T. Uchimarui and Y. Hayashi, *Chem.–Eur. J.*, 2015, **21**, 12337.
- 21 The structure of **11** was determined by converting to the crystalline substrate and X-ray crystallographic analysis was conducted, see the ESI† for details. This reaction was conducted at $-30\text{ }^{\circ}\text{C}$. The reaction also proceeded at rt, but the yield was lower because of the side reactions. On the other hand, a catalytic reaction (Table 1, entry 4) proceeded at room temperature, which was very slow at $-30\text{ }^{\circ}\text{C}$.
- 22 Another pathway for the formation of **10** is the ene reaction of **7** and **8**.
- 23 When the reaction was stopped in a short reaction time, similar enantioselectivity has been obtained. Thus, the Michael product was obtained kinetically.
- 24 (a) E. Nakamura, M. Shimizu, I. Kuwajima, J. Sakata, K. Yokoyama and R. Noyori, *J. Org. Chem.*, 1983, **48**, 932; (b) R. Noyori, I. Nishida and J. Sakata, *J. Am. Chem. Soc.*, 1983, **105**, 1598.
- 25 F. G. Bordwell and H. E. Fried, *J. Org. Chem.*, 1991, **56**, 4218.
- 26 As we cannot find the pK_{a} of ammonium ions of $i\text{-Pr}_2\text{NET}$, we will discuss the pK_{a} of ammonium ions of a similar tertiary amine such as Et_3N . (a) I. M. Kolthoff, M. K. Chantooni and S. Bhowmik, *J. Am. Chem. Soc.*, 1968, **90**, 23; (b) M. R. Crampton and I. A. Robotham, *J. Chem. Res.*, 1997, 22.
- 27 (a) J. P. Guthrie and P. A. Cullimore, *Can. J. Chem.*, 1979, **57**, 240; (b) J. P. Guthrie, *Can. J. Chem.*, 1979, **57**, 1177.
- 28 For the reaction using H_2^{18}O to determine the reaction mechanism, see; Y. Hayashi, T. Mukaiyama, M. Benohoud, N. R. Gupta, T. Ono and S. Toda, *Chem.–Eur. J.*, 2016, **22**, 5868.
- 29 See the general organic textbook, for instance, J. McMurry, *Organic Chemistry*, Physical Sciences, Belmont, 7th edn, 2008, p. 842.
- 30 The generation of an enol (not enolate) from an aldehyde (not ketone) using pyrrolidine was observed to proceed through a cyclic replay mechanism, see ref. 11.
- 31 As the solvent of the reaction is EtOH/toluene (4/1), we discuss pK_{a} in MeOH. The pK_{a} of o -nitrophenol and Et_3NH^+ (in water) is 7.1 and 10.75, respectively, based on Evans' pK_{a} table (ref. 2). The pK_{a} of o -nitrophenol and Et_3NH^+ in MeOH is estimated to be 11.1 and 10.95, respectively, by the use of the empirical conversion method by Knapp. See, E. Rossini, A. D. Bochevarov and E. W. Knapp, *ACS Omega*, 2018, **3**, 1653.
- 32 The enol O–H of the cyclohexanone is acidic enough to be deprotonated by a tertiary amine. The pK_{a} of the enol O–H of the cyclohexanone is 12.1 in water, see; (a) J. P. Guthrie and P. A. Cullimore, *Can. J. Chem.*, 1979, **57**, 240; (b) J. P. Guthrie, *Can. J. Chem.*, 1979, **57**, 1177. The calculated pK_{a} using Advanced Chemistry Development (ACD/Labs) Software V11.02 is 11.5 (gas phase).
- 33 The deprotonation of enol–OH is very fast (often diffusion controlled), see H. O. House, *Modern Synthetic Reactions*, Benjamin, Philippines, 2nd edn, 1972, p. 495.
- 34 The pK_{a} of phenol and p -methoxyphenol in MeOH is estimated to be 13.95 and 14.2, respectively, see ref. 2 and 31.
- 35 The value of pK_{a} in MeOH is estimated to be 4.75, respectively, see ref. 2 and 31.

