

Cite this: *Chem. Sci.*, 2019, 10, 5539

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 31st January 2019

Accepted 25th April 2019

DOI: 10.1039/c9sc00561g

rsc.li/chemical-science

Tuning ligand field strength with pendent Lewis acids: access to high spin iron hydrides†

John J. Kiernicki,^a James P. Shanahan,^a Matthias Zeller^b
and Nathaniel K. Szymczak^{id}*^a

Geometrically flexible 9-borabicyclo[3.3.1]nonyl units within the secondary coordination sphere enable isolation of high-spin Fe(II)-dihydrides stabilized by boron–hydride interactions and a rare example of an isolable $S = 3/2$ reduction product. The borane-capped Fe(II)-dihydride: (1) rapidly deprotonates E–H (E = N, O, P, S) bonds to afford borane-stabilized Fe adducts and (2) releases H₂ upon exposure to π -acids. The Lewis acids provide an avenue for redox-leveling in analogy to the near constant operating potential for N₂ reduction in nitrogenase.

Iron-dihydrides are intermediates in myriad homogenous catalytic reactions and the recent push to develop earth-abundant transition metal catalysts has fueled many research groups to explore their reactivity.¹ Beyond relevance to homogenous catalysis, iron-dihydrides have been implicated in biological reduction sequences. Whereas strong-field hydride ligands typically enforce low-spin configurations, metal-locofactors including nitrogenase contain high-spin hydrides.² To account for the modest conditions used by nitrogenase enzymes for N₂ reduction, one proposal to accumulate reducing power at a near constant potential is to store reducing equivalents as Fe– μ -H–Fe intermediates.^{2,3} For example, the E₄ state of the FeMoco center of nitrogenase is proposed to eliminate H₂ from accumulated bridging hydrides concomitant with N₂ binding/reduction (Fig. 1).

Well characterized synthetic examples of open-shell Fe– μ -H–Fe complexes are rare,⁴ and thus, despite their relevance in biology, the synthesis and reactivity of such species remain largely unknown. This disparity is likely due to the mismatched requirements of the ligand/metal combination – strong-field hydride donor ligands rarely afford high-spin electronic configurations.

Our group is working to evaluate how the precise structural, electronic, and cooperative modes in the secondary coordination sphere can be used to regulate reactivity.⁵ One way in which the ligand-field strength of otherwise strong-field hydride ligands can be attenuated (accommodating high-spin states) is by introducing acidic groups to form bridging hydrides. The

multiple Fe-centers in the E₄ state may serve this role.⁶ To model these intermediates with redox-inactive acids, we targeted the synthesis of ferrous-dihydride compounds in the presence of appended boron Lewis acids. Herein, we report Lewis acid enabled isolation of a high-spin Fe(II)-di(boro)hydride as well as its reduction product, an $S = 3/2$ Fe-di(boro)hydride, and subsequent reactivity.

The complex, (^{BBN}PDP^{tBu})FeBr₂, contains a pair of moderately acidic 9-borabicyclo[3.3.1]nonyl (9-BBN) substituents that are capable of interacting with nitrogenous substrates independent of the metal center (N₂H₄) or cooperatively with the metal center (NH₂[−]).⁷ Treatment of a freshly-thawed orange THF solution of (^{BBN}PDP^{tBu})FeBr₂ with two equiv. KBHET₃ affords an olive-tan powder, assigned as (^{BBN}PDP^{tBu})FeH₂ (**1**) (70%, Fig. 2). **1** is modestly stable with a half-life of ~24 h in THF at room temperature. Investigation by ¹H-NMR spectroscopy (THF) revealed a paramagnetically shifted spectrum with resonances ranging −16.9–63.6 ppm. The decrease in solution symmetry



Fig. 1 FeMo-cofactor and our approach to hydride accumulation with boron Lewis acids.

^aUniversity of Michigan, 930 N. University, Ann Arbor, MI 48109, USA. E-mail: nszym@umich.edu

^bH. C. Brown Laboratory, Purdue University, 560 Oval Dr, West Lafayette, IN 47907, USA

† Electronic supplementary information (ESI) available. CCDC 1884220–1884232 and 1903500. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc00561g






Fig. 2 Synthesis of borane-capped metal–dihydrides.

from C_{2v} to C_2 is consistent with the trialkylboranes interacting with the hydride ligands at the Fe-center. Infrared spectroscopy (KBr) supported this formulation with a broad Fe–H–B stretch at $\sim 1839\text{ cm}^{-1}$.⁸ Notably, the energy of this absorption is at significantly lower energy than bis(tris(mercaptoimidazolyl)hydroborato)-Fe(II) species that display a bridging borohydride.⁹ Solution magnetic susceptibility studies (25 °C, THF) establish **1** as high-spin Fe(II) ($\mu_{\text{eff}} = 4.6 \pm 0.2\ \mu_{\text{B}}$).

Single-crystal X-ray diffraction (XRD) experiments confirmed **1** as an iron-dihydride with each hydride capped by pyramidalized trialkylboranes (Fe–B = 2.970 Å; $\sum B_x = 318.6^\circ$;




Fig. 3 (A) Reduction of **1**. (B) Molecular structures of **1** and **1-K(crypt)** (50% probability ellipsoids). For clarity, the 9-BBN substituents are displayed in wireframe. (C) Electrochemical comparison between **1** and $(\text{BBNPDP}^{\text{tBu}})\text{FeCl}_2$. (D) X-band EPR of **1-K(crypt)** at 10 K in 1 : 1 toluene/THF. (E and F) Spin-density isosurface plots (0.002 a.u.) of **1** and **1-K(crypt)**, respectively.

Fig. 3B). The geometry about iron is best described as square pyramidal ($\tau_5 = 0.31$). Complex **1** is a rare example of a mono-metallic high-spin Fe(II)-dihydride (or Fe(II)-diborohydride).^{1a,6c,9}

The requirements of the intramolecular acids to stabilize **1** were assessed with a control ligand, $\text{B}^{\text{u}}\text{PDP}^{\text{tBu}}$, where the $-(\text{CH}_2)_3\text{BBN}$ fragments are replaced with *n*-butyl groups (Fig. 2).⁷ Treating $(\text{B}^{\text{u}}\text{PDP}^{\text{tBu}})\text{FeBr}_2$ with 2 equiv. KBHET_3 afforded intractable mixtures, precluding characterization (see ESI[†]). We propose that the appended trialkylboranes in **1** serve two important roles: (1) to stabilize the hydride ligands, and (2) to decrease the entropic penalty for stabilization (intramolecular 9-BBN vs. exogenous BET_3). Attempts to install another donor ligand to **1** and enforce a low-spin configuration were unsuccessful: **1** was unreactive toward stoichiometric 1,4-diazabicyclo[2.2.2]octane, NMe_3 , and PMe_3 , but decomposed when treated with 4-dimethylaminopyridine.

To assess the requirements of the metal center and provide additional spectroscopic characterization, we synthesized the zinc analogue, $(\text{BBNPDP}^{\text{tBu}})\text{ZnH}_2$, from $(\text{BBNPDP}^{\text{tBu}})\text{ZnI}_2$ (Fig. 2). ^{11}B -NMR spectroscopy (THF) of $(\text{BBNPDP}^{\text{tBu}})\text{ZnH}_2$ revealed an upfield resonance at 5.43 ppm, consistent with a tetrahedral boron center.¹⁰ IR spectroscopy (KBr) revealed a broad Zn–H–B stretch at $\sim 1775\text{ cm}^{-1}$ that shifts upon deuterium labeling to $\sim 1300\text{ cm}^{-1}$.¹¹ ^2H -NMR spectroscopy established the assignment of the hydride resonance at 0.65 ppm, which is comparable to other $\kappa^1\text{-Zn}(\text{BH}_4)/\text{Zn}(\text{BH}_3\text{R})$ complexes.¹² The molecular structure of $(\text{BBNPDP}^{\text{tBu}})\text{ZnH}_2$ displays bonding metrics analogous to **1** (see ESI[†]).

Electrochemical investigation of **1** using cyclic voltammetry (0.2 M $[\text{Bu}_4\text{N}][\text{PF}_6]$, THF) revealed a quasi-reversible reductive event at -2.06 V vs. Fc/Fc^+ which is minimally shifted from $(\text{BBNPDP}^{\text{tBu}})\text{FeCl}_2$ ($\Delta = +10\text{ mV}$),¹³ despite their different X-type donors (Fig. 3C).¹⁴ The similar redox potentials suggest that, in analogy to the E_4 state of nitrogenase, reducing equivalents can be delivered at a near constant potential to a Fe(II) state, when stored as bridging hydride equivalents.¹⁵ To prepare the reduced complex, 1 equiv. of KC_8 was added to a freshly thawed THF solution of **1** in the presence of 2,2,2-cryptand, resulting in an immediate color change to a vibrant green species assigned as $[\text{K}(2,2,2\text{-cryptand})][(\text{BBNPDP}^{\text{tBu}})\text{FeH}_2]$ (**1-K(crypt)**; Fig. 3A).¹⁶ Investigation of **1-K(crypt)** by ^1H -NMR spectroscopy (THF) revealed a paramagnetically shifted spectrum with resonances ranging -70.6 – 95.0 ppm with solution C_2 symmetry. Samples of **1-K(crypt)** are less stable than **1** and have an approximate half-life of 12 h in THF at room temperature. Upon reduction, the Fe–H–B infrared absorption shifts to higher energy ($\sim 1866\text{ cm}^{-1}$; KBr). Solution magnetic susceptibility studies (25 °C, THF) of **1-K(crypt)** are consistent with an $S = 3/2$ complex ($\mu_{\text{eff}} = 4.1 \pm 0.1\ \mu_{\text{B}}$).¹⁷ X-band EPR spectroscopy was employed to confirm the spin-state. Regardless of coordination environment or geometry, $S = 1/2$ iron complexes, typically exhibit *g* values near the free electron value,¹⁸ while high-spin complexes exhibit *g*_x-tensors > 3.5 .¹⁹ **1-K(crypt)** displays a broad rhombic signal with *g* values of 5.6, 3.97, and 1.82 at 10 K in 1 : 1 THF/toluene glass (Fig. 3D), which suggests **1-K(crypt)** is best described as high-spin.²⁰ Reduced iron complexes with low coordination numbers (2, 3, occasionally 4) are often high-spin,^{19c,19e,21} while



those stabilized by strong-field ligands (*i.e.* NHC, porphyrin, phosphines) are low-spin. **1-K(crypt)** is a rare example of a high-spin complex with a coordination number ≥ 5 .^{19b}

For structural comparison, **1-K(crypt)** was examined by XRD. Data refinement revealed the anionic portion of **1-K(crypt)** to be geometrically similar ($\tau_5 = 0.36$) to **1** (Fig. 3B). The interacting trialkylboranes are equidistant from the metal center in **1** and **1-K(crypt)**. Upon reduction, the Fe–N_{pyridine} bond distance decreases from 2.178(5) to 2.021(6) consistent with enhanced π -backbonding.²² DFT optimized geometries for the high-spin configuration of **1** and **1-K(crypt)** are in agreement with their crystallographic structures.²³ The spin-density for **1** is localized on iron, consistent with a high-spin Fe(II) description (Fig. 3E). For **1-K(crypt)**, significant spin-density is localized on the pyridyl-moiety of the chelate. The calculated β -SOMO for **1-K(crypt)** corresponding to reduction is primarily comprised of ligand- π^* orbitals on the pyridyl moiety with minimal Fe contribution (19%). This analysis is consistent with reduction of **1** affording an $S = 3/2$ system through antiferromagnetic coupling of high-spin Fe(II) with [^{BBN}PDP^{tBu}]¹⁻.²⁴ The calculated electronic structure is consistent with crystallographic bond metrics.²⁵ Pyridyl- π^* population is reflected by C₃–C₄ bond elongation (**1**: 1.377(6); **1-K(crypt)**: 1.433(12)_{ave} Å; Fig. S85[†]).

We sought to examine the generality of the trialkylborane Lewis acids to stabilize both the accumulated hydrides and other small molecules. Addition of E–H substrates (E = NH₂, NHMe, NHPH, OH, PPh, SPh) to freshly thawed THF solutions of **1** afford **2-E** as orange-tan powders with production of H₂ (Fig. 4).²⁶ **2-E** are high-spin Fe(II)-species with solution magnetic susceptibilities ranging $\mu_{\text{eff}} = 4.5\text{--}5.5 \mu_{\text{B}}$ (THF, 25 °C). **2-OH** and **2-PHPh** provide diagnostic infrared handles with sharp $\nu_{\text{(OH)}}$ / $\nu_{\text{(PH)}}$ absorptions observed at 3630 and 2340 cm⁻¹ (KBr), respectively—each consistent with previously reported M–OH–BR₃²⁷ and Fe-phosphides.²⁸ ¹H-NMR spectroscopy revealed **2-NHMe**, **2-NHPH**, **2-OH**, and **2-PHPh** are C₂ symmetric in solution—consistent with Fe–E–B interactions remaining intact in solution. In contrast, **2-SPh** exhibits a solution C_{2v} symmetric spectrum that broadens upon cooling to –80 °C, consistent with reversible B–S binding; likely a consequence of a weaker B–S interaction.²⁹ This observation was supported by DFT analyses. In contrast to the strong B–O interaction in **2-OH** (favored by ~10 kcal per mol per interaction), the calculated B–S interaction is thermodynamically disfavored by ~9 kcal per mol per interaction. In solution, the B–SPh binding equilibria in **2-SPh** were arrested by treating **2-SPh** with two equiv. NH₃. Competitive binding with NH₃ afforded the ammonia-borane species, (^{BBN}PDP^{tBu})Fe(SPh)₂(NH₃)₂ (**3**), highlighting the utility of moderately acidic groups to reversibly interact with small molecule substrates (Fig. 5).

To assess the structural similarities, **2-E** were examined by XRD (Fig. 4). Each displays a pentacoordinate iron best described as distorted square-pyramidal with interacting trialkylboranes. The B-heteroatom distances range from 1.592(2)–2.0504(19) Å following the trend B–OH < B–NHMe < B–NHPH < B–PPh = B–SPh. The same trend is observed for the Fe-heteroatom distance with **2-OH** displaying the shortest bond (1.9812(13) Å) and **2-PHPh**/**2-SPh** displaying the longest bond



Fig. 4 Formation of **2-E** and molecular structures (50% probability ellipsoids) and averaged bond distances. For clarity, the 9-BBN substituents are displayed in wireframe.




Fig. 5 Reversible acid/base interaction in **2-SPh** and molecular structure of **3** (50% probability ellipsoids). For clarity, the 9-BBN and phenyl substituents are displayed in wireframe.






Fig. 6 π -acid induced H_2 extrusion from **1**.

(2.408(4) and 2.4134(5) Å, respectively). The Fe–E bonds are longer than most reported examples, and we attribute this difference to due to quenching the ligand lone-pair by the trialkylborane.^{4d,19b,28,30} Upon substituting the B–S bond of **2-SPH** with NH_3 to afford **3**, the Fe–S bond distance slightly decreases (0.02 Å) while the Fe–S–Ph angle expands (90.06(6) to 110.5(3)_{ave}°). The ability of the appended trialkylborane to stabilize various sizes of small molecules at the metal center is highlighted by comparing the Fe–B distances of **1** and **2-PHPH** which vary by nearly 0.9 Å (2.97 vs. 3.91 Å). The isolation of **2-E** highlights the utility of secondary coordination sphere interactions to not only stabilize hydrides at a metal center, but also to stabilize intermediate species following H_2 loss. Attempts to synthesize most compounds analogous to **2-E** from (^{Bu}PDP^{tBu})FeBr₂ through established methods were unsuccessful (see ESI†).^{19b,28,30b,31} The (^{Bu}PDP^{tBu})Fe(SPh)₂ complex was the only isolable compound, which is consistent with the observed stability of **3** in the absence of Lewis acid/base interactions.

The acid/base interactions can be viewed as an avenue for redox-leveling: the reduction potentials vary by 340 mV across the series of compounds (**1**, **2-E**, and (^{Bu}PDP^{tBu})FeX₂ (X = Cl, Br); Table S3†) despite their disparate ligand donor properties.³² Upon sequestering the Lewis acid of **2-SPH** with NH_3 to form **3**, the redox potential shifts more negative by 80 mV (−2.21 V vs. Fe/Fc⁺). This modest shift is consistent with the weak or dynamic B–S interaction of **2-SPH**. Further description of the Fe–E–B interactions were obtained from their DFT optimized structures.³³ The LUMOs of the β manifold for **2-E** provide insight into their reduction potentials.³⁴ The boron Lewis acids temper the donor properties of the X-type donor ligand, lowering their energy and percent composition in the frontier orbitals (see ESI† for details). The LUMOs of **1** and **2-E** span a modest range of 0.51 eV. In contrast, their computationally investigated truncated (^{Me}PDP^{tBu})Fe(E)₂ counterparts range 1.72 eV. In analogy to the near constant operating potential for N_2 reduction in nitrogenase where $\text{H}^-/\text{N}_x\text{H}_y$ components are present throughout the sequence, the [(^{Bu}PDP^{tBu})Fe] system accommodates small molecule substrates whose donor properties are regulated through Lewis acid/base interactions.

Given the unusual electronic and structural properties of **1**, **1-K(cript)**, and (^{Bu}PDP^{tBu})ZnH₂, Wiberg bond indices for were investigated for each of these complexes to gauge the bonding contribution of the M–H–B interaction (*i.e.* metal–hydride or metal–borohydride character). For each, the interaction is

predominantly borohydride in character with B–H indices ranging 0.65–0.78 and M–H indices ranging 0.14–0.22.³⁵ (^{Bu}PDP^{tBu})ZnH₂ displays the most M–H character whereas **1-K(cript)** displays the most B–H character—consistent with IR spectroscopy. While the data suggest the electronic structures are best described as borohydride, the M–H character varies across the series, indicating M–H reactivity may be accessible.

A key difference between metal–hydrides and borohydrides is the ability of the former to undergo reductive elimination. To test the hydride accumulation hypothesis, *i.e.* whether the bridging hydrides in **1** provide access to Fe(0) by eliminating H_2 , **1** was treated with a π -acid (Fig. 6). Addition of two equiv. 2,4,6-tri-*tert*-butylphenylisocyanide to **1** expels H_2 in 90(2)% yield and affords (^{Bu}PDP^{tBu})Fe(CNAr)₂.³⁶ In contrast, when subjecting the redox-inactive zinc variant, (^{Bu}PDP^{tBu})ZnH₂, to analogous conditions, only 7(3)% H_2 was detected. These results implicate that **1** contains accessible Fe–H character that is available for reactions. In contrast, the reactivity of (^{Bu}PDP^{tBu})ZnH₂ illustrates the inability to extrude H_2 from a borohydride when access to a low-valent metal is not possible.

Conclusions

In summary, we have described a system capable of accumulating hydride equivalents at iron. Intramolecular borane Lewis acid serve a prominent role to not only impart stability to Fe-dihydrides, but also enable previously unobserved electronic structures at iron by regulating the Fe–X-type ligand interactions. Similarities can be drawn between our system and the E₄ state of nitrogenase, where hydrides are accumulated at a near constant potential and addition of small molecule substrates induce H_2 elimination concomitant with substrate stabilization. The geometric flexibility of the appended Lewis acids were previously shown to enable interactions both independent and cooperatively with a metal; here the geometric flexibility is highlighted by their ability to stabilize small molecules of varied sizes. Work is ongoing to interrogate how Lewis acids can be used to further regulate redox transformations at the metal.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the NIH (1R01GM111486-01A1). N. K. S. is a Camille Dreyfus Teacher-Scholar. JJK is supported by the NIH NIGMS F32GM126635. We thank Suzanne Bart for EPR assistance. X-ray diffractometers were funded by the NSF (CHE 1625543).

Notes and references

- (a) H. Dai and H. Guan, Iron Dihydride Complexes: Synthesis, Reactivity, and Catalytic Applications, *Isr. J. Chem.*, 2017, 57, 1170–1203; (b) D. E. Prokopchuk, G. M. Chambers, E. D. Walter, M. T. Mock and



- R. M. Bullock, H₂ Binding, Splitting, and Net Hydrogen Atom Transfer at a Paramagnetic Iron Complex, *J. Am. Chem. Soc.*, 2019, **141**, 1871–1876.
- 2 R. Y. Igarashi, M. Laryukhin, P. C. Dos Santos, H.-I. Lee, D. R. Dean, L. C. Seefeldt and B. M. Hoffman, Trapping H-Bound to the Nitrogenase FeMo-Cofactor Active Site during H₂ Evolution: Characterization by ENDOR Spectroscopy, *J. Am. Chem. Soc.*, 2005, **127**, 6231–6241.
- 3 B. M. Hoffman, D. Lukoyanov, D. R. Dean and L. C. Seefeldt, Nitrogenase: A Draft Mechanism, *Acc. Chem. Res.*, 2013, **46**, 587–595.
- 4 (a) A. Jablonskytė, J. A. Wright, S. A. Fairhurst, J. N. T. Peck, S. K. Ibrahim, V. S. Oganessian and C. J. Pickett, Paramagnetic Bridging Hydrides of Relevance to Catalytic Hydrogen Evolution at Metallosulfur Centers, *J. Am. Chem. Soc.*, 2011, **133**, 18606–18609; (b) W. Wang, M. J. Nilges, T. B. Rauchfuss and M. Stein, Isolation of a Mixed Valence Diiron Hydride: Evidence for a Spectator Hydride in Hydrogen Evolution Catalysis, *J. Am. Chem. Soc.*, 2013, **135**, 3633–3639; (c) R. A. Kinney, C. T. Saouma, J. C. Peters and B. M. Hoffman, Modeling the Signatures of Hydrides in Metalloenzymes: ENDOR Analysis of a Di-iron Fe(μ-NH)(μ-H)Fe Core, *J. Am. Chem. Soc.*, 2012, **134**, 12637–12647; (d) J. M. Smith, R. J. Lachicotte and P. L. Holland, NN Bond Cleavage by a Low-Coordinate Iron(II) Hydride Complex, *J. Am. Chem. Soc.*, 2003, **125**, 15752–15753; (e) J. Rittle, C. C. L. McCrory and J. C. Peters, A 106-Fold Enhancement in N₂-Binding Affinity of an Fe₂(μ-H)₂ Core upon Reduction to a Mixed-Valence Fe^{II}Fe^I State, *J. Am. Chem. Soc.*, 2014, **136**, 13853–13862.
- 5 (a) E. W. Dahl, J. J. Kiernicki, M. Zeller and N. K. Szymczak, Hydrogen Bonds Dictate O₂ Capture and Release within a Zinc Tripod, *J. Am. Chem. Soc.*, 2018, **140**, 10075–10079; (b) E. W. Dahl, H. T. Dong and N. K. Szymczak, Phenylamino derivatives of tris(2-pyridylmethyl)amine: hydrogen-bonded peroxodicopper complexes, *Chem. Commun.*, 2018, **54**, 892–895; (c) C. M. Moore and N. K. Szymczak, Redox-induced fluoride ligand dissociation stabilized by intramolecular hydrogen bonding, *Chem. Commun.*, 2015, **51**, 5490–5492; (d) O. Tutusaus, C. Ni and N. K. Szymczak, A Transition Metal Lewis Acid/Base Triad System for Cooperative Substrate Binding, *J. Am. Chem. Soc.*, 2013, **135**, 3403–3406; (e) E. W. Dahl and N. K. Szymczak, Hydrogen Bonds Dictate the Coordination Geometry of Copper: Characterization of a Square-Planar Copper(I) Complex, *Angew. Chem., Int. Ed.*, 2016, **55**, 3101–3105.
- 6 (a) D.-H. Manz, P.-C. Duan, S. Dechert, S. Demeshko, R. Oswald, M. John, R. A. Mata and F. Meyer, Pairwise H₂/D₂ Exchange and H₂ Substitution at a Bimetallic Dinickel(II) Complex Featuring Two Terminal Hydrides, *J. Am. Chem. Soc.*, 2017, **139**, 16720–16731; (b) K. J. Anderton, B. J. Knight, A. L. Rheingold, K. A. Abboud, R. García-Serres and L. J. Murray, Reactivity of hydride bridges in a high-spin [Fe₃(μ-H)₃]³⁺ cluster: reversible H₂/CO exchange and Fe–H/B–F bond metathesis, *Chem. Sci.*, 2017, **8**, 4123–4129; (c) Y. Yu, A. R. Sadique, J. M. Smith, T. R. Dugan, R. E. Cowley, W. W. Brennessel, C. J. Flaschenriem, E. Bill, T. R. Cundari and P. L. Holland, The Reactivity Patterns of Low-Coordinate Iron–Hydride Complexes, *J. Am. Chem. Soc.*, 2008, **130**, 6624–6638.
- 7 J. J. Kiernicki, M. Zeller and N. K. Szymczak, Hydrazine Capture and N–N Bond Cleavage at Iron Enabled by Flexible Appended Lewis Acids, *J. Am. Chem. Soc.*, 2017, **139**, 18194–18197.
- 8 (a) M. A. Nesbit, D. L. M. Suess and J. C. Peters, E–H Bond Activations and Hydrosilylation Catalysis with Iron and Cobalt Metalloboranes, *Organometallics*, 2015, **34**, 4741–4752; (b) D. L. M. Suess and J. C. Peters, H–H and Si–H Bond Addition to Fe≡NNR₂ Intermediates Derived from N₂, *J. Am. Chem. Soc.*, 2013, **135**, 4938–4941; (c) E. Alberico, P. Sponholz, C. Cordes, M. Nielsen, H.-J. Drexler, W. Baumann, H. Junge and M. Beller, Selective Hydrogen Production from Methanol with a Defined Iron Pincer Catalyst under Mild Conditions, *Angew. Chem., Int. Ed.*, 2013, **52**, 14162–14166; (d) R. Bau, H. S. H. Yuan, M. V. Baker and L. D. Field, An X-ray study of FeH(dmpe)₂(BH₄): a compound containing a singly-bridged BH₄ ligand with a bent Fe–H–B linkage, *Inorg. Chim. Acta*, 1986, **114**, L27–L28; (e) D. L. M. Suess and J. C. Peters, A CO-Derived Iron Dicarbyne That Releases Olefin upon Hydrogenation, *J. Am. Chem. Soc.*, 2013, **135**, 12580–12583.
- 9 (a) J. S. Figueroa, J. G. Melnick and G. Parkin, Reactivity of the Metal → BX₃ Dative σ-Bond: 1,2-Addition Reactions of the Fe → BX₃ Moiety of the Ferraboratrane Complex [κ⁴-B(mimBut)₃]Fe(CO)₂, *Inorg. Chem.*, 2006, **45**, 7056–7058; (b) C. Kimblin, D. G. Churchill, B. M. Bridgewater, J. N. Girard, D. A. Quarless and G. Parkin, Tris(mercaptoimidazolyl)hydroborato complexes of cobalt and iron, [TmPh]₂M (M=Fe, Co): structural comparisons with their tris(pyrazolyl)hydroborato counterparts, *Polyhedron*, 2001, **20**, 1891–1896; (c) See ESI Table S21† for further literature comparisons to Fe–B bond distances and ν(Fe–H–B) vibrational data.
- 10 B–H coupling is not observed presumably due to low symmetry at boron.
- 11 Attempts at deuterium labeling **1** with LiBDEt₃ were unsuccessful.
- 12 S. Marks, R. Köppe, T. K. Panda and P. W. Roesky, Unprecedented Zinc–Borane Complexes, *Chem.–Eur. J.*, 2010, **16**, 7096–7100.
- 13 Potentials were assessed by square wave voltammetry.
- 14 A typical shift of 200–300 mV is expected per Cl/H substitution, see: (a) T. Liu, D. L. DuBois and R. M. Bullock, An iron complex with pendent amines as a molecular electrocatalyst for oxidation of hydrogen, *Nat. Chem.*, 2013, **5**, 228; (b) M. Tilset, I. Fjeldahl, J.-R. Hamon, P. Hamon, L. Toupet, J.-Y. Saillard, K. Costuas and A. Haynes, Theoretical, Thermodynamic, Spectroscopic, and Structural Studies of the Consequences of One-Electron Oxidation on the Fe–X Bonds in 17- and 18-Electron Cp*Fe(dppe)X Complexes (X = F, Cl, Br, I, H, CH₃), *J. Am. Chem. Soc.*, 2001, **123**, 9984–10000; (c) Y. Hu, L. Li, A. P. Shaw, J. R. Norton, W. Sattler and Y. Rong,



- Synthesis, Electrochemistry, and Reactivity of New Iridium(III) and Rhodium(III) Hydrides, *Organometallics*, 2012, **31**, 5058–5064; (d) A. P. Shaw, J. R. Norton, D. Buccella, L. A. Sites, S. S. Kleinbach, D. A. Jarem, K. M. Bocage and C. Nataro, Synthesis, Electrochemistry, and Reactivity of Half-Sandwich Ruthenium Complexes Bearing Metallocene-Based Bisphosphines, *Organometallics*, 2009, **28**, 3804–3814.
- 15 B. J. McNicholas, R. H. Grubbs, J. R. Winkler, H. B. Gray and E. Despagnet-Ayoub, Tuning the formal potential of ferrocyanide over a 2.1 V range, *Chem. Sci.*, 2019, **10**, 3623–3626.
- 16 Crown ethers are not required to isolate the reduced Fe-di(boro)hydride, see ESI.†
- 17 Y. Nakajima, Y. Nakao, S. Sakaki, Y. Tamada, T. Ono and F. Ozawa, Electronic Structure of Four-Coordinate Iron(I) Complex Supported by a Bis(phosphaethenyl)pyridine Ligand, *J. Am. Chem. Soc.*, 2010, **132**, 9934–9936.
- 18 (a) A. V. Polezhaev, C. J. Liss, J. Telser, C.-H. Chen and K. G. Caulton, A PNNH Pincer Ligand Allows Access to Monovalent Iron, *Chem.–Eur. J.*, 2018, **24**, 1330–1341; (b) C. E. MacBeth, S. B. Harkins and J. C. Peters, Synthesis and characterization of cationic iron complexes supported by the neutral ligands NPi-Pr₃, NArPi-Pr₃, and NSt-Bu₃, *Can. J. Chem.*, 2005, **83**, 332–340; (c) C. V. Thompson, I. Davis, J. A. DeGayner, H. D. Arman and Z. J. Tonzetich, Iron Pincer Complexes Incorporating Bipyridine: A Strategy for Stabilization of Reactive Species, *Organometallics*, 2017, **36**, 4928–4935; (d) N. Ehrlich, M. Kreye, D. Baabe, P. Schweyen, M. Freytag, P. G. Jones and M. D. Walter, Synthesis and Electronic Ground-State Properties of Pyrrolyl-Based Iron Pincer Complexes: Revisited, *Inorg. Chem.*, 2017, **56**, 8415–8422.
- 19 (a) A. McSkimming and W. H. Harman, A Terminal N₂ Complex of High-Spin Iron(I) in a Weak, Trigonal Ligand Field, *J. Am. Chem. Soc.*, 2015, **137**, 8940–8943; (b) J. S. Anderson, M.-E. Moret and J. C. Peters, Conversion of Fe–NH₂ to Fe–N₂ with release of NH₃, *J. Am. Chem. Soc.*, 2013, **135**, 534–537; (c) P. P. Samuel, K. C. Mondal, N. Amin Sk, H. W. Roesky, E. Carl, R. Neufeld, D. Stalke, S. Demeshko, F. Meyer, L. Ungur, L. F. Chibotaru, J. Christian, V. Ramachandran, J. van Tol and N. S. Dalal, Electronic Structure and Slow Magnetic Relaxation of Low-Coordinate Cyclic Alkyl(amino) Carbene Stabilized Iron(I) Complexes, *J. Am. Chem. Soc.*, 2014, **136**, 11964–11971; (d) S. A. Stoian, Y. Yu, J. M. Smith, P. L. Holland, E. L. Bominaar and E. Münck, Mössbauer, Electron Paramagnetic Resonance, and Crystallographic Characterization of a High-Spin Fe(I) Diketimate Complex with Orbital Degeneracy, *Inorg. Chem.*, 2005, **44**, 4915–4922; (e) K. P. Chiang, C. C. Scarborough, M. Horitani, N. S. Lees, K. Ding, T. R. Dugan, W. W. Brennessel, E. Bill, B. M. Hoffman and P. L. Holland, Characterization of the Fe–H Bond in a Three-Coordinate Terminal Hydride Complex of Iron(I), *Angew. Chem., Int. Ed.*, 2012, **51**, 3658–3662.
- 20 *g*-Values were obtained by simulation, see ESI.†
- 21 (a) Z. Ouyang, J. Du, L. Wang, J. L. Kneebone, M. L. Neidig and L. Deng, Linear and T-Shaped Iron(I) Complexes Supported by N-Heterocyclic Carbene Ligands: Synthesis and Structure Characterization, *Inorg. Chem.*, 2015, **54**, 8808–8816; (b) M. I. Lipschutz, T. Chantarojsiri, Y. Dong and T. D. Tilley, Synthesis, Characterization, and Alkyne Trimerization Catalysis of a Heteroleptic Two-Coordinate Fe^I Complex, *J. Am. Chem. Soc.*, 2015, **137**, 6366–6372.
- 22 The pyridine(dipyrazole) framework has not been observed to act as a redox-active ligand, see B. J. Cook, C.-H. Chen, M. Pink, R. L. Lord and K. G. Caulton, *Inorg. Chim. Acta*, 2016, **451**, 82–91. It has been proposed as an intermediate species before, see T. J. Sherbow, J. C. Fettinger and L. A. Berben, *Inorg. Chem.*, 2017, **56**, 8651–8660.
- 23 See ESI† for comparison between alternative spin configurations.
- 24 (a) A. M. Tondreau, S. C. E. Stieber, C. Milsmann, E. Lobkovsky, T. Weyhermüller, S. P. Semproni and P. J. Chirik, Oxidation and Reduction of Bis(imino)pyridine Iron Dinitrogen Complexes: Evidence for Formation of a Chelate Trianion, *Inorg. Chem.*, 2013, **52**, 635–646; (b) S. K. Russell, A. C. Bowman, E. Lobkovsky, K. Wieghardt and P. J. Chirik, Synthesis and Electronic Structure of Reduced Bis(imino)pyridine Manganese Compounds, *Eur. J. Inorg. Chem.*, 2012, **2012**, 535–545.
- 25 Q. Knijnenburg, S. Gambarotta and P. H. M. Budzelaar, Ligand-centred reactivity in diiminepyridine complexes, *Dalton Trans.*, 2006, 5442–5448.
- 26 Protonolysis reactions with **1-K(crypt)** resulted in disproportionation. Treating **1** with a diatom surrogate, Me₃SiCN, affords dimeric [(^{BBN}PDP^{tBu})Fe(NC)₂]₂, see ESI.†
- 27 (a) B. L. Conley and T. J. Williams, Thermochemistry and Molecular Structure of a Remarkable Agostic Interaction in a Heterobifunctional Ruthenium–Boron Complex, *J. Am. Chem. Soc.*, 2010, **132**, 1764–1765; (b) R. Choukroun, C. Lorber, L. Vendier and C. Lepetit, Vanadocene-Mediated Ionization of Water in the Aqua Species [H₂O·B(C₆F₅)₃]: Structural Characterization of the Hydride and Hydroxide Complexes [Cp₂V(μ-H)B(C₆F₅)₃] and [Cp₂V(μ-OH)B(C₆F₅)₃], *Organometallics*, 2006, **25**, 1551–1553; (c) N. Tsoureas and A. A. Danopoulos, Pyridyl- and diphenylphosphinoethyl-functionalised N-heterocyclic carbene platinum methyl complexes, *J. Organomet. Chem.*, 2015, **775**, 178–187.
- 28 A. K. Hickey, S. B. Muñoz, S. A. Lutz, M. Pink, C.-H. Chen and J. M. Smith, Arrested α-hydride migration activates a phosphido ligand for C–H insertion, *Chem. Commun.*, 2017, **53**, 412–415.
- 29 **2-SPh** and the zinc analogue can be prepared from the reaction of (^{BBN}PDP^{tBu})MH₂ with PhSSPh. Both reactions extrude H₂, see ESI.†
- 30 (a) S. E. Creutz and J. C. Peters, Exploring secondary-sphere interactions in Fe–N_xH_y complexes relevant to N₂ fixation, *Chem. Sci.*, 2017, **8**, 2321–2328; (b) Y. M. Badiei, M. A. Siegler and D. P. Goldberg, O₂ Activation by Bis(imino)pyridine Iron(II)–Thiolate Complexes, *J. Am. Chem. Soc.*, 2011, **133**, 1274–1277.



- 31 N. A. Eckert, J. M. Smith, R. J. Lachicotte and P. L. Holland, Low-Coordinate Iron(II) Amido Complexes of β -Diketiminates: Synthesis, Structure, and Reactivity, *Inorg. Chem.*, 2004, **43**, 3306–3321.
- 32 (a) R. D. Bemowski, A. K. Singh, B. J. Bajorek, Y. DePorre and A. L. Odom, Effective donor abilities of E-*t*-Bu and EPh (E = O, S, Se, Te) to a high valent transition metal, *Dalton Trans.*, 2014, **43**, 12299–12305; (b) S. A. DiFranco, N. A. Maciulis, R. J. Staples, R. J. Batrice and A. L. Odom, Evaluation of Donor and Steric Properties of Anionic Ligands on High Valent Transition Metals, *Inorg. Chem.*, 2012, **51**, 1187–1200.
- 33 Analysis of the LUMO's were used in lieu of the optimized reduced complexes of **2-E** and truncated variants because (1) the reorganization of **1** upon reduction is minimal, (2) the high correlation between the β -LUMO of **1** and the β -SOMO of **1-K(crypt)**, see Fig. S88.† An analysis LUMO energies and calculated reduction potentials or a subset of **1** and **2-E** is included in the ESI.†
- 34 (a) D. D. Méndez-Hernández, J. G. Gillmore, L. A. Montano, D. Gust, T. A. Moore, A. L. Moore and V. Mujica, Building and testing correlations for the estimation of one-electron reduction potentials of a diverse set of organic molecules, *J. Phys. Org. Chem.*, 2015, **28**, 320–328; (b) J. R. Levin, W. L. Dorfner, A. X. Dai, P. J. Carroll and E. J. Schelter, Density Functional Theory as a Predictive Tool for Cerium Redox Properties in Nonaqueous Solvents, *Inorg. Chem.*, 2016, **55**, 12651–12659.
- 35 (a) R. Hoffmann, P. von Ragué Schleyer and H. F. Schaefer III, Predicting Molecules—More Realism, Please!, *Angew. Chem., Int. Ed.*, 2008, **47**, 7164–7167; (b) The truncated analogue, (^{Me}PDP^{tBu})FeH₂, is calculated to have a M–H bond index of 0.65.
- 36 The assignment of (^{BBN}PDP^{tBu})Fe(CNAr)₂ is based on ¹H NMR and IR spectroscopies as well as MALDI mass spectrometry. We have been unable to isolate analytically pure samples of this species due to its high solubility.

