

Cite this: *Chem. Sci.*, 2021, 12, 8131

All publication charges for this article have been paid for by the Royal Society of Chemistry

# The origin of unidirectional charge separation in photosynthetic reaction centers: nonadiabatic quantum dynamics of exciton and charge in pigment–protein complexes†

Hiroyuki Tamura, \*<sup>ab</sup> Keisuke Saito <sup>ab</sup> and Hiroshi Ishikita <sup>ab</sup>

Exciton charge separation in photosynthetic reaction centers from purple bacteria (PbRC) and photosystem II (PSII) occurs exclusively along one of the two pseudo-symmetric branches (active branch) of pigment–protein complexes. The microscopic origin of unidirectional charge separation in photosynthesis remains controversial. Here we elucidate the essential factors leading to unidirectional charge separation in PbRC and PSII, using nonadiabatic quantum dynamics calculations in conjunction with time-dependent density functional theory (TDDFT) with the quantum mechanics/molecular mechanics/polarizable continuum model (QM/MM/PCM) method. This approach accounts for energetics, electronic coupling, and vibronic coupling of the pigment excited states under electrostatic interactions and polarization of whole protein environments. The calculated time constants of charge separation along the active branches of PbRC and PSII are similar to those observed in time-resolved spectroscopic experiments. In PbRC, Tyr-M210 near the accessory bacteriochlorophyll reduces the energy of the intermediate state and drastically accelerates charge separation overcoming the electron–hole interaction. Remarkably, even though both the active and inactive branches in PSII can accept excitons from light-harvesting complexes, charge separation in the inactive branch is prevented by a weak electronic coupling due to symmetry-breaking of the chlorophyll configurations. The exciton in the inactive branch in PSII can be transferred to the active branch *via* direct and indirect pathways. Subsequently, the ultrafast electron transfer to pheophytin in the active branch prevents exciton back transfer to the inactive branch, thereby achieving unidirectional charge separation.

Received 15th March 2021  
Accepted 3rd May 2021

DOI: 10.1039/d1sc01497h

rsc.li/chemical-science

## 1. Introduction

Light reactions of photosynthesis achieve an extremely high internal quantum efficiency from photoabsorption to separated electrons and holes<sup>1</sup> through ingeniously regulated pathways of energy and charge transfers in pigment–protein complexes. Light-harvesting (antenna) complexes, which contain a number of pigments, absorb a photon to create an electronically excited state characterized as a bound electron–hole pair, *i.e.* exciton.<sup>1–7</sup> Exciton charge separation necessitates a sufficient potential difference between the donor and acceptor of electrons for overcoming the electron–hole Coulomb binding energy.

Photosystem II (PSII) consists of core antenna complexes (CP43 and CP47) and a reaction center (RC).<sup>1–4</sup> Chlorophyll *a* (Chl) molecules in CP43 and CP47 mediate exciton transfers to

the RC consisting of Chls ( $P_{D1}$ ,  $P_{D2}$ ,  $Chl_{D1}$ , and  $Chl_{D2}$ ), pheophytin *a* ( $Pheo_{D1}$  and  $Pheo_{D2}$ ), and plastoquinone ( $Q_A$  and  $Q_B$ ) (Fig. 1).<sup>8–38</sup> Charge separation occurs in the RC, where the electron reduces plastoquinone and the hole eventually oxidizes water at the  $Mn_4CaO_5$  cluster.<sup>11–13</sup> Similarly, bacteriochlorophyll *a* (BChl) molecules in the light harvesting complex I (LHI) from purple bacteria, *Rhodobacter sphaeroides*, transfer an exciton to the RC (PbRC) consisting of BChl ( $P_L$ ,  $P_M$ ,  $B_L$ , and  $B_M$ ), bacterioopheophytin *a* (BPheo,  $H_L$  and  $H_M$ ), and ubiquinone ( $Q_A$  and  $Q_B$ ) (Fig. 1).<sup>11</sup>

Charge separation in PSII and PbRC occurs only along the active branch of the pseudo- $C_2$  symmetric pigment–protein complexes, *i.e.*, D1- and L-branches, respectively (Fig. 1).<sup>11</sup> The D2- and M-branches are referred to as inactive branches. PSII and PbRC may have been evolved from a common ancestor and classified as type II RCs.<sup>11–13</sup> In type II RCs,  $Q_B$  in the inactive branch accepts an electron from  $Q_A$  in the active branch while it does not directly accept an electron from (B) Pheo in the inactive branch (Fig. 1).

In PbRC, the strong electronic coupling between the special pair BChls,  $P_L$  and  $P_M$ , leads to stabilization of the delocalized

<sup>a</sup>Department of Applied Chemistry, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8654, Japan. E-mail: tamura@protein.rcast.u-tokyo.ac.jp

<sup>b</sup>Research Center for Advanced Science and Technology, The University of Tokyo, 4-6-1 Komaba, Meguro-ku, Tokyo 153-8904, Japan

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1sc01497h





Fig. 1 Exciton transfer<sup>5-7,9,10</sup> and charge separation<sup>11</sup> pathways in (a) PbRC and (b) PSII. Dashed lines indicate the rotation axis of pseudo- $C_2$  symmetry. Blue and red arrows indicate exciton and charge transfer pathways, respectively. Phe and Tyr denote phenylalanine and tyrosine.

exciton,  $(P_L P_M)^*$ .<sup>11,34</sup> The  $P_L P_M$  can accept an exciton from LHI, which absorbs a near infrared photon.<sup>5-7</sup> Time-resolved spectroscopic measurements indicated that the excited electron in  $(P_L P_M)^*$  is transferred to  $H_L$  via  $B_L$  along the L-branch on a time scale of a few ps.<sup>39-50</sup> Despite the pseudo- $C_2$  symmetric cofactor arrangement, the difference in the amino acid sequences between the L- and M-branches leads to the difference in the redox potentials of the pigments via electrostatic interactions and polarization.<sup>51-53</sup>

A previous study using time-dependent density functional theory (TDDFT) with the quantum mechanics/molecular mechanics/polarizable continuum model (QM/MM/PCM) method indicated that the intermediate states of charge separation along the L- and M-branches, i.e.,  $[P_L P_M]^+ B_L^-$  and  $[P_L P_M]^+ B_M^-$ , are lower and higher in energy than that of  $(P_L P_M)^*$ , respectively.<sup>34</sup>

In contrast to PbRC, the excitation energies of  $P_{D1}$  and  $P_{D2}$  in PSII are higher than those of  $Chl_{D1}$  and  $Chl_{D2}$ ,<sup>9,34</sup> where the exciton tends to be localized on a single pigment owing to a weak excitonic coupling. Charge separation in PSII creates a hole localized on  $P_{D1}^+$ , which is the nearest pigment to the  $Mn_4CaO_5$  cluster located on the D1 side.<sup>54-57</sup> The localized nature of a hole on  $P_{D1}^+$  is important for PSII to keep a high oxidation potential.<sup>58</sup>

In PSII, CP43 and CP47 transfer an exciton to the RC, presumably, via the peripheral Chls on the D1 ( $Chl_{D1}$ ) and D2 ( $Chl_{D2}$ ) sides.<sup>3,9,10</sup> Time-resolved spectroscopic measurements on PSII suggested that the primary electron transfer occurs from an exciton on  $Chl_{D1}^*$  to  $Phe_{O_{D1}}$  on a time scale of a few hundred fs.<sup>14,22</sup> The hole on  $Chl_{D1}^{+}$  is, in turn, transferred to  $P_{D1}$  on a time scale of a few ps.<sup>14</sup>

Because the potential for electron transfer is energetically downhill along both the D1- and D2-branches toward  $Phe_{O_{D1}}$  and  $Phe_{O_{D2}}$ , respectively,<sup>33,34</sup> the energetics alone cannot explain unidirectional charge separation in PSII. Given that both  $Chl_{D1}$  and  $Chl_{D2}$  can accept an exciton from the core antenna complexes, the mechanism that leads to charge separation exclusively along the D1-branch is of particular interest. The charge separation pathways in pigment-protein complexes can be determined by various factors including energetics, electronic coupling, vibronic coupling, and quantum effects.<sup>2-4,18-21,31,44-46</sup>

In this study, we address the long-standing open question as to how PbRC and PSII achieve unidirectional charge separation exclusively along the active branch, by means of nonadiabatic quantum dynamics calculations<sup>59-62</sup> parametrized on the basis of TDDFT in the framework of the QM/MM/PCM method.<sup>63,64</sup> First, we show that the experimentally observed kinetics of charge separation along the active branches of PbRC and PSII are fairly well reproduced by nonadiabatic quantum dynamics calculations, which is based on the energetics and electronic coupling of the pigments, accounting for electrostatic interactions and polarization of whole protein environments from the X-ray crystal structures. On this basis, we clarify the essential factors which regulate the charge separation pathways in the reaction centers.

## 2. Methods

The energetics and electronic couplings in PbRC and PSII are analyzed by means of the polarizable QM/MM/PCM method, using the QuanPol code<sup>63</sup> implemented in the GAMESS code.<sup>65</sup> The electronic states in the QM regions are calculated using DFT and TDDFT with the CAMB3LYP functional<sup>66</sup> with the range separation parameter  $\mu$  of 0.14,  $\alpha$  of 0.19, and  $\beta$  of 0.46, which is well suited for the present systems including charge separated states.<sup>34</sup> The quantitative values of excitation energies may depend on functionals and parameters.<sup>32,38,67</sup> The 6-31G(d) basis set is used for all the QM calculations.

The QM region comprises pigments, ligands, hydrogen bonded water, and residues which interact directly with pigments as detailed in a previous report.<sup>34</sup> A polarizable amber-02 force field<sup>68</sup> is applied for proteins in the MM region, where induced dipoles of the MM atoms are taken into account to reproduce the dielectric screening. The PCM with a dielectric constant of 80 is applied to reproduce the polarization of water, which surrounds the proteins and fills the cavities. The PCM in the QuanPol code is based on a conductor-like screening model,<sup>63,64</sup> where the polarization points are put on spheres of radius 3.0 Å from the atom positions.<sup>34</sup> All atoms from the X-ray crystal structures are explicitly considered, where each MM atom contains an induced dipole in addition to the permanent charge. The induced dipole of each MM atom is determined iteratively together with the self-consistent field calculation of



electronic states, considering the electrostatic interactions with the electrons and nuclei in the QM region as well as the permanent charges and induced dipoles of other MM atoms.<sup>63</sup> The molecular orbital levels of the cofactors calculated using QM/MM reproduce the redox potential values calculated solving the Poisson–Boltzmann equation.<sup>33,34,57,69</sup> While the dielectric constant for the membrane region may be lower than 80 (*e.g.* 20),<sup>70</sup> a small dielectric constant makes the electrostatic interactions with the charged groups in the membrane-extrinsic region overestimated for membrane proteins. The optimal values for the dielectric constant depend on the protein model used.<sup>71,72</sup> The dielectric constant of 80 for the bulk region appears to be optimal for the present models, as suggested previously.<sup>33,34</sup>

The atomic coordinates of PSII and PbRC are obtained from the X-ray crystal structures from *Thermosynechococcus vulcanus* at 1.9 Å resolution (PDB code, 3ARC)<sup>73</sup> and from *Rhodobacter sphaeroides* at 2.01 Å resolution (PDB code, 3I4D),<sup>74</sup> respectively. The intramolecular reorganization energies of pigments are calculated through geometry optimization with QM/MM, where DFT with the CAMB3LYP functional plus Grimme's dispersion correction<sup>75</sup> is used for the QM region. The atomic coordinates of the MM region are fixed to the X-ray crystal structures. The reorganization energy of the MM region is not taken into account.

The electronic coupling between excited states is evaluated on the basis of the diabaticization scheme for TDDFT<sup>76</sup> in the framework of the QM/MM/PCM method.<sup>34,77</sup> The protocol of diabaticization is summarized below.

(1) We prepare a set of reference wavefunctions,  $\Phi_I$ , that possess pure characters of the excited states such as an exciton on a single molecule (*i.e.*, Frenkel exciton) and charge separated states for decoupled molecules.

(2) We calculate adiabatic electronic states in the pigment–protein complexes using TDDFT-QM/MM/PCM.

(3) The diabatic wavefunctions are expressed as a linear combination of the adiabatic wavefunctions,  $\Psi_J$ , by evaluating the overlap integrals between the reference and adiabatic wavefunctions:

$$\Phi_I = \sum_J C_{IJ} \Psi_J, \quad C_{IJ} = \langle \Psi_J | \Phi_I \rangle \quad (1)$$

That is, the adiabatic states from the TDDFT-QM/MM/PCM calculations are considered as basis functions for expanding the diabatic states. We consider 10 adiabatic states for expanding the diabatic states. The diabatic coupling is then evaluated as follows:

$$H_{IJ} = \langle \Phi_I | H | \Phi_J \rangle, \quad (2)$$

where  $H$  is the electronic Hamiltonian. The excitonic coupling in the present scheme includes both the Coulomb (Förster) and electron exchange (Dexter) contributions.<sup>76</sup>

For the nonadiabatic quantum dynamics calculations, we consider the following linear vibronic coupling Hamiltonian in the diabatic representation:

$$H = \sum_I h_I(x) |I\rangle \langle I| + \sum_{I \neq J} H_{IJ} (|I\rangle \langle J| + |J\rangle \langle I|) \quad (3)$$

$$h_I(x) = \sum_i \frac{\omega_i}{2} (p_i^2 + x_i^2) + \sum_i \kappa_i^I x_i + H_{II} \quad (4)$$

$H_{IJ}$  is the diabatic coupling (electronic coupling) between the states  $I$  and  $J$ .  $H_{II}$  is the vertical excitation energy of the  $I$ th electronic states.  $\omega_i$ ,  $x_i$ , and  $p_i$  are the frequency, position, and momentum of the  $i$ th vibrational mode (harmonic oscillator) in the dimensionless coordinate.  $\kappa_i^I$  is the vibronic coupling of the  $i$ th vibrational mode in the  $I$ th electronic state.

The exciton on the special pair, (P<sub>L</sub>P<sub>M</sub>)\*, is considered for the initial conditions of the quantum dynamics calculations of charge separation in PbRC. For PSII, in addition to the exciton localized on Chl<sub>D1</sub>\* in the D1-branch, Chl<sub>D2</sub>\* in the D2-branch is also considered for the initial conditions of the quantum dynamics calculations of charge separation. For the exciton transfer between Chl<sub>D2</sub>\* and Chl<sub>D1</sub>\*, the direct pathway and the indirect pathway *via* P<sub>D1</sub>\* and P<sub>D2</sub>\* are considered, where the quantum dynamics calculations account for the interference of the phase factors from several pathways. The initial vibrational wave packet is put on the Franck–Condon region of the initial electronic state.

$\omega_i$  and  $\kappa_i^I$  in eqn (4), *i.e.*, spectral density, are determined on the basis of the normal mode analysis and the geometry optimization of the pigments using the QM/MM/PCM method, where the atomic displacements from the Franck–Condon region to the potential bottom on the respective electronic states are projected onto the normal modes. The present model explicitly considers the vibronic couplings of the pigments and axial ligands, which are relevant to the dynamics of charge separation on a time scale of a few ps, whereas slow vibrational modes from surrounding proteins are neglected. The vibrational modes are reduced to a limited number of effective modes which reproduce the short-time dynamics and the reorganization energy of the system (see ESI†).<sup>60–62</sup> We consider 25 effective modes for each pigment, unless otherwise noted. For charge separation in PSII *via* indirect exciton transfer from Chl<sub>D1</sub>\* to Chl<sub>D2</sub>\*, 10 effective modes are considered for the respective intermediate states, P<sub>D1</sub>\* and P<sub>D2</sub>\*. The multi-configuration time-dependent Hartree (MCTDH) method<sup>59</sup> is used for the nonadiabatic quantum dynamics calculations, which properly consider correlations among the nuclear degrees of freedom, the Franck–Condon factor of vibrational wavefunctions, and vibrational energy redistribution along with electronic state transitions.

For analyzing the time constants of the first ( $\tau_1$ ) and second ( $\tau_2$ ) charge transfers along the active branches,  $\tau_1$  and  $\tau_2$  in the following rate equations are determined *via* curve fitting against the populations of the exciton ( $P_{EX}$ ), and the first ( $P_{CS1}$ ) and second ( $P_{CS2}$ ) charge separated states in the quantum dynamics calculations:

$$\frac{\partial P_{EX}}{\partial t} = -\frac{P_{EX}}{\tau_1}, \quad \frac{\partial P_{CS1}}{\partial t} = \frac{P_{EX}}{\tau_1} - \frac{P_{CS1}}{\tau_2}, \quad \frac{\partial P_{CS2}}{\partial t} = \frac{P_{CS1}}{\tau_2}, \quad (5)$$



where the corresponding kinetic scheme is expressed as follows:



### 3. Results and discussion

#### 3.1. Charge separation in PbRC

$(P_L P_M)$  in PbRC can be regarded as a single molecular site owing to the strong electronic coupling.<sup>34</sup> The electron transfers from  $(P_L P_M)^*$  to  $(P_L P_M)^{+} B_L^{-}$  and  $(P_L P_M)^{+} B_M^{-}$  are exothermic (downhill) and endothermic (uphill), respectively (Fig. 2a).<sup>34</sup> As a benchmark, we first compare the calculated time constants of charge separation along the L-branch with the corresponding experimental values. The quantum dynamics calculations

indicate that  $(P_L P_M)^*$  initially transfers the excited electron to  $B_L^{-}$  on a time scale of  $\tau_1 \approx 3.2$  ps.  $B_L^{-}$ , in turn, transfers the electron to  $H_L$  on a time scale of  $\tau_2 \approx 1.8$  ps (Fig. 2c and f). A similar order of time constants was observed in time-resolved spectroscopic measurements on charge separation in PbRC ( $\tau_1 = 3.5 \pm 0.4$  ps and  $\tau_2 = 1.2 \pm 0.3$  ps).<sup>41</sup>

The electronic coupling of the  $B_L^{-} \rightarrow H_L$  transfer (16 meV) is stronger than that of the  $(P_L P_M)^* \rightarrow (P_L P_M)^{+} B_L^{-}$  transfer (5 meV) (Fig. 2e). Thus, the population of the intermediate  $(P_L P_M)^{+} B_L^{-}$  state is kept small (Fig. 2c). The fast electron transfer from  $B_L^{-}$  to  $H_L$  is advantageous for preventing charge recombination, because  $(P_L P_M)^{+} H_L^{-}$  is difficult to decay to the ground state owing to a negligibly small orbital overlap between  $(P_L P_M)^{+}$  and  $H_L^{-}$ . Charge separation along the M-branch is negligibly slow, because the intermediate  $(P_L P_M)^{+} B_M^{-}$  state is substantially higher in energy than




Fig. 2 Bottom-to-bottom (adiabatic) excitation energies of the electronic states considering the intramolecular reorganization energies in (a) wild type and (b) Y(M210)F mutant PbRC. Dotted lines indicate the destabilized charge separated states in Y(M210)F mutant PbRC. Population of electronic states during quantum dynamics calculations of charge separation in (c) wild type and (d) Y(M210)F mutant PbRC, where the  $(P_L P_M)^*$ ,  $(P_L P_M)^{+} B_L^{-}$ , and  $(P_L P_M)^{+} H_L^{-}$  states along the L-branch are considered. Dotted lines indicate curve fitting by using eqn (5). (e) Diagram of charge transfer pathways (red lines) with electronic coupling (meV). Tyr-M210 is shown in yellow. (f) Diagram of the electron and hole locations in the  $(P_L P_M)^{+} B_L^{-}$  and  $(P_L P_M)^{+} H_L^{-}$  states with  $\tau_1$  and  $\tau_2$  (ps) for wild type PbRC ( $\sim 5$  ps in total).  $\tau_1$  and  $\tau_2$  for the mutant PbRC are 110 and 8 ps, respectively ( $\sim 118$  ps in total).





$(P_L P_M)^*$ , even though  $(P_L P_M)^+ H_M^-$  is lower in energy than  $(P_L P_M)^*$  (Fig. 2a).

The previous time-resolved spectroscopic measurements of mutant PbRC suggested that some specific residues especially contribute to unidirectional charge separation.<sup>47–50,52</sup> We have extensively analyzed the contribution of each residue to the potential shift on the pigments one by one, and concluded that Tyr-M210 near  $B_L$  has the largest contribution to the stabilization of  $B_L^-$ ,<sup>33,34</sup> where Phe-L181 is located at the counterpart position near  $B_M$ .

To verify the essential role of Tyr-M210, we consider the mutation of Tyr-M210 to phenylalanine, Y(M210)F, and investigate charge separation in the mutant PbRC by means of quantum dynamics calculations. The present TDDFT-QM/MM/PCM calculations indicate that the Y(M210)F mutation, in which the hydroxyl group is replaced with hydrogen, makes the intermediate  $(P_L P_M)^+ B_L^-$  state energetically uphill with respect to  $(P_L P_M)^*$ , even though the final  $(P_L P_M)^+ H_L^-$  state remains downhill (Fig. 2b). The quantum dynamics calculation indicates that the destabilization of the intermediate  $(P_L P_M)^+ B_L^-$  state drastically slows charge separation along the L-branch through the superexchange mechanism (Fig. 2d). This trend is qualitatively consistent with the experimental observations for the mutant PbRC,<sup>47–50,52</sup> where the calculated time constant ( $\sim 118$  ps) is quantitatively larger than the experimental values ( $\sim 16$

ps).<sup>47</sup> Here, only the local geometry of Phe-M210 was optimized in QM/MM, fixing surrounding proteins at the original geometry of wild type, although the mutation may also affect the surrounding geometry. Overall, the present analysis highlights the impact of the electrostatic interaction of Tyr-M210 on the efficient charge separation along the L-branch.

### 3.2. Charge separation in PSII

In PSII,  $Chl_{D1}$  and  $Chl_{D2}$  are supposed to accept an exciton from CP43 and CP47 *via*  $Chl_{D1}^*$  and  $Chl_{D2}^*$ , respectively (Fig. 3a).<sup>9,10</sup> The present calculations indicate that the bottom-to-bottom excitation energy of  $Chl_{D1}^*$  (1991 meV) lies between those of  $Chl_{D1}^*$  (1965 meV) and  $P_{D1}^*$  (2032 meV, Fig. 3b). Similarly, the  $Chl_{D2}^*$  energy (2015 meV) lies between those of  $Chl_{D2}^*$  (1992 meV) and  $P_{D2}^*$  (2038 meV, Fig. 3b). Thus,  $Chl_{D1}^*$  and  $Chl_{D2}^*$  can accept an exciton from  $Chl_{D1}^*$  and  $Chl_{D2}^*$ , respectively, in terms of energetics.

The absolute values of the calculated excitonic couplings in PSII are in the range of 7 to 15 meV (Table 1). The lowest and second lowest excitons obtained by diagonalizing the coupling matrix are localized on  $Chl_{D1}^*$  and  $Chl_{D2}^*$ , respectively (Fig. S2†), which can be regarded as Frenkel excitons. The quantitative values of the exciton energies in PSII calculated using TDDFT-QM/MM/PCM with the CAMB3LYP functional tend to be blue-shifted as compared to the experimental values,<sup>25–27</sup> where the




Fig. 3 (a) Arrangement of Chl molecules in PSII. Calculated bottom-to-bottom (adiabatic) excitation energies considering the intramolecular reorganization energies of the (b) exciton and (c) exciton and charge separated states in PSII. The intermediate and final charge separated states are indicated in red and blue, respectively.



Table 1 Electronic coupling in PSII (meV)

	$P_{D1}^*$	$P_{D2}^*$	$Chl_{D1}^*$	$Chl_{D2}^*$	$Chl_{D1}^{*+}Pheo_{D1}^{*-}$
$P_{D2}^*$	-10.1				
$Chl_{D1}^*$	7.4	-13.6			
$Chl_{D2}^*$	-14.4	7.0	-1.8		
$Chl_{D1}^{*+}Pheo_{D1}^{*-}$			-21.6		
$P_{D1}^{*+}Pheo_{D1}^{*-}$					6.4
$P_{D1}^{*+}Chl_{D1}^{*-}$	-6.5		5.4		
$P_{D1}^{*+}Chl_{D2}^{*-}$	0.7			-0.3	
$P_{D2}^{*+}Chl_{D1}^{*-}$		-0.5	-5.4		

calculated lowest exciton energy of 632 nm is blue-shifted as compared to the experimental value of 680 nm.<sup>25–27</sup>

We analyze charge separation from an exciton on  $Chl_{D1}^*$  by means of quantum dynamics calculations. The initial electron transfer from  $Chl_{D1}^*$  to  $Chl_{D1}^{*+}Pheo_{D1}^{*-}$  occurs on an ultrafast time scale ( $\tau_1 \approx 0.15$  ps) (Fig. 4a and c) owing to a strong electronic coupling ( $\sim 22$  meV, Fig. 4c and Table 1). The subsequent hole transfer to  $P_{D1}^{*+}Pheo_{D1}^{*-}$  occurs on a time scale of  $\tau_2 \approx 3.7$  ps (Fig. 4a and c). Thus, once  $Chl_{D1}^*$  accepts an exciton, charge separation occurs efficiently along the D1-branch. Similar time constants of charge separation in PSII were observed in the time-resolved spectroscopy measurements.<sup>14,22</sup> Another charge separation pathway,  $Chl_{D1}^* \rightarrow P_{D1}^{*+}Chl_{D1}^{*-}$ , is endothermic (Fig. 3c) and thus cannot compete with  $Chl_{D1}^* \rightarrow Chl_{D1}^{*+}Pheo_{D1}^{*-}$ . Although other charge separation pathways from an exciton on  $P_{D1}^*$  and  $P_{D2}^*$  were also proposed,<sup>23,24</sup> the quantum dynamical analysis for these pathways is beyond the scope of the present study. Overall, we can conclude that charge separation along the D1-branch proceeds via two-step  $Chl_{D1}^* \rightarrow Chl_{D1}^{*+}Pheo_{D1}^{*-}$  and  $Chl_{D1}^{*+} \rightarrow P_{D1}^{*+}$  transfers, considering the quantum dynamical analysis based on the energetics and electronic couplings from the QM/MM/PCM method.

The electronic coupling of the  $Chl_{D1}^* \rightarrow Chl_{D1}^{*+}Pheo_{D1}^{*-}$  transfer ( $\sim 22$  meV) is stronger than that of the  $Chl_{D1}^* \rightarrow P_{D1}^{*+}$  transfer ( $\sim 6$  meV, Fig. 4c and Table 1). The strong electronic coupling between the accessory Chl/BChl and the Pheo/BPheo is a common feature of PSII/PbRC. Nevertheless, the  $B_L^{*-} \rightarrow H_L^{*-}$  electron transfer ( $\sim 1.8$  ps) in PbRC is slower than the  $Chl_{D1}^* \rightarrow Chl_{D1}^{*+}Pheo_{D1}^{*-}$  transfer ( $\sim 0.15$  ps) in PSII, because the population of the  $(P_L P_M)^{*+}B_L^{*-}$  intermediate state in PbRC is kept small.

Because  $Chl_{D2}$  can also accept an exciton from CP47 on the D2 side,<sup>3,9,10</sup> the question arises as to how the exciton on  $Chl_{D2}^*$  eventually undergoes charge separation in the D1-branch. To analyze charge separation mechanisms from an exciton in the D2-branch, we carried out quantum dynamics calculations considering the initial exciton localized on  $Chl_{D2}^*$ . The charge separated state in the D2-branch,  $P_{D1}^{*+}Pheo_{D2}^{*-}$ , is less stable than that in the D1-branch,  $P_{D1}^{*+}Pheo_{D1}^{*-}$  (Fig. 3c), owing mainly to a difference in the potentials between  $Pheo_{D1}^{*-}$  and  $Pheo_{D2}^{*-}$ .<sup>33,34,36</sup>

The most stable charge separated state in the D2-branch is  $P_{D1}^{*+}Chl_{D2}^{*-}$  (Fig. 3c). However, PSII can avoid charge separation from  $Chl_{D2}^*$  to  $P_{D1}^{*+}Chl_{D2}^{*-}$  (Fig. 4b) because of a weak electronic coupling ( $\sim 0.3$  meV, Fig. 4d and Table 1), which is

significantly weaker than that between  $Chl_{D1}^*$  and  $P_{D2}^{*+}Chl_{D1}^{*-}$  ( $\sim 5.4$  meV) on the counterpart side (Fig. 5a and Table 1). The difference originates from the difference in the vinyl-group orientation between  $P_{D1}$  and  $P_{D2}$  (Fig. 5). The vinyl group is rather in plane for  $P_{D1}$  and out of plane for  $P_{D2}$  (Fig. 6).<sup>57</sup> The present results indicate that the in-plane  $P_{D1}$  vinyl group interferes with the  $\pi$ - $\pi$  interaction between  $P_{D1}$  and  $Chl_{D2}$  (Fig. 5b), thereby preventing the charge transfer to form  $P_{D1}^{*+}Chl_{D2}^{*-}$ .

The out-of-plane orientation of the  $P_{D2}$  vinyl group is caused by the relatively large steric hindrance from the  $P_{D1}$  phytol chain as compared with that between the  $P_{D1}$  vinyl group and the  $P_{D2}$  phytol chain (Fig. 5c and 6). The potential curves calculated using QM/MM indicate that the rotations of the  $P_{D1}$  and  $P_{D2}$  vinyl groups are hindered in the protein environments (Fig. 6 and S1†). Umena *et al.* reported that the conformations of vinyl groups were determined unambiguously from the corresponding electron density distributions and most of the vinyl groups are located in or near the same plane of the chlorine ring,<sup>73</sup> which suggests that the out-of-plane vinyl orientation for  $P_{D2}$  is exceptional. The same conformations of the  $P_{D1}$  and  $P_{D2}$  vinyl groups have also been observed in the X-ray free electron laser (XFEL) structure.<sup>78</sup> Thus, the observed vinyl orientations of  $P_{D1}$  and  $P_{D2}$  are considered to be robust in the protein environments. The phytol chains of  $P_{D1}$  and  $P_{D2}$  are less flexible due to the presence of the highly packed protein environment of D1/D2/CP43/CP47, as compared to those exposed to the protein surface in antenna proteins (*e.g.*, LH1 and the Fenna-Matthews-Olson protein). Umena *et al.* also confirmed that all of the C8 and C13 positions in the phytol chains have a (*R,R*) configuration as indicated by the low *B*-factor values.<sup>73</sup> Notably, the difference in the phytol-chain conformation also contributes to the asymmetric hole distribution on  $P_{D1}$  and  $P_{D2}$ , *i.e.*,  $P_{D1}^{*+} > P_{D2}^{*+}$ .<sup>57</sup> These results suggest that the symmetry-breaking of the  $P_{D1}P_{D2}$  geometry not only increases the  $P_{D1}^{*+}$  population, which facilitates water oxidation at the  $Mn_4CaO_5$  moiety on the D1 side, but also prevents charge separation along the D2-branch via a weak electronic coupling between  $Chl_{D2}^*$  and  $P_{D1}^{*+}Chl_{D2}^{*-}$ .

The quantum dynamics calculations indicate that the exciton on  $Chl_{D2}^*$  can be transferred to  $Chl_{D1}$  via the direct pathway and the indirect pathway mediated by  $P_{D1}^*$  and  $P_{D2}^*$  owing to adequately strong excitonic couplings (Table 1 and Fig. 4g) and small energy differences (Fig. 4h). Note that among the residues near  $Chl_{D1}$  and  $Chl_{D2}$ , D1-Met172 adjacent to  $Chl_{D1}$  contributes to the difference in excitation energy between  $Chl_{D1}^*$  and  $Chl_{D2}^*$ .<sup>34</sup> Previous calculations by Sirohiwal *et al.*<sup>38</sup> using other DFT functionals and an equation-of-motion coupled cluster method also indicated that  $Chl_{D1}$  exhibits the lowest excitation energy in the protein environment of PSII. Once the exciton is transferred to  $Chl_{D1}$ , subsequent charge separation to  $Chl_{D1}^{*+}Pheo_{D1}^{*-}$  occurs rapidly. The present analysis indicates that an overall time scale of charge separation from  $Chl_{D2}^*$  to  $Chl_{D1}^{*+}Pheo_{D1}^{*-}$  is in the order of a few tens ps (Fig. 4e and f), where exponential fitting indicates a  $\tau$  of  $\sim 50$  ps. It is highly likely that the exciton on  $Chl_{D2}^*$  can eventually undergo charge separation in the D1-branch without charge separation in the






Fig. 4 Population of excited states during quantum dynamics calculations of charge separation in PSII: (a)  $\text{Chl}_{\text{D1}}^*$ ,  $\text{Chl}_{\text{D1}}^+\text{Pheo}_{\text{D1}}^-$ , and  $\text{P}_{\text{D1}}^+\text{Pheo}_{\text{D1}}^-$  along the D1-branch and (b) from  $\text{Chl}_{\text{D2}}^*$  to  $\text{P}_{\text{D1}}^+\text{Chl}_{\text{D2}}^-$ . Dotted lines indicate curve fitting by using eqn (5). Diagram of charge separation with the absolute value of electronic coupling (meV): (c) from  $\text{Chl}_{\text{D1}}^*$  and (d) from  $\text{Chl}_{\text{D2}}^*$ . Population of excited states during quantum dynamics calculations from  $\text{Chl}_{\text{D2}}^*$ : (e) simultaneously considering indirect  $\text{Chl}_{\text{D2}}^* \rightarrow (\text{P}_{\text{D1}}^*, \text{P}_{\text{D2}}^*) \rightarrow \text{Chl}_{\text{D1}}^* \rightarrow \text{Chl}_{\text{D1}}^+\text{Pheo}_{\text{D1}}^-$  ( $\tau \approx 50$  ps) and direct  $\text{Chl}_{\text{D2}}^* \rightarrow \text{Chl}_{\text{D1}}^* \rightarrow \text{Chl}_{\text{D1}}^+\text{Pheo}_{\text{D1}}^-$  pathways, and (f)  $\text{Chl}_{\text{D1}}^+\text{Pheo}_{\text{D1}}^-$  population considering only the indirect (black) or the direct (blue) pathway. The  $\text{P}_{\text{D1}}^+\text{Chl}_{\text{D2}}^-$  population (green) is shown again for comparison. (g) Exciton transfer pathways (blue arrows) with the excitonic coupling (meV). Orange arrows indicate the direction of transition dipole moment. Solid and dotted arrows indicate indirect and direct exciton transfers, respectively. (h) Exciton transfer and charge separation pathways with the excitation energy.

D2-branch. Even though the energy difference between  $\text{Chl}_{\text{D1}}^*$  and  $\text{Chl}_{\text{D2}}^*$  is small (Fig. 4h), the ultrafast charge separation from  $\text{Chl}_{\text{D1}}^*$  to  $\text{Chl}_{\text{D1}}^+\text{Pheo}_{\text{D1}}^-$  prevents exciton back transfer to

$\text{Chl}_{\text{D2}}$ , enhancing the robustness of unidirectional charge separation along the D1-branch. The charge separation pathway via the exciton transfer from the D2- to D1-branches may








Fig. 5 Equilibrium geometry and the highest occupied molecular orbital (HOMO) of the (a)  $P_{D2}Chl_{D1}$  dimer and (b)  $P_{D1}Chl_{D2}$  dimer. Red circles indicate the vinyl groups. Electronic coupling (meV) between  $Chl_{D1}^*$  and  $P_{D2}^+Chl_{D1}^{*-}$  and that between  $Chl_{D2}^*$  and  $P_{D1}^+Chl_{D2}^{*-}$  are shown. (c) Configuration of the vinyl (red circles) and phytol (blue circles) groups of  $P_{D1}$  and  $P_{D2}$ .




Fig. 6 Potential energy curve as a function of the C-C-C=C torsion angle (degree) of the vinyl group of (a)  $P_{D1}$  and (b)  $P_{D2}$  in PSII, where the geometry of chlorophyll is optimized by QM/MM, fixing the torsion angle. The vinyl group is in the same plane as the chlorin ring at 0 degree. The black arrows indicate the equilibrium angle.

correspond to the delayed component observed in the time-resolved spectroscopic measurements apart from the ultrafast  $Chl_{D1}^* \rightarrow Chl_{D1}^+Pheo_{D1}^{*-}$  charge separation within the D1-branch.<sup>14</sup>

The direct excitonic coupling between  $Chl_{D1}^*$  and  $Chl_{D2}^*$  is relatively weak ( $\sim 2$  meV) owing to the long distance ( $\sim 20$  Å) as compared with the coupling between neighboring Chls, *i.e.*,  $P_{D1}^* - Chl_{D1}^*$ ,  $P_{D1}^* - Chl_{D2}^*$ ,  $P_{D2}^* - Chl_{D1}^*$  and  $P_{D2}^* - Chl_{D2}^*$  pairs (Fig. 4g and Table 1). Consequently, charge separation considering only the direct  $Chl_{D2}^* \rightarrow Chl_{D1}^* \rightarrow Chl_{D1}^+Pheo_{D1}^{*-}$  pathway is slower than charge separation considering only the indirect  $Chl_{D2}^* \rightarrow (P_{D1}^*, P_{D2}^*) \rightarrow Chl_{D1}^* \rightarrow Chl_{D1}^+Pheo_{D1}^{*-}$  pathway in the quantum dynamics calculations (Fig. 4f). The excitonic coupling between  $P_{D1}^*$  and  $Chl_{D2}^*$  ( $-14$  meV) is stronger than that between  $P_{D1}^*$  and  $Chl_{D1}^*$  (7 meV, Table 1). The former and latter are characterized as J- and H-aggregates (minus and plus signs), respectively, considering the directions of the transition dipole moments (Fig. 4g). The excitonic coupling is relatively insensitive to the orbital overlap as compared with the case of the charge transfer coupling. The direct and indirect  $Chl_{D2}^* \rightarrow Chl_{D1}^*$  exciton transfers exhibit the destructive interference of the quantum phase factor, which is dictated by the signs of excitonic couplings, *i.e.*, relative orientation of the transition dipole moments. Consequently, the exciton transfer rate considering all pathways is slower than the rate considering only the indirect pathway (Fig. 4f). Thus, in terms of the phase factor, the configuration of Chls in PSII is not necessarily optimal for accelerating the  $Chl_{D2}^* \rightarrow Chl_{D1}^*$  transfer, while the configuration is optimal for charge separation along the D1-branch.

Overall, it can be concluded that the  $Chl_{D2}^* \rightarrow Chl_{D1}^*$  exciton transfer followed by charge separation to  $Chl_{D1}^+Pheo_{D1}^{*-}$  in the D1-branch is overwhelmingly faster than charge separation in the D2-branch (Fig. 4f). The irreversible  $Chl_{D2}^* \rightarrow Chl_{D1}^*$  exciton transfer allows PSII to utilize the excitation energy from both the CP43 and CP47 antenna complexes for charge separation in the active branch.

### 3.3. Role of $Mn_4CaO_5$ in the charge separation pathway in PSII

The localized electronic states on  $P_{D1}$  in PSII are advantageous to maintain a high oxidation potential for water splitting in contrast to the strongly coupled  $(P_L P_M)^*$  and  $(P_L P_M)^{+}$  in PbRC.<sup>34</sup> The hole on  $P_{D1}^{+}$  is largely stabilized by acidic residues near the  $Mn_4CaO_5$  cluster, namely D1-Asp61, D1-Glu189, and D1-Asp170.<sup>34,57</sup> This may explain why the  $Mn_4CaO_5$  cluster is located on the D1 side, because the electrostatic potential, which attracts a hole toward the D1 side, also enhances charge separation to  $P_{D1}^+Pheo_{D1}^{*-}$ .<sup>33,34,58</sup> Because the difference in the redox potential between  $P_{D1}$  and  $Chl_{D1}$  is small,<sup>34</sup>  $P_{D1}^+Chl_{D1}^{*-}$  is substantially higher in energy than  $Chl_{D1}^+Pheo_{D1}^{*-}$  (Fig. 3c). Thus, the exciton funneling to  $Chl_{D1}^*$  rather than  $P_{D1}^*$  is a reasonable design principle for efficient charge separation to use excitons from the antenna complexes in PSII.

## 4. Conclusion

Quantum dynamics calculations indicated that two-step  $(P_L P_M)^* \rightarrow (P_L P_M)^{+}B_L^{*-}$  and  $B_L^{*-} \rightarrow H_L^{*-}$  electron transfers occur on a time scale of  $\sim 3.2$  and  $\sim 1.8$  ps, respectively (Fig. 2c). The population of the intermediate  $(P_L P_M)^{+}B_L^{*-}$  state is kept





small, owing to a strong  $B_L^{\cdot-} \rightarrow H_L^{\cdot-}$  coupling ( $\sim 16$  meV, Fig. 2e). The rapid electron transfer to  $H_L$  is advantageous for preventing charge recombination, because the orbital overlap between  $(P_L P_M)^{++}$  and  $H_L^{\cdot-}$  is negligibly small owing to a long molecular distance. The electrostatic interaction with the hydroxyl group of Tyr-M210 near  $B_L$  stabilizes the intermediate  $(P_L P_M)^{++} B_L^{\cdot-}$  state and accelerates charge separation along the L-branch, highlighting the essential role of Tyr-M210 in efficient unidirectional charge separation.

In PSII, both  $Chl_{D1}$  and  $Chl_{D2}$  can accept an exciton from CP43 and CP47, respectively. The  $Chl_{D1}^* \rightarrow Chl_{D1}^{*+} Pheo_{D1}^{\cdot-}$  electron transfer occurs on an ultrafast time scale ( $\sim 0.15$  ps), followed by the  $Chl_{D1}^{*+} \rightarrow P_{D1}^{*+}$  hole transfer on a time scale of  $\sim 3.7$  ps (Fig. 4a), as suggested by time-resolved spectroscopic measurements.<sup>14</sup> Charge separation in the D2-branch is unlikely to occur despite the relatively stable  $P_{D1}^{*+} Chl_{D2}^{\cdot-}$  state, because the in-plane  $P_{D1}$  vinyl group interferes with the  $\pi$ - $\pi$  interaction between  $P_{D1}$  and  $Chl_{D2}$ , thereby weakening the electronic coupling. The exciton on  $Chl_{D2}^*$  can be transferred to  $Chl_{D1}$  via the direct and indirect pathways. Subsequently, the ultrafast  $Chl_{D1}^* \rightarrow Chl_{D1}^{*+} Pheo_{D1}^{\cdot-}$  charge separation prevents exciton back transfer to  $Chl_{D2}$ , thereby enhancing the robustness of unidirectional charge separation in the D1-branch. Thus, PSII efficiently utilizes excitons not only from CP43 (D1 side) but also from CP47 (D2 side) for charge separation in the D1-branch, which leads to electron transfer to  $Q_B$  via  $Q_A$  and hole transfer to the  $Mn_4CaO_5$  cluster on the D1 side.

## Author contributions

H. T. designed the research. H. T., K. S., and H. I. performed the research. H. T. wrote the main part of the manuscript. All the authors were involved in the discussion of the results and contributed to the final version of the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This research was supported by JST CREST (JPMJCR1656 to H. I.), JSPS KAKENHI (JP18H01937 to H. T. and H. I., JP18H05155, JP20H03217, and JP20H05090 to H. I., JP18H01186 to K. S., and JP16H06560 to K. S.), and the Interdisciplinary Computational Science Program in CCS, University of Tsukuba. The authors acknowledge valuable discussions with prof. Akihiko Ishizaki.

## References

- 1 E. Wientjes, H. van Amerongen and R. Croce, *J. Phys. Chem. B*, 2013, **117**, 11200–11208.
- 2 G. D. Scholes, G. R. Fleming, A. Olaya-Castro and R. van Grondelle, *Nat. Chem.*, 2011, **3**, 763–774.
- 3 C. Kreisbeck and A. Aspuru-Guzik, *Chem. Sci.*, 2016, **7**, 4174–4183.
- 4 J. Pan, A. Gelzinis, V. Chorošajev, M. Vengris, S. S. Senlik, J.-R. Shen, L. Valkunas, D. Abramavicius and J. P. Ogilvie, *Phys. Chem. Chem. Phys.*, 2017, **19**, 15356–15367.
- 5 R. van Grondelle and V. I. Novoderezhkin, *Phys. Chem. Chem. Phys.*, 2006, **8**, 793–807.
- 6 F. Ma, L.-J. Yu, Z.-Y. Wang-Otomo and R. van Grondelle, *Biochim. Biophys. Acta Bioenerg.*, 2016, **1857**, 408–414.
- 7 L.-M. Tan, J. Yu, T. Kawakami, M. Kobayashi, P. Wang, Z.-Y. Wang-Otomo and J.-P. Zhang, *J. Phys. Chem. Lett.*, 2018, **9**, 3278–3284.
- 8 M. Germano, A. Y. Shkuropatov, H. Permentier, R. de Wijn, A. J. Hoff, V. A. Shuvalov and H. J. van Gorkom, *Biochemistry*, 2001, **40**, 11472–11482.
- 9 B. A. Diner and F. Rappaport, *Annu. Rev. Plant Biol.*, 2002, **53**, 551–580.
- 10 S. Vasil'ev, J.-R. Shen, N. Kamiya and D. Bruce, *FEBS Lett.*, 2004, **561**, 111–116.
- 11 T. Cardona, A. Sedoud, N. Cox and A. W. Rutherford, *Biochim. Biophys. Acta Bioenerg.*, 2012, **1817**, 26–43.
- 12 T. Cardona and A. W. Rutherford, *Trends Plant Sci.*, 2019, **24**, 1008–1021.
- 13 T. Cardona, J. W. Murray and A. W. Rutherford, *Mol. Biol. Evol.*, 2015, **32**, 1310–1328.
- 14 M. L. Groot, N. P. Pawlowicz, L. J. G. W. van Wilderen, J. Breton, I. H. M. van Stokkum and R. van Grondelle, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 13087.
- 15 M.-L. Groot, F. van Mourik, C. Eijkelhoff, I. H. M. van Stokkum, J. P. Dekker and R. van Grondelle, *Proc. Natl. Acad. Sci. U. S. A.*, 1997, **94**, 4389.
- 16 A. R. Holzwarth, M. G. Müller, M. Reus, M. Nowaczyk, J. Sander and M. Rögner, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 6895.
- 17 M. Hasegawa, H. Nagashima, R. Minobe, T. Tachikawa, H. Mino and Y. Kobori, *J. Phys. Chem. Lett.*, 2017, **8**, 1179–1184.
- 18 J. A. Myers, K. L. M. Lewis, F. D. Fuller, P. F. Tekavec, C. F. Yocum and J. P. Ogilvie, *J. Phys. Chem. Lett.*, 2010, **1**, 2774–2780.
- 19 F. D. Fuller, J. Pan, A. Gelzinis, V. Butkus, S. S. Senlik, D. E. Wilcox, C. F. Yocum, L. Valkunas, D. Abramavicius and J. P. Ogilvie, *Nat. Chem.*, 2014, **6**, 706–711.
- 20 E. Romero, R. Augulis, V. I. Novoderezhkin, M. Ferretti, J. Thieme, D. Zigmantas and R. van Grondelle, *Nat. Phys.*, 2014, **10**, 676–682.
- 21 E. Romero, V. I. Novoderezhkin and R. van Grondelle, *Nature*, 2017, **543**, 355–365.
- 22 G. Raszewski and T. Renger, *J. Am. Chem. Soc.*, 2008, **130**, 4431–4446.
- 23 E. Romero, I. H. M. van Stokkum, V. I. Novoderezhkin, J. P. Dekker and R. van Grondelle, *Biochemistry*, 2010, **49**, 4300–4307.
- 24 V. I. Novoderezhkin, E. Romero, J. P. Dekker and R. van Grondelle, *ChemPhysChem*, 2011, **12**, 681–688.
- 25 G. Raszewski, W. Saenger and T. Renger, *Biophys. J.*, 2005, **88**, 986–998.
- 26 V. I. Novoderezhkin, J. P. Dekker and R. van Grondelle, *Biophys. J.*, 2007, **93**, 1293–1311.



- 27 K. Acharya, B. Neupane, V. Zazubovich, R. T. Sayre, R. Picorel, M. Seibert and R. Jankowiak, *J. Phys. Chem. B*, 2012, **116**, 3890–3899.
- 28 G. Raszewski, B. A. Diner, E. Schlodder and T. Renger, *Biophys. J.*, 2008, **95**, 105–119.
- 29 F. Müh, M. Plöckinger and T. Renger, *J. Phys. Chem. Lett.*, 2017, **8**, 850–858.
- 30 A. Gelzinis, L. Valkunas, F. D. Fuller, J. P. Ogilvie, S. Mukamel and D. Abramavicius, *New J. Phys.*, 2013, **15**, 075013.
- 31 Y. Fujihashi, M. Higashi and A. Ishizaki, *J. Phys. Chem. Lett.*, 2018, **9**, 4921–4929.
- 32 M. A. Kavanagh, J. K. G. Karlsson, J. D. Colburn, L. M. C. Barter and I. R. Gould, *Proc. Natl. Acad. Sci. U. S. A.*, 2020, **117**, 19705.
- 33 K. Kawashima and H. Ishikita, *Chem. Sci.*, 2018, **9**, 4083–4092.
- 34 H. Tamura, K. Saito and H. Ishikita, *Proc. Natl. Acad. Sci. U. S. A.*, 2020, **117**, 16373.
- 35 M. Mandal, K. Kawashima, K. Saito and H. Ishikita, *J. Phys. Chem. Lett.*, 2020, **11**, 249–255.
- 36 H. Ishikita, J. Biesiadka, B. Loll, W. Saenger and E.-W. Knapp, *Angew. Chem., Int. Ed.*, 2006, **45**, 1964–1965.
- 37 L. Zhang, D.-A. Silva, H. Zhang, A. Yue, Y. Yan and X. Huang, *Nat. Commun.*, 2014, **5**, 4170.
- 38 A. Sirohiwal, F. Neese and D. A. Pantazis, *J. Am. Chem. Soc.*, 2020, **142**, 18174–18190.
- 39 M. Ziolek, N. Pawłowicz, R. Naskrecki and A. Dobek, *J. Phys. Chem. B*, 2005, **109**, 18171–18176.
- 40 H. Wang, S. Lin and N. W. Woodbury, *J. Phys. Chem. B*, 2008, **112**, 14296–14301.
- 41 C. Lauterwasser, U. Finklele, H. Scheer and W. Zinth, *Chem. Phys. Lett.*, 1991, **183**, 471–477.
- 42 W. Zinth and J. Wachtveitl, *ChemPhysChem*, 2005, **6**, 871–880.
- 43 A. Niedringhaus, V. R. Policht, R. Sechrist, A. Konar, P. D. Laible, D. F. Bocian, D. Holten, C. Kirmaier and J. P. Ogilvie, *Proc. Natl. Acad. Sci. U. S. A.*, 2018, **115**, 3563.
- 44 P. Huppman, T. Arlt, H. Penzkofer, S. Schmidt, M. Bibikova, B. Dohse, D. Oesterhelt, J. Wachtveitl and W. Zinth, *Biophys. J.*, 2002, **82**, 3186–3197.
- 45 F. Ma, E. Romero, M. R. Jones, V. I. Novoderezhkin and R. van Grondelle, *J. Phys. Chem. Lett.*, 2018, **9**, 1827–1832.
- 46 F. Ma, E. Romero, M. R. Jones, V. I. Novoderezhkin and R. van Grondelle, *Nat. Commun.*, 2019, **10**, 933.
- 47 U. Finklele, C. Lauterwasser, W. Zinth, K. A. Gray and D. Oesterhelt, *Biochemistry*, 1990, **29**, 8517–8521.
- 48 H. Wang, Y. Hao, Y. Jiang, S. Lin and N. W. Woodbury, *J. Phys. Chem. B*, 2012, **116**, 711–717.
- 49 C. Kirmaier, C. He and D. Holten, *Biochemistry*, 2001, **40**, 12132–12139.
- 50 P. D. Laible, D. K. Hanson, J. C. Buhmaster, G. A. Tira, K. M. Faries, D. Holten and C. Kirmaier, *Proc. Natl. Acad. Sci. U. S. A.*, 2020, **117**, 865.
- 51 M. A. Steffen, K. Lao and S. G. Boxer, *Science*, 1994, **264**, 810.
- 52 T. P. Treynor, C. Yoshina-Ishii and S. G. Boxer, *J. Phys. Chem. B*, 2004, **108**, 13523–13535.
- 53 M. Saggi, S. D. Fried and S. G. Boxer, *J. Phys. Chem. B*, 2019, **123**, 1527–1536.
- 54 S. E. J. Rigby, J. H. A. Nugent and P. J. O'Malley, *Biochemistry*, 1994, **33**, 10043–10050.
- 55 B. A. Diner, E. Schlodder, P. J. Nixon, W. J. Coleman, F. Rappaport, J. Lavergne, W. F. J. Vermaas and D. A. Chisholm, *Biochemistry*, 2001, **40**, 9265–9281.
- 56 T. Okubo, T. Tomo, M. Sugiura and T. Noguchi, *Biochemistry*, 2007, **46**, 4390–4397.
- 57 K. Saito, T. Ishida, M. Sugiura, K. Kawakami, Y. Umena, N. Kamiya, J. R. Shen and H. Ishikita, *J. Am. Chem. Soc.*, 2011, **133**, 14379–14388.
- 58 H. Ishikita, W. Saenger, J. Biesiadka, B. Loll and E.-W. Knapp, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 9855.
- 59 M. H. Beck, A. Jäckle, G. A. Worth and H. D. Meyer, *Phys. Rep.*, 2000, **324**, 1–105.
- 60 H. Tamura, E. R. Bittner and I. Burghardt, *J. Chem. Phys.*, 2007, **127**, 034706.
- 61 H. Tamura, J. G. S. Ramon, E. R. Bittner and I. Burghardt, *Phys. Rev. Lett.*, 2008, **100**, 107402.
- 62 H. Tamura, *J. Chem. Phys.*, 2009, **130**, 214705.
- 63 N. M. Thellamurege, D. Si, F. Cui, H. Zhu, R. Lai and H. Li, *J. Comput. Chem.*, 2013, **34**, 2816–2833.
- 64 N. M. Thellamurege and H. Li, *J. Chem. Phys.*, 2012, **137**, 246101.
- 65 M. W. Schmidt, K. K. Baldrige, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. Su, T. L. Windus, M. Dupuis and J. A. Montgomery Jr, *J. Comput. Chem.*, 1993, **14**, 1347–1363.
- 66 T. Yanai, D. P. Tew and N. C. Handy, *Chem. Phys. Lett.*, 2004, **393**, 51–57.
- 67 L. Cupellini, S. Caprasecca, C. A. Guido, F. Müh, T. Renger and B. Mennucci, *J. Phys. Chem. Lett.*, 2018, **9**, 6892–6899.
- 68 P. Cieplak, J. Caldwell and P. Kollman, *J. Comput. Chem.*, 2001, **22**, 1048–1057.
- 69 K. Saito and H. Ishikita, *Biophys. J.*, 2011, **101**, 2018–2025.
- 70 G. M. Ullmann and E.-W. Knapp, *Eur. Biophys. J.*, 1999, **28**, 533–551.
- 71 C. N. Schutz and A. Warshel, *Proteins*, 2001, **44**, 400–417.
- 72 A. Warshel, P. K. Sharma, M. Kato and W. W. Parson, *Biochim. Biophys. Acta*, 2006, **1764**, 1647–1676.
- 73 Y. Umena, K. Kawakami, J.-R. Shen and N. Kamiya, *Nature*, 2011, **473**, 55–60.
- 74 A. W. Roszak, V. Moulisová, A. D. P. Reksodipuro, A. T. Gardiner, R. Fujii, H. Hashimoto, N. W. Isaacs and R. J. Cogdell, *Biochem. J.*, 2012, **442**, 27–37.
- 75 S. Grimme, *J. Comput. Chem.*, 2006, **27**, 1787–1799.
- 76 H. Tamura, *J. Phys. Chem. A*, 2016, **120**, 9341–9347.
- 77 K. Mitsunashi, H. Tamura, K. Saito and H. Ishikita, *J. Phys. Chem. B*, 2021, **125**, 2879–2885.
- 78 J. Kern, R. Chatterjee, I. D. Young, F. D. Fuller, L. Lassalle, M. Ibrahim, S. Gul, T. Fransson, A. S. Brewster, R. Alonso-Mori, R. Hussein, M. Zhang, L. Douthit, C. de Lichtenberg, M. H. Cheah, D. Shevela, J. Wersig, I. Seuffert, D. Sokaras, E. Pastor, C. Weninger, T. Kroll, R. G. Sierra, P. Aller, A. Butryn, A. M. Orville, M. Liang, A. Batyuk, J. E. Koglin, S. Carbajo, S. Boutet, N. W. Moriarty, J. M. Holton, H. Dobbek, P. D. Adams, U. Bergmann, N. K. Sauter, A. Zouni, J. Messinger, J. Yano and V. K. Yachandra, *Nature*, 2018, **563**, 421–425.

