

Cite this: *Chem. Sci.*, 2015, 6, 3816

Assembly line termination in cylindrocyclophane biosynthesis: discovery of an editing type II thioesterase domain in a type I polyketide synthase†

H. Nakamura,^a J. X. Wang^b and E. P. Balskus^{*a}

The termination step is an important source of structural diversity in polyketide biosynthesis. Most type I polyketide synthase (PKS) assembly lines are terminated by a thioesterase (TE) domain located at the C-terminus of the final module, while other PKS assembly lines lack a terminal TE domain and are instead terminated by a separate enzyme *in trans*. In cylindrocyclophane biosynthesis, the type I modular PKS assembly line is terminated by a freestanding type III PKS (CylI). Unexpectedly, the final module of the type I PKS (CylH) also possesses a C-terminal TE domain. Unlike typical type I PKSs, the CylH TE domain does not influence assembly line termination by CylI *in vitro*. Instead, this domain phylogenetically resembles a type II TE and possesses activity consistent with an editing function. This finding may shed light on the evolution of unusual PKS termination logic. In addition, the presence of related type II TE domains in many cryptic type I PKS and nonribosomal peptide synthetase (NRPS) assembly lines has implications for pathway annotation, product prediction, and engineering.

Received 12th October 2014

Accepted 11th April 2015

DOI: 10.1039/c4sc03132f

www.rsc.org/chemicalscience

Introduction

Polyketides are a large group of secondary metabolites that display a wide range of important biological activities.¹ Many polyketides are synthesized by type I polyketide synthases (PKSs), which generate structural diversity and complexity by employing combinations of domains and modules organized into enzymatic assembly lines.² The core domains of a type I PKS module are the ketosynthase (KS), acyltransferase (AT), and acyl-carrier protein (ACP) domains. The AT domain selects an initial substrate or an extender unit, typically malonyl-coenzyme A (CoA) or methylmalonyl-CoA, and loads it onto the phosphopantetheinyl (ppant) arm of the ACP domain. The KS domain then catalyses a decarboxylative Claisen condensation between the ACP-tethered extender unit and a polyketide intermediate covalently bound to an active site cysteine. The initial β -keto thioester product can be further modified by tailoring domains present in the module (typically ketoreductase (KR), dehydratase (DH), and enoyl reductase (ER) domains). Type I PKSs can either be modular or iterative: each

module of a type I modular PKS catalyses one round of elongation and tailoring, while the single module of a type I iterative PKS catalyses multiple rounds of elongation and tailoring.

At the end of an assembly line, the nascent polyketide product must be released from the PKS by a terminating enzyme. The type of reaction used for termination has a large influence on the architecture of the final natural product and is therefore an important source of structural diversity in polyketide biosynthesis. There are three major types of termination logic used by type I PKS assembly lines: release by C-terminal thioesterase (TE) domains, release by alternate C-terminal domains, and release by freestanding enzymes *in trans*. The canonical and most well-studied strategy is C-terminal TE domain-mediated termination. The TE domains that catalyse assembly line terminations are referred to as type I TEs and are typically found at the C-terminus of the final module. TE domains release the nascent polyketide product from the PKS assembly lines by catalysing either hydrolysis or macrocyclization using a conserved Ser-His-Asp catalytic triad (Fig. 1a, top).³ The second termination logic involves product release by C-terminal domains other than TEs. For example, reductase (R) domains catalyse reductive release in multiple PKS assembly lines, including those in xenovulene⁴ and azanigerone⁵ biosyntheses. Furthermore, a special type of TE domain called TE/CLC (Claisen-like cyclase) catalyses Claisen-type condensations to form a new ring system in the biosynthesis of the fungal polyketides naphthopyrone⁶ and aflatoxin.⁷ Both of these approaches involve the use of C-terminal domains *in cis*.

^aDepartment of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138, USA. E-mail: balskus@chemistry.harvard.edu

^bSmall Molecule Mass Spectrometry Facility, FAS Division of Science, Cambridge, Massachusetts 02138, USA

† Electronic supplementary information (ESI) available: Fig. S1–S12; Tables S1–S8, full experimental details and procedures, ¹H and ¹³C NMR spectral data and HRMS data of compounds 10 and 11 and the internal standards. See DOI: 10.1039/c4sc03132f




Fig. 1 Cylirocyclophane biosynthesis involves unusual type I polyketide synthase (PKS) termination logic. (a) Biosynthetic logic for termination of type I PKS assembly lines. (b) Structures of cylirocyclophanes isolated from *Cylindrospermum licheniforme* ATCC 29412. (c) Overview of the hypothesized cylirocyclophane biosynthetic pathway and proposed termination chemistry. KS = ketosynthase, AT = acyltransferase, ACP = acyl carrier protein, TE = thioesterase. (d) Structure of the *N*-acetylcysteamine thioester (SNAC) mimic of β-keto ACP thioester 7.

The third termination logic involves PKSs that lack a C-terminal domain for product release, instead employing a separate, freestanding enzyme *in trans* (Fig. 1a, bottom). For example, a freestanding TE promotes hydrolysis in the biosynthesis of polyether ionophores,⁸ and a discrete acyltransferase catalyses macrolactamization in rifamycin biosynthesis.⁹ More recently, a freestanding ketosynthase was postulated to condense two β-ketothioester intermediates, each tethered to a separate type I modular PKS, to form a pyrone ring in myxopyronin biosynthesis.¹⁰ *In trans* termination chemistry is frequently found in type I iterative PKS systems. A discrete thioesterase releases a polyene intermediate from a type I iterative PKS in enediyne biosynthesis,¹¹ and an acyltransferase releases an intermediate from an iterative PKS in lovastatin synthesis.¹² Other types of enzymes that catalyse product release from PKS assembly lines *in trans* include a pyridoxal 5'-phosphate (PLP)-dependent enzyme in fumonisins biosynthesis,¹³ a lactamase in atrochrysone carboxylic acid biosynthesis¹⁴ and a Baeyer-Villiger monooxygenase in aurafuron biosynthesis.¹⁵ The variety of enzymes that participate in assembly line termination further illustrates the importance of this step to polyketide structural diversity.

The cylirocyclophanes (1–6) are a family of cyanobacterial polyketides with a rare [7.7]paracyclophane carbon scaffold (Fig. 1b). The biosynthesis of these natural products appears to involve an unusual assembly line termination mechanism.¹⁶ In

our previous investigation of cylirocyclophane biosynthesis, we hypothesized that putative biosynthetic precursor 8 would be generated by *in trans* termination of the type I modular PKS assembly line (CylD–CylH) *via* freestanding type III PKS CylII (Fig. 1c). In support of this hypothesis, purified CylII converted 9, an *N*-acetylcysteamine thioester (SNAC) mimic of the CylH-bound β-ketothioester 7 (Fig. 1d), to resorcinol 8. This result implied that CylII accepts acyl-ACP 7 as a substrate *in vivo*. Type III PKSs typically use acyl-CoA electrophiles and form aromatic products by catalysing iterative decarboxylative Claisen condensations directly with malonyl-CoA extender units.¹⁷ Recently several type III PKSs have been shown to accept acyl-ACPs as electrophiles, including ArsB and ArsC from *Azobacter vinelandii*¹⁸ and BpsA from *Bacillus subtilis*,¹⁹ which use ACP-thioesters from fatty acid biosynthesis. Additionally, FtpA from *Myxococcus xanthus* accepts an acyl-ACP substrate produced by a fatty acyl-AMP ligase FtpD,²⁰ a type III PKS domain fused to a type I fatty acid synthase (FAS) uses an acyl-ACP electrophile in the synthesis differentiation factor DIF-1 by *Dictyostelium discoideum*,²¹ and gramicidin synthase from *Streptomyces coelicolor* accepts both acyl-CoA and acyl-ACP electrophiles *in vitro*.²² While several type III PKSs are known to utilize acyl-ACP substrates, to our knowledge CylII is the first type III PKS predicted to use an acyl-ACP substrate generated by a type I modular PKS.¹⁶

Although there are many examples of PKS termination catalysed by discrete enzymes *in trans*, all previously characterized pathways with *in trans* termination logic, to the best of our knowledge, have assembly lines that lack C-terminal TE domains. Cylirocyclophane biosynthesis, however, is unique in possessing both a freestanding terminating enzyme and a C-terminal TE domain on the final module of its type I modular PKS assembly line (CylH). This is surprising, as termination *via* a type III PKS should not require a type I TE domain. To understand the unusual termination chemistry involved in this pathway and the role of the TE domain, we have characterized the activity of CylH and CylII *in vitro*. We confirm that type III PKS CylII accepts a CylH-tethered substrate. We employ mutagenesis to show that the CylH TE domain activity does not influence assembly line termination by CylII or catalyse competing hydrolysis of acyl-ACP 7. Rather, biochemical experiments and bioinformatic analysis of the CylH TE domain reveal similarity to type II TEs, freestanding enzymes that typically play editing roles in polyketide biosynthesis.^{2,23} Further searches of bacterial genomes reveal that similar editing TE domains are widely incorporated into cryptic PKS and non-ribosomal peptide synthetase (NRPS) assembly lines.

These discoveries have broad implications for how we think about the roles of TE domains in biosynthesis. Previous work has revealed the existence of type II-like TE domains in type I PKS assembly lines, including those that make ajudazole and jerangolid,^{24–26} but their biochemical functions have been largely unexplored. The discovery of an *in cis* TE domain with a dedicated editing role raises the possibility that similar domains are found in other assembly lines. This prospect overturns canonical enzymatic assembly line biosynthetic logic, which assumes that *in cis* TE domains mediate only termination



chemistry, and should influence how we annotate newly discovered assembly lines. Our findings also raise the possibility that *in cis* editing is important for optimal function of certain assembly lines, an observation that could be exploited to better engineer these systems. Overall, the discovery of an editing C-terminal TE domain in a type I PKS assembly line has implications both for the evolution of unusual termination logic and our understanding of the functions of these important domains.

Results and discussion

Assessment of the CylH TE domain's role in assembly line termination

We previously showed that type III PKS CylI converted acyl-SNAC substrate **9** to resorcinol **8** *via* two extension cycles using malonyl-CoA as an extender unit, followed by cyclization, decarboxylation and aromatization.¹⁶ The direct use of **9** by CylI suggested that the C-terminal TE domain of the final PKS module of CylH was not necessary for assembly line termination. In this work, we aimed to investigate whether the CylH TE domain is catalytically active and to elucidate the role of the TE domain in the context of the type I PKS assembly line. We cloned, heterologously expressed and purified the final PKS module of CylH (CylH_{PKS} = KS-AT-ACP-TE) and the excised CylH TE domain (CylH_{TE}), as well as their corresponding TE active site serine to alanine mutants (CylH_{PKS}-S1201A and CylH_{TE}-S95A) using *E. coli* as an expression host (Fig. S1–S3, Tables S1 and S2†). Production of CylH_{PKS} and CylH_{PKS}-S1201A required co-expression of chaperones GroES and GroEL. Successful posttranslational modification of the ACP domain of CylH_{PKS} and the CylH_{PKS}-S1201A mutant by Sfp, a promiscuous phosphopantetheinyl transferase,²⁷ was verified using the BODIPY-CoA loading assay (Fig. S4 and Table S3†).

We performed *in vitro* assays to examine the elongation activity of CylH_{PKS} (Fig. 2a). *Holo* CylH_{PKS} was prepared *in situ* by incubation with CoA and Sfp, and the resulting *holo* enzyme was incubated with acyl-SNAC **10** and malonyl-CoA. The assay mixtures were quenched by the addition of 1 M KOH, followed by heating at 65 °C to ensure complete hydrolysis of acyl-ACP **7** to β -keto acid **11**. We did not observe significant amounts of product decarboxylation under these conditions (Fig. S5†). CylH_{PKS} catalysed elongation of substrate **10**, leading to the

formation of **11** in full assays but not negative controls lacking either enzyme or substrate (Fig. S6 and Table S4†). Addition of type III PKS CylI to the assay mixture resulted in the production of resorcinol **8** (Fig. 2b). We observed a small amount of resorcinol formation in the absence of CylH_{PKS} due to promiscuous activity of CylI toward substrate **10**, which undergoes three extension cycles prior to cyclization, decarboxylation and aromatization to give resorcinol **8**. The substantial increase in the rate of resorcinol formation in the presence of CylH_{PKS}, however, shows that CylI uses the β -keto ACP thioester **7** as a preferred substrate. Overall, these assays confirmed that CylH_{PKS} catalysed elongation and that CylI could utilize a CylH-tethered substrate.

To determine whether TE domain activity impacted the rate of resorcinol formation, we performed the elongation assay with CylH_{PKS}-S1201A. The TE domain mutation did not appear to affect the qualitative rate of resorcinol formation, suggesting that the TE domain is not directly involved in either aromatic ring formation or a competing hydrolysis (Fig. 3a and S7†). If the TE domain is catalytically active and has a competing hydrolytic activity for acyl-ACP **7**, we should observe accumulation of the hydrolysis product **11**. We therefore quantified the amount of resorcinol **8** and acyl-ACP hydrolysis product **11** in assays containing CylI and either CylH_{PKS} or CylH_{PKS}-S1201A by LC-MS (Fig. 3b, S8 and S9†). In the full assays containing CylI, very small quantities of hydrolysis product **11** were detected (~ 0.3 μ M for both CylH_{PKS} and CylH_{PKS}-S1201A) compared to




Fig. 3 The activity of the CylH TE domain does not impact assembly line termination by CylI. (a) HPLC assay of CylI-catalysed resorcinol formation with CylH_{PKS}-S1201A mutant (monitored at 210 nm). (b) LC-MS quantitation of resorcinol **8** and hydrolysis product **11** in assays containing CylI and either wild type CylH_{PKS} or the CylH_{PKS}-S1201A mutant. Mean values \pm standard deviation of assays done in triplicate are shown, and *p*-values < 0.05 are marked with *.




Fig. 2 The final module of the cylindrocyclophane PKS assembly line generates a substrate for type III PKS CylI. (a) Assays used to study the elongation activity of CylH_{PKS} and the resorcinol formation by CylI using acyl-ACP substrate **7**. (b) HPLC assays of CylI-catalysed resorcinol formation show that CylH increases reaction efficiency (monitored at 210 nm).



resorcinol **8**, indicating that the CylH TE domain does not have a competing hydrolytic activity in assembly line termination. The trace amount of hydrolysis product **11** observed in these assays can also result from background hydrolysis of acyl-ACP **7**, if the transfer of nascent polyketide to CylI is inefficient. This is likely the reason for the higher concentrations of **11** present in assays lacking CylI.

We also tested whether addition of CylH_{TE} impacted the efficiency of CylI-catalysed resorcinol formation using SNAC substrate **9**, and observed no qualitative difference in reaction efficiency in the presence of CylH_{TE} or the CylH_{TE}-S95A mutant (Fig. S10†). These results support the hypothesis that the CylH TE domain does not influence termination in the PKS assembly line.

Bioinformatic and phylogenetic analyses of the CylH TE domain

To determine whether the apparent lack of CylH TE domain activity arose from mutations to key active site residues, specifically the conserved Ser–His–Asp catalytic triad, we analysed this domain using bioinformatics. When we aligned the amino acid sequence of CylH_{TE} with those of characterized type I TEs, the catalytic Asp residue appeared to be absent (Fig. S1†). However, when we performed the alignment with sequences of characterized type II TEs, the catalytic triad of CylH_{TE} was conserved. This discrepancy between the two alignments suggested that despite its position at the C-terminus of a type I modular PKS, the CylH TE domain might be more closely related to type II TEs. Additional support for this hypothesis included an HHPred structure homology search, which identified type II TEs RifR²⁸ and RedJ²⁹ as the top two hits. Using the crystal structure of RifR as a template, we constructed a CylH_{TE} homology model, which possesses a well-aligned catalytic triad (Fig. 4a).

As in the case of CylH_{TE}, C-terminal TE domains in the ajudazole (AjuTE) and jerangolid (JerTE) biosyntheses were previously reported to resemble type II TEs.²⁴ In fact, CylH_{TE} appears to have high sequence homology to AjuTE (53% similarity, 33% identity) and JerTE (55% similarity, 39% identity). Interestingly, the hydrolase signature sequence (GX₂SG) of CylH_{TE} matches that of type II TEs perfectly (GHSMG), while the His residue in the signature sequence of AjuTE and JerTE are replaced by Ser and Cys respectively.²⁴ This His residue has been shown to stabilize the active site by forming a hydrogen bond with the backbone carbonyl of the catalytic His residue in the crystal structure of RifR.²⁸ The hydrolase signature sequence of CylH_{TE} provides another evidence for its closer homology to type II TEs.

Finally, we constructed a phylogenetic tree following the method used by Buntin and co-workers²⁴ and found that CylH_{TE} clusters with freestanding type II TEs, not C-terminal type I TEs (Fig. 4c). Overall, these analyses show that the CylH TE domain has the catalytic residues needed for activity, but is more closely related to type II TEs rather than the type I TEs more typically associated with modular PKSs.

In all previous studies that have characterized *in cis* type II-like TE domains in enzymatic assembly lines, the TE domain is



Fig. 4 Bioinformatic analysis of the CylH TE domain reveals similarity to type II thioesterases. (a) Homology model of the CylH TE domain (green) overlaid with type II TE RifR (pink) shows conservation of the catalytic triad (Asp–His–Ser). (b) Multiple sequence alignment of the hydrolase signature sequence of characterized TEs shows that the CylH TE domain has the GHSMG motif common in type II TEs. (c) Phylogenetic tree of characterized TE domains using Bayesian estimation reveals clustering of the CylH TE domain with type II TEs (all bootstrap values were greater than 50. Ones with values above 75 are marked with *). TEI = type I TE, TEII = type II TE, P = PKS, N = NRPS, PN = hybrid PKS–NRPS, uk = unknown. TEs that have not been previously characterized are marked with *. The accession numbers of the proteins used in the analysis are listed in Table S7†).

thought to play a role in product release. The C-terminal TE domain in ajudazol has been found to catalyse isochromanone ring formation,²⁴ and the homologous TE domains in jerangolid³⁰ and legiolulin²⁵ biosyntheses have been implicated in promoting lactonization and isochromanone ring formation, respectively. The excised AjuTE and JerTE domains also possess hydrolytic activity toward short chain acyl *p*-nitrophenyl esters, substrates commonly used to assess the editing function of type II TEs.²⁴ More recently, the NRPS assembly line that constructs 4-methyloxazoline-containing natural products was found to contain an unusual module (FmoA5) with two tandem internal



This journal is © The Royal Society of Chemistry 2015

Identification of *in cis* type II TE domains in other PKS and NRPS assembly lines

We performed additional bioinformatic analyses to determine the prevalence of *in cis* type II TE domains in type I PKS or NRPS assembly lines. A BLASTP search of sequenced bacterial genomes using the CylH TE domain as a query identified 84 unique type I PKS and NRPS assembly lines containing *in cis* type II TE domains, including AjuTE and JerTE (Table S8†). Many of the hits have unusual domains or domain organizations, such as a hybrid PKS–NRPS from *Burkholderia mallei* that contains TE domains at both its N- and C-termini. The ubiquity of *in cis* type II TE domains, as well as the diverse activities of those that have been biochemically characterized, suggests that these domains may possess new, unexplored functions and could be important new tools for combinatorial biosynthesis. These TE domains may also display editing activity at particular steps in biosynthesis that could be important for maintaining the efficiency of these assembly line enzymes. While the function of the TE/CLC domain in maintaining assembly line fidelity of fungal type I iterative PKS from aflatoxin biosynthesis is well-characterised,³¹ the editing roles of TE domains in type I modular PKS assembly lines have not been widely explored. A better understanding of the functions of these domains could be applied to improve the efficiency of engineered assembly lines through the strategic introduction of editing activity. Finally, the wide distribution of C-terminal type II TE domains has implications for future characterization and annotation of cryptic multimodular assembly line pathways in bacterial genomes, as one can no longer assume that these domains follow canonical termination logic.

Conclusions

In summary, we have discovered and characterized the role of a C-terminal *in cis* type II TE domain in cylindrocyclophane biosynthesis. We have shown that the activity of this domain does not influence assembly line termination by type III PKS CylI. However, the CylH TE domain is catalytically active and can hydrolyze acetyl-SNAC and acetyl-ACP, as well as other short-chain acyl-SNACs, which is consistent with a role in preserving assembly line function *via* editing. The activity of freestanding type II TEs are often important for maintaining the efficiency of enzymatic assembly lines. Likewise, the CylH TE domain may also be necessary for maintaining the productivity of the type I modular PKS in cylindrocyclophane biosynthesis. Bioinformatic analyses suggest that such atypical *in cis* type II TE domains are widespread in other multimodular assembly lines. Further efforts to understand the functions of these domains will improve both our understanding of modular enzymatic assembly line function and our ability to engineer and utilize the chemical capabilities of these systems.

Acknowledgements

We acknowledge Li Zha for help with the autoradiography experiments, Zebulon Levine for the synthesis of acetyl-SNAC,

and Jonathan Marks for help with the bioinformatics. We received financial support for Harvard University and the Searle Scholars Program. H.N. acknowledges fellowship support from the NSF GRFP (DGE1144152).

Notes and references

- (a) J. Staunton and K. J. Weissman, *Nat. Prod. Rep.*, 2001, **18**, 380; (b) B. J. Rawlings, *Nat. Prod. Rep.*, 2001, **18**, 190; (c) B. J. Rawlings, *Nat. Prod. Rep.*, 2001, **18**, 231.
- M. A. Fischbach and C. T. Walsh, *Chem. Rev.*, 2006, **106**, 3468.
- L. Du and L. Lou, *Nat. Prod. Rep.*, 2010, **27**, 255.
- A. M. Bailey, R. J. Cox, K. Harley, C. M. Lazarus, T. J. Simpson and E. Skellam, *Chem. Commun.*, 2007, 4053.
- A. O. Zabala, W. Xu, Y.-H. Chooi and Y. Tang, *Chem. Biol.*, 2012, **19**, 1049.
- I. Fujii, A. Watanabe, U. Sankawa and Y. Ebizuka, *Chem. Biol.*, 2001, **8**, 189.
- T. P. Korman, J. M. Crawford, J. W. Labonte, A. G. Newman, J. Wong, C. A. Townsend and S.-C. Tsai, *Proc. Natl. Acad. Sci. U. S. A.*, 2010, **107**, 6246.
- (a) T. Liu, D. You, C. Valenzano, Y. Sun, J. Li, Q. Yu, X. Zhou, D. E. Cane and Z. Deng, *Chem. Biol.*, 2006, **13**, 945; (b) B. M. Harvey, H. Hong, M. A. Jones, Z. A. Hughes-Thomas, R. M. Goss, M. L. Heathcote, V. M. Bolanos-Garcia, W. Kroutil, J. Staunton, P. F. Leadlay and J. B. Spencer, *ChemBioChem*, 2006, **7**, 1435; (c) T. Liu, X. Lin, X. Zhou, Z. Deng and D. E. Cane, *Chem. Biol.*, 2008, **15**, 449.
- T.-W. Yu, Y. Shen, Y. Doi-Katayama, L. Tang, C. Park, B. S. Moore, C. R. Hutchinson and H. G. Floss, *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 9051.
- H. Sucipto, S. C. Wenzel and R. Müller, *ChemBioChem*, 2013, **14**, 1581.
- (a) R. Kong, L. P. Goh, C. W. Liew, Q. S. Ho, E. Murugan, B. Li, K. Tang and Z.-X. Liang, *J. Am. Chem. Soc.*, 2008, **130**, 8142; (b) J. Zhang, S. G. Van Lanen, J. Ju, W. Liu, P. C. Dorrestein, W. Li, N. L. Kelleher and B. Shen, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 1460; (c) M. Kotaka, R. Kong, I. Qureshi, Q. S. Ho, H. Sun, C. W. Liew, L. P. Goh, P. Cheung, Y. Mu, J. Lescar and Z.-X. Liang, *J. Biol. Chem.*, 2009, **284**, 15739; (d) K. Belecki, J. M. Crawford and C. A. Townsend, *J. Am. Chem. Soc.*, 2009, **131**, 12564.
- X. Xie, M. J. Meehan, W. Xu, P. C. Dorrestein and Y. Tang, *J. Am. Chem. Soc.*, 2009, **131**, 8388.
- R. Gerber, L. Lou and L. Du, *J. Am. Chem. Soc.*, 2009, **131**, 3148.
- T. Awakawa, K. Yokota, N. Funa, F. Doi, N. Mori, H. Watanabe and S. Horinouchi, *Chem. Biol.*, 2009, **16**, 613.
- B. Frank, S. C. Wenzel, H. B. Bode, M. Scharfe, H. Blöcker and R. Müller, *J. Mol. Biol.*, 2007, **374**, 24.
- H. Nakamura, H. A. Hamer, G. Sirasani and E. P. Balskus, *J. Am. Chem. Soc.*, 2012, **134**, 18518.
- M. B. Austin and J. P. Noel, *Nat. Prod. Rep.*, 2003, **20**, 79.
- A. Miyanaga, N. Funa, T. Awakawa and S. Horinouchi, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 871.



- 19 C. Nakano, H. Ozawa, G. Akanuma, N. Funa and S. Horinouchi, *J. Bacteriol.*, 2009, **191**, 4916.
- 20 T. Hayashi, Y. Kitamura, N. Funa, Y. Ohnishi and S. Horinouchi, *ChemBioChem*, 2011, **12**, 2166.
- 21 M. B. Austin, T. Saito, M. E. Bowman, S. Haydock, A. Kato, B. S. Moore, R. R. Kay and J. P. Noel, *Nat. Chem. Biol.*, 2006, **2**, 494.
- 22 (a) L. Song, F. Barona-Gomez, C. Corre, L. Xiang, D. W. Udvary, M. B. Austin, J. P. Noel, B. S. Moore and G. L. Challis, *J. Am. Chem. Soc.*, 2006, **128**, 14754; (b) S. Gruschow, T. J. Buchholz, W. Seufert, J. S. Dordick and D. H. Sherman, *ChemBioChem*, 2007, **8**, 863; (c) J. A. Chemler, T. J. Buchholz, T. W. Geders, D. L. Akey, C. M. Rath, G. E. Chlipala, J. L. Smith and D. H. Sherman, *J. Am. Chem. Soc.*, 2012, **134**, 7359.
- 23 (a) A. R. Butler, N. Bate and E. Cundliffe, *Chem. Biol.*, 1999, **6**, 287; (b) D. Schwarzer, H. D. Mootz, U. Linne and M. A. Marahiel, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 14083; (c) M. L. Heathcote, J. Staunton and P. F. Leadlay, *Chem. Biol.*, 2001, **8**, 207; (d) B. S. Kim, T. A. Cropp, B. J. Beck, D. H. Sherman and K. A. Reynolds, *J. Biol. Chem.*, 2002, **277**, 48028; (e) E. Yeh, R. M. Kohli, S. D. Bruner and C. T. Walsh, *ChemBioChem*, 2004, **5**, 1290; (f) M. Kotowska and K. Pawlik, *Appl. Microbiol. Biotechnol.*, 2014, **98**, 7735.
- 24 K. Buntin, K. J. Weissman and R. Müller, *ChemBioChem*, 2010, **11**, 1137.
- 25 T. Ahrendt, M. Miltenberger, I. Haneburger, F. Kirchner, M. Kronenwerth, A. O. Brachmann, H. Hilbi and H. B. Bode, *ChemBioChem*, 2013, **14**, 1415.
- 26 A. Muliandi, Y. Katsuyama, K. Sone, M. Izumikawa, T. Moriya, J. Hashimoto, I. Kozono, M. Takagi, K. Shin-ya and Y. Ohnishi, *Chem. Biol.*, 2014, **21**, 923.
- 27 L. E. N. Quadri, P. H. Weinreb, M. Lei, M. M. Nakano, P. Zuber and C. T. Walsh, *Biochemistry*, 1998, **37**, 1585.
- 28 H. B. Claxton, D. L. Akey, M. K. Silver, S. J. Admiraal and J. L. Smith, *J. Biol. Chem.*, 2009, **284**, 5021.
- 29 J. R. Whicher, G. Florova, P. K. Sydor, R. Singh, M. Alhamadsheh, G. L. Challis, K. A. Reynolds and J. L. Smith, *J. Biol. Chem.*, 2011, **286**, 22558.
- 30 B. Julien, Z.-Q. Tian, R. Reid and C. D. Reeves, *Chem. Biol.*, 2006, **13**, 1277.
- 31 A. L. Vagstad, S. B. Bumpus, K. Belecki, N. L. Kelleher and C. A. Townsend, *J. Am. Chem. Soc.*, 2012, **134**, 6865.

