

## REVIEW

[View Article Online](#)  
[View Journal](#) | [View Issue](#)Cite this: *Chem. Sci.*, 2024, 15, 13Received 26th September 2023  
Accepted 22nd November 2023

DOI: 10.1039/d3sc05056d

[rsc.li/chemical-science](https://rsc.li/chemical-science)

## f-Element heavy pnictogen chemistry

Jingzhen Du,<sup>a</sup> Philip J. Cobb,<sup>b</sup> Junru Ding,<sup>a</sup> David P. Mills<sup>\*b</sup>  
and Stephen T. Liddle<sup>\*b</sup>

The coordination and organometallic chemistry of the f-elements, that is group 3, lanthanide, and actinide ions, supported by nitrogen ligands, e.g. amides, imides, and nitrides, has become well developed over many decades. In contrast, the corresponding f-element chemistry with the heavier pnictogen analogues phosphorus, arsenic, antimony, and bismuth has remained significantly underdeveloped, due largely to a lack of suitable synthetic methodologies and also the inherent hard(f-element)–soft(heavier pnictogen) acid–base mismatch, but has begun to flourish in recent years. Here, we review complexes containing chemical bonds between the f-elements and heavy pnictogens from phosphorus to bismuth that spans five decades of endeavour. We focus on complexes whose identity has been unambiguously established by structural authentication by single-crystal X-ray diffraction with respect to their synthesis, characterisation, bonding, and reactivity, in order to provide a representative overview of this burgeoning area. By highlighting that much has been achieved but that there is still much to do this review aims to inspire, focus and guide future efforts in this area.

## 1. Introduction

Due to their widespread and important applications in magnetic materials,<sup>1,2</sup> electronic devices,<sup>3,4</sup> bioimaging,<sup>5,6</sup> synthesis,<sup>7,8</sup> catalysis,<sup>9,10</sup> materials science<sup>11,12</sup> and nuclear

technologies,<sup>13,14</sup> there has been burgeoning interest in the fundamental chemistry of the f-elements over the past few decades. Since f-element metal ions, that is group 3, lanthanide, and actinide ions, are hard Lewis acids with typically large radii and high coordination numbers, they preferentially bind with hard bases (by the hard–soft-acid-base definition); the chemical bonds of these ions are understood to be predominantly ionic, thus their solution chemistry is dominated by N-, O-, and halide-donor ligands.<sup>15</sup> With ever-developing synthetic methods

<sup>a</sup>College of Chemistry, Zhengzhou University, Zhengzhou, 450001, China<sup>b</sup>Department of Chemistry and Centre for Radiochemistry Research, The University of Manchester, Oxford Road, Manchester, M13 9PL, UK. E-mail: david.mills@manchester.ac.uk; steve.liddle@manchester.ac.uk

Jingzhen Du

Jingzhen Du received his BSc in 2013 from Shangqiu Normal University. Before moving to the University of Manchester as a President's Doctoral Scholarship PhD student in 2016 to investigate actinide nitride chemistry with Prof. Steve Liddle, he had a few years research training in Prof. Liang Deng's Group working on organometallic d-block compounds at Shanghai Institute of Organic Chemistry. In 2020, he received

his PhD degree from the University of Manchester and continued as a Postdoctoral researcher in the Liddle group. In 2023, he started his independent career at Zhengzhou University. His research interests are organometallic f-block complexes with unusual structures, properties, and reactivities.



Philip J. Cobb

Philip Cobb gained his PhD in 2018 from the University of Manchester after conducting a research project on the chemistry of uranyl chemistry under the supervision of Prof. Steve Liddle. He then spent 2 years researching f-element complexes at Manchester before moving into the nuclear industry.




Fig. 1 Bar graph summarising the number of molecular structures deposited into the CCDC by 10-08-2023 for any type of f-element pnictogen bond. Total numbers: M–N = 29 713, M–P = 817, M–As = 67, M–Sb = 21, M–Bi = 28.

and characterisation techniques, molecular non-aqueous f-element chemistry has developed in recent years, and under non-aqueous conditions f-element complexes with novel linkages involving softer donor atoms can be accessed and investigated.<sup>16</sup> More peripherally, but still relevant, it is known that softer donor ligands can effect better selectivity in extraction processes, so the study of such linkages can provide bonding

benchmarks of wider relevance.<sup>17</sup> For group 15, the pnictogens, f-element chemistry is well developed for nitrogen ligands, *e.g.* amides, imides, and nitrides, but this is not the case for the heavier congeners phosphorus, arsenic, antimony, and bismuth.<sup>18–20</sup> To illustrate the point, a search of the Cambridge Structural Database (CSD, 10th August 2023†)<sup>21</sup> for any type of crystallographically characterised chemical bond between the f-elements and any pnictogen reveals a stark picture, Fig. 1. There are almost thirty thousand complexes with f-element nitrogen bonds, over 30 times the total number of 933 for f-element heavier pnictogen complexes. Furthermore, below phosphorus there are only 67, 21, and 28 examples of f-element bonds to arsenic, antimony, and bismuth, respectively, with most of those examples reported in the last decade.

The above data parallel transition metal chemistry in many regards, though are a more extreme picture reflecting that stabilisation of the heavier and softer pnictogen ligands multiply bonded at large and hard Lewis acidic f-element metal ions is certainly more challenging. However, f-element pnictinidene and pnictido complexes remain of interest since in addition to being heavy amide, imide, and nitride analogues, they are isoelectronic congeners of transition metal alkyls, carbenes, and carbynes, respectively, that have developed into excellent catalysts for various organic transformations<sup>22</sup> or as precursors to inorganic materials.<sup>23</sup> Furthermore, whilst f-element phosphorus and arsenic multiple bonding isprecedented, f-element antimony or bismuth multiple bonds are conspicuous by their absence.<sup>20</sup> Indeed, as group 15 is descended the pnictide ions become increasingly electropositive and metal-like, which increases the challenges of pairing electropositive f-element and increasingly large pnictogen metal ions together to form weak and highly polarised metal–metal bonds. Thus, well-defined molecules are of vital importance to study the inherent physicochemical properties and nature of covalency in f-element ligand bonds. This in turn could benefit the development of new synthetic methods, ligand design, catalytic



Junru Ding

*Junru Ding received her MSc degree at Zhengzhou University in 2023, and then began her PhD investigating f-block organometallic chemistry under the supervision of Prof. Jingzhen Du at the College of Chemistry, Zhengzhou University.*



David P. Mills

*David P. Mills is a Professor of Inorganic Chemistry in the Department of Chemistry at The University of Manchester, where he started his independent career as a Lecturer in 2012. His current research interests are focussed on the synthesis, characterisation, and applications of coordinatively unsaturated complexes with unusual electronic structures and bonding motifs, mainly with the f-block elements.*



Stephen T. Liddle

*Steve Liddle is Professor and Head of Inorganic Chemistry and Co-Director of the Centre for Radiochemistry Research at the University of Manchester Department of Chemistry, which are positions he has held since 2015. His research interests broadly encompass the physical-inorganic chemistry of the f-elements, with particular interests in synthesis, reactivity, magnetism, small molecule activation, and catalysis.*

† Shortly after the census date of this review, reports of a yttrium-bismoly complex and a yttrium-bismuth cluster were published by Demir and co-workers, and an account of rare-earth phosphinidene complexes was published by Chen and co-workers. See ref. 156–158 for details.



Table 1  $^{31}\text{P}$  NMR chemical shifts of reported f-element complexes with phosphorus ligands in this review


Complex name	$^{31}\text{P}$ NMR (ppm)	Solvent	Ref.
$[\text{Y}(\text{Cp}^{\prime\prime})_2(\text{THF})(\text{PHSi}^i\text{Bu}_3)]$ (5)	−181.1	$\text{C}_6\text{D}_6$	44
$[\text{Y}(\text{Cp}^{\prime\prime})_2\{(\mu\text{-PH}_2)(\mu\text{-Li}[\text{tmada}])_2(\text{Cl})\}]$ (6)	−218.5	Toluene- $d_8$	45
$[\text{Yb}(\text{Cp}^*)_2\{(\text{PCHCMeCMeCHC})_2\}]$ (8c)	191.6	THF- $d_8$	48
$[\text{Yb}(\text{Cp}^*)_2\{(\text{PCHCMeCMeCHC})_2\}]$ (8c)	178.2	Toluene- $d_8$	48
$[\{\text{Lu}(\text{PNP}^{\text{iPr}})(\mu\text{-PMes})_2\}]$ (9)	186.8	$\text{C}_6\text{D}_6$	50
$[\{\text{Lu}(\text{PNP}^{\text{iPr}})(\mu\text{-PMes})_2\}]$ (9)	18.1	$\text{C}_6\text{D}_6$	50
$[\{\text{Nd}(\mu\text{-PDipp})(\text{I})(\text{THF})_3\}_2]$ (10)	−168	$\text{C}_6\text{D}_6$	51
$[\{\text{Sc}(\text{PNP}^{\text{iPr}})(\mu\text{-PTripp})_2\}]$ (13)	227.4	$\text{C}_6\text{D}_6$	53
$[\{\text{Sc}(\text{PNP}^{\text{iPr}})(\mu\text{-PTripp})_2\}]$ (13)	7.0	$\text{C}_6\text{D}_6$	53
$[\text{Sc}(\text{PNP}^{\text{iPr}})(\mu\text{-PDmp})(\mu\text{-Br})\text{Li}]$ (14)	9.8	$\text{C}_6\text{D}_6$	53
$[\text{Sc}(\text{PNP}^{\text{iPr}})(\mu\text{-PDmp})(\mu\text{-Br})\text{Li}]$ (14)	8.0/13.2	$\text{C}_6\text{D}_6$	53
$[\text{Sc}(\text{PNP}^{\text{iPr}})(\mu\text{-PDmp})(\mu\text{-Br})\text{Li}(\text{DME})]$ (15)	56.1	$\text{C}_6\text{D}_6$	53
$[\text{Sc}(\text{PNP}^{\text{iPr}})(\mu\text{-PDmp})(\mu\text{-Br})\text{Li}(\text{DME})]$ (15)	10.8/5.6	$\text{C}_6\text{D}_6$	53
$[\{\text{Sc}(\text{NCCN}^{\text{iPr}})(\mu\text{-PXyl})_2\}]$ (16)	183.8	$\text{C}_6\text{D}_6$	54
$[\{\text{Sc}(\text{NCCN}^{\text{iPr}})(\mu\text{-PXyl})(\text{DMAP})_2\}]$ (17)	181.3	$\text{C}_6\text{D}_6$	54
$[\text{Sc}(\text{NCCN}^{\text{iPr}})(2,2'\text{-bipy})\{\eta^2\text{-P}_2(\text{Xyl})_2\}]$ (18)	30.0/25.9	$\text{C}_6\text{D}_6$	54
$[\{\text{Sc}(\text{NCCN}^{\text{iPr}})_2(\mu\text{-S})\{\mu\text{-}\eta^2\text{-P}_2(\text{Xyl})_2\}]$ (19S)	−79.0	$\text{C}_6\text{D}_6$	54
$[\{\text{Sc}(\text{NCCN}^{\text{iPr}})_2(\mu\text{-Se})\{\mu\text{-}\eta^2\text{-P}_2(\text{Xyl})_2\}]$ (19Se)	−72.2	$\text{C}_6\text{D}_6$	54
$[\{\text{Sc}(\text{NCCNiPr})_2(\mu\text{-Te})\{\mu\text{-}\eta^2\text{-P}_2(\text{Xyl})_2\}]$ (19Te)	−60.1	$\text{C}_6\text{D}_6$	54
$[\text{Sc}(\text{NCCN}^{\text{Dipp}})(\text{Me})\{\text{P}(\text{H})\text{Dipp}\}]$ (20)	−90.9	$\text{C}_6\text{D}_6$	55
$[\{\text{Sc}(\text{NCCN}^{\text{Dipp}})_2(\mu\text{-CH}_2)(\mu\text{-PDipp})]$ (21)	84.1	$\text{C}_6\text{D}_6$	55
$[\{\text{Y}[\text{PhC}(\text{NDipp})_2](\mu_2\text{-Me})_3(\mu_3\text{-Me})(\mu_3\text{-PPh})]$ (26Y)	138.8	$\text{C}_6\text{D}_6$	56
$[\{\text{Lu}[\text{PhC}(\text{NDipp})_2](\mu_2\text{-Me})_3(\mu_3\text{-Me})(\mu_3\text{-PPh})]$ (26Lu)	103.4	$\text{C}_6\text{D}_6$	56
$[\{\text{Y}[\text{PhC}(\text{NDipp})_2](\mu_2\text{-Me})_3(\mu_3\text{-Me})(\mu_2, \eta^2: \eta^3\text{-PC}_6\text{H}_4)]$ (27Y)	262.48	$\text{C}_6\text{D}_6$	56
$[\{\text{Lu}[\text{PhC}(\text{NDipp})_2](\mu_2\text{-Me})_3(\mu_3\text{-Me})(\mu_2, \eta^2: \eta^3\text{-PC}_6\text{H}_4)]$ (27Lu)	192.52	$\text{C}_6\text{D}_6$	56
$[\{\text{Y}(\text{Cp}^{\text{Me}})_2\}_3(\mu\text{-PMes})_3\text{Li}][\text{Li}(\text{THF})_4]_2$ (28Y)	57.24	$\text{C}_6\text{D}_6$	58
$[\text{Sc}(\text{NCCN}^{\text{Dipp}})\{\text{PP}(\text{EDA}^{\text{Dipp}})\}]$ (29)	412.0	$\text{C}_6\text{D}_6$	59
$[\text{Sc}(\text{NCCN}^{\text{Dipp}})\{\text{PP}(\text{EDA}^{\text{Dipp}})\}]$ (29)	157.2	$\text{C}_6\text{D}_6$	59
$[\text{Sc}(\text{NCCN}^{\text{Dipp}})\{\text{PP}(\text{EDA}^{\text{Dipp}})\}]$ (29)	402.3	THF- $d_8$	59
$[\text{Sc}(\text{NCCN}^{\text{Dipp}})\{\text{PP}(\text{EDA}^{\text{Dipp}})\}]$ (29)	158.5	THF- $d_8$	59
$[\text{Sc}(\text{NCCN}^{\text{Me}})\{\text{PP}(\text{EDA}^{\text{Dipp}})\}]$ (30)	324.8	$\text{C}_6\text{D}_6$	59
$[\text{Sc}(\text{NCCN}^{\text{Me}})\{\text{PP}(\text{EDA}^{\text{Dipp}})\}]$ (30)	169.0	$\text{C}_6\text{D}_6$	59
$[\text{Sc}(\text{NCCN}^{\text{Me}})\{\text{PP}(\text{EDA}^{\text{Dipp}})\}]$ (30)	312.2	THF- $d_8$	59
$[\text{Sc}(\text{NCCN}^{\text{Me}})\{\text{PP}(\text{EDA}^{\text{Dipp}})\}]$ (30)	166.8	THF- $d_8$	59
$[\text{Sc}(\text{NCCN}^{\text{Me}})\{(\mu\text{-PB})[\text{N}(\text{Dipp})\text{CHCHN}(\text{Dipp})](\mu\text{-Cl})\text{K}\}]_2$ (35)	11.4	Toluene- $d_8$	60
$[\text{Sc}(\text{NCCN}^{\text{Me}})\{(\mu\text{-PB})[\text{N}(\text{Dipp})\text{CHCHN}(\text{Dipp})](\mu\text{-Cl})\{\text{K}(\text{DB18C6})\}]$ (36)	19.6	Toluene- $d_8$	60
$[\text{Sc}(\text{NCCN}^{\text{Me}})\{\text{N}(\text{Pr})\text{C}(\text{PB}[\text{N}(\text{Dipp})\text{CHCHN}(\text{Dipp})]\text{N}(\text{Pr})\}]$ (37)	−103.1	$\text{C}_6\text{D}_6$	60
$[\text{Y}(\text{Tp}^{\text{tBu, Me}})(\text{Me})(\text{HPDipp})]$ (38Y)	−117.8	$\text{C}_6\text{D}_6$	61
$[\text{Y}(\text{Tp}^{\text{tBu, Me}})(\text{PDipp})(\text{DMAP})_2]$ (40)	−5.5	$\text{C}_6\text{D}_6$	61
$[\{\text{Y}(\text{I})\}\{\text{Y}[\mu_3\text{-P}(\text{Dipp})](\mu\text{-I})(\text{THF})_4(\mu_6\text{-P})\{\text{K}(\text{C}_7\text{H}_8)\}]$ (41)	347.4	$\text{C}_6\text{D}_6$	62
$[\{\text{Y}(\text{I})\}\{\text{Y}[\mu_3\text{-P}(\text{Dipp})](\mu\text{-I})(\text{THF})_4(\mu_6\text{-P})\{\text{K}(\text{C}_7\text{H}_8)\}]$ (41)	154.7	$\text{C}_6\text{D}_6$	62
$[\{\text{Y}(\text{I})\}\{\text{Y}[\mu_3\text{-P}(\text{Dipp})](\mu\text{-I})(\text{THF})_4(\mu_6\text{-P})\{\text{K}(\text{THF})\}]$ (42)	346.6	$\text{C}_6\text{D}_6$	62
$[\{\text{Y}(\text{I})\}\{\text{Y}[\mu_3\text{-P}(\text{Dipp})](\mu\text{-I})(\text{THF})_4(\mu_6\text{-P})\{\text{K}(\text{THF})\}]$ (42)	154.3	$\text{C}_6\text{D}_6$	62
$[\{\text{Y}(\text{I})\}\{\text{Y}(\text{I})(\text{THF})_2\}\{\text{Y}(\text{THF})_2\}_2(\mu\text{-I})[\mu_3\text{-P}(\text{Dipp})]_4(\mu_5\text{-P})]$ (43)	358.8	THF- $d_8$	62
$[\{\text{Y}(\text{I})\}\{\text{Y}(\text{I})(\text{THF})_2\}\{\text{Y}(\text{THF})_2\}_2(\mu\text{-I})[\mu_3\text{-P}(\text{Dipp})]_4(\mu_5\text{-P})]$ (43)	148.4	THF- $d_8$	62
$[\{\text{Y}(\text{I})\}\{\text{Y}(\text{I})(\text{THF})_2\}\{\text{Y}(\text{NCCN}^{\text{iPr}})_2\}_2(\mu\text{-I})[\mu_3\text{-P}(\text{Dipp})]_4(\mu_5\text{-P})]$ (44)	400.7	THF- $d_8$	62
$[\{\text{Y}(\text{I})\}\{\text{Y}(\text{I})(\text{THF})_2\}\{\text{Y}(\text{NCCN}^{\text{iPr}})_2\}_2(\mu\text{-I})[\mu_3\text{-P}(\text{Dipp})]_4(\mu_5\text{-P})]$ (44)	176.3/172.9	THF- $d_8$	62
$[\{\text{Y}(\text{THF})\}\{\text{Y}(\mu\text{-I})(\text{THF})_2\}\{\text{Y}(\text{THF})\}\{\text{Y}(\text{Cp}^*)\}(\mu_3\text{-I})(\mu\text{-I})[\mu_3\text{-P}(\text{Dipp})]_4(\mu_6\text{-P})\{\text{K}(\text{THF})\}]$ (45)	300.9	THF- $d_8$	62
$[\{\text{Y}(\text{THF})\}\{\text{Y}(\mu\text{-I})(\text{THF})_2\}\{\text{Y}(\text{THF})\}\{\text{Y}(\text{Cp}^*)\}(\mu_3\text{-I})(\mu\text{-I})[\mu_3\text{-P}(\text{Dipp})]_4(\mu_6\text{-P})\{\text{K}(\text{THF})\}]$ (45)	72.8 to 130.4	THF- $d_8$	62
$[\{\text{Sc}(\text{NN}^{\text{fC}})_3\text{P}_7]$ (47Sc)	−131.4 to 23.2	$\text{C}_6\text{D}_6$	64
$[\{\text{Y}(\text{NN}^{\text{fC}})(\text{THF})_3\text{P}_7]$ (47Y)	−130.4 to −20.4	$\text{C}_6\text{D}_6$	64
$[\{\text{Sm}(\text{DippForm})_2\}_2(\mu^2\text{-}\eta^4\text{-}\eta^4\text{-P}_4)]$ (48)	453	$\text{C}_6\text{D}_6$	67
$[\{\text{Y}(\text{DippDBD})(\text{THF})_2(\text{P}_3)]\text{K}(\text{18C6}(\text{toluene}))]$ (49)	−273.01	THF- $d_8$	68
$[\text{Th}(\text{Cp}^*)_2(\text{PPh}_2)_2]$ (50)	143	$\text{C}_6\text{D}_6$	79
$[\text{Th}(\text{Cp}^*)_2(\mu\text{-PPh}_2)_2\text{Ni}(\text{CO})_2]$ (51)	177	$\text{C}_6\text{D}_6$	79
$[\text{Th}(\text{Cp}^*)_2(\mu\text{-PPh}_2)_2\text{Pt}(\text{PMe}_3)]$ (52)	149.3	Toluene- $d_8$	80
$[\text{Th}(\text{Cp}^*)_2(\mu\text{-PPh}_2)_2\text{Pt}(\text{PMe}_3)]$ (52)	−3.3	Toluene- $d_8$	80
$[\text{Th}(\text{Cp}^*)_2(\text{PHTripp})_2]$ (53)	1.66	$\text{C}_6\text{D}_6$	81
$[\text{Th}(\text{Tren}^{\text{TIPS}})(\text{PH}_2)]$ (54Th)	−144.08	$\text{C}_6\text{D}_6$	83
$[\text{U}(\text{Tren}^{\text{TIPS}})(\text{PH}_2)]$ (54U)	595.07	$\text{C}_6\text{D}_6$	82
$[\text{Th}(\text{Tren}^{\text{TCHS}})(\text{PH}_2)]$ (55Th)	−133.01	THF- $d_8$	84
$[\text{U}(\text{Tren}^{\text{TCHS}})(\text{PH}_2)]$ (55U)	605.91	THF- $d_8$	84
$[\{\text{Th}(\text{Cp}^*)_2\}_2\{\mu\text{-P}[(2,6\text{-CH}_2\text{CHCH}_3)_2\text{-4-}^i\text{PrC}_6\text{H}_2]\}]$ (57)	161.9	$\text{C}_6\text{D}_6$	81



Complex name	<sup>31</sup> P NMR (ppm)	Solvent	Ref.
[{Th(Cp*) <sub>2</sub> (μ-PTripp)(μ-PHTripp)(K)} <sub>2</sub> ] ( <b>58</b> )	171.91	THF- <i>d</i> <sub>8</sub>	87
[{Th(Cp*) <sub>2</sub> (μ-PTripp)(μ-PHTripp)(K)} <sub>2</sub> ] ( <b>58</b> )	−110.54	THF- <i>d</i> <sub>8</sub>	87
[Th(Cp*) <sub>2</sub> (PTripp)](PTripp)[K(2,2,2-cryptand)] ( <b>59</b> )	177.86	THF- <i>d</i> <sub>8</sub>	87
[Th(Cp*) <sub>2</sub> (PTripp)](PTripp)[K(2,2,2-cryptand)] ( <b>59</b> )	−106.99	THF- <i>d</i> <sub>8</sub>	87
[{Th(Cp <sup>tt</sup> ) <sub>2</sub> (PMes*){ClK}} <sub>2</sub> ] ( <b>60</b> )	108.79	C <sub>6</sub> D <sub>6</sub>	93
[Th(Cp <sup>tt</sup> ) <sub>2</sub> (PMes*)(μ-Cl){K(18C6)}] ( <b>61</b> )	133.5	C <sub>6</sub> D <sub>6</sub>	93
[U(Cp*) <sub>2</sub> (PMes*)(OPMe <sub>3</sub> )] ( <b>62</b> )	71.06	C <sub>6</sub> D <sub>6</sub>	95
[U(Cp*) <sub>2</sub> (PMes*)(OPMe <sub>3</sub> )] ( <b>62</b> )	−59.84	C <sub>6</sub> D <sub>6</sub>	95
[Th(Cp <sup>ttt</sup> ) <sub>2</sub> (PMes*)] ( <b>63</b> )	145.7	C <sub>6</sub> D <sub>6</sub>	97
[{U(Tren <sup>TIPS</sup> )(μ-PH)}{K(2,2,2-cryptand)}] ( <b>66</b> )	2460.4	C <sub>6</sub> D <sub>6</sub>	82
[Th(Tren <sup>TIPS</sup> )(PH)][Na(12C4) <sub>2</sub> ] ( <b>67</b> )	198.8	C <sub>6</sub> D <sub>6</sub>	83
[Th(Tren <sup>TCHS</sup> )(PH)][Na(2,2,2-cryptand)] ( <b>69Th</b> )	266.16	THF- <i>d</i> <sub>8</sub>	84
[U(Tren <sup>TCHS</sup> )(PH)][Na(2,2,2-cryptand)] ( <b>69U</b> )	2628.50	THF- <i>d</i> <sub>8</sub>	84
[{Th(Tren <sup>TIPS</sup> ) <sub>2</sub> (μ-P)][Na(12C4) <sub>2</sub> ] ( <b>70</b> )	553.5	THF- <i>d</i> <sub>8</sub>	83
[{Th(Tren <sup>TIPS</sup> ) <sub>2</sub> (μ-PH)] ( <b>71Th</b> )	145.7	C <sub>6</sub> D <sub>6</sub>	83
[Th(Tren <sup>TIPS</sup> )(OCP)] ( <b>74Th</b> )	−339.91	C <sub>6</sub> D <sub>6</sub>	110
[U(Tren <sup>TIPS</sup> )(OCP)] ( <b>74U</b> )	−319.96	C <sub>6</sub> D <sub>6</sub>	109
[{Th(Tren <sup>TIPS</sup> ) <sub>6</sub> (μ-OC <sub>2</sub> P <sub>3</sub> H) <sub>2</sub> (μ-OC <sub>2</sub> P <sub>3</sub> H) <sub>2</sub> Rb <sub>4</sub> }] ( <b>76</b> )	217.99 to 261.14	C <sub>6</sub> D <sub>6</sub>	110
[{Th(Cp <sup>tt</sup> ) <sub>2</sub> } <sub>2</sub> (μ <sup>2</sup> -η <sup>4</sup> -P <sub>6</sub> )] ( <b>77</b> )	−41.9 to 125.3	Toluene- <i>d</i> <sub>8</sub>	111
[{Th(Cp <sup>tt</sup> ) <sub>2</sub> } <sub>2</sub> (μ <sup>2</sup> -η <sup>3</sup> -P <sub>3</sub> ){Th(Cp <sup>tt</sup> ) <sub>2</sub> Cl}] ( <b>78</b> )	−94.5 to −69.7	CD <sub>2</sub> Cl <sub>2</sub>	111
[{U(Cp*)(C <sub>6</sub> H <sub>6</sub> (Si <sup>†</sup> Pr <sub>3</sub> ) <sub>2</sub> ) <sub>2</sub> (μ <sup>2</sup> -η <sup>4</sup> -P <sub>4</sub> )] ( <b>79</b> )	718	C <sub>6</sub> D <sub>6</sub>	112
[{Th(Cp <sup>tt</sup> ) <sub>3</sub> } <sub>2</sub> (μ <sup>2</sup> -η <sup>2</sup> -P <sub>4</sub> )] ( <b>80</b> )	−246.55 to 323	C <sub>6</sub> D <sub>6</sub>	113

This review highlights the most notable achievements in the field of f-element heavy pnictogen chemistry from phosphorus to the heaviest abundant main group element bismuth up to August 2023. In line with the criteria for reviews, a representative selection, rather than a complete literature survey, is presented, and discussions concentrate on structurally characterised molecules. We aim to highlight the major advances involving all heavy pnictogen ligand types, with the exception of phospholyl and arsolyl ligands, which were reviewed in 2021,<sup>30</sup> and (OCE)<sup>−</sup> (E = P, As) ligands, which were reviewed in 2019 and are normally O-bound unless the E centre decisively directs the chemistry;<sup>31</sup> several other previous reviews have separately covered the ligand classes that comprise this review.<sup>20,24–29</sup> Here we present current challenges to inspire researchers and focus and guide future efforts of the field to develop f-element heavy pnictogen chemistry more rapidly in the future. In this review, we include the group 3 elements scandium, yttrium, and lanthanum under the heading of lanthanide sections for convenience. <sup>31</sup>P NMR chemical shifts for P-bound complexes covered in this review are compiled in Table 1.

Metal heavy pnictogen nomenclature depends upon the pnictogen identity, charge and binding mode. The prefix is



© 2024 The Author(s). Published by the Royal Society of Chemistry





bond distance is 2.770(1) Å. Complex 5 is diamagnetic and the  $^{31}\text{P}$  NMR spectrum exhibits a doublet of doublets resonance at  $-181.1$  ppm with  $^1J_{\text{YP}}$  and  $^1J_{\text{PH}}$  values of 144.0 and 201.0 Hz, respectively. Phosphanide complexes featuring the  $(\text{PH}_2)^-$  ligand are exceedingly rare for f elements. Although actinide phosphanide complexes with terminal An-PH<sub>2</sub> linkages are known (see below), isostructural analogues still remain rare for lanthanides. The only relevant example is the yttrium complex  $[\text{Y}(\text{Cp}^*)_2\{(\mu\text{-PH}_2)(\mu\text{-Li}[\text{tmeda}])\}_2(\text{Cl})\}]$  (6), Fig. 3, which contains two  $(\text{PH}_2)^-$  groups that bridge to two lithium cations.<sup>45</sup> It was found that 6 is unstable in both solid and solution states, and decomposes under argon atmosphere at room temperature to produce PH<sub>3</sub> as well as a small amount of H<sub>2</sub>PSiMe<sub>3</sub>, reflecting the synthetic challenge of stabilising terminal lanthanide-PH<sub>2</sub> species.

## 4.2 Lanthanide phosphorin and biphosphinine complexes

In 1997, Cloke and co-workers reported the synthesis and molecular structure of the bis(2,4,6-tri-*tert*-butyl-phosphorin) holmium(0) complex  $[\text{Ho}(\eta^6\text{-Ttp})_2]$  (7), Fig. 4, prepared by co-condensation of holmium vapor with an excess of 2,4,6-tri-*tert*-butyl-phosphorin at  $-196$  °C followed by further work-up and recrystallisation.<sup>46</sup> The remarkable thermostability of 7 ( $T_{\text{sublimation}} = 160$  °C,  $10^{-5}$  mbar, 90% recovery) arises from the better  $\pi$ -acceptor capability for phosphorin over arenes, which was confirmed by optical and magnetic data for 7. The structure of 7 was found to exhibit extensive disorder, with the P-atoms equally disordered over the three possible positions of each phosphorin ligand, so no preference for *syn* or *anti* conformations could be inferred.

During 2014 to 2016 Nocton, Clavaguéra, and co-workers reported biphosphinine complexes of the general formula  $[\text{Ln}(\text{L})_2\{(\text{PCHCMCMeCHC})_2\}]$  (Ln = Tm, L = {P(CBu<sup>t</sup>CMe)<sub>2</sub>}, 8a; Ln = Sm or Yb, L = Cp\*, C<sub>5</sub>Me<sub>5</sub>, 8b, 8c),<sup>47,48</sup> Fig. 4. In these complexes the biphosphinine ligands are formally radical anions but the extent of electron transfer for the Yb complex was ambiguous, with characterisation data intermediate to closed or fully open shell formulations. The Tm-P, Sm-P, and Yb-P distances were found to be 2.825/2.862(2), 2.909(2)/2.927(2), and 2.872(2)/2.938(2) Å, consistent with the radii of the lanthanide ions. We note that this work followed on from prior work on P-methylated phosphinine ligands with chelating side arms from Arliguie, Mézailles, and co-workers;



Fig. 4 Lanthanide complexes containing phosphorin ligands 7 and 8.

however, in those complexes the anion charge partially delocalises into the C<sub>5</sub>P rings, resulting in rather long M-P bonds ( $\sim 3$  Å for Ce, Nd, and U) that are between dative phosphines and covalent phosphides,<sup>49</sup> thus we do not discuss them further.

## 4.3 Lanthanide phosphinidiide complexes

Unlike their d-transition metal counterparts, lanthanide-pnictogen multiple bonds are relatively rare as a result of the valence orbital spatial and energy mismatch of 4f metal ions and pnictogen ligands. Most lanthanide pnictinidenes form bridging dimeric pnictinidiide complexes where the “Ln=Pn” moiety is stabilised through additional interactions with adjacent rare earth metal centres or electropositive alkali metal cations.<sup>26</sup> This section describes the progresses made in isolation of a handful of lanthanide pnictinidiide complexes before the terminal phosphinidene species was finally secured very recently.

In 2008, Kiplinger and co-workers synthesised the first bridging phosphinidiide lanthanide complex  $[\text{Lu}\{(\text{PNP}^{\text{iPr}})(\mu\text{-PMes})\}_2]$  (9,  $\text{PNP}^{\text{iPr}} = [\{2\text{-}(\text{iPr}_2\text{P})\text{C}_6\text{H}_4\}_2\text{N}\}]^-$ ) by protonolysis of  $[(\text{PNP}^{\text{iPr}})\text{Lu}(\text{CH}_2\text{SiMe}_3)_2]$  with MesPH<sub>2</sub>, in 52% yield, Scheme 1.<sup>50</sup> Complex 9 exhibits an asymmetric Lu<sub>2</sub>P<sub>2</sub> core, with two short [2.5973(15)/2.6031(16) Å] and two long [2.6527(16)/2.6724(14) Å] Lu-P bonds. The sum of the angles about the phosphorus atoms range from 358.9 to 356.5°, which indicates that the phosphorus lone pairs possibly  $\pi$ -donate to the Lu ions. The authors concluded that the structural data for 9 suggested




Scheme 1 Synthesis of Lu-phosphinidiide complex 9.



Scheme 2 Synthesis of Nd-phosphinidiide complexes 10–12.

In 2010, Mindiola and co-workers reported that the reaction of the sterically demanding phosphide precursor,  $\text{LiPH}(\text{Tripp})$  ( $\text{Tripp} = 2,4,6\text{-iPr}_3\text{C}_6\text{H}_2$ ), with  $[\text{Sc}(\text{PNP}^{\text{iPr}})(\text{Me})(\text{Br})]$  yielded the bridging phosphinidide dinuclear scandium complex



Mindiola and co-workers subsequently disclosed the first mononuclear rare earth phosphinidiide complexes. The Li-capped complexes [Sc(PNP<sup>iPr</sup>)(μ-PDmp)(μ-Br)Li(DME)<sub>*n*</sub>] (Dmp = 2,6-Mes<sub>2</sub>C<sub>6</sub>H<sub>3</sub>; *n* = 0, **14**; *n* = 1, **15**) were prepared from the concomitant salt metathesis and protonolysis reactions of the bulky primary phosphide lithium salt, LiPHDmp with [(PNP<sup>iPr</sup>)Sc(Me)(Br)], eliminating methane as the driving force, to give **14** (55%) and **15** (47%), Scheme 4.<sup>53</sup> The exact product depends on the reaction solvent; complex **15** was also synthesised by the addition of a stoichiometric amount of DME to **14** (21%). Single crystal X-ray diffraction studies revealed that both **14** and **15** have short Sc–P bond lengths of 2.338(2) and 2.3732(18) Å, respectively. Calculations performed on **15** indicated that the Sc=P bond has significant multiple bond character with a Mayer bond order of 1.46. The authors varied the reaction



conditions for the synthesis of **14** and **15** to investigate if the elimination of LiBr was possible. However, heating reaction mixtures up to 100 °C did not liberate the occluded LiBr.


In 2013, Maron, Chen and co-workers utilised a similar one-pot salt metathesis and protonolysis methodology with the reaction of KPXyl (Xyl = 2,6-Me<sub>2</sub>C<sub>6</sub>H<sub>3</sub>) with [Sc(NCCN<sup>iPr</sup>)(Me)(Cl)] (NCCN<sup>iPr</sup> = MeC(NDipp)CHC(Me)(NCH<sub>2</sub>CH<sub>2</sub>N<sup>iPr</sup>Pr<sub>2</sub>) yielding the bridging phosphinidide dinuclear scandium complex [{Sc(NCCN<sup>iPr</sup>)(μ-PXyl)}<sub>2</sub>] (**16**, 74%), *via* the elimination of KI and methane, Scheme 5.<sup>54</sup> In contrast with the previously reported Sc-phosphinidide complex **13**, the Sc<sub>2</sub>P<sub>2</sub> core in **16** is more symmetric with Sc–P distances of 2.522(1) and 2.528(1) Å. They also attempted to synthesise a terminal Sc-phosphinidene complex by reacting **16** with a strongly donating ligand, DMAP (DMAP = 4-NMe<sub>2</sub>-C<sub>5</sub>H<sub>4</sub>N) to break the dimer up; however, this resulted in the formation of the adduct [{Sc(NCCN<sup>iPr</sup>)(μ-PXyl)(DMAP)}<sub>2</sub>] (**17**) in a 90% yield. The authors noted that **17** was the major product if an excess of DMAP (four equivalents) was added to **16**. Whilst **17** retains its Sc<sub>2</sub>P<sub>2</sub> core, the Sc–P bonds are longer than **16** at 2.540(1) and 2.589(1) Å, reflecting the increased Sc coordination numbers.

Maron, Chen and co-workers showed that **16** exhibits notable redox chemistry; for example, the addition of three equivalents of 2,2'-bipyridine (2,2'-bipy) results in the oxidative coupling of two phosphinidide fragments to form a diphosphanide Sc-complex [Sc(NCCN<sup>iPr</sup>)(2,2'-bipy){η<sup>2</sup>-P<sub>2</sub>(Xyl)<sub>2</sub>}] (**18**), with the single electron reduction of two molecules of 2,2'-bipy


affording [Sc(NCCN<sup>iPr</sup>)(2,2'-bipy)<sub>2</sub>] as a side product, Scheme 6. Similarly, the oxidative coupling of the two (PXyl)<sup>2-</sup> ligands to (P<sub>2</sub>(Xyl)<sub>2</sub>)<sup>2-</sup> promotes the reduction of elemental selenium/tellurium or Ph<sub>3</sub>P=E (E = Se or S) to S<sup>2-</sup>, Se<sup>2-</sup> or Te<sup>2-</sup>, yielding the bridging chalcogenido complexes [{Sc(NCCN<sup>iPr</sup>)}<sub>2</sub>(μ-E){μ-η<sup>2</sup>-P<sub>2</sub>(Xyl)<sub>2</sub>}] (**19E**, E = S, Se or Te). The authors also found that **16** readily undergoes nucleophilic addition chemistry with a range of unsaturated allene, nitrile, isocyanide, and CS<sub>2</sub> substrates to generate the corresponding organophosphorus scandium complexes. In follow-up work they reported that **16** could readily cleave the boron–oxygen bonds in pinacol–borane and catecholborane to give complexes featuring a newly-generated ligand (HB{P(Xyl)}<sub>2</sub>)<sup>2-</sup>, demonstrating the reactive nature of the Sc-phosphinidide species due to the highly ionic Sc–P bonding interaction.<sup>54</sup>

In 2015, Maron, Chen and co-workers found that thermal decomposition of the scandium phosphide precursor [Sc(NCCN<sup>Dipp</sup>)(Me){P(H)Dipp}] (**20**, NCCN<sup>Dipp</sup> = {MeC(NDipp)}<sub>2</sub>CH) supported by the NCCN<sup>Dipp</sup> scaffold afforded the scandium phosphinidide complex [{(NCCN<sup>Dipp</sup>)Sc}<sub>2</sub>(μ-CH<sub>2</sub>)(μ-PDipp)] (**21**). However, this reaction also gave a diphosphide by-product [(NCCN<sup>Dipp</sup>){P(H)Dipp}<sub>2</sub>] in a 1 : 1 ratio with **21**. An optimal route was then implemented by reacting one equivalent of **20** with [Sc(NCCN<sup>Dipp</sup>)(Me)<sub>2</sub>], giving **21** in a 77% yield, Scheme 7.<sup>55</sup> Complex **21** exhibits Sc–P bond lengths of 2.495(1) and 2.508(1) Å, which are shorter than other scandium bridging phosphinidide complexes. Interestingly, there is an up-field shift of the phosphinidide moiety resonances in the <sup>31</sup>P NMR spectrum of **21** to 84.1 ppm, *cf.* **13** (227.4 ppm) and **16** (183.8 ppm), which is presumably due to one of the bridging phosphinidide units being replaced by a methylenide fragment.

In the same publication Maron, Chen and co-workers conducted a reactivity study of **21**; this revealed that, in contrast with the previously reported complex **16**, the phosphinidide ligand in **21** is relatively unreactive, with all small molecules reacting at the adjacent methylenide centre, Scheme 8. The reaction of **21** with either CO<sub>2</sub>, PhCN, <sup>t</sup>BuNC or CS<sub>2</sub> resulted in



Scheme 6 Selected redox reactions of **16**.



Scheme 7 Synthesis of the Sc-phosphinidide complex **21**.



Scheme 8 Reactivity of Sc-phosphinidide complex **21**.





insertion of the unsaturated functional group into the Sc-CH<sub>2</sub> bonds to give the bridging phosphinidiide complexes  $[\{\text{Sc}(\text{NCCN}^{\text{Dipp}})\}_2(\text{L})\{\mu\text{-P}(\text{Dipp})\}]$  (L =  $\mu\text{-O-COCH}_2\text{CO}_2$ , **22**;  $\mu\text{-N-C(Ph)(CH}_2\text{)}$ , **23**;  $\mu\text{-C(CH}_2\text{)(N}^t\text{Bu)}$ , **24**; or  $(\mu\text{-S})_2\text{CCH}_2$ , **25**). Complexes **22–25** all demonstrate similar Sc-P distances between 2.4922(13) and 2.588(2) Å. DFT studies of **21** indicated that the lack of reactivity of the Sc-P bond was possibly due to its increased covalency when compared to the Sc-C bonds in the same complex.<sup>55</sup>

Zhou, Luo, Zhang and co-workers later expanded the number of structurally characterised lanthanide phosphinidiide complexes *via* the protonolysis reactions of  $[\{\text{Ln}[\text{PhC}(\text{NDipp})_2](\mu_2\text{-Me})\}_3(\mu_3\text{-Me})(\mu_3\text{-CH}_2)]$  (Ln = Y, Lu) and one equivalent of PhPH<sub>2</sub> to give the clusters  $[\{\text{Ln}[\text{PhC}(\text{NDipp})_2](\mu_2\text{-Me})\}_3(\mu_3\text{-Me})(\mu_3\text{-PPh})]$  (Ln = Lu, **26Lu**, 92%; Ln = Y, **26Y**, 90%), Scheme 9.<sup>56</sup> These clusters exhibit a rare  $\mu^3$ -bridging mode of the phosphinidiide between three rare earth metal ions, which is in contrast to most other rare earth phosphinidiide complexes where a  $\mu^2$ -bridging mode is observed. Complex **26Y** exhibits one long [2.9432(12) Å] and two short [2.7142(11) and 2.7317(12) Å] Y-P bond lengths. The longer distance is within the range of previously reported dative R<sub>2</sub>P: → Ln(III) interactions. Once the change in metal radii is accounted for, this rationale can be applied to **26Lu**, which exhibits similar asymmetric bonding between the phosphinidiide and three bonded Ln(III) ions, with Lu-P distances of 2.902(2), 2.684(2) and 2.639(2) Å. Both complexes contain P-C bonds that are bent out of the Ln<sub>3</sub> plane (Lu: 49.3°; Y: 59.1°); this differs from analogous transition metal bridging phosphinidiide complexes where the P-C bond is approximately perpendicular to the M<sub>3</sub> plane. In common with other rare earth phosphinidiide complexes, the authors reported that complexes **26Ln** exhibited reactivity towards unsaturated molecules such as ketones, thiones, or isothiocyanates, where the complex undergoes phospho-Wittig chemistry, with the exchange of the phosphinidene for an oxo or a sulfido group. Upon heating complexes **26Ln** in toluene, an

additional molecule of methane was eliminated to yield the  $\mu_2$ -bridging phosphinidiide complexes  $[\{\text{Ln}[\text{PhC}(\text{NDipp})_2](\mu_2\text{-Me})_2(\mu_3\text{-Me})(\mu_2, \eta^2: \eta^3\text{-PC}_6\text{H}_4)]$ , **27Ln**, in high yields (Ln = Lu, 94%; Y, 91%). Complex **27Lu** exhibits Lu-P distances of 2.649(4)/2.644(4) Å, which is a similar range to that of the two shorter Lu-P bonds in **26Lu**, whilst **27Y** displays a range of Y-P distances [2.698(2)/2.692(2) Å] that are shorter than those seen for **26Y**. The authors have additionally reported the crystal structure of the yttrium bridging phosphinidiide complex  $[\{\text{Y}(\text{PhC}(\text{NDipp})_2)(\mu\text{-Me})\}_2(\mu_3\text{-CCCPh})(\mu, \eta^2: \eta^3\text{-PC}_6\text{H}_4)]$ ,<sup>57</sup> which exhibits Y-P bond distances of 2.701(3) and 2.700(3) Å; these are statistically indistinguishable from the corresponding distances seen in **27Y**. However, the synthetic route and additional characterisation data for this complex have not been reported to date.

In 2015, Layfield and co-workers synthesised the bridging phosphinidiide lanthanide complexes  $[\{\text{Ln}(\text{Cp}^{\text{Me}})_2\}_3(\mu\text{-PMes})_3\text{-Li}][\text{Li}(\text{THF})_4]_2$  (**28Ln**; Ln = Y or Dy; Cp<sup>Me</sup> = C<sub>5</sub>H<sub>4</sub>Me), Scheme 10.<sup>58</sup> The deprotonation reactions of the bridging lanthanide phosphides,  $[\{\text{Ln}(\text{Cp}^{\text{Me}})_2(\mu\text{-PHMes})\}_3]$  (Ln = Y or Dy), with three equivalents of *n*-butyl-lithium gave **28Y** and **28Dy**, in yields of 56% and 64%, respectively, in addition to three equivalents of butane gas. The solid-state structures of **28Ln** exhibit central Ln<sub>3</sub>P<sub>3</sub> cores in a chair-like configuration, and the phosphinidiide units are capped with a single lithium ion. The authors reported longer Y-P bond distances for **28Y** [2.7869(12)–2.8268(13) Å] than the yttrium complex **27Y**, which is possibly due to the coordination a lithium ion to the three



Scheme 10 Synthesis of Y- and Dy-phosphinidiide complexes **28Ln**.



Scheme 9 Synthesis of Lu- and Y-phosphinidiide complexes **26Ln–27Ln**.



Scheme 11 Synthesis of the Sc-phosphinophosphinidene complexes **29** and **30**.




Fig. 5 Reactivity of Sc-phosphinophosphinidene complexes **29** and **30** to give **31–34**.


phosphinidide units. Complex **28Dy** exhibits Dy–P bond distances of 2.7850(15)–2.8249(15) Å, which are shorter than the range of distances seen for the phosphide precursor [ $\text{Dy}(\text{Cp}^{\text{Me}})_2(\mu\text{-PHMe})_3$ ] [2.926(6)–2.951(6) Å], which is likely due to increasing negative charge localised at the phosphorous atom resulting in stronger electrostatic bonding.

In 2020, Maron, Chen and co-workers utilised the phosphinophosphinidene ligand  $[\text{PP}\{\text{N}(\text{Dipp})\text{CH}_2\text{CH}_2\text{N}(\text{Dipp})\}]^{2-}$  ( $\text{PP}\{\text{EDA}^{\text{Dipp}}\}^{2-}$ ) in the one-pot salt metathesis and protonolysis reactions of  $[\text{Sc}(\text{NCCN}^{\text{R}})(\text{Me})(\text{Cl})]$  ( $\text{NCCN}^{\text{R}} = \text{DippNC}(\text{Me})\text{CHC}(\text{Me})\text{NR}'$ ;  $\text{R} = \text{Me}$  (see **33** below)  $\text{R}' = \text{CH}_2\text{CH}_2\text{NMe}_2$  or  $\text{Dipp}$ ) with one equivalent of  $\text{K}[\text{HPP}\{\text{EDA}^{\text{Dipp}}\}]$  to afford the Sc phosphinophosphinidene complexes  $[\text{Sc}(\text{NCCN}^{\text{R}})\{\text{PP}(\text{EDA}^{\text{Dipp}})\}]$  ( $\text{R} = \text{Dipp}$ , **29**, 54%;  $\text{R} = \text{Me}$ , **30**, 67%), with elimination of one equivalent each of methane and potassium chloride, Scheme 11.<sup>59</sup> Due to the poor stability of the  $\text{K}[\text{HPP}\{\text{EDA}^{\text{Dipp}}\}]$  ligand precursor, it was synthesised *in situ* from the phosphine derivative and benzyl potassium. Complex **29** exhibits a shorter Sc–P phosphinidene bond distance [2.448(1) Å] when compared to **30** [2.484(1) Å], which the authors attributed to the increase in coordination number of the scandium centre from five to six.

Complex **29** displays interesting reactivity towards unsaturated small molecules such as alkynes when compared to **30**.<sup>59</sup> The reaction of **29** with one equivalent of  $\text{PhC}\equiv\text{CR}$  ( $\text{R} = \text{H}$  or  $\text{Me}$ ) gave  $[\text{Sc}(\text{NCCN}^{\text{Dipp}})\{\eta^2\text{-P}=\text{P}(\text{EDA}^{\text{Dipp}})\text{CR} = \text{CPh}\}]$ , Fig. 5, ( $\text{R} = \text{H}$ , **31**;  $\text{R} = \text{Me}$ , **32**), which was surprising as reactions with unsaturated molecules typically occur at the more nucleophilic centre, *i.e.* the phosphinidene ( $\text{P}^{\text{a}}$ -phosphorus). Indeed, the authors reported that **30** reacts as expected at the phosphinidene centre with the same alkynes  $\text{PhC}\equiv\text{CR}$  to yield  $[(\text{NCCN}^{\text{Me}})\text{Sc}\{\text{P}(\text{H})\text{P}(\text{EDA}^{\text{Dipp}})\}(\text{C}\equiv\text{CPh})]$  (**33**) and  $[(\text{NCCN}^{\text{Me}})\text{Sc}\{\eta^2\text{-PP}(\text{EDA}^{\text{Dipp}})\text{MeC} = \text{CPh}\}]$  (**34**). The calculated reaction pathways for **29** and **30** with  $\text{PhC}\equiv\text{CH}$  illustrated that the interesting reactivity at the  $\text{P}^{\text{b}}$ -phosphorus in the case of **29** was a consequence of the coordinated THF blocking the access of reactants to the  $\text{P}^{\text{a}}$ -phosphinidene centre.

#### 4.4 Lanthanide phosphinidene complexes

Whilst there were many lanthanide metal phosphinidide complexes reported in the past two decades, a bona fide terminal phosphinidene complex for any lanthanide metal remained elusive for decades. A breakthrough in this field was achieved very recently; in 2021, Maron, Chen and co-workers



Scheme 12 Synthesis and reactivity of a terminal Sc-phosphinidene complexes **36**.

reported the synthesis and molecular structure of the first terminal scandium phosphinidene complex, Scheme 12.<sup>60</sup>

Building on their Sc-phosphinophosphinidene work, Maron, Chen and co-workers reacted the boronylphosphine  $\text{H}_2\text{PB}\{\text{N}(\text{Dipp})\text{CHCHN}(\text{Dipp})\}$  with  $\text{KCH}_2\text{Ph}$  to give *in situ*-generated  $\text{K}[\text{HPB}\{\text{N}(\text{Dipp})\text{CHCHN}(\text{Dipp})\}]$ , which was treated with  $[\text{Sc}(\text{NCCN}^{\text{Me}})(\text{Me})(\text{Cl})]$  in a THF/toluene mixture to afford a K/Sc heterometallic phosphinidide complex **35** after heating at 50 °C for 24 h. Dimeric **35** could be converted to a terminal monomeric boronylphosphinidene complex **36** as a dark purple solid in 83% yield by reacting with dibenzo-18-crown-6 in toluene. The solid-state structure confirmed the boronylphosphinidene ligand of **36** adopts an end-on coordination even though the chloride ligand is still bridged between the Sc and K metal centres. The Sc–P bond length in **36** (2.381(1) Å) is close to that in **35** (2.397(2) Å) but shorter than that in the scandium phosphinophosphinidene complex **30** (2.484(1) Å). The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum for **36** at 25 °C shows a very broad signal, but data recorded at –30 °C gave a sharper resonance at 19.6 ppm. DFT studies on **36** revealed a three-centre two-electron (3c-2e) Sc–P–B  $\sigma$  bond with a strong Sc–P  $\pi$ -interaction. In line with the nucleophilic nature of the phosphinidene ligand, a preliminary reactivity study showed that **36** reacted with  $N,N'$ -diisopropylcarbodiimide at room temperature *via* a  $[2 + 2]$ -addition fashion to give a four-membered scandium metallaheterocycle complex,  $[\text{Sc}(\text{NCCN}^{\text{Me}})\{\text{N}(\text{Pr})\text{C}(\text{PB}\{\text{N}(\text{Dipp})\}\text{CHCHN}(\text{Dipp}))\}\text{N}(\text{Pr})]$  (**37**) in 82% yield.

Very recently, Sirsch, Anwender and co-workers reported the synthesis and molecular structure of the first terminal yttrium phosphinidene complex, Scheme 13.<sup>61</sup> The reactions of  $\text{H}_2\text{PDipp}$  with the lanthanide dimethyl complexes  $[\text{Ln}(\text{Tp}^{\text{tBu,Me}})(\text{Me})_2]$  ( $\text{Ln} = \text{Y, Dy, Ho}$ ;  $\text{Tp}^{\text{tBu,Me}} = \text{HB}(2\text{-Me-4-}^t\text{Bu-N}_2\text{C}_3\text{H})_3$ ) afforded the corresponding phosphide complexes  $[(\text{Tp}^{\text{tBu,Me}})\text{Ln}(\text{Me})(\text{HPDipp})]$  (**38Ln**,  $\text{Ln} = \text{Y, Dy, Ho}$ ). Addition of DMAP to **38Ln** gave the corresponding DMAP adducts  $[(\text{Tp}^{\text{tBu,Me}})\text{Ln}(\text{Me})(\text{HPDipp})(\text{DMAP})]$  (**39Ln**,  $\text{Ln} = \text{Y, Dy, Ho}$ );




Scheme 13 Synthesis of a terminal Y-phosphinidene complex 40.

addition of a further equivalent of DMAP to **38Y** produced  $[\text{Y}(\text{Tp}^{\text{tBu,Me}})(\text{PDipp})(\text{DMAP})_2]$  (**40**). Overall, **40** was prepared *via* a double-deprotonation approach previously established by Anwander for terminal lanthanide imido complexes. However, attempts to synthesise the Dy and Ho phosphinidene analogues using the similar strategy were unsuccessful. Moreover, the Lewis acid-stabilised yttrium phosphinidide  $[\text{Y}(\text{Tp}^{\text{tBu,Me}})[(\mu\text{-PDipp})(\mu\text{-Me})\text{AlMe}_2]]$  was prepared by addition of  $\text{H}_2\text{PDipp}$  to the solution of  $[\text{Y}(\text{Tp}^{\text{tBu,Me}})(\text{Me})(\text{AlMe}_4)]$  in toluene. However, cleavage of trimethylaluminum within this phosphinidide complex to target a terminal phosphinidene was unsuccessful as well. The solid-state structure of **40** revealed a short Y–P distance of 2.4855(7) Å, which is in between the sum of single and double covalent bond radii of Y and P (2.74 and 2.32 Å, respectively), indicating multiple bonding character. The  $^{31}\text{P}$  NMR spectrum of **40** exhibits a very weak signal at  $-5.5$  ppm with a maximum  $^1J_{\text{YP}}$  coupling constant of 282 Hz, which is shifted to lower frequency compared to that of  $[\text{Y}(\text{Tp}^{\text{tBu,Me}})[(\mu\text{-PDipp})(\mu\text{-Me})\text{AlMe}_2]]$  at  $-52.2$  ppm, suggesting the increase of electron density at the yttrium centre in **40**. A computational study of **40** confirmed the highly polarised covalent multiple bond of Y=PDipp linkage and revealed one  $\sigma$ - and two  $\pi$ -type Y–P interactions with a Wiberg bond index (WBI) of 1.37. The successful isolation and characterisation of terminal phosphinidenes **36** and **40** indicates that it should be possible to access terminal phosphinidene species with other lanthanides with the right supporting ligands combined with suitable phosphinidene group transfer methodologies and reagents.

#### 4.5 Lanthanide phosphido complexes

In common with their pnictinidene derivatives, there are relatively few examples of f-element, and hence lanthanide, heavy pnictido complexes. To date the only structurally authenticated complexes to feature f-element pnictido bonding are with phosphorus or arsenic, with the pnictogen bridging between



Scheme 14 Synthesis of yttrium phosphido complexes 41–45.

two or more metal centres.<sup>20</sup> There have been no structurally authenticated f-element stibido or bismuthido derivatives to date, therefore they are highly sought-after synthetic targets. For lanthanide metal complexes, only a few clusters were reported containing phosphido ligand.

In 2011, Chen and co-workers reported the first structurally authenticated rare earth phosphido complex,  $[\{\text{Y}(\text{I})\}\{\text{Y}[\mu_3\text{-P}(\text{Dipp})](\mu\text{-I})(\text{THF})_4(\mu_6\text{-P})\{\text{K}(\text{C}_7\text{H}_8)\}]\}$  (**41**) *via* a salt elimination/protonolysis protocol, Scheme 14.<sup>62</sup> The reaction of  $[\text{Y}\{\text{P}(\text{SiMe}_3)(\text{Dipp})\}(\text{I})_2(\text{THF})_3]$  with one equivalent of  $\text{K}[\text{P}(\text{H})\text{Dipp}]$  gave **41** in a 33% yield. The product formed depends upon the reaction solvent; dissolution of **41** in THF results in the loss of half an equivalent of potassium iodide and the conversion of **41** to half an equivalent each of the THF adduct,  $[\{\text{Y}(\text{I})\}\{\text{Y}[\mu_3\text{-P}(\text{Dipp})](\mu\text{-I})(\text{THF})_4(\mu_6\text{-P})\{\text{K}(\text{THF})\}]\}$  (**42**), and the potassium-free complex  $[\{\text{Y}(\text{I})\}\{\text{Y}(\text{I})(\text{THF})_2\}_2\{\text{Y}(\text{THF})_2\}_2(\mu\text{-I})[\mu_3\text{-P}(\text{Dipp})]_4(\mu_5\text{-P})]$  (**43**). The authors attempted to displace the yttrium-bound THF molecules by exposing **41** to potassium salts of  $\text{NCCN}^{\text{iPr}}$  and  $\text{Cp}^*$ , yielding the corresponding phosphido complexes  $[\{\text{Y}(\text{I})\}\{\text{Y}(\text{I})(\text{THF})_2\}_2\{\text{Y}(\text{NCCN}^{\text{iPr}})\}_2(\mu\text{-I})[\mu_3\text{-P}(\text{Dipp})]_4(\mu_5\text{-P})]$  (**44**) and  $[\{\text{Y}(\text{THF})\}\{\text{Y}(\mu\text{-I})(\text{THF})_2\}_2\{\text{Y}(\text{THF})\}\{\text{Y}(\text{Cp}^*)\}_3(\mu\text{-I})[\mu_3\text{-P}(\text{Dipp})]_4(\mu_6\text{-P})\{\text{K}(\text{THF})\}]\}$  (**45**), respectively.<sup>62</sup> The phosphido centres in complexes **41**–**45** are stabilised through interactions with five yttrium ions where bridging phosphinidide ligands


defined *cyclo*-P<sub>3</sub> and -P<sub>4</sub> lanthanide complexes have been reported more recently. For example, in 2018, Roesky and co-workers reported the lanthanide *cyclo*-P<sub>4</sub> complex [Sm(DippForm)<sub>2</sub>]<sub>2</sub>(μ<sup>2</sup>-η<sup>4</sup>:η<sup>4</sup>-P<sub>4</sub>) (**48**, DippForm = {(NDipp)<sub>2</sub>-CH}) by the reduction of P<sub>4</sub> with the divalent precursor [Sm(DippForm)<sub>2</sub>(THF)<sub>2</sub>], Fig. 6.<sup>67</sup> In 2019, Zhang, Zhou and co-workers reported the lanthanide *cyclo*-P<sub>3</sub> complex [K(18C6)(toluene)][{Y(DippDBD)(THF)<sub>2</sub>(P<sub>3</sub>)}] (**49**, DippDBD = *N,N'*-2,6-diisopropylphenyl-1,4-diazabutadiene), with the central P<sub>3</sub><sup>3-</sup> trianion bridging two Y(III) metal ions, prepared by alkyl migration of an organosubstituted *cyclo*-P<sub>4</sub>R<sub>2</sub> precursor followed by encapsulation of K<sup>+</sup> cation with 18-crown-6 reagent, Fig. 6.<sup>68</sup> Very recently, using a redox synthetic strategy, Roesky and co-workers reported another two *cyclo*-P<sub>3</sub> and -P<sub>4</sub> inverse sandwich complexes for lanthanides supported by a xanthene-diamide ligand.<sup>69</sup> Lastly, diphosphorus (P<sub>2</sub>) complexes, which are heavy N<sub>2</sub> analogues, remain elusive for lanthanides because of the synthetic challenges of making and stabilising P<sub>2</sub>.<sup>69</sup>

A few years later, Diaconescu and co-workers reported another rare earth metal P<sub>8</sub> cluster [Sc(NN<sup>fc</sup>)<sub>4</sub>P<sub>8</sub>] (NN<sup>fc</sup> = 1,1-fc(NSi<sup>t</sup>BuMe<sub>2</sub>)<sub>2</sub>, fc = ferrocenylene) prepared by the reaction of the scandium naphthalendiide complex [Sc(NN<sup>fc</sup>)<sub>2</sub>(μ-C<sub>10</sub>H<sub>8</sub>)] with P<sub>4</sub>.<sup>64</sup> The core P<sub>8</sub><sup>4-</sup> ligand in this molecule is very similar to that in **46**. Interestingly, using excess P<sub>4</sub> in the reactions with [Sc(NN<sup>fc</sup>)<sub>2</sub>(μ-C<sub>10</sub>H<sub>8</sub>)] and [Y(NN<sup>fc</sup>)(THF)<sub>2</sub>(μ-C<sub>10</sub>H<sub>8</sub>)] yielded the P<sub>7</sub> complexes [(NNfc)Ln(THF)<sub>*n*</sub>]<sub>3</sub>P<sub>7</sub>] (**47Ln**, Ln = Sc, *n* = 0; Ln = Y, *n* = 1), respectively, Fig. 6. Complexes **47Ln** are the first



© 2024 The Author(s). Published by the Royal Society of Chemistry





Scheme 15 Synthesis of Th phosphide complexes 50–52.

### 5.1 Actinide phosphide complexes

In terms of actinide complexes containing metal–phosphorus single bond interactions, monoanionic charged phosphide ligands can have stronger interactions with actinide metal centres than neutral phosphine ligands, which only form dative bonds to actinide ions.<sup>20</sup> Furthermore, some actinide phosphide complexes are useful precursors to novel linkages such as actinide–metal and -ligand multiple bonds (see below), low-valent U–P bonds,<sup>76</sup> and hydrophosphination catalysts.<sup>77,78</sup>

The first examples of actinide phosphide complexes were reported in 1985 by Ryan and co-workers. The mononuclear complex  $[\text{Th}(\text{Cp}^*)_2(\text{PPh}_2)_2]$  (**50**) was prepared by a salt metathesis reaction between  $[\text{Th}(\text{Cp}^*)_2(\text{Cl})_2]$  and potassium diphenylphosphide, Scheme 15.<sup>79</sup> Importantly, the phosphide ligands are able to coordinate to transition metals as well, supporting actinide and transition metal interactions, making it possible to investigate actinide–metal bonding. Treatment of **50** with  $[\text{Ni}(\text{COD})_2]$  ( $\text{COD} = 1,5\text{-cyclooctadiene}$ ) under a CO atmosphere, or  $[\text{Pt}(\text{COD})_2]$  in the presence of  $\text{PMe}_3$ , led to the formation of the heterobimetallic compounds  $[\text{Th}(\text{Cp}^*)_2(\mu\text{-PPh}_2)_2\text{Ni}(\text{CO})_2]$  (**51**)<sup>79</sup> and  $[\text{Th}(\text{Cp}^*)_2(\mu\text{-PPh}_2)_2\text{Pt}(\text{PMe}_3)_2]$  (**52**)<sup>80</sup> in moderate yields, respectively, Scheme 15. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum for **51** exhibits a signal at 177 ppm, which is shifted downfield from the resonance for **50** found at 143 ppm. In contrast, the  $^{31}\text{P}$  NMR spectrum for **52** shows a doublet at 149.3 ppm, and a triplet at  $-3.3$  ppm. These were attributed to the  $[\text{PPh}_2]^-$  and  $\text{PMe}_3$  ligands, respectively. Both resonances contain coupling to Pt, suggesting a direct interaction between the phosphorus atoms of both ligands and the transition metal. The solid-state structure of **51** revealed the Th–P bond lengths to be 2.869(4) and 2.900(4) Å, which are close to the Th–P distance of 2.866(7) Å found in the mononuclear starting material **50**. The Th–P bond lengths in **52** are unexceptional and are similar to those in **50**. The distance between the thorium and nickel atoms in **51** was found to be 3.206(2) Å, which is longer than the sum of the covalent single bond radii of Th and Ni (2.85 Å). In contrast, in **52** the distance between the thorium and platinum centres was




Fig. 7 Actinide phosphide and phosphanide complexes 53–55An.

found to be 2.984(1) Å, which is similar to the sum of the covalent single bond radii of Th and Pt (2.98 Å). The metal–metal bonding interactions in **51** and **52** were interpreted as a weak, donor–acceptor dative bonds from the low-valent electron-rich Ni(0) and Pt(0) ions to the electron-poor thorium(IV) ions.

As well as actinide phosphide complexes with secondary phosphide ligands, there are also some actinide complexes containing primary phosphide ligands, which could potentially be used to access actinide phosphorus multiple bonds. For example, in 2015 Walensky and co-workers reported the thorium bisphosphide complex  $[\text{Th}(\text{Cp}^*)_2(\text{PHTripp})_2]$  (**53**,  $\text{Tripp} = 2,4,6\text{-}^i\text{Pr}_3\text{C}_6\text{H}_2$ ), prepared by the reaction of  $\text{H}_2\text{PTripp}$  with the thorium dimethyl precursor  $[\text{Th}(\text{Cp}^*)_2(\text{CH}_3)_2]$  in a 2 : 1 stoichiometric ratio *via* an alkane elimination method, Fig. 7.<sup>81</sup> The Th–P bond distances in **53** (2.8754(6) and 2.8830(6) Å) are very close to those in **50–52**, indicative of typical single bond interactions. However, the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of **53** has a resonance at 1.66 ppm, whereas the  $^{31}\text{P}\{^1\text{H}\}$  resonances in **50–52** range from 143 to 177 ppm. Complex **53** is a useful precursor to thorium phosphinidide and phosphinidene complexes (see below). Unlike complexes **50–53**, which contain bulky phosphide substituents, by using bulky triamidoamine ligand frameworks Liddle, Scheer, and co-workers synthesised the first actinide parent phosphanide complexes  $[\text{An}(\text{Tren}^{\text{R}})(\text{PH}_2)]$  ( $\text{Tren}^{\text{R}} = \{\text{N}(\text{CH}_2\text{CH}_2\text{NR})_3\}^{3-}$ ,  $\text{R} = \text{TIPS}$ , triisopropylsilyl,  $\text{An} = \text{Th}$  and  $\text{U}$ , **54An**;  $\text{R} = \text{TCHS}$ , tricyclohexylsilyl,  $\text{An} = \text{Th}$  and  $\text{U}$ , **55An**), Fig. 7.<sup>82–84</sup> Complexes **54An** were prepared by salt metathesis reactions of the corresponding actinide precursors with  $\text{NaPH}_2$ ,<sup>82,84</sup> whilst **55An** were synthesised by protonation of their respective terminal parent phosphinidene complexes due to the lack of suitable actinide precursors for salt metathesis reactions.<sup>84</sup> These four terminal phosphanide complexes are the only known examples to date for any f-element. The solid-state structures of **54An** and **55An** revealed trigonal-bipyramidal metal geometries with the parent phosphide group well-protected by the bulky silyl substituents. The An–P bond distances are 2.982(2), 2.883(2), 3.0360(15), and 2.8725(13) Å for **54Th**, **54U**, **55Th**, and **55U**, respectively, which are slightly longer than the respective sum of the single bond covalent radii for thorium and phosphorus (2.86 Å) and uranium and phosphorus (2.81 Å). Due to the P–H coupling, the  $^{31}\text{P}$  NMR spectra for **54Th**, **54U**, **55Th**, and **55U** exhibit triplet resonances at  $-144.1$ , 595.0,  $-133.0$  and 605.9 ppm, respectively. As expected,

Scheme 16 Synthesis of the U-phosphinidiide complex **56**.

the resonances for uranium complexes are significantly shifted owing to the paramagnetic nature of  $5f^2$  uranium(IV). These phosphanide complexes have proven to be key precursors to access actinide-phosphorus multiple bonds (see below).

## 5.2 Actinide phosphinidiide complexes

Phosphinidiide complexes can be viewed as polynuclear forms of phosphinidene complexes where the phosphinidene ligands are bridged between two or more electropositive metal ions to enhance the stability of the reactive moiety. As discussed above, phosphinidiide complexes are also more common than terminal phosphinidenes for both lanthanides and actinides, but some phosphinidiide complexes can still have metal-phosphorus multiple bonding character.

The first crystallographically characterised actinide phosphinidiide complex was isolated in 1984 by Marks, Day and co-workers, as a parent  $(HP)^{2-}$  unit bridging two uranium(IV) centres. Mixing three equivalents of  $[U(Cp^*)_2(Me)_2]$  with one

equivalent of  $P(OCH_3)_3$  and excess hydrogen gave  $[U(Cp^*)_2(OMe)]_2(\mu-PH)$  (**56**) in yields of 42%, Scheme 16.<sup>85</sup> The authors additionally prepared  $[Th(Cp^*)_2(OMe)]_2(\mu-PH)$  in an analogous manner. Marks and Day postulated that the mechanism for the synthesis of **56** proceeded *via* an actinide hydride. To confirm this, additional reactions were conducted in the absence of hydrogen or utilising  $[An(Cp^*)_2(H)_2]_2$  as starting materials. The former showed no detectable reaction, whilst the latter gave **56**. Complex **56** exhibits two U–P distances of 2.743(1) Å and a U–P–U angle of 157.7(2)°. The IR spectrum of **56** confirmed the presence of the parent phosphinidiide, with a PH stretching mode at 2193  $cm^{-1}$  ( $\nu_{P-H}/\nu_{P-D} = 1.39$ ). Liddle, Scheer, and co-workers also reported diactinide parent phosphinidiide complexes supported by triamidoamine ligands,<sup>86</sup> which will be discussed in the phosphido section below for comparison purposes.

In 2015 Walensky and co-workers utilised the bis- $Cp^*$  framework to synthesise the bridging thorium phosphinidiide complex  $[Th(Cp^*)_2]_2[\mu-P\{(2,6-CH_2CHCH_3)_2-4-^iPrC_6H_2\}]$  (**57**, 63%) by the protonolysis of two equivalents of  $[Th(Cp^*)_2(Me)_2]$  with one equivalent of  $H_2PTripp$  at 90–95 °C, Scheme 17.<sup>81</sup> Complex **57** exhibits Th–P bond distances of 2.8083(9) and 2.8186(9) Å, which are marginally shorter than those observed for the thorium phosphide complex **53** (2.8755(6) and 2.8829(7) Å) due to the bridging mode of the phosphinidiide.

Walensky and co-workers also derivatised **53** to synthesise the bridging phosphide/phosphinidiide thorium complex  $[Th(Cp^*)_2(\mu-PTripp)(\mu-PHTripp)(K)]_2$  (**58**), Scheme 17,<sup>87</sup> which paved the way to a number of derivatives.<sup>88–92</sup> The reaction of **53** with one equivalent of  $KN(SiMe_3)_2$  gave **58** in a yield of 64% *via* the elimination of one equivalent of  $HN(SiMe_3)_2$ . It was found that if the deprotonation reaction was conducted in the presence of 2.2.2-cryptand the monomeric phosphide/phosphinidene complex  $[K(2.2.2-cryptand)][Th(Cp^*)_2(PTripp)(PHTripp)]$  (**59**) was obtained.<sup>87</sup> Complex **58** exhibits Th–P distances of 2.6957(10) Å, whilst **59** demonstrates a shorter Th–P bond length of 2.6024(9) Å, which is a result of the encapsulation of the potassium ion allowing more electron

Scheme 17 Synthesis of Th-phosphinidiide complexes **57** and **58** and phosphinidene **59**.Scheme 18 Synthesis of thorium phosphinidiide and phosphinidene complexes **60** and **61**.

density to be donated from the phosphorous atom to the thorium centre.

Recently, Walter, Ding, Zi and co-workers reported the salt metathesis/protonolysis reaction of one equivalent each of  $[\text{Th}(\text{Cp}^{\text{tt}})_2(\text{Me})(\text{Cl})]$  ( $\text{Cp}^{\text{tt}} = 1,3\text{-}^t\text{Bu}_2\text{-C}_5\text{H}_3$ ) and  $\text{KPH}(\text{Mes}^*)$  to synthesise the dinuclear phosphinidide complex  $[\{\text{Th}(\text{Cp}^{\text{tt}})_2(\text{P}(\text{Mes}^*)\text{Cl})\}_2]$  (**60**) in 78% yield, Scheme 18, with elimination of one equivalent each of methane and potassium chloride.<sup>93</sup> The authors reported that **60** could be converted to the mononuclear terminal phosphinidene complex  $[\text{Th}(\text{Cp}^{\text{tt}})_2(\text{P}(\text{Mes}^*)(\mu\text{-Cl})\{\text{K}(\text{18C6})\})]$  (**61**, 92%) by the addition of 18-crown-6.<sup>93</sup> The solid-state structure of **60** and **61** revealed that **60** exhibits a slightly shorter Th–P distance of 2.560(1) Å and a more acute Th–P–C<sup>Ar</sup> angle of 162.6(2)° than found in **61** (2.582(1) Å; Th–P–C<sup>Ar</sup> angles = 171.3(1)°) which is most likely a result of the removal of the interaction between phosphorus and the potassium cation from **60** to **61**. The authors reported that all four complexes **58–61** exhibit  $^{31}\text{P}\{\text{H}\}$  NMR resonances in the range 108.8 to 177.85 ppm.

### 5.3 Actinide phosphinidene complexes


Although uranium imido chemistry is well-developed,<sup>18</sup> actinide complexes containing heavier pnictogen An = PnR (An = actinide; Pn = P, As, Sb, Bi) multiple bonds are scarce.<sup>20</sup> To date, there are few complexes containing An=PnR double bonds outside of cryogenic matrix isolation conditions, but only with P or As ligands, not for Sb and Bi ligands. Also, there is no example of a terminal actinide heavy pnictido An≡Pn triple bond isolated under ambient conditions. Even for d-block metals there are relatively few structurally characterised terminal metal pnictinidene complexes (~50), far fewer than their imido counterparts (>3000).<sup>94</sup> Actinide pnictinidene complexes normally require kinetic stabilisation by sterically demanding ancillary ligands, coupled with bulky pnictinidene substituents (*cf.* **59** and **61**).

In 1996, Burns and co-workers reported the first example of a terminal uranium phosphinidene complex utilising the sterically demanding Cp\* supporting ligand. The salt elimination/protonolysis reaction of one equivalent each of

$[\text{U}(\text{Cp}^*)_2(\text{Me})(\text{Cl})]$  and  $\text{KPH}(\text{Mes}^*)$  in the presence of  $\text{OPMe}_3$  gave  $[\text{U}(\text{Cp}^*)_2(\text{P}(\text{Mes}^*)(\text{OPMe}_3))]$  (**62**), in yields of up to 62%, Scheme 19 (top).<sup>95</sup> Complex **62** exhibits a U–P distance of 2.562(3) Å, which is shorter than the bridging phosphinidide U–P distance in **56** or, for example, the U–P distance in the phosphide complex  $[\text{U}(\text{Cp}^*)_2\{\text{P}(\text{SiMe}_3)_2\}(\text{Cl})]$  (2.789(4) Å).<sup>96</sup> The authors postulated that the U–P–C<sup>Ar</sup> angle of 143.7(3)° in **62** was due to crystal packing forces, or a result of the combination of heavy main group elements generally adopting bent geometries in addition to the preferred linear geometry required to minimise P–U  $\pi$ -overlap.

In 2018, using sterically demanding cyclopentadienyl ligands, Walter, Ding, Zi and co-workers reported a base-free terminal thorium phosphinidene  $[\text{Th}(\text{Cp}^{\text{ttt}})_2(\text{P}(\text{Mes}^*))]$  (**63**,  $\text{Cp}^{\text{ttt}} = 1,2,4\text{-}^t\text{Bu}_3\text{-C}_5\text{H}_2$ ), Scheme 19 (bottom).<sup>97</sup> Complex **63** was prepared by a similar salt metathesis/protonolysis reaction that was employed to prepare **62**; reaction of one equivalent each of  $[\text{Th}(\text{Cp}^{\text{ttt}})_2(\text{Me})(\text{I})]$  and  $\text{KPH}(\text{Mes}^*)$  produced **63** in high yield (80%), with elimination of one equivalent each of methane and potassium iodide. The molecular structure of **63** further confirmed a base-free terminal phosphinidene species with a short Th–P bond distance (2.536(2) Å). The  $^{31}\text{P}\{\text{H}\}$  NMR spectrum of **63** shows one resonance at 145.7 ppm, which is close to those for **59** and **61**. Calculations probing the Th=P bonding of **63** and the theoretical monomeric model of **63** suggest more covalency in this linkage than related imido complexes. The authors additionally treated the thorium complexes **63** with a wide range of unsaturated substrates to probe the reactivity of Th=P double bond. In general, the actinide metallocene phosphinidene motif has proven practicable to effect with a range of cyclopentadienyl substituents.<sup>98–105</sup>

From the above examples, it is clear that sterically demanding cyclopentadienyl ligands have proven effective at stabilising actinide phosphinidide and phosphinidene complexes, but sterically demanding triamidoamine ligands



Scheme 19 Synthesis of terminal uranium and thorium phosphinidene complexes **62** (top) and **63** (bottom).



Scheme 20 Synthesis of uranium phosphinidene and phosphinidide complexes **64M** and **66**.


have in parallel extended this chemistry into a new regime. Following the successful isolation of terminal uranium nitrides supported by the  $\text{Tren}^{\text{TIPS}}$  ligand ( $\text{Tren}^{\text{TIPS}} = \{\text{N}(\text{CH}_2\text{CH}_2\text{NSi}^{\text{iPr}}\text{Pr}_3)_3\}^{3-}$ ) in 2012 and 2013,<sup>106,107</sup> Liddle, Scheer, and co-workers utilised this bulky ligand framework to stabilise the first terminal uranium parent phosphinidene in 2014, Scheme 20. The reaction of **54U** with one equivalent of benzyl potassium and two equivalents of benzo-15-crown-5 ether (B15C5) yielded the terminal parent uranium(IV) phosphinidene complex  $[\text{K}(\text{B15C5})_2][\text{U}(\text{Tren}^{\text{TIPS}})(\text{PH})]$  (**64K**).<sup>82</sup> The synthesis of the sodium analogue  $[\text{Na}(\text{12C4})_2][\text{U}(\text{Tren}^{\text{TIPS}})(\text{PH})]$  (**64Na**) was achieved by reaction of the uranium cyclometallate complex  $[\text{U}\{\text{N}(\text{CH}_2\text{CH}_2\text{NSi}^{\text{iPr}}\text{Pr}_3)_2(\text{CH}_2\text{CH}_2\text{NSi}^{\text{iPr}}\text{Pr}_2\text{C}(\text{H})\text{MeCH}_2)\}]$  (**65U**) with one equivalent of  $\text{NaPH}_2$  and two equivalents of 12-crown-4 ether (12C4).<sup>86</sup> The authors reported that the treatment of **54U** with one equivalent each of  $\text{KCH}_2\text{Ph}$  and 2.2.2-cryptand gave the contact ion pair phosphinidide complex  $\{[\text{U}(\text{Tren}^{\text{TIPS}})(\mu\text{-PH})]\{\text{K}(2.2.2\text{-cryptand})\}\}$  (**66**).<sup>82</sup> The solid-state structures revealed that the terminal phosphinidene complexes **64M** and phosphinidide **66** exhibit U–P distances ranging from 2.613(2) to 2.685(2) Å, which are shorter than that in the parent phosphide precursor **54U** (2.883(2) Å). The authors reported that the IR spectrum of **64K** exhibits a P–H stretch of  $2360\text{ cm}^{-1}$ . The U–P bond lengths in **64M** are longer than those observed for **62** (2.562(3) Å) lying between the sum of the covalent single and double bond radii for uranium and phosphorus (2.81 Å and 2.36 Å, respectively); this reflects the sterically demanding nature of  $\text{Tren}^{\text{TIPS}}$  and indicates polarised covalent  $\text{U}=\text{P}$  interactions that is confirmed by DFT calculations.

In 2016, Liddle, Scheer, and co-workers utilised the  $\text{Tren}^{\text{TIPS}}$  ligand framework to stabilise terminal thorium parent phosphinidene analogues. The two methodologies used to synthesise the uranium complexes **64M** were adapted to prepare the terminal phosphinidene thorium complex  $[\text{Th}(\text{Tren}^{\text{TIPS}})(\text{PH})]$  (**67**), Scheme 21.<sup>83</sup> Complex **67** could be synthesised either through deprotonation of **54Th** with one equivalent of  $\text{NaCH}_2\text{Ph}$  and two equivalents of 12C4, or by the

reaction of the thorium cyclometallate complex  $[\text{Th}\{\text{N}(\text{CH}_2\text{CH}_2\text{NSi}^{\text{iPr}}\text{Pr}_3)_2(\text{CH}_2\text{CH}_2\text{NSi}^{\text{iPr}}\text{Pr}_2\text{C}(\text{H})\text{MeCH}_2)\}]$  (**65Th**) with one equivalent of  $\text{NaPH}_2$  and two equivalents of 12C4 in yields of up to 38%. Complex **67** is isostructural with the uranium phosphinidene complex **64Na**.

The Th–P bond length of 2.758(2) Å in **67** is *ca.* 0.22 Å shorter than that of the Th– $\text{PH}_2$  bond in **54Th** (2.982(2) Å), but is longer than the  $\text{U}=\text{P}$  double distance of 2.613(2) Å in **64K**, suggesting a more polarised double bond interaction for  $\text{Th}=\text{PH}$  linkage. This is in accord with a smaller Th–P Mayer bond order of 1.67 in **67** than that of 1.92 in **64K**. The  $\text{Th}=\text{P}-\text{H}$  angle of  $67.45(8)^\circ$  in **67** indicates an ‘agostic-type’ interaction between the metal ion and the electron density of the P–H bond, whereas this interaction was not observed in **64K**, which has a  $\text{U}=\text{P}-\text{H}$  angle of  $118.8(9)^\circ$ . The  $^{31}\text{P}$  NMR spectrum for **67** has a doublet resonance at 198.8 ppm due to P–H coupling, further confirming the presence of  $[\text{PH}]^{2-}$  group at thorium. By contrast, because of the strong paramagnetic shielding from the uranium(IV) centre, no resonance was observed in the  $^{31}\text{P}$  NMR spectra for **64M**.

Protonation of reactive actinide–carbon bonds has proven to be an effective strategy for constructing actinide–pnictogen multiple bonds. In 2022, Liddle, Scheer, and co-workers developed a bulky  $\text{Tren}^{\text{TCHS}}$  ligand ( $\{\text{N}(\text{NCH}_2\text{CH}_2\text{NSiCy}_3)_3\}^{3-}$ ) to prepare two new terminal actinide phosphinidene complexes, Scheme 22. Reaction of the cyclometallate actinide complexes  $[\text{An}\{\text{N}(\text{CH}_2\text{CH}_2\text{NSiCy}_3)_2(\text{CH}_2\text{CH}_2\text{NSiCy}_2[\text{CHCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}])\}]$  (**68An**, An = Th, U) with  $\text{NaPH}_2$  in the presence of 2.2.2-cryptand in THF afforded  $[\text{Na}(2.2.2\text{-cryptand})][\text{An}(\text{Tren}^{\text{TCHS}})(\text{PH})]$  (**69An**, An = U, Th).<sup>84</sup> The molecular structures of **69An** confirmed the presence of terminal phosphinidenes, with the  $(\text{HP})^{2-}$  ligand well-protected by the super bulky tricyclohexylsilyl groups. The An–P bond distances of 2.7237(9) and 2.6381(12) Å for **69Th** and **69U**, respectively are statistically invariant to those in **67** and **64M**, indicating multiple bonding interactions in these  $\text{An}=\text{PH}$  linkages with polarised covalent interactions supported by DFT studies. The ‘agostic-type’ interaction between the An ion and the



Scheme 21 Synthesis of thorium phosphinidene complex **67**.



Scheme 22 Synthesis of thorium and uranium phosphinidene complexes **69An** and phosphanide complexes **55An**.





phosphinidene ligand was observed in both structures, with An–P–H angles of 65.83(17)° and 65.23(13)°, respectively. In addition, absorptions corresponding to P–H stretches at 2072 and 2070 cm<sup>−1</sup> for **69Th** and **69U**, respectively, were observed in their ATR-IR spectra. The An=P vibrations are also observed in the Raman spectra of **69Th** and **69U** at 306 and 296 cm<sup>−1</sup>, respectively. The <sup>31</sup>P NMR spectrum for **69U** exhibits a broad resonance at 2629 ppm due to the paramagnetic uranium(IV) centre, while this is not observed in **64M**. Similar to **67**, **69Th** exhibits a doublet resonance at 266.2 ppm in its <sup>31</sup>P NMR spectrum due to P–H coupling.

Reflecting the basic and nucleophilic nature of **69An**, Liddle, Scheer, and co-authors additionally found that treatment of **69An** with [HNEt<sub>3</sub>][BPh<sub>4</sub>], as a proton source, in THF resulted in the isolation of the phosphanide complexes **55An** in good yields, Scheme 22,<sup>84</sup> which have similar bond metrics to **54An**.<sup>82</sup> Alternatively, **55U** could be prepared by oxidation of **69U** with AgBPh<sub>4</sub> in benzene. The formation of **55U** in these oxidation reactions may involve a transient U(V)=PH species (or valence isomer, e.g. U(IV)=P'H), which then abstracts a proton (or H<sup>+</sup>) in the reaction mixture due to the HSAB mismatch of U and P. This reactivity contrasts to the disproportionation observed for Tren<sup>TIPS</sup>-supported U(V)=NH chemistry, which produces U(IV)–NH<sub>2</sub> amide and U(VI)≡N nitride products.<sup>108</sup> These reactivity outcomes reflect the periodic differences between nitrogen and phosphorus, and that when the latter is paired with electro-positive metals P-based electrons can become involved in redox reactions as found in the reactivity of **16**.

#### 5.4 Actinide phosphido complexes

Actinide phosphido complexes remain exceedingly rare. There are only a few bridging dinuclear complexes isolated in recent years with no examples of terminal actinide heavy pnictido An≡Pn triple bond isolated under ambient conditions to date.<sup>20</sup>

In 2016, Liddle, Scheer, and co-workers reported the first example of an actinide–phosphido complex [Na(12C4)<sub>2</sub>]



Scheme 23 Synthesis of bridging thorium phosphinidide and phosphido complex **70** and **71Th**.

[{Th(Tren<sup>TIPS</sup>)<sub>2</sub>(μ-P)] (**70**), where the phosphido ligand bridges two thorium centres, Scheme 23.<sup>83</sup> Complex **70** was also the first such f-element phosphido species, and was prepared by the reaction of two equivalents of **65Th** with NaPH<sub>2</sub> in the presence of two equivalents of 12C4 in up to 57% yield. Alternatively, **70** can be synthesised by the stoichiometric reaction of **67** with **65Th**. The authors additionally reported the synthesis of the bridging phosphinidide complex [{Th(Tren<sup>TIPS</sup>)<sub>2</sub>(μ-PH)] (**71Th**) in a 40% yield, either by treatment of **54Th** with one equivalent of **65Th**, or the reaction of two equivalents each of **65Th** with NaPH<sub>2</sub>, eliminating one equivalent of 'Na<sub>2</sub>PH'. However, attempts to prepare **70** by deprotonation of **71Th** were unsuccessful. Complex **70** has a symmetrical ThPTh core with Th–P bond distances of 2.740(2) and 2.735(2) Å, which are shorter than those in **71Th** (2.898(2) Å) but compares well to that of the terminal phosphinidene **67** (2.7584(18) Å), and lies between the sum of the covalent single and double bond radii for thorium and phosphorus (2.86 and 2.45 Å, respectively), suggesting multiple bonding interactions in the ThPTh linkage. The <sup>31</sup>P NMR spectra for **71Th** and **70** exhibit doublet and singlet resonances at 145.7, and 553.5 ppm, confirming the presence of (HP)<sup>2−</sup> and P<sup>3−</sup>, respectively.

Subsequently, in 2017 Liddle, Scheer, and co-authors reported that addition of one equivalent of KCH<sub>2</sub>Ph to a mixture of **54U** and [U(Tren<sup>TIPS</sup>)(THF)][BPh<sub>4</sub>] gave the bridging diuranium phosphinidide complex [{U(Tren<sup>TIPS</sup>)<sub>2</sub>(μ-PH)] (**71U**)<sup>86</sup> that is isostructural to **71Th**, Scheme 24. Importantly, the authors found that deprotonation of **71U** with benzyl potassium in the presence of two equivalents of B15C5 produced the diuranium(IV/IV) phosphido compound [K(B15C5)<sub>2</sub>][{U(Tren<sup>TIPS</sup>)<sub>2</sub>(μ-P)] (**72K**), Scheme 24;<sup>86</sup> recall that this method did not work for **71Th**.<sup>84</sup> Complex **72K** was found to rapidly decompose, which was attributed to steric overload, and so,



Scheme 24 Synthesis of bridging uranium phosphinidide and phosphido complexes **71U**–**73Na**.

seeking a more stable combination, the slightly less bulky and asymmetric sodium analogue of **72K**,  $[\text{Na}(\text{12C}_4)_2][\{\text{U}(\text{Tren}^{\text{TIPS}})\}\{\text{U}(\text{Tren}^{\text{DMBS}})\}(\mu\text{-P})]$  (**73Na**;  $\text{Tren}^{\text{DMBS}} = \{\text{N}(\text{CH}_2\text{CH}_2\text{NSiMe}_2\text{-}^t\text{Bu})_3\}^{3-}$ ) was prepared<sup>86</sup> by the reaction of one equivalent each of **64Na** with the cyclometallate complex  $[\text{U}\{\text{N}(\text{CH}_2\text{CH}_2\text{-NSiMe}_2^t\text{Bu})_2(\text{CH}_2\text{CH}_2\text{NSi}(\text{Me})(\text{CH}_2)(^t\text{Bu}))\}]$ , which is a less sterically demanding Tren ligand compared to  $\text{Tren}^{\text{TIPS}}$ .

Unlike **70** which is stable, **72K** and **73Na** are to various extents unstable in solution, in line with the paucity of such phosphido species, which resulted in both complexes being isolated in relatively low yields of 29% and <5% for **73Na** and **72K**, respectively. Interestingly, **73Na** exhibits an asymmetric UPU core, with two U–P bond distances of 2.657(2) and 2.713(2) Å, and the shorter U–P distance is the  $\text{Tren}^{\text{TIPS}}$  ligated unit. In contrast, **72K** has a symmetric core with statistically indistinguishable U–P bond lengths of 2.653(4) and 2.665(4) Å, which are shorter than those in the phosphinidide **71U** (2.8187(12) and 2.8110(12) Å). Calculations on **72K** and **73Na** suggested that the U–P bond are less covalent than the terminal phosphinidene species **64M**. This is reflected by smaller U–P Mayer bond orders of 1.41/1.43 and 1.44/1.66 for **72K** and **73Na**, respectively, when compared to **[64K]<sup>−</sup>** (1.92). DFT calculations also suggest that the Th–P bonds in **70** are more polarised and ionic than the U–P bonds in **72K** and **73Na**, as evidenced by smaller Th–P Mayer bond orders of 1.26 and 1.28 for **[70]<sup>−</sup>**. However, these An–P Mayer bond orders are nearly twice that of the single U–N<sub>amide</sub> bonds (~0.71), suggesting that these An–P bonds are polarised multiple bond interactions.

### 5.5 Actinide 2-phosphaethynolate complexes

Transition metal 2-phosphaethynolate (OCP)<sup>−</sup> complexes have proven to be useful precursors to prepare phosphido complexes *via* redox or photolysis process.<sup>31</sup> Recently, Liddle, Scheer, and co-workers reported the two new actinide–OCP complexes

$[\text{An}(\text{Tren}^{\text{TIPS}})(\text{OCP})]$  (**74An**, An = Th and U) and their reduction chemistry.<sup>109,110</sup> Reduction of **74U** with  $\text{KC}_8$  in the presence of 2.2.2-cryptand gives  $[\text{K}(2.2.2\text{-cryptand})][\{\text{U}(\text{Tren}^{\text{TIPS}})\}_2\{\mu\text{-}\eta^2(\text{OP})\text{:}\eta^2(\text{CP})\text{-OCP}\}]$  (**75**), Scheme 25.<sup>109</sup> Although there was no phosphido species isolated from this reduction, the coordination mode of this trapped OCP-ligand is unique, and derives from a novel highly reduced and bent carbene-like form of this ligand with a bridging P-centre and the most acute P–C–O angle of ~127° in any complex to date. The mixed valence diuranium(III/IV) formulation is supported by the characterisation data and DFT calculations, where back-bonding from uranium gives a highly reduced form of the OCP unit that is perhaps best described as a uranium stabilised  $(\text{OCP})^{2-}$  radical dianion. In contrast, reduction of **74Th** with  $\text{KC}_8$  or  $\text{CsC}_8$  produced the phosphinidide C–H bond activation product  $[\{\text{Th}(\text{Tren}^{\text{TIPS}})\}\text{Th}\{\text{N}(\text{CH}_2\text{CH}_2\text{NSiPr}_3)_2[\text{CH}_2\text{CH}_2\text{SiPr}_2\text{CH}(\text{Me})\text{CH}_2\text{C}(\text{O})\mu\text{-P}]\}]$  and the oxo complex  $[\{\text{Th}(\text{Tren}^{\text{TIPS}})(\mu\text{-OCs})\}_2]$ . Surprisingly, using  $\text{RbC}_8$  for the reduction afforded a hexathorium complex,  $[\{\text{Th}(\text{Tren}^{\text{TIPS}})\}_6(\mu\text{-OC}_2\text{P}_3)_2(\mu\text{-OC}_2\text{P}_3\text{H}_2)\text{Rb}_4]$  (**76**), which contains four five-membered  $[\text{C}_2\text{P}_3]$  phosphorus heterocycles *via* a  $[2 + 2 + 1]$  cycloaddition, Scheme 25.<sup>110</sup> In addition, this hexathorium complex can be converted to the oxo complex  $[\{\text{Th}(\text{Tren}^{\text{TIPS}})(\mu\text{-ORb})\}_2]$  and the known cyclometallated complex **65Th** at 80 °C, *via* an otherwise hidden example of reductive cycloaddition reactivity in the chemistry of 2-phosphaethynolate. From the above examples it can be seen that the reduction chemistry of 2-phosphaethynolate for actinides can be quite complicated.

### 5.6 Actinide polyphosphorus complexes

Similar to most polyphosphorus complexes, actinide polyphosphorus complexes are most often prepared from  $\text{P}_4$  and low-valent actinide compounds.<sup>20</sup> In 1991, Scherer and co-workers reported the first examples of actinide polyphosphorus complexes,  $[\{\text{Th}(\text{Cp}^{\text{tt}})_2\}_2(\mu^2\text{-}\eta^4\text{-P}_6)]$  (**77**) and  $[\{\text{Th}(\text{Cp}^{\text{tt}})_2\}(\mu^2\text{-}\eta^3\text{-P}_3)\{\text{Th}(\text{Cp}^{\text{tt}})_2\text{Cl}\}]$  (**78**), prepared by treatment of  $[\text{Th}(\text{Cp}^{\text{tt}})_2(\eta^4\text{-C}_4\text{H}_6)]$  with  $\text{P}_4$  at 100 °C in the absence, or presence of,  $\text{MgCl}_2$ , respectively, Fig. 8.<sup>111</sup> The solid-state structures of **77** and **78** revealed distinct structural differences: a bicyclic  $\text{P}_6^{4-}$  ligand is bridged by two thorium(IV) centres in **77**, whilst *cyclo*- $\text{P}_3^{3-}$  is bridged in **78**, with Th–P bond distances between 2.840(7) and 2.921(7) Å. The <sup>31</sup>P NMR



Scheme 25 Reduction of actinide–OCP complexes **74An** to produce actinide complexes **75** and **76**.




Fig. 8 Core structures for reported actinide polyphosphorus complexes **78–82** prepared from  $\text{P}_4$ .

spectrum for **77** exhibits resonances at 125.3, 18.4 and  $-41.9$  ppm. In contrast, the  $^{31}\text{P}$  NMR spectrum for **78** shows temperature-dependent features; at room temperature, there is one resonance at  $-75.7$  ppm, however, at 193 K, two resonances at  $-69.7$  and  $-94.5$  ppm were observed, indicating two unique phosphorus environments at low temperature.

This area progressed little over 20 years until in 2011 Cloke, Green and co-workers isolated the *cyclo*- $\text{P}_4$  uranium complex  $[\{\text{U}(\text{Cp}^*)(\text{C}_8\text{H}_6(\text{Si}^i\text{Pr}_3)_2)_2(\mu^2-\eta^4-\text{P}_4)\}]$  (**79**), Fig. 8, from the reaction of the U(III) complex  $[\text{U}(\text{Cp}^*)(\text{C}_8\text{H}_6(\text{Si}^i\text{Pr}_3)_2)(\text{THF})]$  with half an equivalent of  $\text{P}_4$ .<sup>112</sup> The molecular structure of **79** revealed a diuranium structure where each of the uranium metal centres interacts with the *cyclo*- $\text{P}_4$  ligand in an  $\eta^2$ -fashion. The  $\text{P}_4$  ligand forms a chair-like structure with the two uranium atoms, with U–P bond distances of 2.9763(12) and 2.9773(12) Å, respectively, which are typical single bond character. In addition, the P–P bond distances of 2.152(2) and 2.149(2) Å within the bridging *cyclo*- $\text{P}_4$  unit suggest the dianionic charge of the  $\text{P}_4$  ligand. There is only a single resonance at 718 ppm in the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of **79**. In 2016, Mills and co-workers reported another actinide *cyclo*- $\text{P}_4$  complex,  $[\{\text{Th}(\text{Cp}^*)_3\}_2(\mu^2-\eta^2-\text{P}_4)]$  (**80**), Fig. 8, by reacting the Th(III) complex  $[\text{Th}(\text{Cp}^*)_3]$  with  $\text{P}_4$ .<sup>113</sup> The solid-state structure of **80** revealed a planar  $(\text{P}_4)^{2-}$  ligand bridging two thorium centres *via* two individual  $\eta^1$  bonding interactions. The Th–P bond distances of 2.919(4) Å indicates Th–P single bond interactions and the short P–P bond distances of only 2.051(9) Å are 0.1 Å shorter than those in **79**, suggesting that the bonding within the *cyclo*- $\text{P}_4$  ligand is closer to double bond character, which might be due to the  $\eta^1$  coordination mode and the lack of Th–P  $\pi$ -bonding. The  $^1\text{H}$  NMR spectrum was diagnostic of a diamagnetic complex, reflecting that the reduction of the  $\text{P}_4$  arises from the oxidation of the Th(III) precursor to two Th(IV) centres; the  $^{31}\text{P}$  NMR spectrum of **80** suggests the presence of two different phosphorus environments with two triplet signals at 227.59 and 328.86 ppm, respectively, with a P–P coupling constant of *ca.* 400 Hz.

In 2013, Liddle and co-workers reported that reaction of the diuranium(v) arene complex  $[\{\text{U}(\text{Ts}^{\text{tol}})\}_2(\mu_2-\eta^6:\eta^6-\text{C}_6\text{H}_5\text{CH}_3)]$  ( $\text{Ts}^{\text{tol}} = \text{HC}(\text{SiMe}_2\text{NAr})$ ,  $\text{Ar} = 4\text{-MeC}_6\text{H}_4$ ) with one equivalent of  $\text{P}_4$  resulted in the formation and isolation of the triuranium Zintl cluster  $[\{\text{U}(\text{Ts}^{\text{tol}})\}_3(\mu^3-\eta^3-\text{P}_7)]$  (**81**),<sup>114</sup> Fig. 8. The solid-state structure of **81** revealed a  $\text{P}_7^{3-}$  trianion cluster capped on three of its faces by three  $\text{Ts}^{\text{tol}}$ -uranium cation fragments. The U–P bond lengths of **81** are in the range of 2.949(2)–3.031(2) Å, which are comparable to the U–P cluster distances discussed above. Interestingly, **81** reacts with a variety of halide reagents, such as  $\text{Me}_3\text{-SiCl}$ ,  $\text{LiCl}$ ,  $\text{MeI}$ , and  $\text{PhI}$ , to afford  $\text{P}_7\text{R}_3$  products with subsequent reduction of the U component closing the synthetic cycle.


Extending the reactivity of  $\text{P}_4$  with other U(III) complexes, Liddle and co-workers reported that treatment of  $[\text{U}(\text{Tren}^{\text{TIPS}})]$  with 0.25 equivalents of  $\text{P}_4$  reproducibly gives the actinide inverted sandwich *cyclo*- $\text{P}_5$  complex  $[\{\text{U}(\text{Tren}^{\text{TIPS}})\}_2(\mu-\eta^5:\eta^5\text{-cyclo-P}_5)]$  (**82**),<sup>115</sup> Fig. 8. All previous examples of *cyclo*- $\text{P}_5$  complexes were stabilised by transition metals, but the isolation of **82** indicates that *cyclo*- $\text{P}_5$  can also be stabilised by hard actinide ions. Moreover, the characterisation data are consistent with **82** being a diuranium(IV) complex, and thus the *cyclo*-



Scheme 26 Synthesis of uranium diphosphorus complex **83**.

$\text{P}_5$  unit in **82** is formally a radical dianion rather than the usual monoanion form. The molecular structure of **82** revealed quite long U–P bond distances spanning the range 3.250(6)–3.335(6) Å, which are longer than the sum of the single bond covalent radii of U and P (2.81 Å), perhaps owing to the sterically demanding nature of  $\text{Tren}^{\text{TIPS}}$  ligands combined with the bridging  $\eta^5$ -bound (per U) nature of the *cyclo*- $\text{P}_5$  unit in **82**. DFT studies on **82** indicates the principal bonding in the  $\text{U}(\text{P}_5)\text{U}$  unit is polarised  $\delta$ -bonding, whilst the isolobal cyclopentadienyl ligand normally interacts with metals *via*  $\sigma$ - and  $\pi$ -bonding interactions with minimal  $\delta$ -interaction. In a related study, Zhu, Maron, and co-workers investigated the reactivity of  $\text{P}_4$  with U(III) supported by a tertiary phosphine-appended Tren ligand, resulting in a diuranium product containing a  $\text{P}_4$  chain.<sup>116</sup>

In 2021, Liddle and co-workers reported the synthesis and structure of a side-on bound diphosphorus U(IV) complex,  $[\{\text{U}(\text{Tren}^{\text{TIPS}})\}_2(\mu-\eta^2:\eta^2-\text{P}_2)]$  (**83**), Scheme 26, by reacting the  $7\lambda^3$ -(dimethylamino)phosphadibenzonorborene P-atom transfer reagent (anthracene- $\text{PNMe}_2$ ) with  $[\text{U}(\text{Tren}^{\text{TIPS}})]$ .<sup>117</sup> The by-product,  $[\text{U}(\text{Tren}^{\text{TIPS}})(\text{NMe}_2)]$ , was isolated from the reaction mixture by fractional crystallisation, accounting for the fate of the  $\text{NMe}_2$  unit. Complex **83** is the first diphosphorus complex for any f-element complex, coming after the first f-element dinitrogen complex in 1988.<sup>118</sup> The molecular structure of **83** revealed P–P bond distances of 2.036(2) Å, indicative of  $\text{P}=\text{P}$  double bond character and hence a dianionic  $\text{P}_2^{2-}$ , which is consistent with a diuranium(IV) formulation confirmed by the characterisation data. The U–P distances of 2.9441(12) and 2.9446(12) Å are longer than the sum of the single bond covalent radii of U and P (2.81 Å), reflecting the side-on bridging mode of the  $\text{P}_2$  unit in **83**. Computational results indicated that within the  $\text{UP}_2\text{U}$  motif the in-plane U–P  $\pi$ -bonding dominates with a very weak  $\delta$ -interaction. It was subsequently found that oxidation of **64M** with  $\text{AgBPh}_4$  also produces **83** (along with **54U**), which suggests the formation of a transient  $\text{U}(\text{v})=\text{PH}$  linkage that disproportionates to **54U** and  $\text{U}(\text{vi})\equiv\text{P}$ , the latter of which could dimerise to give the



more stable P–P coupled **83**. In addition, a preliminary reactivity study demonstrated that **83** can be converted to uranium *cyclo*-P<sub>3</sub> complexes [M(arene)<sub>4</sub>][{U(Tren<sup>TIPS</sup>)<sub>2</sub>(μ-η<sup>3</sup>:η<sup>3</sup>-P<sub>3</sub>)}] (M = K, Rb, Cs; arene = toluene or benzene); these reactions are low yielding, implying the presence of reactive phosphido intermediates.<sup>107</sup> DFT calculations indicate that uranium moves from π-bonding to P<sub>2</sub> and *cyclo*-P<sub>3</sub> to δ-bonding with *cyclo*-P<sub>5</sub>, highlighting the flexibility of the chemical bonding of uranium.

## 6. Lanthanide heavier pnictogen complexes

Upon descending group 15, the number of f-element heavier pnictogen bonds falls away rapidly, and lanthanide complexes containing heavier pnictogen ligands from As to Bi are much rarer than P analogues. Furthermore, such complexes tend to form multi-centre clusters with bridging pnictogen ligands. Thus, well-defined mononuclear species are sparse. To the best of our knowledge, there are no lanthanide complexes isolated to date with multiple bonding to As, Sb, or Bi.

### 6.1 Lanthanide arsenic complexes

The first crystallographically characterised complex containing a lanthanide–arsenic bond was reported in 1988 by Schumann and co-workers. It was found that [Lu(Cp)<sub>2</sub>(μ-CH<sub>3</sub>)<sub>2</sub>{Li(TMEDA)}] (TMEDA = *N,N,N',N'*-tetramethylethylenediamine) reacted with diphenylarsine in benzene to afford the lanthanide–arsenide complex [Lu(Cp)<sub>2</sub>(μ-AsPh<sub>2</sub>)<sub>2</sub>{Li(TMEDA)}] (**84**), Fig. 9, *via* methane elimination.<sup>119</sup> The solid-state structure of **84** revealed the Lu–As bond distances are 2.896(2) and 2.870(2) Å, with a As–Lu–As bond angle of 81.14(6)°. By using the reducing nature of Sm(II), Evans and co-worker prepared [Sm(Cp\*)<sub>2</sub>(AsPh<sub>2</sub>)] (**85**), Fig. 9, by the reaction of two equivalents of [Sm(Cp\*)<sub>2</sub>] with Ph<sub>2</sub>AsAsPh<sub>2</sub> *via* reductive cleavage.<sup>120</sup> This strategy also works for making the phosphide analogue. When dissolved in THF, **85** converts to the THF adduct [Sm(Cp\*)<sub>2</sub>(-AsPh<sub>2</sub>)(THF)], which can ring open THF to produce [Sm(Cp\*)<sub>2</sub>{O(CH<sub>2</sub>)<sub>4</sub>AsPh<sub>2</sub>}(THF)] under thermolysis conditions.




Fig. 9 Selected examples of lanthanide complexes containing metal–arsenide bonds **84**–**87**.



Scheme 27 Synthesis of the yttrium and dysprosium arsinide complexes **88Ln**.

In the molecular structure of **85**, there are two independent molecules in unit cell, and the Sm–As bond distances are 2.973(3) and 2.966(3) Å, respectively, which are slightly longer than those seen in **84**. Apart from these Ln(III) arsenide complexes, there are two Ln(II) complexes containing metal–arsenic bonds, where Nief and co-workers reported the preparation of [Sm(AsMes)<sub>2</sub>(THF)<sub>4</sub>] (**86**)<sup>36</sup> and [Tm(Dsas)<sub>2</sub>(THF)] (**87**, Dsas = 2,5-bis(trimethylsilyl)-3,4-dimethylarsolide)<sup>121</sup> *via* salt metathesis, Fig. 9.


During 2015 and 2016, Layfield and co-workers synthesised the bridging arsinide lanthanide complexes [{Ln(Cp<sup>Me</sup>)<sub>2</sub>}(μ-AsMes)<sub>3</sub>Li][Li(THF)<sub>4</sub>]<sub>2</sub> (**88Ln**; Ln = Y, Dy; Cp<sup>Me</sup> = C<sub>5</sub>H<sub>4</sub>Me; Mes = mesityl), Scheme 27, when investigating the effects of arsinide ligands on SMM properties.<sup>122,123</sup> Deprotonation of the bridging lanthanide arsenides [{Ln(Cp<sup>Me</sup>)<sub>2</sub>}(μ-AsHMes)<sub>3</sub>] (Ln = Y or Dy), with three equivalents of *n*-butyl-lithium gave **88Y** and **88Dy**, in yields of 73% and 77%, respectively, in addition to three equivalents of butane gas. Complexes **88Ln** feature bridging pnictide units, similar to the phosphinide clusters **28Ln**. The central Ln<sub>3</sub>As<sub>3</sub> core shows a chair-like arrangement that is analogous to the Ln<sub>3</sub>P<sub>3</sub> cores in **28Ln**. The authors reported that **88Ln** exhibit Ln–As bond distances (Y: 2.8574(6)–2.8893(7) Å; Dy: 2.8515(6)–2.8908(7) Å) that are shorter than in their arsenide counterparts [{Ln(Cp<sup>Me</sup>)<sub>2</sub>}(μ-AsHMes)<sub>3</sub>](Ln = Y,



Scheme 28 Synthesis of lanthanide polyarsenide complexes **90**–**92Ln**.



More recent work from the Roesky group has proved that a redox strategy, using  $[\text{Fe}(\text{Cp}^*)(\eta^5\text{-As}_5)]$  as an arsenic ligand source to react with low-valent lanthanide starting materials, is an effective way to synthesise new lanthanide polyarsenide complexes. Three new 3d/4f polyarsenide complexes in the separate ion pair form  $[\text{K}(\text{18C6})][\text{Ln}(\text{Cp}^*)_2(\mu\text{-}\eta^4\text{:}\eta^4\text{-As}_5)\text{Fe}(\text{Cp}^*)]$  (**92Ln**, Ln = La, Ce, Nd) were prepared by the reduction of  $[\text{Fe}(\text{Cp}^*)(\eta^5\text{-As}_5)]$  with formally low-valent bridging arene lanthanide compounds in moderate yields, Scheme 28.<sup>125</sup> The molecular structures of **92Ln** are very similar to each other, containing highly reduced  $\text{As}_5$  units with an envelope shape. In **92Ln** the Fe centre is  $\eta^4$ -coordinated by the *cyclo*- $\text{As}_5$  unit and the shortest As–As bond distance of 2.3781(6) Å is observed, while the rest of the As–As bonds range from 2.3928(5) Å to 2.4325(5) Å. As expected, Ln–As bond distances vary over a wide range because of the steric constraints and anisotropic charge distribution within the  $\text{As}_5$  ligand. In other work, Roesky and co-workers found that reacting the arsenic source  $[\{\text{Co}(\text{Cp}''')\}_2(\mu\text{-}\eta^2\text{:}\eta^2\text{-As}_2)_2]$  with samarocenes produced two new mixed 3d/4f



The practical difficulties of working with As<sub>4</sub> solutions has spurred the development of new arsenic starting materials for construction of polyarsenic complexes. Roesky and co-workers reported the preparation for arsenic nanoparticles using a reductive synthetic method developed by Feldmann and co-workers.<sup>128</sup> The elemental As<sup>0</sup> nanoparticles (*nano*-As<sup>0</sup>,  $d = 7.2 \pm 1.8$  nm) can be formed by the reduction of AsI<sub>3</sub> with a freshly prepared solution of lithium naphthalenide at 0 °C in THF. The *nano*-As<sup>0</sup> can be isolated in a pure form as the LiI by-product is sufficiently soluble in THF and Et<sub>2</sub>O to be washed away; this has



proven very effective at introducing As oligomers to f-elements, Scheme 30. It was reported that **94** could be prepared by the reaction of  $[\text{Sm}(\text{Cp}^*)_2]$  with *nano-As*<sup>0</sup> at 60 °C, but again with inseparable byproducts and in low yield. Interestingly, under harsh conditions, the tetrasamarium polyarsenide cluster  $[\{\text{Sm}(\text{Cp}^*)_2\}_4(\mu\text{-As}_8)]$  (**95**) was reproducibly isolated in yields of 22% as a crystalline solid after work-up.<sup>128</sup> In the molecular structure of **95**, the Sm–As bond distances fall into the small range of 3.0814(10)–3.1734(10) Å, which are slightly longer than those of **94**, and the As–As bond distances are between 2.4044(12) Å and 2.5003(12) Å, suggesting single bonds with angles of 93.52(4)° to 103.68(4)° within the  $[\text{As}_8]^{4-}$  tetraanionic cage. Very recently, the authors expanded this chemistry, reacting *nano-As*<sup>0</sup> with the dilanthanide inverted arene complexes  $[\text{K}(18\text{-crown-6})][\{\text{Ln}(\text{Cp}^*)_2\}_2(\mu\text{-}\eta^6\text{:}\eta^6\text{-C}_6\text{H}_6)]$  (Ln = La, Ce) and  $[\text{K}(18\text{-crown-6})]_2[\{\text{Ln}(\text{Cp}^*)_2\}_2(\mu\text{-}\eta^6\text{:}\eta^6\text{-C}_6\text{H}_6)]$  (Ln = Ce, Nd) to give a range of lanthanide Zintl anions **96–99** containing  $\text{As}_3^{3-}$ ,  $\text{As}_7^{3-}$ , and  $\text{As}_{14}^{4-}$  ligands, respectively, that were previously not accessible in molecular lanthanide chemistry, Scheme 30.<sup>129</sup> The  $\text{As}_{14}^{4-}$  unit in **97** is the largest organo-lanthanide-polyarsenic complex to date. The synthesis and characterisation of these diverse polyarsenic lanthanide complexes demonstrated the great utility of *nano-As*<sup>0</sup> for accessing novel molecular polyarsenic clusters.

## 6.2 Lanthanide antimony complexes

Although there have not been any reports of structurally authenticated f-element-antimony or -bismuth multiple bonds, a relatively small number of lanthanide complexes featuring single bond interactions with these heavy pnictogens have been crystallographically characterised. Lanthanide–antimony or -bismuth bonds are commonly synthesised as clusters with the polyantimony or polybismuth ligands stabilised between multiple metal centres. Complexes containing metal–antimony or -bismuth multiple bonding interaction are also rare for d-block metals; to date, there has been only one single example of a terminal metal stibido complex,  $[\text{W}^{\text{VI}}(\text{Tren}^{\text{TMS}})(\text{Sb})]$  ( $\text{Tren}^{\text{TMS}} = \{\text{N}(\text{CH}_2\text{CH}_2\text{NSiMe}_3)_3\}^{3-}$ ) with a  $\text{W}\equiv\text{Sb}$  triple bond (2.526(2) Å), prepared by Scheer and co-workers.<sup>130</sup> No structurally authenticated metal–bismuth multiple bonds are known both for d-block and f-block metals. In 1992, Evans and co-workers reported the synthesis of a complex featuring Sm–Sb interactions. The reaction of one equivalent each of  $[\text{Sm}(\text{Cp}^*)_2]$  and  $\text{Sb}(\text{nBu})_3$  afforded the samarium antimony Zintl ion complex  $[\{\text{Sm}(\text{Cp}^*)_2\}_3(\mu\text{-}\eta^2\text{:}\eta^2\text{:}\eta^1\text{-Sb}_3)(\text{THF})]$  (**100**), Scheme 31.<sup>131</sup> Complex **100** exhibits five Sm–Sb bonds in the range of 3.162(1)–3.205(1) Å.




Scheme 31 Synthesis of samarium polyantimony complex **100**.



Scheme 32 Synthesis of lanthanide Zintl polyantimony complexes **101Ln**. Ln–Sb interactions are omitted for clarity.

Surprisingly, there were no further reports of lanthanide–antimony bonds until in 2016 a wide range of lanthanide–polyantimony clusters emerged. Sun, Boldyrev and co-workers reported the synthesis of lanthanide(III) complexes with three Zintl  $\text{Sb}_4^{2-}$  units  $[\text{K}(2.2.2\text{-cryptand})]_3[\text{Ln}(\eta^4\text{-Sb}_4)_3]$  (**101Ln**; Ln = La, Y, Ho, Er, Lu).<sup>132</sup> Complexes **101Ln** were synthesised *via* the reaction of  $[\text{Ln}(\text{CH}_2\text{C}_6\text{H}_5)_3(\text{THF})_3]$  (Ln = La, Y, Ho, Er, Lu) with three equivalents each of  $\text{K}_2\text{Sb}_4$  and 2.2.2-cryptand in pyridine, Scheme 32. The five complexes exhibit two types of Ln–Sb bond distances, with equatorial Ln–Sb<sub>eq</sub> distances of  $\sim 3.4$  Å and shorter non-equatorial Ln–Sb<sub>ne</sub> distances of  $\sim 3.2$  Å. The authors noted that the Ln–Sb bond distances decrease from lanthanum to lutetium due to the lanthanide contraction (Ln–Sb<sub>ne</sub>: 3.2461(5), 3.0932(10), 3.0843(9) and 3.0643(11) Å for La, Ho, Er and Lu respectively). Calculations on **101Ln** indicated that the  $\text{Sb}_4$  units are aromatic in nature, similar to cyclobutadienyl.

In 2017, Layfield and co-workers reported the synthesis of lanthanide stibide complexes which contain either a bridging stibide or a Zintl-like  $[\text{Sb}_4\text{Mes}_3]^{3-}$  moiety. The bridging stibide  $[\{\text{Ln}(\text{Cp}^{\text{Me}})_2\}_3(\mu\text{-SbHMe}_3)]$  (**102Ln**, Ln = Y, Dy) was synthesised *via* the reaction of three equivalents each of  $[\text{Ln}(\text{Cp}^{\text{Me}})_2]$  and  $\text{MesSbH}_2$ , Scheme 33.<sup>133</sup> Complexes  $[\{\text{Ln}(\text{Cp}^{\text{Me}})_2\}_3(\mu\text{-}(\text{SbMes})_3\text{-Sb})]$  (**103Ln**, Ln = Y, Dy) were synthesised *via* two different routes, the first of which involved the reaction of three equivalents each of  $[\text{Ln}(\text{Cp}^{\text{Me}})_2]$  and *n*-butyl lithium with four



Scheme 33 Synthesis of lanthanide polyantimony complexes **102Ln** and **103Ln**.



Scheme 34 Synthesis of samarium polyantimony complexes 104–107.

equivalents of  $\text{MesSbH}_2$ ; however, stibine dehydrocoupling to give the distibane ( $\text{Sb}_2\text{H}_2\text{Mes}_2$ ) and tetrastibetane ( $\text{Sb}_4\text{Mes}_4$ ) can occur. The second reported route to  $103\text{Ln}$  involved the cross-dehydrocoupling reaction of one equivalent of the stibide precursor  $102\text{Ln}$  with one equivalent of  $\text{MesSbH}_2$  to give one equivalent each of  $103\text{Ln}$  ( $\text{Ln} = \text{Y}$ , quantitative; or  $\text{Dy}$ , 45%) and mesitylene, and two equivalents of hydrogen gas. Complexes  $103\text{Ln}$  exhibit solid state structures with  $\text{Ln}_3\text{Sb}_3$  cores akin to the arsinidiide  $88\text{Ln}$  and phosphinidiide  $28\text{Ln}$  analogues, however in  $103\text{Ln}$  the  $\text{Sb}_3$  unit is capped with a  $\text{Sb}^{3-}$  fragment to form a Zintl-like moiety. Complexes  $103\text{Ln}$  feature mean  $\text{Y-Sb}$  and  $\text{Dy-Sb}$  bond distances of 3.1211(15)–3.1420(16) Å, and 3.119(1)–3.138(1) Å, respectively. These bond distances are very similar to that of the stibide precursors  $102\text{Ln}$ , which exhibit  $\text{Ln-Sb}$  bond distances of 3.0987(11)–3.2008(6) and 3.092(6)–3.212(3) Å for  $\text{Y}$  and  $\text{Dy}$ , respectively. This observation differs from the analogous pnictinidiide complexes  $28\text{Ln}$  and  $88\text{Ln}$ , where the  $\text{Ln-Pn}$  distances are shorter than the pnictide precursors; this is likely due to  $103\text{Ln}$  exhibiting  $[\text{Sb}_4\text{Mes}_3]^{3-}$  units as opposed to discrete  $\text{RSb}^{2-}$  stibinidene ligands. The authors reported that both  $102\text{Dy}$  and  $103\text{Dy}$  show SMM behaviour, with  $U_{\text{eff}}$  values of 345 and 270  $\text{cm}^{-1}$ , respectively, and both complexes exhibiting magnetic hysteresis at 1.8 K.

In 2018, Roesky and co-workers utilised  $[\text{Sm}(\text{Cp}^*)_2]$  to stabilise a range of samarium polyantimony complexes, Scheme 34.<sup>134</sup> The reaction of  $[\text{Sm}(\text{Cp}^*)_2]$  with an  $\text{Sb/Hg}$  amalgam in toluene yielded a mixture of the polyantimony complexes  $[\{\text{Sm}(\text{Cp}^*)_2\}_2(\mu\text{-}\eta^2\text{-}\eta^2\text{-Sb}_2)]$  (**104**) and  $[\{\text{Sm}(\text{Cp}^*)_2\}_2\text{Sb}_2(\mu\text{-Hg})]$  (**105**), Scheme 34. The product was dependent upon the temperature of the reaction; whilst at room temperature the primary products were **104** and **105**, if the reaction mixture was heated at 60–70 or 120 °C, the polyantimony complexes  $[\{\text{Sm}(\text{Cp}^*)_2\}_3(\text{Sb}_4)_2\text{Hg}]$  (**106**) or  $[\{\text{Sm}(\text{Cp}^*)_2\}_4(\text{Sb}_8)]$  (**107**) were isolated, respectively. Using a  $\text{Sb/Hg}$  amalgam however resulted in difficult to separate antimony products, so the authors



Fig. 10 Lanthanide-bismuth complexes **108** and **109**.

reported an alternate synthetic route to both **104** and **107** via the reaction of  $[\text{Sm}(\text{Cp}^*)_2]$  with antimony nanoparticles. Complex **104** exhibits an  $\text{Sm-Sb}$  bond distance of 3.2141(9) Å, which is comparable to complex **100**, whereas complex **105** demonstrates significantly shorter  $\text{Sm-Sb}$  distances of 3.0052(14)–3.0158(13) Å. Complexes **106** and **107** exhibit longer  $\text{Sm-Sb}$  bond distances of 3.2238(8)–3.3694(8) Å and 3.3134(8)–3.4119(7) Å, respectively. More recently, Roesky and co-workers have shown that  $[\text{Sm}(\text{Cp}^*)_2]$  can also be used to make 4d/4f polyantimony clusters containing a planar  $\text{Sb}_4$ -unit that is similar to that seen in **91**.<sup>135</sup>

### 6.3 Lanthanide bismuth complexes

The first example of a structurally authenticated lanthanide bismuth interaction was reported in 1991 by Evans and co-workers. The reaction of two equivalents of  $[\text{Sm}(\text{Cp}^*)_2]$  with one equivalent of  $\text{BiPh}_3$  in toluene afforded  $[\{\text{Sm}(\text{Cp}^*)_2\}_2(\mu\text{-}\eta^2\text{-}\eta^2\text{-Bi}_2)]$  (**108**) in yields of up to 60%,<sup>136</sup> Fig. 10. The complex is isostructural to the diantimony complex **104** and exhibits  $\text{Sm-Bi}$  distances ranging from 3.2645(10) to 3.3108(11) Å, which are longer than that the  $\text{Sm-Sb}$  distances in **104** (3.2141(9) Å). More recent work from the Evans group has shown that the reaction of  $\text{BiPh}_3$  with the  $\text{Gd(III)}$  complex  $[\text{K}(2.2.2\text{-cryptand})][\text{Gd}\{\text{N}(\text{SiMe}_3)_2\}_3]$  afforded a mononuclear  $\text{Gd(III)}$ -bismuthide complex  $[\text{K}(2.2.2\text{-cryptand})][\text{Gd}\{\text{N}(\text{SiMe}_3)_2\}_3(\text{BiPh}_2)]$  (**109**),<sup>137</sup> Fig. 10, which is the only example of a mononuclear lanthanide-bismuth compound. In the molecular structure of **109**, the  $\text{Gd-}$




Scheme 35 Synthesis of lanthanide dibismuth complexes **110Ln**–**111Ln**.



Bi bond distance of 3.3516(5) Å is very close to the Sm–Bi bond in **108**.

Very recently, Chilton, Demir and co-workers reported two new sets of lanthanide dibismuth complexes, Scheme 35. The neutral complexes  $[\{\text{Ln}(\text{Cp}^*)_2\}_2(\mu\text{-}\eta^2\text{:}\eta^2\text{-Bi}_2)]$  (**110n**, Ln = Gd, Tb, Dy, Y) were prepared by the one-pot reactions of eight equivalents of  $[\text{Ln}(\text{Cp}^*)_2(\text{BPh}_4)]$  (Ln = Gd, Tb, Dy, Y), two equivalents of triphenylbismuth, and eight equivalents of  $\text{KC}_8$  at room temperature under argon.<sup>138</sup> Complexes **110Ln** are quite soluble in toluene, thus the poor atom-efficiency of the reaction is overcome by separation from the poorly soluble byproducts. Reduction of **110Ln** using  $\text{KC}_8$  in the presence of 2.2.2-cryptand in THF afforded the radical complexes  $[\text{K}(2.2.2\text{-cryptand})][\{\text{Ln}(\text{Cp}^*)_2\}_2(\mu\text{-}\eta^2\text{:}\eta^2\text{-Bi}_2^{\cdot-})]$  (**111Ln**, Ln = Gd, Tb, Dy, Y).<sup>138</sup> Complexes **111Ln** are the first examples containing the heaviest dinitrogen radical analogues of dibismuth for any d- or f-block metal. The radical nature of **111Ln** are confirmed by the characterisation data and DFT calculations. Reflecting the radical electron in the  $\pi$ -antibonding orbital of the  $\text{Bi}_2$  ligand in **111Ln**, the Bi–Bi bond distances of (2.9310(11) to 2.9450(13) Å) are longer than those in neutral **110Ln** (2.8418(10) to 2.8549(9) Å), and the Ln–Bi bond distances are shorter in **111Ln** (3.1865(8) to 3.2064(5) Å) vs. **110Ln** (3.2335(2) to 3.2611(8) Å). Magnetic studies have shown that the  $\text{Bi}_2^{3-}$  radical-bridged **111Tb** and **111Dy** are SMMs with magnetic hysteresis where the  $\text{Bi}_2^{3-}$  radical as the bridge engenders antiferromagnetic exchange coupling with the paramagnetic lanthanide ions, leading to a ferrimagnetic ground state.

Similar to antimony chemistry there have been a range of lanthanide–bismuth interactions stabilised through the isolation of Zintl anions bound to the lanthanide metal ions. In 2011, Dehnen and co-workers utilised this method to synthesise a mini-fullerene-type Zintl-lanthanide complex, Scheme 36. The reaction of  $[\text{K}(2.2.2\text{-cryptand})]_2[\text{Sn}_2\text{Bi}_2]\cdot\text{en}$  (en = 1,2-ethylenediamine) and  $[\text{Eu}(\text{C}_5\text{Me}_4\text{H})_3]$  in toluene gave  $[\text{K}(2.2.2\text{-cryptand})]_4[\text{Eu}@\text{Sn}_6\text{Bi}_8]$  (**112**) in a 11% yield.<sup>139</sup> The solid-state structure of **112** indicates each position of the cage is occupied by either a Sn or Bi atom in a ratio of 0.76–0.11 and 0.24–0.89, respectively. The Eu–(Sn/Bi) bond distances exhibited by **112** span a wide range of 3.3515(8)–3.5770(9) Å. The Sn/Bi atoms in the non-equatorial positions exhibit shorter mean Eu–Sn/Bi




Scheme 37 Synthesis of lanthanide polybismuth Zintl anionic clusters **113Ln** and **114Ln** and representations of the polyhedral architectures. Ln–A interactions are omitted for clarity.

distances (3.4743(11) Å) compared to the equatorial distance (3.5208(15) Å). Complex **112** exhibits a short mean axial Eu–Sn/Bi distance of 3.3418(10) Å.

In 2012, Dehnen and co-workers further expanded the range of lanthanide bismuth clusters, Scheme 37; the separate reactions of  $[\text{K}(2.2.2\text{-cryptand})]_2[\text{Sn}_2\text{Bi}_2]\cdot\text{en}$  with  $[\text{Ln}(\text{C}_5\text{Me}_4\text{H})_3]$  (Ln = La or Ce) in *p*-xylene gave mixtures of tin–bismuth clusters surrounding lanthanide centres in  $[\text{K}(2.2.2\text{-cryptand})]_4[\text{Ln}@\text{Sn}_7\text{Bi}_7]_x[\text{Ln}@\text{Sn}_4\text{Bi}_9]_{1-x}$  (Ln = La,  $x = 0.70$ ; Ln = Ce,  $x = 0.39$ ).<sup>140</sup> The crystallographic data for both complexes demonstrate two types of cluster, the 14-vertex anion  $[\text{Ln}@\text{Sn}_7\text{Bi}_7]^{4-}$  (**113Ln**) and 13-vertex anion  $[\text{Ln}@\text{Sn}_4\text{Bi}_9]^{4-}$  (**114Ln**). Like previous Sn/Bi clusters, anions **113Ln** and **114Ln** exhibit disorder of the Sn and Bi atoms occupying the same atomic positions in the cluster. The clusters **113Ln** have similar structures to **112**, exhibiting longer mean Ln–Sn/Bi equatorial distances of 3.539(2) (La) and 3.493(3) Å (Ce) when compared to the non-equatorial distances of 3.433(2) (La) and 3.416(2) Å (Ce). Clusters **113La** and **113Ce** exhibit mean Ln–Sn/Bi axial distances of 3.426(2) and 3.436(2) Å, respectively. The second cluster type, **114Ln**, features Ln–Sn/Bi bond distances over the range of 3.107(4)–3.557(2) and 3.046(12)–3.543(3) Å for La and



Scheme 36 Synthesis of europium polybismuth Zintl anionic clusters **112**. Eu–A interactions are omitted for clarity.



Scheme 38 Synthesis of lanthanum polybismuth Zintl anionic clusters **115** and representation of the polyhedral architecture. La–Bi interactions are omitted for clarity.





Scheme 39 Synthesis of samarium polybismuth Zintl anionic clusters **116**. Sm–Bi/A interactions are omitted for clarity.

Ce, respectively. The authors postulated that the differences in Ln–Sn/Bi distances between the La and Ce analogues of **114La** and **114Ce** is most likely a result of the different ionic radii of La(III) and Ce(III).

In parallel work in 2012, Dehnen and co-workers reported the synthesis of the indium metalloid bismuth Zintl cluster,  $[K(2.2.2\text{-cryptand})]_6[La@In_2Bi_{11}]_2(\mu\text{-Bi})_2$  (**115**) via the reaction of  $[K(2.2.2\text{-cryptand})]_2[InBi_3]\cdot en$  with  $[La(C_5Me_4H)_3]$  in toluene in a yield of 11%,<sup>141</sup> Scheme 38. Complex **115** features two 13-vertex anions connected via two bridging bismuth atoms. The La–Bi distances exhibited by **115**, 3.1343(8)–3.5018(9) Å, are similar to that of the anion **114Ln** [3.106(4)–3.557(2) Å].

In 2014, Dehnen and co-workers utilised similar protocols to expand the number of 13-vertex lanthanide–bismuth metalloid clusters to include the gallium-substituted cage  $[K(2.2.2\text{-cryptand})]_2[Sm@Ga_2HBi_{11}]_{0.9}[Sm@Ga_3H_3Bi_{10}]_{0.1}$  (**116**) via the reaction of  $[K(2.2.2\text{-cryptand})]_2[GaBi_3]\cdot en$  with  $[Sm(C_5Me_4H)_3]$ , Scheme 39.<sup>142</sup> The authors reported that the two different



Scheme 41 Synthesis of lanthanide polybismuth anionic clusters **121Ln**.

trianionic fragments feature disorder similar to the tin analogue **113Ln**, with several of the atomic positions in the cluster being occupied by Bi, Ga and GaH. Formulation of each anionic cluster was determined by the examination of a combination of techniques, namely single crystal diffraction, energy dispersive X-ray and electrospray ionisation mass spectrometry (ESI-MS), which indicated that the two fragments were present in a ratio of 92% to 8% for  $[Sm@Ga_2HBi_{11}]$  and  $[Sm@Ga_3H_3Bi_{10}]$ , respectively. The Sm–Bi/Ga distances in **116** span 3.0464(6)–3.4134(6) Å, which is similar to the other 13-vertex lanthanide cluster complexes **114Ln** and **115**.

In 2015, Dehnen and co-workers reported the synthesis of a wide range of lead bismuth cages. The separate reactions of the *in situ*-generated Zintl lead precursor  $[K(2.2.2\text{-cryptand})]_2[Pb_2Bi_2]\cdot en$  with  $[Ln(C_5Me_4H)_3]$  (Ln = La, Ce, Nd, Gd, Sm, Tb) in toluene yielded ten complexes,  $[K(2.2.2\text{-cryptand})]_3[Ln@Pb_6Bi_8]_x[Ln@Pb_3Bi_{10}]_{1-x}$  ( $x = 0.038, 0$  or  $0.545$  for La, Ce or Nd respectively) or  $[K(2.2.2\text{-cryptand})]_4[Ln@Pb_7Bi_7]_y[Ln@Pb_4Bi_9]_{1-y}$  (Ln = La;  $y = 0.038, 0.279$  or  $0.458$ ; Ln = Nd, Sm or Tb;  $y = 0$ ), depending on the lanthanide used, Scheme 40.<sup>143</sup> All ten complexes feature a mixture of 14- and 13-vertex anions,  $[Ln@Pb_6Bi_8]_x[Ln@Pb_3Bi_{10}]_{1-x}]^{3-}$  (**117Ln**,  $x=18Ln_{1-x}$ ) and  $[Ln@Pb_7Bi_7]_y[Ln@Pb_4Bi_9]_{1-y}]^{4-}$  (**119Ln**,  $y=120Ln_{1-y}$ ); the structures for the differing anions are analogous to those of **113Ln** and **114Ln**. Like the previously reported lanthanide bismuth cages (**114Ln** and **116**), the anionic clusters **117Ln**–**120Ln** all contain disorder, resulting in both Pb and Bi occupying the same atomic sites.

In 2021, Demir and co-workers reported the synthesis of  $[K(THF)_4]_2[Ln(Cp^*)_2(\mu\text{-Bi}_6)]$  (**121Ln**, Ln = Tb, Dy) by one-pot reactions of  $[Ln(Cp^*)_2(BPh_4)]$  (Ln = Tb, Dy) with triphenylbismuth in THF, followed by reduction with  $KC_8$  at 45 °C, Scheme 41.<sup>144</sup> The authors proposed that the use of  $KC_8$  induced reduction and bismuth cluster formation. The solid-state structures of **121Ln** revealed  $[Ln_2Bi_6]$  cores with lanthanide centres bridged by a rare  $[Bi_6]^{6-}$  Zintl ion. Notably, complexes **121Ln** are the first examples of a cyclic bismuth hexamer in an organometallic complex with any metal. The Ln–Bi bond distances for **121Tb** and **121Dy** are 3.055(1)–3.070(1) and 3.042(1)–3.060(1) Å, respectively, which are approximately 0.2 Å shorter than that the Sm–Bi bonds in **108** (3.2645(10)–3.3108(11) Å). The Bi–Bi distances of 3.029(1)–3.042(1) and



Scheme 40 Synthesis of lanthanide polybismuth Zintl anionic clusters **117Ln**–**120Ln**. Ln–A interactions are omitted for clarity.

3.027(1)–3.036(1) Å for **121Tb** and **121Dy**, respectively, are significantly longer than multiple Bi–Bi bonds (2.82–2.87 Å), and are comparable with Bi–Bi single bonds (>2.99 Å). Magnetic data and quantum calculations indicates strong ferromagnetic interactions between the lanthanide ions facilitated by the Zintl  $[\text{Bi}_6]^{6-}$  ligands, resulting in magnetic blocking and open hysteresis loops that are rarely observed for super exchange-coupled SMMs containing solely lanthanide ions.

## 7. Actinide heavier pnictogen complexes


In parallel with lanthanide chemistry, actinide complexes containing heavier pnictogen ligands (from As to Bi) are much rarer than actinide–phosphorus congeners. Again, these complexes tend to form multi-nuclear clusters with the heavier pnictogen ligands bridged by two or more metal centres, thus well-defined mononuclear species are sparse. However, significant advances have been achieved in actinide arsenic multiple bonding chemistry that are discussed in this section. In contrast, there are currently no examples of actinide complexes featuring multiple bonding interactions with antimony or bismuth ligands.

### 7.1 Actinide arsenic complexes

In 1994, Using a similar synthetic approach used for the preparation of **77**, Scherer and co-workers synthesised the first examples of an actinide polyarsenic complex  $[\{\text{Th}(\text{Cp}^{\text{tt}})_2\}_2(\mu^2\text{-}\eta^3\text{-As}_6)]$  (**122**) by the treatment of  $[(\text{Cp}^{\text{tt}})_2\text{Th}(\eta^4\text{-C}_4\text{H}_6)]$  with elemental  $\text{As}_4$  in boiling xylene,<sup>145</sup> Fig. 11. Complex **122** is also the first structurally characterised actinide complex containing metal–arsenic bonds, which is isostructural to the polyphosphorus analogue **77**. In the molecular structure of **122**, the Th–As bond distances span the range of 2.913(2) to 3.044(2) Å. However, it would be more than 20 years before more actinide–arsenic complexes started to emerge, reflecting the synthetic challenges of the area.



Fig. 11 Examples containing actinide–arsenic bonds **122**–**125**.



Scheme 42 Synthesis of thorium arsinidiide and diarsene complexes **126**–**128**.

In 2015, Liddle, Scheer, and co-workers reported the diuranium complex  $[\{\text{U}(\text{Tren}^{\text{TIPS}})\}_2(\mu\text{-}\eta^2\text{-}\eta^2\text{-HASAsH})]$  (**123**)<sup>146</sup> by the reaction of  $[\text{U}(\text{Tren}^{\text{TIPS}})(\text{THF})][\text{BPh}_4]$  with  $\text{KAsH}_2$  in 1 : 1.4 ratio *via* a dehydrocoupling process, Fig. 11. On one occasion, the uranium diarsenic complex  $[\{\text{U}(\text{Tren}^{\text{TIPS}})\}_2(\mu\text{-}\eta^2\text{-}\eta^2\text{-As}_2)]$  (**124**)<sup>146</sup> was also obtained from the reaction in less than 1% yield, which can be viewed as a fully dehydrocoupled product, Fig. 11. The solid-state structures of **123** and **124** are similar to each other, though the As–As bond distance in **123** (2.4102(13) Å) is longer than that in **124** (2.2568(14) Å), indicating As–As single and As=As double bond characters, respectively. The +4 oxidation state of **123** was confirmed by magnetometry data, so the diarsene ligand has been reduced to its diarsane-1,2-diide form. Complex **123** is the first example of HASAsH complex for any d- or f-block metal. The characterisation data and theoretical calculations supported the presence of back-bonding-type interactions from uranium to the HASAsH  $\pi^*$ -orbital, indicating the strong  $\pi$ -accepting ability of this ligand. In other work, the authors also reported the structure of  $[\{\text{U}(\text{Tren}^{\text{TIPS}})\}_2(\mu\text{-}\eta^3\text{-}\eta^3\text{-As}_3)]$  (**125**),<sup>147</sup> Fig. 11, but the low yield for this compound prevented further characterisation.

In between 2016 and 2021, Walensky and co-workers reported the synthesis of thorium arsenic complexes, Scheme 42. Using methane elimination, the authors prepared the bridging dithorium arsinidiide complex  $[\{\text{Th}(\text{Cp}^*)_2(\mu\text{-AsMes})_2\}]$  (**126**) and the arsenide complexes  $[\text{Th}(\text{Cp}^*)_2(\text{HASAr})_2]$  (Ar = Mes, Tripp) by the reactions of  $[\text{Th}(\text{Cp}^*)_2(\text{Me})_2]$  under different conditions.<sup>148</sup> It



Scheme 43 Synthesis of the uranium arsinidene complexes **130**, arsenido **131**, and arsinidiide **132** from the arsinide **129**.



Scheme 44 Synthesis of thorium arsinidiide and arsenido complexes **133–135**.

was found that  $[\text{Th}(\text{Cp}^*)_2(\text{HAsMes})_2]$  is unstable at room temperature and converts to the dehydrocoupled product  $[\text{Th}(\text{Cp}^*)_2(\mu\text{-As}_2\text{Mes}_2)]$  (**127**) with the release of  $\text{H}_2$  gas. Applying the same synthetic approach, the uranium analogues of **126** and **127** were prepared. With the bulkier substituents on the arsinide group,  $[\text{Th}(\text{Cp}^*)_2(\text{HAsTripp})_2]$  is thermally stable at room temperature and its reactivity with *tert*-butyl isocyanide  $\text{BuNC}$  to give an arsaazaallene product was investigated.<sup>149</sup> The authors reported that deprotonation of  $[\text{Th}(\text{Cp}^*)_2(\text{AsH}(\text{Tripp}))_2]$  with one equivalent of  $\text{KN}(\text{SiMe}_3)_2$  gave  $[\text{Th}(\text{Cp}^*)_2(\mu\text{-As}(\text{Tripp}))\{\mu\text{-AsH}(\text{Tripp})\}\text{K}]_2$  (**128**) in yields of 77%.<sup>86</sup> The Th–As bond distances in **126** (2.8787(6) Å) and **127** (2.923(2)/2.971(3) Å) are quite long; by contrast, **128** exhibits a short Th–As distance of 2.7994(4) Å, indicating a multiple bonding interaction. This was probed computationally, revealing a Th=As Wiberg bond index (1.30) nearly twice that of the Th–As single bonds (0.70).

In 2015, Liddle, Scheer, and co-workers adapted the synthetic protocols used for the preparation of the uranium phosphinidene complex **64K**. Deprotonation of **129** with benzyl potassium in the presence of two equivalents of  $\text{B15C5}$  produced the analogous arsinidene complex  $[\text{K}(\text{B15C5})_2][\text{U}(\text{Tren}^{\text{TIPS}})(\text{AsH})]$  (**130**).<sup>147</sup> Scheme 43. Most surprisingly, double deprotonation of **129** by two equivalents of benzyl potassium afforded the arsenido complex  $[\{\text{U}(\text{Tren}^{\text{TIPS}})(\mu\text{-As})(\mu\text{-K}_2)\}_4]$  (**131**).<sup>147</sup> Attempts to abstract the potassium cations from **131** using 2.2.2-cryptand resulted in formation of the arsinidiide  $[\{\text{U}(\text{Tren}^{\text{TIPS}})(\mu\text{-AsH})\}\{\text{K}(2.2.2\text{-cryptand})\}]$  (**132**),<sup>147</sup> with K/H exchange from solvent. The authors additionally reported an As–H stretch in the ATR-IR spectrum of **130** at  $1857 \text{ cm}^{-1}$ . Complex **131** is a tetramer with an  $\text{As}_4\text{K}_6$  adamantane-type core, with the bridging potassium ions self-evidently playing an indispensable role in stabilising the arsenido  $[\text{As}]^{3-}$  centres, cf. formation of **132**.

The U–As distance of 2.7159(13) Å in the terminal arsinidene complex **130** is shorter than that seen for the potassium-capped arsinidiide complex **132** (2.7489(10) Å) and the arsenido complex **131** (2.730(2)–2.775(2) Å), which is due to the

coordination of the potassium ion (or ions) to the arsenic centre(s). Computational analysis showed highly polarised single, double, and triple uranium arsenic bonding interactions for  $\text{U-AsH}_2$  (**129**),  $\text{U=AsH}$  (**130**) and  $\text{U}\equiv\text{As}$  (**131**), with substantial 5f orbital contributions to these bonds.

In 2017, Liddle, Scheer, and co-workers adapted the protonolysis chemistry used to access **67**, **70**, and **71**, performing protonolysis reactions of **65Th** with different ratios of  $\text{KAsH}_2$  both with and without 15C5 to synthesise a range of Th–As complexes, Scheme 44, including the parent arsinidiide  $[\{\text{Th}(\text{Tren}^{\text{TIPS}})\}_2(\mu\text{-AsH})]$  (**133**), the arsinidiide  $[\{\text{Th}(\text{Tren}^{\text{TIPS}})(\mu\text{-AsH})\}\{\text{K}(15\text{C5})\}]$  (**134**), and the bridging arsenido  $[\text{K}(15\text{C5})_2][\{\text{Th}(\text{Tren}^{\text{TIPS}})\}_2(\mu\text{-As})]$  (**135**), Scheme 44.<sup>150</sup> The authors reported that attempts to prepare a terminal arsinidene  $[\text{K}(\text{L})_n][\text{Th}(\text{Tren}^{\text{TIPS}})(\text{AsH})]$  ( $\text{L} = \text{crown ethers or 2.2.2-cryptand}$ ) under various conditions were unsuccessful. Complex **134** could also be readily prepared in a 70% yield *via* the deprotonation of the parent arsinide  $[\text{Th}(\text{Tren}^{\text{TIPS}})\text{AsH}_2]$  (the Th analogue of **129**) with one equivalent of benzyl potassium in the presence of one equivalent of 15C5. The ATR-IR spectra of **133** and **134** exhibited As–H stretches of  $1930$  and  $1922 \text{ cm}^{-1}$ , respectively. Consistent with the arsenido nature of **135**, no As–H stretch was observed in the ATR-IR spectrum of **135**.

Complexes **133–135** have respective Th–As distances of 2.9619(6)/3.0286(6), 2.8565(7) and 2.8063(14)/2.8060(14) Å, which are shorter than that of the parent arsinide  $[\text{Th}(\text{Tren}^{\text{TIPS}})\text{AsH}_2]$  [3.065(3) Å]. When compared to the sum of the covalent single and double bond radii of Th and As of 2.96 and 2.57 Å, respectively, these Th–As bonds are relatively long and clearly polarised. An acute Th–As–H bond angle of  $79.1(2)^\circ$  in **134** suggested that a Th $\cdots$ H ‘agostic-type’ interaction could be present. The solid state structure of **135** is analogous to the thorium phosphido complex **70**, with a symmetrical ThAsTh core and near linear Th–As–Th angle of  $177.04(6)^\circ$ . In line with the multiple bonding interaction in the ThAsTh linkage, the Th–As distances exhibited by **135** (2.8063(14)/2.8060(14) Å) are





Scheme 45 Synthesis of terminal actinide parent arsinidene and arsenide complexes **136An** and **137An**.

shorter than those in **133** (2.9619(6)/3.0286(6) Å). The authors reported that attempts to prepare a terminal  $\text{Tren}^{\text{TIPS}}$  thorium arsenido  $\text{Th}=\text{As}$  species using the similar double deprotonation method for **132** proved unsuccessful; indeed, closely related work attempting to construct thorium nitrides supported by  $\text{Tren}^{\text{TIPS}}$  consistently resulted in the formation of parent imido complexes.

In 2022, Liddle, Scheer, and co-workers showed that a terminal parent arsinidene at thorium (and uranium for comparison) could be isolated using the bulky  $\text{Tren}^{\text{TCHS}}$  ligand, Scheme 45. Specifically, reaction of **68An** with  $\text{KAsH}_2$  in the presence of 2.2.2-cryptand in THF afforded directly, in good yields,  $[\text{K}(2.2.2\text{-cryptand})][\text{An}(\text{Tren}^{\text{TIPS}})(\text{AsH})]$  (**136An**), Scheme 45.<sup>84</sup> The molecular structures of **136An** confirmed the presence of terminal arsinidene ( $\text{AsH}^{2-}$ ) units that are well-protected by the tricyclohexylsilyl groups. The An–As bond distances of 2.8521(8) Å and 2.7581(6) Å for **136Th** and **136U**, respectively, are statistically invariant to those in **134** and **130**, indicating multiple bonding interactions in these An = AsH linkages that were confirmed by DFT calculations. ‘Agostic-type’ interactions between the metal ions and arsinidene ligands are observed in both structures, with An–As–H angles of 60.50(12)° and 61.18(12)°, respectively. In addition, ATR-IR spectroscopy revealed As–H stretches at 1867 and 1875  $\text{cm}^{-1}$  for **136Th** and **136U**, respectively. The authors found that treatment of **136An** with  $[\text{HNEt}_3][\text{BPh}_4]$  in THF results in the isolation of the parent arsenide complexes  $[\text{An}(\text{Tren}^{\text{TCHS}})(\text{AsH}_2)]$  (**137An**) in good yields, Scheme 45; oxidation of **136U** also gave the protonated product **137U**.<sup>84</sup>

## 7.2 Actinide 2-arsaethynolate complexes

Following the aforementioned developments in f-element 2-phosphethynolate chemistry, the chemistry of the corresponding 2-arsaethynolate anion ( $\text{OCAs}^-$ ) is beginning to emerge. In 2018, Meyer and co-workers reported reactions of the U(III)



Scheme 46 Synthesis of the actinide–antimony complexes **138An**.

complex  $[\text{U}\{(\text{Ad},\text{MeArO})_3\text{N}\}(\text{DME})][\{(\text{Ad},\text{MeArO})_3\text{N}\}^{3-}]$  = trianion of tris(2-hydroxy-3-(1-adamantyl)-5-methylbenzyl)amine) with one equivalent of  $[\text{Na}(\text{OCAs})(\text{dioxane})_3]$ , and 2.2.2-cryptand giving  $[\text{Na}(2.2.2\text{-cryptand})][\{(\text{Ad},\text{MeArO})_3\text{N}\}(\text{THF})\{\mu\text{-O}\}\{(\text{Ad},\text{MeArO})_3\text{N}\}(\text{OCAs})]$ .<sup>151</sup> In contrast, using two equivalents of  $[\text{Na}(\text{OCAs})(\text{dioxane})_3]$  yielded the binuclear,  $\mu$ -oxo bridged diuranium(IV/IV) complex  $[\text{Na}(2.2.2\text{-cryptand})]_2[\{(\text{Ad},\text{MeArO})_3\text{N}\}(\mu\text{-O})(\mu\text{-AsCAs})]$ , which contains a  $\mu\text{-}\eta^1\text{-}\eta^1$ -coordinated ( $\text{AsCAs}^{2-}$ ) ligand.<sup>151</sup> In 2019, Liddle, Scheer, and co-workers reported the first U–OCAs complex  $[\text{U}(\text{Tren}^{\text{TIPS}})(\text{OCAs})]$ , and treatment of this complex with  $\text{KC}_8$  and 2.2.2-cryptand as an *in situ* electrode mixture yielded  $[\text{K}(2.2.2\text{-cryptand})][\{(\text{U}(\text{Tren}^{\text{TIPS}}))_2\{\mu\text{-}\eta^2(\text{OAs})\}(\mu\text{-OCAs})\}]$  which is the As analogue of **75**, with a highly reduced bent, carbene-like OCAs ligand.<sup>152</sup> In contrast, reduction or photolysis of  $[\text{U}(\text{Tren}^{\text{TIPS}})(\text{OCAs})]$  with  $[\text{U}(\text{Tren}^{\text{TIPS}})]$  gave the mixed-valence arsenido complex  $[\{(\text{U}(\text{Tren}^{\text{TIPS}}))_2(\mu\text{-As})\}]$ , in very low yield, or **123**, respectively.<sup>152</sup> All of these results demonstrate the challenges of using the OCAs ligand to synthesise actinide–arsenic bonds and also that the synthetic methods and ancillary ligands drive the ( $\text{OCAs}^-$ ) bond cleavage chemistry in very different directions.

## 7.3 Actinide antimony complexes

In 2017, Liddle, Scheer, and co-workers reported the isolation and characterisation of the first discrete (*i.e.* not multi-centre) An–Sb bonds (An = U and Th) from the reactions of  $[\text{An}(\text{Tren}^{\text{TIPS}})(\text{L})][\text{BPh}_4]$  (An = U, L = THF; An = Th, L = DME) and  $\text{KSb}(\text{SiMe}_3)_2$  in THF, yielding  $[\text{An}(\text{Tren}^{\text{TIPS}})\{\text{Sb}(\text{SiMe}_3)_2\}]$  (**138An**), Scheme 46.<sup>153</sup> The closely related uranium complex  $[\text{U}(\text{Tren}^{\text{DMBS}})\{\text{Sb}(\text{SiMe}_3)_2\}]$  with a less sterically bulky supporting ligand could also be prepared using the same synthetic approach.<sup>153</sup>

Complexes **138Th** and **138U** exhibit An–Sb distances of 3.2849(3) and 3.2089(6) Å, respectively. The authors observed a shorter U–Sb bond in **138U** (3.2089(6) Å) compared to  $[\text{U}(\text{Tren}^{\text{DMBS}})\{\text{Sb}(\text{SiMe}_3)_2\}]$  (3.2437(8) Å), likely a result of the differing sterics between the  $\text{Tren}^{\text{TIPS}}$  and  $\text{Tren}^{\text{DMBS}}$  ligand



Scheme 47 Synthesis of the uranium–bismuthide complex **139**.





Scheme 48 Synthesis of the uranium–bismuth Zintl anionic clusters 140–142. U–Bi bonds are omitted for clarity.

systems; the latter results in more orthogonal, and presumably weaker, binding of the stibide ligand. The analogous U–P and U–As complexes were also prepared in that study, revealing increasingly pyramidalised pnictide centres as the group is descended.

#### 7.4 Actinide bismuth complexes

In the same publication describing the work in Section 7.3, Liddle, Scheer, and co-workers also reported the synthesis and characterisation of the first two-centre-two-electron (2c–2e) U–Bi bond,  $[U(\text{Tren}^{\text{DMBS}})\{Bi(\text{SiMe}_3)_2\}]$  (**139**), by the reaction of  $[U(\text{Tren}^{\text{DMBS}})(\text{THF})][\text{BPh}_4]$  with  $\text{KBi}(\text{SiMe}_3)_2$  in THF, Scheme 47.<sup>153</sup> It was found that the U–Bi bond could not be stabilised using the bulkier  $\text{Tren}^{\text{TIPS}}$  ligands, likely due to steric overload. Complex **139** exhibits U–Bi distances of 3.3208(4) Å which is longer than the above An–Sb bonds. Whilst the U–Bi bond in **139** was isolable, no Th–Bi bond could be isolated with  $\text{Tren}^{\text{TIPS}}$  or  $\text{Tren}^{\text{DMBS}}$ , underscoring the fragility of these linkages. Complex **139** is the only monomeric actinide complex containing a discrete bond to Bi to date.

Dehnen and co-workers reported actinide bismuth clusters in 2016. Reaction of  $[U(\text{C}_5\text{Me}_4\text{H})_3]$  with  $[K(2.2.2\text{-cryptand})]_2[EE'Bi_2]\cdot\text{en}$  (E = Ga, Tl, E' = Bi; E = E' = Pb) in 1,2-ethylenediamine gave either  $[K(2.2.2\text{-cryptand})]_3[U@Bi_{12}]$  (**140**),  $[K(2.2.2\text{-cryptand})]_2[K(2.2.2\text{-cryptand})(\text{en})][U@Ti_2Bi_{11}]$  (**141**) or  $[K(2.2.2\text{-cryptand})]_3[U@Pb_7Bi_7]_{0.66}[U@Pb_4Bi_9]_{0.34}$  (**142**), Scheme 48.<sup>154</sup> The structure of the trianionic fragment of **140** is analogous to the antimony cluster **101Ln**, exhibiting longer equatorial U–Bi bond distances (3.463(3)–3.545(3) Å) than the non-equatorial U–Bi bond distances (3.119(3)–3.167(3) Å). Complex **141** exhibits a 13-vertex cluster that is analogous to **114Ln**, with U–Bi



Scheme 49 Synthesis of the thorium–bismuth Zintl anionic clusters **143**. The Th–Bi bonds are omitted for clarity.

distances ranging from 3.068(1) to 3.4515(6) Å. Complex **142** has an isomorphous structure to **118Ln/120Ln**, exhibiting U–Bi bond distances ranging from 2.885(9) to 3.6885(12) Å.

In 2021, Dehnen and co-workers extended this Bi cluster chemistry to include the first example of a thorium bismuth cluster containing Th–Bi bonds, Scheme 49. The authors reported that reaction of  $[C_5Me_4H_3Th(Cl)]$  with  $K_5Ga_2Bi_4$ , which can be used as an *in situ* source of  $[GaBi_3]^{2-}$  and  $Bi_4^{2-}$ , in the presence of 2.2.2-cryptand in 1,2-ethylenediamine afforded  $[K(2.2.2\text{-cryptand})]_4[Th@Bi_{12}]\cdot 2\text{en}$  (**143**) as black, prismatic crystals, Scheme 49.<sup>155</sup> The structure of **143** was confirmed by single-crystal X-ray diffraction and the Th:Bi ratio within the cluster was verified by micro-X-ray fluorescence spectroscopy. The molecular structure revealed Bi–Bi bond distances over a relatively small range (3.0420(14)–3.132(1) Å) and Th–Bi bond lengths (3.2104(11)–3.5908(9) Å) that are comparable to the An–Bi bonds in **139–142**. Magnetic data and theoretical studies on **143** confirm the formal assignment as  $Th^{4+}$  and  $Bi_{12}^{8-}$  with a remarkable ring current strength of 24.8 nAT<sup>−1</sup> for  $[Th@Bi_{12}]^{4-}$  (and 23.7 nAT<sup>−1</sup> for  $Bi_{12}^{8-}$ ). This is much larger than in 6π-aromatic benzene (11.4 nAT<sup>−1</sup>), but close to that in 26π-aromatic porphine (25.3 nAT<sup>−1</sup>), despite the much smaller number of 2π-electrons involved. The aromatic nature of  $[Th@Bi_{12}]^{4-}$  extends aromaticity to the heaviest all-metal inorganic system.

## 8. Conclusions and outlook

Although f-element heavy pnictogen chemistry initially progressed quite slowly for many decades, with the resurgence of non-aqueous f-element chemistry momentum in this area has increased significantly in recent years, as evidenced by the burgeoning array of novel metal-heavy-pnictogen bond types that are now known. These well-characterised compounds have enabled us to secure new structural motifs and probe the nature of f-element ligand chemical bonds to further deepen our

understanding of chemical bonding with increasingly heavy ions in non-relativistic to relativistic regimes. Overall, f-element heavier pnictogen chemistry is developing well but there are certainly numerous opportunities to advance knowledge and understanding in this burgeoning field.

Looking forward, there are several appealing directions for researchers to explore in this area: (1) in general terms, f-element-phosphorus chemistry is maturing, but arsenic and especially antimony and bismuth are poorly developed – bringing the latter three to the same level of maturity as phosphorus will do much to elucidate periodic trends; (2) there are still relatively few actinide–pnictidene/ido multiple bond complexes and even fewer lanthanide analogues – however, the reports of isolated complexes to date suggests that there is ample scope to secure new Ln and An double and triple bonds to P, As, Sb, and Bi if the right supporting ligands can be identified and coupled with suitable synthetic approaches; (3) heavy analogues of dinitrogen are now known and even as radical species, but examples remain few in number – developing better synthetic approaches would open the area up and provide interesting electronic and physicochemical properties and potential atom-transfer methodologies; and, (4) though relatively few in number, it is already clear that f-element pnictogen clusters can exhibit novel magnetic and aromaticity properties – expanding the range of such compounds can only enhance our understanding of these fundamental phenomena.

## Author contributions

J. D., D. P. M. and S. T. L. designed the structure of the review. All authors contributed to the writing of the article.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We gratefully acknowledge funding and support from the Royal Society (UF071260, UF110005, RG080285, RG110238), Engineering and Physical Sciences Research Council (EP/K024000/1, EP/M027015/1, EP/P001386/1, EP/P002560/1, EP/S033181/1, EP/T011289/1), European Research Council (CoG-239621, CoG-612724), the Universities of Nottingham, Manchester, Regensburg, and Zhengzhou. S. T. L. thanks the Alexander von Humboldt Foundation for a Friedrich Wilhelm Bessel Research Award.

## References

- J. P. Liu, M. Willard, W. Tang, E. Brück, F. de Boer, E. Liu, J. Liu, C. Felser, G. Fecher, L. Wollmann, O. Isnard, E. Burzo, S. Liu, J. F. Herbst, F. Hu, Y. Liu, J. Sun, B. Shen and A. de Visser, *Metallic magnetic materials*, in *Handbook of Magnetism and Magnetic Materials*, ed. Coey M. and Parkin S., Springer, Cham, 2021.
- O. Gutfleisch, M. A. Willard, E. Brück, C. H. Chen, S. G. Sankar and J. P. Liu, *Adv. Mater.*, 2011, **23**, 821–842.
- Y. Hasegawa and Y. Kitagawa, Lanthanide-based materials for electroluminescence, in *Modern Applications of Lanthanide Luminescence*, ed. de Bettencourt-Dias A., Springer, Cham, 2021, Springer Series on Fluorescence, vol. 19.
- R. Marin and D. Jaque, *Chem. Rev.*, 2021, **121**, 1425–1462.
- H. Dong, S.-R. Du, X.-Y. Zheng, G.-M. Lyu, L.-D. Sun, L.-D. Li, P.-Z. Zhang, C. Zhang and C.-H. Yan, *Chem. Rev.*, 2015, **115**, 10725–10815.
- A. S. Hyre and L. H. Doerr, *Coord. Chem. Rev.*, 2020, **404**, 213098.
- G. A. Molander, *Chem. Rev.*, 1992, **92**, 29–68.
- F. Ortu, *Chem. Rev.*, 2022, **122**, 6040–6116.
- M. Scharfe, P. A. Lira-Parada, A. P. Amrute, S. Mitchell and J. Pérez-Ramírez, *J. Catal.*, 2016, **344**, 524–534.
- Z. Zhang, S. Liu, X. Li, T. Qin, L. Wang, X. Bo, Y. Liu, L. Xu, S. Wang, X. Sun, Y. Lu, F. Luo and S. Liu, *ACS Appl. Mater. Interfaces*, 2018, **10**, 22023–22030.
- S. G. McAdams, A. M. Ariciu, A. K. Kostopoulos, J. P. S. Walsh and F. Tuna, *Coord. Chem. Rev.*, 2017, **346**, 216–239.
- X. Cheng, J. Zhou, J. Yue, Y. Wei, C. Gao, X. Xie and L. Huang, *Chem. Rev.*, 2022, **122**, 15998–16050.
- A. Leoncini, J. Huskens and W. Verboom, *Chem. Soc. Rev.*, 2017, **46**, 7229.
- A. V. Gelis, P. Kozak, A. T. Breshears, M. A. Brown, C. Launier, E. L. Campbell, G. B. Hall, T. G. Levitskaia, V. E. Holfeltz and G. J. Lumetta, *Sci. Rep.*, 2019, **9**, 1–11.
- D. A. Atwood, *The Rare Earth Elements: Fundamentals and Applications*, John Wiley & Sons, Ltd, Chichester, UK, 2012.
- S. T. Liddle, *Angew. Chem., Int. Ed.*, 2015, **54**, 8604–8641.
- N. P. Bessen, J. A. Jackson, M. P. Jensen and J. C. Shafer, *Coord. Chem. Rev.*, 2020, **421**, 213446.
- D. Schädle and R. Anwender, *Chem. Soc. Rev.*, 2019, **48**, 5752.
- M. Keener, L. Maria and M. Mazzanti, *Chem. Sci.*, 2023, **14**, 6493–6521.
- S. P. Vilanova and J. R. Walensky, *Encyclopedia of Inorganic and Bioinorganic Chemistry*, John Wiley & Sons, Ltd, 2018.
- C. R. Groom, I. J. Bruno, M. P. Lightfoot and S. C. Ward, *Acta Crystallogr., Sect. B: Struct. Sci., Cryst. Eng. Mater.*, 2016, **72**, 171–179.
- L.-W. Ye, X.-Q. Zhu, R. L. Sahani, Y. Xu, P.-C. Qian and R.-S. Liu, *Chem. Rev.*, 2021, **121**, 9039–9112.
- C. Zhu and H. Xia, *Acc. Chem. Res.*, 2018, **51**, 1691–1700.
- H. Aktaş, J. C. Slootweg and K. Lammertsma, *Angew. Chem., Int. Ed.*, 2010, **49**, 2102–2113.
- O. T. Summerscales and J. C. Gordon, *RSC Adv.*, 2013, **3**, 6682–6692.
- T. Li, S. Kaercher and P. W. Roesky, *Chem. Soc. Rev.*, 2014, **43**, 42–57.
- M. E. García, D. García-Vivó, A. Ramos and M. A. Ruiz, *Coord. Chem. Rev.*, 2017, **330**, 1–36.
- L. Qiao, C. Zhang, X.-W. Zhang, Z.-C. Wang, H. Yin and Z.-M. Sun, *Chin. J. Chem.*, 2020, **38**, 295–304.



- 29 L. Giusti, V. R. Landaeta, M. Vanni, J. A. Kelly, R. Wolf and M. Caporali, *Coord. Chem. Rev.*, 2021, **441**, 213927.
- 30 D. P. Mills and P. Evans, *Chem.–Eur. J.*, 2021, **27**, 6645–6665.
- 31 L. N. Grant and D. J. Mindiola, *Chem.–Eur. J.*, 2019, **25**, 1–9.
- 32 M. D. Fryzuk, T. S. Haddad and D. J. Berg, *Coord. Chem. Rev.*, 1990, **99**, 137–212.
- 33 B. Ramirez, P. Sharma, R. J. Eisenhart, L. Gagliardi and C. C. Lu, *Chem. Sci.*, 2019, **10**, 3375–3384.
- 34 B. Ramirez and C. C. Lu, *J. Am. Chem. Soc.*, 2020, **142**, 5396–5407.
- 35 K. Izod, P. O'Shaughnessy, J. M. Sheffield, W. Clegg and S. T. Liddle, *Inorg. Chem.*, 2000, **39**, 4741–4748.
- 36 K. Izod, S. T. Liddle, W. McFarlane and W. Clegg, *Organometallics*, 2004, **23**, 2734–2743.
- 37 K. Izod, S. T. Liddle and W. Clegg, *Chem. Commun.*, 2004, 1748–1749.
- 38 G. W. Rabe, J. Riede and A. Schier, *J. Chem. Soc., Chem. Commun.*, 1995, 577–578.
- 39 G. W. Rabe and J. W. Ziller, *Inorg. Chem.*, 1995, **34**, 5378–5379.
- 40 G. W. Rabe, J. Riede and A. Schier, *Organometallics*, 1996, **15**, 439–441.
- 41 S. Atlan, F. Nief and L. Ricard, *Bull. Soc. Chim. Fr.*, 1995, **132**, 649.
- 42 F. Nief and L. Ricard, *J. Organomet. Chem.*, 1997, **529**, 357–360.
- 43 G. W. Rabe, I. A. Guzei and A. L. Rheingold, *Inorg. Chem.*, 1997, **36**, 4914–4915.
- 44 M. Westerhausen, S. Schneiderbauer, M. Hartmann, M. Warchhold and H. Nöth, *Z. Anorg. Allg. Chem.*, 2002, **628**, 330–332.
- 45 M. Westerhausen, S. Schneiderbauer, N. Makropoulos, M. Warchhold, H. Nöth, H. Piotrowski and K. Karaghiosoff, *Organometallics*, 2002, **21**, 4335–4341.
- 46 P. L. Arnold, F. G. N. Cloke and P. B. Hitchcock, *Chem. Commun.*, 1997, 481–482.
- 47 L. Jacquot, M. Xémard, C. Clavaguéra and G. Nocton, *Organometallics*, 2014, **33**, 4100–4106.
- 48 A. Jaoul, C. Clavaguéra and G. Nocton, *New J. Chem.*, 2016, **40**, 6643–6649.
- 49 T. Arliguie, M. Doux, N. Mézailles, P. Thuéry, P. Le Floch and M. Ephritikhine, *Inorg. Chem.*, 2006, **45**, 9907–9913.
- 50 J. D. Masuda, K. C. Jantunen, O. V. Ozerov, K. J. T. Noonan, D. P. Gates, B. L. Scott and J. L. Kiplinger, *J. Am. Chem. Soc.*, 2008, **130**, 2408–2409.
- 51 P. Cui, Y. Chen, X. Xu and J. Sun, *Chem. Commun.*, 2008, 5547–5549.
- 52 P. Cui, Y. Chen and M. V. Borzov, *Dalton Trans.*, 2010, **39**, 6886–6890.
- 53 B. F. Wicker, J. Scott, J. G. Andino, X. Gao, H. Park, M. Pink and D. J. Mindiola, *J. Am. Chem. Soc.*, 2010, **132**, 3691–3693.
- 54 Y. Lv, C. E. Kefalidis, J. Zhou, L. Maron, X. Leng and Y. Chen, *J. Am. Chem. Soc.*, 2013, **135**, 14784–14796.
- 55 J. Zhou, T. Li, L. Maron, X. Leng and Y. Chen, *Organometallics*, 2015, **34**, 470–476.
- 56 K. Wang, G. Luo, J. Hong, X. Zhou, L. Weng, Y. Luo and L. Zhang, *Angew. Chem., Int. Ed.*, 2014, **53**, 1053–1056.
- 57 H. Tian, J. Hong, K. Wang, I. de Rosal, L. Maron, X. Zhou and L. Zhang, *CSD Communications*, Private Communications, 2017.
- 58 T. Pugh, F. Tuna, L. Ungur, D. Collison, E. J. L. McInnes, L. F. Chibotaru and R. A. Layfield, *Nat. Commun.*, 2015, **6**, 7492.
- 59 B. Feng, L. Xiang, K. N. McCabe, L. Maron, X. Leng and Y. Chen, *Nat. Commun.*, 2020, **11**, 2916.
- 60 B. Feng, L. Xiang, A. Carpentier, L. Maron, X. Leng and Y. Chen, *J. Am. Chem. Soc.*, 2021, **143**, 2705–2709.
- 61 T. E. Rieser, P. Wetzel, C. Maichle-Mössmer, P. Sirsch and R. Anwender, *J. Am. Chem. Soc.*, 2023, **145**, 17720–17733.
- 62 Y. Lv, X. Xu, Y. Chen, X. Leng and M. V. Borzov, *Angew. Chem., Int. Ed.*, 2011, **50**, 11227–11229.
- 63 S. N. Konchenko, N. A. Pushkarevsky, M. T. Gamer, R. Köppe, H. Schnöckel and P. W. Roesky, *J. Am. Chem. Soc.*, 2009, **131**, 5740–5741.
- 64 W. Huang and P. L. Diaconescu, *Chem. Commun.*, 2012, **48**, 2216–2218.
- 65 T. Li, J. Wiecko, N. A. Pushkarevsky, M. T. Gamer, R. Koepp, S. N. Konchenko, M. Scheer and P. W. Roesky, *Angew. Chem., Int. Ed.*, 2011, **50**, 9491–9495.
- 66 T. Li, M. T. Gamer, M. Scheer, S. N. Konchenko and P. W. Roesky, *Chem. Commun.*, 2013, **49**, 2183–2185.
- 67 C. Schoo, S. Bestgen, R. Köppe, S. N. Konchenko and P. W. Roesky, *Chem. Commun.*, 2018, **54**, 4770–4773.
- 68 F. Zhang, J. Zhang, Z. Chen, L. Weng and X. Zhou, *Inorg. Chem.*, 2019, **58**, 8451–8459.
- 69 A. Hauser, L. Münzfeld, S. Schlittenhardt, R. Köppe, C. Uhlmann, U.-C. Rauska, M. Ruben and P. W. Roesky, *Chem. Sci.*, 2023, **14**, 2149–2158.
- 70 P. G. Edwards, R. A. Andersen and A. Zalkin, *Organometallics*, 1984, **3**, 293–298.
- 71 B. S. Newell, T. C. Schwaab and M. P. Shores, *Inorg. Chem.*, 2011, **50**, 12108–12115.
- 72 J. W. Napoline, S. J. Kraft, E. M. Matson, P. E. Fanwick, S. C. Bart and C. M. Thomas, *Inorg. Chem.*, 2013, **52**, 12170–12177.
- 73 A. L. Ward, W. W. Lukens, C. C. Lu and J. Arnold, *J. Am. Chem. Soc.*, 2014, **136**, 3647–3654.
- 74 A. J. Ayres, M. Zegke, J. P. A. Ostrowski, F. Tuna, E. J. L. McInnes, A. J. Wooles and S. T. Liddle, *Chem. Commun.*, 2018, **54**, 13515–13518.
- 75 A. J. Ayres, A. J. Wooles, M. Zegke, F. Tuna and S. T. Liddle, *Inorg. Chem.*, 2019, **58**, 13077–13089.
- 76 D. Perales, R. Bhowmick, M. Zeller, P. Miro, B. Vlasisavljevich and S. C. Bart, *Chem. Commun.*, 2022, **58**, 9630–9633.
- 77 M. E. Garner and J. Arnold, *Organometallics*, 2017, **36**, 4511–4514.
- 78 M. E. Garner, B. F. Parker, S. Hohloch, R. G. Bergman and J. Arnold, *J. Am. Chem. Soc.*, 2017, **139**, 12935–12938.
- 79 J. M. Ritchey, A. J. Zozulin, D. A. Wroblewski, R. R. Ryan, H. J. Wasserman, D. C. Moody and R. T. Paine, *J. Am. Chem. Soc.*, 1985, **107**, 501–503.
- 80 P. J. Hay, R. R. Ryan, K. V. Salazar, D. A. Wroblewski and A. P. Sattelberger, *J. Am. Chem. Soc.*, 1986, **108**, 313–315.



- 81 A. C. Behrle, L. Castro, L. Maron and J. R. Walensky, *J. Am. Chem. Soc.*, 2015, **137**, 14846–14849.
- 82 B. M. Gardner, G. Balázs, M. Scheer, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2014, **53**, 4484–4488.
- 83 E. P. Wildman, G. Balázs, A. J. Wooles, M. Scheer and S. T. Liddle, *Nat. Commun.*, 2016, **7**, 12884–12894.
- 84 J. Du, G. Balázs, J. A. Seed, J. D. Cryer, A. J. Wooles, M. Scheer and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2022, **61**, e202211627.
- 85 M. R. Duttera, V. W. Day and T. J. Marks, *J. Am. Chem. Soc.*, 1984, **106**, 2907–2912.
- 86 T. M. Rookes, B. M. Gardner, G. Balázs, M. Gregson, F. Tuna, A. J. Wooles, M. Scheer and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2017, **56**, 10495–10500.
- 87 S. P. Vilanova, P. Alayoglu, M. Heidarian, P. Huang and J. R. Walensky, *Chem.–Eur. J.*, 2017, **23**, 16748–16752.
- 88 P. Runghanaphatsophon, T. J. Duignan, A. J. Myers, S. P. Vilanova, C. L. Barnes, J. Autschbach, E. R. Batista, P. Yang and J. R. Walensky, *Inorg. Chem.*, 2018, **57**, 7270–7278.
- 89 M. L. Tarlton, Y. Yang, S. P. Kelley, L. Maron and J. R. Walensky, *Organometallics*, 2021, **40**, 2701–2708.
- 90 M. L. Tarlton, X. Yu, R. J. Ward, S. P. Kelley, J. Autschbach and J. R. Walensky, *Chem.–Eur. J.*, 2021, **27**, 14396–14400.
- 91 M. L. Tarlton, S. P. Vilanova, M. G. Kaumini, S. P. Kelley, P. Huang and J. R. Walensky, *Inorg. Chem.*, 2021, **60**, 14932–14943.
- 92 M. L. Tarlton, Y. Yang, S. P. Kelley, L. Maron and J. R. Walensky, *Organometallics*, 2021, **40**, 2701.
- 93 C. Zhang, G. Hou, G. Zi, W. Ding and M. D. Walter, *Inorg. Chem.*, 2019, **58**, 1571–1590.
- 94 CSD Search by 10-08-2023.
- 95 D. S. J. Arney, R. C. Schnabel, B. C. Scott and C. J. Burns, *J. Am. Chem. Soc.*, 1996, **118**, 6780–6781.
- 96 S. W. Hall, J. C. Huffman, M. M. Miller, L. R. Avens, C. J. Burns, A. P. Sattelberger, D. S. J. Arney and A. F. England, *Organometallics*, 1993, **12**, 752–758.
- 97 C. Zhang, G. Hou, G. Zi, W. Ding and M. D. Walter, *J. Am. Chem. Soc.*, 2018, **140**, 14511–14525.
- 98 C. Zhang, G. Hou, G. Zi and M. D. Walter, *Dalton Trans.*, 2019, **48**, 2377–2387.
- 99 Y. Wang, C. Zhang, G. Zi, W. Ding and M. D. Walter, *New J. Chem.*, 2019, **43**, 9257–9539.
- 100 D. Wang, W. Ding, G. Hou, G. Zi and M. D. Walter, *Chem.–Eur. J.*, 2020, **26**, 16888–16899.
- 101 D. Wang, S. Wang, G. Hou, G. Zi and M. D. Walter, *Inorg. Chem.*, 2020, **59**, 14549–14563.
- 102 D. Wang, G. Hou, G. Zi and M. D. Walter, *Organometallics*, 2020, **39**, 4085–4101.
- 103 D. Wang, G. Hou, G. Zi and M. D. Walter, *Organometallics*, 2021, **40**, 383–396.
- 104 S. Wang, T. Li, Y. Heng, G. Hou, G. Zi and M. D. Walter, *Organometallics*, 2021, **40**, 2149–2165.
- 105 D. Wang, S. Wang, T. Li, Y. Heng, G. Hou, G. Zi and M. D. Walter, *Dalton Trans.*, 2021, **50**, 8349–8363.
- 106 D. M. King, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Science*, 2012, **337**, 717–720.
- 107 D. M. King, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Nat. Chem.*, 2013, **5**, 482–488.
- 108 D. M. King, J. McMaster, F. Tuna, E. J. L. McInnes, W. Lewis, A. J. Blake and S. T. Liddle, *J. Am. Chem. Soc.*, 2014, **136**, 5619–5622.
- 109 R. Magnall, G. Balázs, E. Lu, F. Tuna, A. J. Wooles, M. Scheer and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2019, **58**, 10215–10219.
- 110 J. Du, G. Balázs, A. J. Wooles, M. Scheer and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2021, **60**, 1197–1202.
- 111 O. J. Scherer, B. Werner, G. Heckmann and G. Wolmershäuser, *Angew. Chem., Int. Ed. Engl.*, 1991, **30**, 553–555.
- 112 A. S. P. Frey, F. G. N. Cloke, P. B. Hitchcock and J. C. Green, *New J. Chem.*, 2011, **35**, 2022–2026.
- 113 A. Formanui, F. Ortu, R. Beekmeyer, A. Kerridge, R. W. Adams and D. P. Mills, *Dalton Trans.*, 2016, **45**, 2390–2393.
- 114 D. Patel, F. Tuna, E. J. L. McInnes, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2013, **52**, 13334–13337.
- 115 B. M. Gardner, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2015, **54**, 7068–7072.
- 116 W. Fang, I. Douair, A. Hauser, K. Li, Y. Zhao, P. W. Roesky, S. Wang, L. Maron and C. Zhu, *CCS Chem.*, 2022, **4**, 2630–2638.
- 117 J. Du, D. Hunger, J. A. Seed, J. D. Cryer, D. M. King, A. J. Wooles, J. van Slageren and S. T. Liddle, *J. Am. Chem. Soc.*, 2021, **143**, 5343–5348.
- 118 W. J. Evans, T. A. Ulibarri and J. W. Ziller, *J. Am. Chem. Soc.*, 1988, **110**, 6877–6879.
- 119 H. Schumann, E. Palamidis, J. Loebel and J. Plckardt, *Organometallics*, 1988, **7**, 1008–1010.
- 120 W. J. Evans, J. T. Leman, J. W. Ziller and S. I. Khan, *Inorg. Chem.*, 1996, **35**, 4283–4291.
- 121 F. Nief, D. Turcitu and L. Ricard, *Chem. Commun.*, 2002, 1646–1647.
- 122 T. Pugh, A. Kerridge and R. A. Layfield, *Angew. Chem., Int. Ed.*, 2015, **54**, 4255–4258.
- 123 T. Pugh, V. Vieru, L. F. Chibotaru and R. A. Layfield, *Chem. Sci.*, 2016, **7**, 2128–2137.
- 124 N. Arleth, M. T. Gamer, R. Köppe, S. N. Konchenko, M. Fleischmann, M. Scheer and P. W. Roesky, *Angew. Chem., Int. Ed.*, 2016, **55**, 1557–1560.
- 125 N. Reinfandt, N. Michenfelder, C. Schoo, R. Yadav, S. Reichl, S. N. Konchenko, A. N. Unterreiner, M. Scheer and P. W. Roesky, *Chem.–Eur. J.*, 2021, **27**, 7862–7871.
- 126 C. Schoo, R. Köppe, M. Piesch, M. T. Gamer, S. N. Konchenko, M. Scheer and P. W. Roesky, *Chem.–Eur. J.*, 2018, **24**, 7890–7895.
- 127 C. Schöttle, P. Bockstaller, R. Popescu, D. Gerthsen and C. Feldmann, *Angew. Chem., Int. Ed.*, 2015, **54**, 9866–9870.





- 128 C. Schoo, S. Bestgen, A. Egeberg, J. Seibert, S. N. Konchenko, C. Feldmann and P. W. Roesky, *Angew. Chem., Int. Ed.*, 2019, **58**, 4386–4389.
- 129 N. Reinfandt, A. Hauser, L. Münzfeld and P. W. Roesky, *Chem. Sci.*, 2022, **13**, 3363–3368.
- 130 G. Balázs, M. Sierka and M. Scheer, *Angew. Chem., Int. Ed.*, 2005, **44**, 4920–4924.
- 131 W. J. Evans, S. L. Gonzales and J. W. Ziller, *J. Chem. Soc., Chem. Commun.*, 1992, 1138–1139.
- 132 X. Min, I. A. Popov, F.-X. Pan, L.-J. Li, E. Matito, Z.-M. Sun, L.-S. Wang and A. I. Boldyrev, *Angew. Chem., Int. Ed.*, 2016, **55**, 5531–5535.
- 133 T. Pugh, N. F. Chilton and R. A. Layfield, *Chem. Sci.*, 2017, **8**, 2073–2080.
- 134 C. Schoo, S. Bestgen, A. Egeberg, S. Klementyeva, C. Feldmann, S. N. Konchenko and P. W. Roesky, *Angew. Chem., Int. Ed.*, 2018, **57**, 5912–5916.
- 135 N. Reinfandt, C. Schoo, L. Dütsch, R. Köppe, S. N. Konchenko, M. Scheer and P. W. Roesky, *Chem.–Eur. J.*, 2021, **27**, 3974–3978.
- 136 W. J. Evans, S. L. Gonzales and J. W. Ziller, *J. Am. Chem. Soc.*, 1991, **113**, 9880–9882.
- 137 A. B. Chung, A. J. Ryan, M. Fang, J. W. Ziller and W. J. Evans, *Inorg. Chem.*, 2021, **60**, 15635–15645.
- 138 P. Zhang, R. Nabi, J. K. Staab, N. F. Chilton and S. Demir, *J. Am. Chem. Soc.*, 2023, **145**, 9152–9163.
- 139 F. Lips, R. Clérac and S. Dehnen, *Angew. Chem., Int. Ed.*, 2011, **50**, 960–964.
- 140 F. Lips, M. Hołyńska, R. Clérac, U. Linne, I. Schellenberg, R. Pöttgen, F. Weigend and S. Dehnen, *J. Am. Chem. Soc.*, 2012, **134**, 1181–1191.
- 141 B. Weinert, F. Weigend and S. Dehnen, *Chem.–Eur. J.*, 2012, **18**, 13589–13595.
- 142 B. Weinert, F. Müller, K. Harms, R. Clérac and S. Dehnen, *Angew. Chem., Int. Ed.*, 2014, **53**, 11979–11983.
- 143 R. Ababei, W. Massa, B. Weinert, P. Pollak, X. Xie, R. Clérac, F. Weigend and S. Dehnen, *Chem.–Eur. J.*, 2015, **21**, 386–394.
- 144 P. Zhang, F. Benner, N. F. Chilton and S. Demir, *Chem.*, 2022, **8**, 717–730.
- 145 O. J. Scherer, J. Schulze and G. Wolmershäuser, *J. Organomet. Chem.*, 1994, **484**, C5–C7.
- 146 B. M. Gardner, G. Balázs, M. Scheer, A. J. Wooles, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2015, **54**, 15250–15254.
- 147 B. M. Gardner, G. Balázs, M. Scheer, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Nat. Chem.*, 2015, **7**, 582–590.
- 148 M. L. Tarlton, O. J. Fajen, S. P. Kelley, A. Kerridge, T. Malcomson, T. L. Morrison, M. P. Shores, X. Xhani and J. R. Walensky, *Inorg. Chem.*, 2021, **60**, 10614–10630.
- 149 A. C. Behrle and J. R. Walensky, *Dalton Trans.*, 2016, **45**, 10042–10049.
- 150 E. P. Wildman, G. Balázs, A. J. Wooles, M. Scheer and S. T. Liddle, *Nat. Commun.*, 2017, **8**, 14769–14777.
- 151 C. J. Hoerger, F. W. Heinemann, E. Louyriac, M. Rigo, L. Maron, H. Grützmacher, M. Driess and K. Meyer, *Angew. Chem., Int. Ed.*, 2019, **131**, 1693–1697.
- 152 R. Magnall, G. Balázs, E. Lu, M. Kern, J. van Slageren, F. Tuna, A. J. Wooles, M. Scheer and S. T. Liddle, *Chem.–Eur. J.*, 2019, **25**, 14246–14252.
- 153 T. M. Rookes, E. P. Wildman, G. Balázs, B. M. Gardner, A. J. Wooles, M. Gregson, F. Tuna, M. Scheer and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2018, **57**, 1332–1336.
- 154 N. Lichtenberger, R. J. Wilson, A. R. Eulenstein, W. Massa, R. Clérac, F. Weigend and S. Dehnen, *J. Am. Chem. Soc.*, 2016, **138**, 9033–9036.
- 155 A. R. Eulenstein, Y. J. Franzke, N. Lichtenberger, R. J. Wilson, H. L. Deubner, F. Kraus, R. Clérac, F. Weigend and S. Dehnen, *Nat. Chem.*, 2021, **13**, 149–155.
- 156 E. R. Pugliese, F. Benner and S. Demir, *Chem.–Eur. J.*, 2023, e202302687.
- 157 E. R. Pugliese, F. Benner and S. Demir, *Chem. Commun.*, 2023, DOI: [10.1039/d3cc04520j](https://doi.org/10.1039/d3cc04520j).
- 158 Q. Wen, B. Feng and Y. Chen, *Acc. Chem. Res.*, 2023, DOI: [10.1021/acs.accounts.3c00429](https://doi.org/10.1021/acs.accounts.3c00429).

